Genomic Landscape and Molecular Subtypes of Primary Central Nervous System Lymphoma

Shengjie Li[1,2,3,4,5,6*#], Danhui Li[7#], Zuguang Xia[8,9#], Jianing Wu[6], Jun Ren[6], Yingzhu Li[6], Jiazhen Cao[10], Ying Sun[11], Liyang Zhang[12], Hongwei Ye[12], Chengxun Li[13*], Wenjun Cao[6*], Xingtao Zhou[5*], Ying Mao[1,2,3,4,14*]

1. Department of Neurosurgery, Huashan Hospital, Department of Clinical Laboratory, Eye & ENT Hospital, Shanghai Medical College, Fudan University, Shanghai, China

2. National Center for Neurological Disorders, Shanghai, China

3. Shanghai Key Laboratory of Brain Function Restoration and Neural Regeneration, Shanghai, China

4. Neurosurgical Institute of Fudan University, Shanghai, China

5. Eye Institute and Department of Ophthalmology, Eye & ENT Hospital, Fudan University, Shanghai, China

6. Department of Clinical Laboratory, Eye & ENT Hospital, Shanghai Medical College, Fudan University, Shanghai, China

7. Department of Pathology, RenJi Hospital, School of Medicine, Shanghai JiaoTong University, Shanghai, 200127, China

8. Department of Medical Oncology, Fudan University Shanghai Cancer Center, Fudan University, Shanghai, China

9. Department of Oncology, Shanghai Medical College, Fudan University, Shanghai, China

10. Department of Clinical Laboratory, Huashan Hospital, Fudan University, Shanghai, China

11. GenomiCare Biotechnology (Shanghai) Co. Ltd., Shanghai, China

12. Guangzhou KingMed Center for Clinical Laboratory Co. Ltd., Guangzhou, China

13. Institutes of Biomedical Sciences, Fudan University, Shanghai, 200032, China

14. Lead contact

#Shengjie Li, Danhui Li, and Zuguang Xia contributed equally to this work.

Corresponding Authors:

Wenjun Cao, Department of Clinical Laboratory, Eye & ENT Hospital, Fudan University, No.

83 Fenyang Road, Shanghai, China; Electronic address: wgkjyk@aliyun.com.

Chengxun Li, Institutes of Biomedical Sciences, Fudan University, Shanghai, 200032, China

Electronic address: licx21@m.fudan.edu.cn

Xingtao, Zhou, Eye Institute and Department of Ophthalmology, Eye & ENT Hospital, Fudan

University, Shanghai, 200031, China. Electronic address: doctzhouxingtao@163.com.

Ying Mao, Department of Neurosurgery, Huashan Hospital, Fudan University, No. 12

Wulumuqi Road, Shanghai, 200040, China; Electronic address: maoyin@fudan.edu.cn

Abstract

Primary central nervous system lymphoma (PCNSL) is a rare and aggressive brain tumor with a poor prognosis and is almost exclusively diffuse large B-cell lymphoma (DLBCL). Its genetic characteristics and molecular subtypes in Chinese patients remain poorly understood, which in turn makes developing effective new therapies challenging. By enrolling 176 newly diagnosed Chinese PCNSLs at five tertiary care centers with extensive follow-up, we performed a genomic study aimed at expanding the genomic landscape and developing new molecular subtypes. We first confirmed that the molecular subtyping of DLBCL, as previously published, is not applicable to Chinese PCNSLs. We then identified (n = 58) and validated (n = 82) three prominent genetic subtypes related to different clinical and molecular features of PCNSL, and further confirmed by an independent external Chinese PCNSL cohort (n = 36). We called these BMIs (from the co-occurrence of mutations in two genes among BTG1, MYD88, and IRF4), which are associated with favorable outcomes; E3s (so-called EP300 mutations), which are associated with unfavorable outcomes; and UCs (unclassified, without characteristic mutations). Importantly, EP300 was mutated in more Asians (16.98%) than in Western PCNSLs (< 4.53%), resulting in unfavorable outcomes independent of the specific mutation site. Our analysis comprehensively reveals the genomic landscape of Chinese PCNSL and emphasizes the clinical value of molecular classification for improving precision medicine strategies.

Keywords: Primary central nervous system lymphoma; Genomic landscape; Molecular subtypes; EP300; Precision medicine

Introduction

Primary central nervous system lymphoma (PCNSL), a rare subtype of extranodal non-Hodgkin lymphoma, accounts for >60% of PCNSL cases and has an unfavorable prognosis even when the patient receives standard treatment, which is based on a high-dose methotrexate (HD-MTX) regimen[1,2]. PCNSL is a unique lymphoma that differs from other tumors and is characterized as a genetically heterogeneous disorder marked by a variety of low-frequency mutations, somatic copy number alterations, and structural variants[3–6]. Thus, combining standard immunochemotherapy with promising novel agents that target specific pathways via different molecular clusters may improve patient prognosis[7,8].

Pathologically, nearly 95% of PCNSLs are diffuse large B-cell lymphomas (DLBCLs). Recently, DLBCL has been categorized into various molecular clusters on the basis of genomic sequencing and RNA sequencing, including quartet categorization by *L.M. Staudt*'s team[9], quintet categorization by *M.A. Shipp*'s team[10], *L.M. Staudt*'s team's refined septet method[11], and the LymphPlex classification proposed by *W.L. Zhao*'s team[12]. However, PCNSL has been proven to be a biological entity that is molecularly distinct from DLBCL[13], so these molecular clusters of DLBCL may not be applicable to PCNSL. Notably, the integration of genome-wide data from multiomic studies by *A. Alentorn* et al.[14] showed four molecular patterns of PCNSL mainly use gene expression data. These patterns have a distinctive prognostic impact, providing a basis for future clinical stratification and subtype-based targeted interventions. Owing to the molecular heterogeneity between various ethnic groups of PCNSL patients[3,15], the applicability of the proposed classifications for PCNSL to other populations requires further confirmation. No effective biomarker or molecular classification scheme exists to tailor therapies for individual PCNSL patients. Furthermore, the issue of PCNSL molecular heterogeneity in Chinese patients has not been adequately addressed, primarily because the data have come from small, single-center studies of Chinese PCNSL patients[16,17].

To address these issues, we conducted whole-exome sequencing (WES) on specimens obtained from 140 patients with newly diagnosed PCNSL who were seen at one of three tertiary care cancer centers. These patients were divided into a discovery cohort and a

validation cohort. Among the 140 patients, 94.29% received treatment with an HD-MTX-

based chemotherapy regimen followed by consolidation therapy (detailed in Table S1), and

all 140 patients underwent long-term follow-up. Additionally, WES/Whole Genome

Sequencing (WGS) data were obtained from 36 Chinese PCNSL patients at two other tertiary

care cancer centers[17], serving as an independent external cohort. This representative and

clinically annotated PCNSL cohort was utilized to comprehensively detect mutations.

Importantly, to facilitate routine clinical implementation, we identified three molecular

subtypes composed only of single-nucleotide variations (SNVs); these SNVs were associated

with different outcomes, and this was confirmed in the validation cohort and in an independent

external Chinese PCNSL cohort (n=36)[17]. Our results comprehensively reveal the genomic

landscape of PCNSLs in Chinese patients and provide a clinically actionable PCNSL

classification system, thereby improving precision medicine strategies.

METHODS

Sample collection and clinicopathological features of PCNSL patients

In this study, a total of 140 newly diagnosed PCNSL patients were retrospectively enrolled

from January 2010 to December 2020 from three independent medical centers, namely,

Huashan Hospital of Fudan University, Renji Hospital of Shanghai Jiao Tong University, and

Shanghai Cancer Center of Fudan University. The study comprised two cohorts: the

discovery cohort, which included 58 patients from Huashan Hospital of Fudan University,

Shanghai; and the validation cohort (n = 82), which consisted of 25 patients from Huashan

Hospital of Fudan University, 51 patients from Renji Hospital of Shanghai Jiao Tong University,

and 6 patients from Shanghai Cancer Center of Fudan University. In the discovery cohort,

paired blood samples were also obtained for analysis in conjunction with each tumor sample.

In the validation cohort, only unpaired FFPE tumor tissue samples were obtained for analysis.

Additionally, WES/WGS data of 36 patients newly diagnosed with PCNSL from Fujian

Cancer Hospital and The First Affiliated Hospital of Fujian Medical University between

February 2012 and October 2020 were included in this study, as an independent external

Chinese PCNSL cohort. The general clinical characteristics of these 36 PCNSL patients and

the sequencing protocol have been previously described[17].

None of the patients had received steroid treatment for PCNSL, and PCNSL tumors were

diagnosed on the basis of the World Health Organization criteria[13]. All patients underwent a

2-deoxy-2[F-18] fluoro-D-glucose positron emission tomography/computed tomography

(FDG PET/CT) scan and bone marrow aspiration to exclude systemic tumor manifestation

on the basis of the guidelines of the International PCNSL Collaborative Group[18] and the

European Association of Neuro-Oncology[19]. PCNSL patients for whom ocular involvement

was present were excluded. Each tumor tissue sample was independently reviewed by at

least 2 pathologists to confirm that the tumor sample was histologically consistent with the

PCNSL. All patients underwent long-term follow-up, which ran to December 30, 2022.

Ethics statement

This study was conducted with the approval of the Institutional Review Boards at Huashan

Hospital of Fudan University (Approval No. 2022-529), Shanghai Cancer Center of Fudan

University (Approval No. 1612167-18), Renji Hospital of Shanghai Jiao Tong University

(Approval No. LY2024-112-C), and Fujian Cancer Hospital (Approval no. SQ 2020-106-01).

All participants provided written informed consent before participating in the clinical study and

before the collection of tumor tissues. The research was carried out in strict adherence to the

ethical principles of the Declaration of Helsinki and conformed to international norms of good

clinical practice.

Treatment Regimens

All PCNSL patients were treated with HD-MTX-based combination immunochemotherapy,

followed by consolidation therapy (detailed in Table S1). The following chemotherapy

protocols were used for induction: HD-MTX combined with IDA, HD-MTX combined with R,

HD-MTX combined with IDA and R, or a combination of BTK inhibitors. Whole-brain radiation

therapy or stem cell transplantation is a form of consolidated therapy. The detailed therapeutic

schedule was the same as that previously described[20].

Sample processing

The collection and processing of samples for this study were carried out by the Department

of Pathology or Department of Neurosurgery at Huashan Hospital of Fudan University, Renji

Hospital of Shanghai Jiao Tong University, and Shanghai Cancer Center of Fudan University. In the discovery cohort, surplus tumor tissues and paired blood samples obtained during surgical procedures were collected. To guarantee the quality of these samples, each specimen was promptly and accurately labeled with relevant patient information within 30 minutes of collection and then immediately snap-frozen in liquid nitrogen. These samples were stored at -80°C until being shipped to GenomiCare Biotechnology (Shanghai, China). In the validation cohort, the residual tumor tissue specimens were subjected to FFPE. These FFPE samples were transported in a dry ice container to Sinotech Genomics Technologies in Shanghai, China, accompanied by a time and temperature tracker to ensure proper monitoring during shipment. Upon arrival, the samples were promptly stored in liquid nitrogen to maintain their integrity until further processing.

DNA extraction

For frozen fresh tissue and matched peripheral blood samples, total genomic DNA (gDNA) was extracted via the Maxwell RSC Blood DNA Kit (AS1400, Promega) on a Maxwell RSC system (AS4500, Promega) following the manufacturer's instructions. For FFPE tissue, total gDNA was extracted via the QIAamp DNA FFPE Tissue Kit (56404, Qiagen) following the manufacturer's instructions. The integrity and concentration of the gDNA were determined via the Qsep100 System (BIOptic, China) and a Qubit 3.0 fluorometer (Thermo Fisher Scientific). The OD_{260} was measured with a NanoDrop One (Thermo Fisher Scientific).

WES library preparation and sequencing

For frozen tissue and matched peripheral blood samples, exome DNA was extracted via the SureSelect Human All Exon V7 Kit (5991-9039EN, Agilent). The library was subsequently prepared via the SureSelectXT Low Input Target Enrichment and Library Preparation system (G9703-90000, Agilent). For the FFPE sample, the exome DNA was captured via the SureSelect Human All Exon V8 Kit (5191-6874, Agilent) and prepared into a library via the SureSelect XT HS2 DNA Reagent Kit (G9983A, Agilent). The library was validated via the use of an Agilent 2100 Bioanalyzer and a Qubit 3.0 fluorometer. Paired-end 150 bp read sequencing was performed on an Illumina NovaSeq 6000. Image analysis and base calling were performed via onboard RTA3 software (Illumina).

Somatic mutation calling

Quality control was conducted via FastQC software. The cleaned and trimmed FASTQ files were aligned to the UCSC human reference genome (hg19) via the Burrows–Wheeler Aligner (BWA) with default parameters. For each paired blood sample, SNVs and short insertions/deletions (INDELs) were identified via Sentieon TNseq[21] with default parameters. The identified somatic mutations were removed if they did not satisfy any one of the following criteria: a variant allele frequency (VAF) of at least 0.05, support from a minimum of three reads, annotation by the Variant Effect Predictor (VEP) package[22], and transformation into a mutation annotation format (MAF) file for further analysis with maftools[23].

When a paired normal sample (FFPE sample) was not available, a panel of normal (PON) files was used as a paired control to call the SNVs and InDels. A panel of normal files was created using the reads from clinical blood samples that showed no evidence of tumor contamination from 100 different individuals collected by Genomicare Biotechnology (Shanghai). The reads were retained when they passed the standard WES QC, as described in the WES subsection. The reads for reference alleles and alternative alleles were gathered for each candidate site in all 100 samples, except for false positives, which were those exhibiting a candidate allele frequency exceeding 0.05 in more than 5 samples.

We applied an additional PON mask for all the candidate somatic SNV and INDEL sites to exclude germline mutations, as previously reported[10,24]. In general, in addition to the standard retention rules, mutations were retained if they were known driver alterations highlighted as being biologically significant in the COSMIC database. Mutations with a tumor mutation frequency ≥20% were also retained. Other germline mutations were retained if their existence ratio in the Exome Aggregation Consortium (ExAC) exac_all was < 0.00005, their existence ratio in the Asian population exac_eas was < 0.0003, and their existence ratio in previous GenomiCare samples germline_gc was < 0.0006. Retained mutations were considered somatic mutations.

Thus, the PON file and PON mask limit possible recurrent artifacts of sequencing and the presence of germline mutations in somatic mutations that are falsely selected owing to a lack of paired blood data.

Somatic copy number alteration calling and profiling

Using the method described in the ExomeCNV package[25], we implemented a normalized depth–coverage ratio method to identify CNVs in paired samples. To correct for five potential biases (the size of exonic regions, batch effects, both the quantity and quality of sequencing data, local GC content, and genomic mappability) that affect the raw read counts, we utilized a standard normal distribution model. Genes with a haploid copy number ≥3 or ≤1.2 were defined as amplified or deleted, respectively, and a minimum tumor content (purity) of 20% was needed.

Significance analysis of recurrent somatic copy number alterations

Sequenza[26] was used to generate the segment files. The segment files were used as the input for GISTIC2.0[27] to detect recurrent arm-level and focal peaks in copy number alterations via the parameters -js 40 -conf 0.99 -ta 0.2 -td 0.2 -brlen 0.8, with other parameters set to the defaults.

Cancer-driver gene mutations filtering

After the aforementioned steps of calling SNVs and CNVs, we further filtered the resulting mutated genes by intersecting them with a designated group of genes recognized as cancer-related genes. This group of genes was sourced from two prominent public cancer gene databases. Part 1 of the group was obtained from the OncoKB curated cancer gene list (https://www.oncokb.org/cancerGenes)[28]. These genes are classified as cancer genes by OncoKB on the basis of their inclusion in various sequencing panels, the Sanger Cancer Gene Census, or the criteria established by *Vogelstein* et al.[29]. Part 2 was derived from the ranked CIViC gene candidates table (https://github.com/griffithlab/civic-server/blob/master/public/downloads/RankedCivicGeneCandidates.tsv)[30]. The final list of cancer-related genes consisted of all genes in Part 1 and genes in Part 2 only when its 'panel_count' value (in the CIViC tsv file listed above) was ≥ 2.

Bioinformatic analysis

The mutational signature classification was based on all single-base substitution (SBS) signatures of the COSMIC mutational signatures[31]. TMB was defined as the aggregate count of somatic nonsynonymous mutations, including SNVs or INDELs, within the tumor exome

235 for each patient[32]. This count was then divided by the overall size of the targeted regions (35

236 for WES), resulting in the TMB, expressed in counts per megabase (counts/Mb). The MATH

237 score[33] was calculated on the basis of the width of the variant allele frequency (VAF)

238 distribution, utilizing maftools for the analysis[23]. The pathway map was generated via

239 maftools as previously described[34].

Classification model for identifying molecular types

CoxNet survival analyses and least absolute shrinkage and selection operator (LASSO)

regressions were performed via the scikit-learn package in Python (version 3.9.7). K–M

survival curves (log-rank test) were used for OS analysis via the survival and survminer

packages in R (version 4.1.2).

(1) Selection of 24 initial features in the discovery cohort

Molecular features derived from SNVs were selected to develop subsequent classification

models.

First, to identify genes with high variability for use in clustering, those with low mutation

frequencies were filtered out. This process led to the selection of genes with a mutation

frequency greater than 5% for clustering purposes. A total of 150 genes were screened for

further analysis.

In the second phase, OS was analyzed as the dependent variable, with the aforementioned

150 genes serving as independent variables. The feature selection steps were as follows: (1)

a comprehensive approach involving CoxNet survival analyses and LASSO regressions was

employed to identify characteristic genes; (2) the importance score of each feature in the

model was calculated and ranked in descending order; (3) features that ranked in the top 50%

were recorded as potential features once; (4) processes 1-3 were repeated 500 times, and

all potential features were recorded; and (5) among the feature sets obtained in the above

process, those recorded more than 250 times were selected.

Ultimately, 24 feature genes, namely, EP300, KMT2D, MPEG1, TUSC3, MYD88, PTCH1,

DST, ZNF521, BTG1, NOTCH2, PRDM1, ADGRL3, CDH11, CREBBP, ATRX, TCL1A,

GNA13, ETV6, IRF4, HIST2H3D, CD79B, GRM8, MYH11, and SPTA1, were selected.

(2) Differential feature selection

To determine the most suitable features among the initial 24 for the classification model, Kaplan–Meier analyses were conducted. Features qualifying for further analysis met the following criteria: a log-rank test P value of less than 0.2 and any group with more than 5 patients. Ultimately, 6 DEGs, namely, EP300, CREBBP, DST, MYD88, BTG1, and IRF4, were identified.

(3) Model training utilizing random combinations of 6 differential features

To adhere to clinical practice guidelines for predicting patient outcomes and implementing individualized management strategies for PCNSL patients, patients with PCNSL were categorized into three subgroups (those with good, poor, and intermediate prognoses) in the present study. Six differential features were ultimately selected for building a classification model to classify patients into the three subtypes. These features were randomly combined to form three types, yielding a total of 288 combinations. To identify the most appropriate classification model among these combinations, Kaplan–Meier analyses were performed. Models were chosen on the basis of a log-rank test P value of less than 0.05 and subgroup sample sizes greater than 8. Ultimately, seven candidate molecular subtype combinations met these criteria: (molecular markers shown in the format of subtype 1 vs. subtype 2 vs. subtype 3): ① EP300 (predominant) vs. MYD88/BTG1/IRF4≥2 vs. others; ② EP300 vs. MYD88/BTG1/IRF4≥2 (predominant) vs. others; ③ BTG1 (predominant) vs. CREBP/DST≥1 vs. others; ④ BTG1 vs. CREBP/DST≥1 (predominant) vs. others; ⑤ BTG1/IRF4≥1 vs. EP300/CREBP/DST≥1 vs. others; ⑥ BTG1/IRF4≥1 vs. EP300/DST≥1 vs. others; and ⑦ CREBP/DST≥1 vs. MYD88/BTG1/IRF4≥2 vs. others.

(4) Validation of candidate molecular subtypes

The WES data and OS data of 82 patients in the validation cohort were included to validate the candidate molecular subtypes. The mutation frequencies of EP300, CREBBP, DST, MYD88, BTG1, and IRF4 in the validation cohort were greater than 5%. Kaplan–Meier survival analysis was used to validate the seven candidate molecular subtype combinations. The performance of [EP300 (predominant) vs. MYD88/BTG1/IRF4≥2 vs. others] (3-way P = 0. 0202) and [EP300 vs. MYD88/BTG1/IRF4≥2 vs. others (predominant)] (3-way P = 0. 0346) was better than those of the other five molecular subtypes (3-way P > 0.05 for both).

Additionally, compared with the [EP300 vs. MYD88/BTG1/IRF4≥2 vs. others (predominant)]

molecular subtype, the Kaplan–Meier survival curve of [EP300 (predominant) vs.

MYD88/BTG1/IRF4≥2 vs. others] showed no crossover. Finally, the performance of [EP300

(predominant) vs. MYD88/BTG1/IRF4≥2 vs. others] was the best, so we selected it as the

molecular subtype of Chinese PCNSL patients. In other words, PCNSL patients with both

EP300 and BMI (BTG1, IRF4, and MYD88) variants were categorized into the E3 subtype.

Hematoxylin & eosin and immunohistochemical staining

PCNSL tumor tissue was fixed with 10% paraformaldehyde and embedded in paraffin. The

paraffin-embedded tissues were cut into 4-µm-thick sections. These sections were then

subjected to staining with hematoxylin and eosin (H&E) or for immunohistochemistry. For

H&E staining, the sections were stained with hematoxylin and eosin following standard

protocols (ab245880, Abcam). Immunohistochemistry was performed for ki-67 (ab15580,

Abcam), CD2 (ab314761, Abcam), CD5 (ab75877, Abcam), CD10 (ab256494, Abcam), CD19

(ab134114, Abcam), CD20 (ab78237, Abcam), CD79a (ab79414, Abcam), BCL2 (MA5-11757,

Thermo Scientific), BCL6 (PA5-14259, Thermo Scientific), MUM1 (ab247079, Abcam), c-Myc

(MA1-980, Thermo Scientific), and p53 (MA5-14516, Thermo Scientific) via an automated

Leica BOND-III staining system. H&E and immunohistochemistry images were obtained with

a KFBIO scanner (KF-PRO-005-EX, Zhejiang, China) and visualized via KFBIO SlideViewer

software. Each staining was independently reviewed by at least 2 pathologists.

In situ hybridization of fluorescence

PCNSL tumor tissue was fixed with 10% paraformaldehyde and embedded in paraffin. The

paraffin-embedded tissues were cut into 4-µm-thick sections. Fluorescence in situ

hybridization was performed in accordance with standard protocols via the following

commercial probes: MYC break-apart, BCL-2 break-apart, and BCL-6 break-apart probes

(LBP Medicine Science & Technology Co., Ltd., Guangzhou, China). Staining was

independently evaluated by two hematopathologists, and discrepancies were resolved by

another hematopathologist.

HIV, HBV, and EBV detection

Antibody and antigen levels of HIV were measured via an Elecscy HIV combi PT assay kit

(Roche, Germany) with a Cobas e 601 analyzer (Roche). Hepatitis B virus (HBV)-specific

PCR was performed via an HBV virus nucleic acid test kit (Sansure Biotech, China) with an

Applied Biosystems 7500 Real-Time PCR System (Thermo Scientific). EBV-encoded RNA in

situ hybridization (EBER-ISH) was performed to assess the presence of Epstein-Barr virus

(EBV) in tumor tissue. The testing utilized a fluorescein isothiocyanate -coupled specific

peptide nucleic acid probe (Roche Diagnostics, Mannheim, Germany).

Statistical analysis

All the statistical analyses were performed via R version 4.1.2 and GraphPad Prism version

9. The specific statistical analyses employed for each figure are detailed in the respective

legends. We used PASS software to calculate the sample size for our study, with a

significance level of 0.05 and a power of 0.80. The observed group proportions were 0.6, 0.2,

and 0.2, whereas the expected proportions were 0.33 for each group. The sample size

required to meet these criteria was 25 participants. Consequently, the sample size utilized in

this study is deemed adequate. Whole-exome sequencing data were processed via nf-

core/sarek v3.4.2[35] of the nf-core collection of workflows[36], which uses reproducible software

environments from the Bioconda[37] and Biocontainers[38] projects. The pipeline was executed

with Nextflow v24.04.2[39].

Results

PCNSL genomic landscape

Using WES, we detected mutations in 140 patients who were newly diagnosed with PCNSL

at one of three cancer centers; in 59% of cases, blood samples were lacking. Because EBV-

positive PCNSL is recognized as an immunodeficiency-associated PCNSL in the current

WHO CNS5 classification[40,41], all participants in the study who tested negative for EBV were

included. Additionally, DLBCL patients with HBV and HIV infection may have different

mutational profiles[42,43]; thus, all participants in the study who tested negative for HIV and HBV

were included. The key demographic and clinical characteristics of the patients are

summarized in **Figure S1** and **Table S1**. The study design and flow chart are displayed in

Figure 1A. After filtering, we identified 115,938 genetic variation events in the 140 PCNSL

samples analyzed (median = 19.06 mutations/Mb, range = 0.24–91.2 mutations/Mb; median

number of variants = 949.5/sample) (**Figure S2A, Figure S3A**).

We applied a filter to the 140 PCNSLs on the basis of the genes curated in the CIViC and

OncoKB databases to identify candidate cancer genes with hallmark mutations in PCNSL.

Some of the mutated genes were PIM1 (in 62.1% of PCNSLs), MYD88 (55.0%), KMT2D

(48.6%), CD79B (45.7%), BTG2 (38.6%), LRP1B (37.9%), FAT4 (37.1%), RNF213 (35%),

FAT1 (34.3%), ZFHX3 (33.6%), DTX1 (31.4%), KMT2C (31.4%), SETD1B (31.4%), MPEG1

(29.3%), CREBBP (27.9%), ANKRD11 (27.1%), KAT6B (27.1%), BTG1 (25%), and PTEN

(25%) (**Figure 1B**), which are involved in chromatin histone modification, BCR-TLR-mediated

NF-κB signaling, immune signaling, the cell cycle, PI3 kinase signaling, MAPK signaling, and

Wnt/β-catenin signaling (**Figure S2B**)[3,5,9–11,44–51]. In this study, we also identified several

other pathways in PCNSL, such as pathways related to genome integrity, RTK signaling, RNA

abundance, TGFB signaling, TOR signaling, and apoptosis (**Figure S2B**), many of which

have defined roles in other cancers[34]. The mutually exclusive or cooccurring relationships of

these genes are shown in **Figure S3D**. Survival associated with candidate cancer genes with

mutation frequencies greater than 15% plus the NOTCH2 and EZH2 genes[9] is shown in

Figure S4.

In the search for focal copy number alterations (CNAs) in the 140 PCNSLs, we detected

significant recurrent amplifications at chromosomal locations 1q21.1, 4q16.3, 11p15.5,

11q12.2, 17q12, 19p13.3, 22q11.1, Yq11.221, and Yq12. We also detected deletions at 11q11,

11q12.1, 14q32.33, 17q11.2, 18p11.21, 19p12, 19q13.42, and 22q11.21 (**Figure 1C**). The

arm-level CNAs among all 140 PCNSL samples are displayed in **Figure S2C**. The pattern of

somatic mutations caused by the different mutational processes in the genome, termed

mutational signatures, was calculated and compared to the well-established signatures in

COSMIC[52]. The mutational signature contributions are shown in **Figure S2D**. The COSMIC

5, COSMIC 45, and COSMIC 1 signatures contributed the most to the PCNSL genome

(**Figure S2E**). These factors are associated with aging, tobacco smoking, NER deficiency,

the regulation of oxidative DNA damage repair, and the spontaneous deamination of 5-

methylcytosine. Taken together, these results establish a comprehensive genomic landscape

380 of PCNSL in Chinese individuals.

Comparison of the genomic landscapes of the discovery and validation cohorts

Owing to differences in sample type between the discovery and validation cohorts (fresh tumor [discovery cohort] vs. formalin-fixed paraffin-embedded [FFPE] [validation cohort]) and the presence or absence of matched samples (paired [discovery cohort] vs. tumor-only [validation cohort]), we applied further filtering and recalibration steps to reduce false-positive calls in the validation cohort (for details, see the Methods subsection on somatic mutation calling) and compared the genomic landscapes of the discovery and validation cohorts. The key demographic and clinical characteristics of the patients in the discovery and validation cohorts are summarized in **Table S2**. We identified 12,222 genetic variation events (median = 3.84 mutations/Mb, median variants = 159/sample, **Figure S3B, Figure S5A**) in the analyzed discovery cohort of PCNSL samples and 103,716 genetic variation events (median = 6.85 mutations/Mb, median variants = 1597/sample, **Figure S3C, Figure S5B**) in the analyzed validation cohort of PCNSL samples. The genetic variation profiles of the discovery and validation cohort of PCNSL samples were similar, as revealed by principal component analysis (PCA) (**Figure 2A**), but the mutation rate was higher in the validation cohort of PCNSL samples (**Figure 2B**).

PIM1, MYD88, CD79B, and KMT2D were the most frequently mutated candidate cancer genes of PCNSL. PIM1 (70.7%), MYD88 (63.8%), CD79B (51.7%), and KMT2D (39.7%) (**Figure S5C**) were identified in the discovery cohort, and PIM1 (56.1%), MYD88 (48.8%), CD79B (41.5%), and KMT2D (54.6%) were also observed at similar mutation frequencies in the validation cohort (**Figure S5D**). However, several of the less frequently mutated candidate cancer genes in PCNSL whose mutation frequencies were significantly different when the discovery and validation cohorts were compared included ZFHX3, TRRAP, SZT2, RNF213, RANBP2, PRKDC, NUMA1, NOTCH1, NOTCH2, KMT2C, KMT2B, KAT6B, FLT3, FAT4, FAT1, and ANKRD11 (all $P < 0.05$) (**Figure 2C**). The arm-level CNAs between the discovery and validation cohort of PCNSL samples were mostly similar and are displayed in **Figure 2D**. Significant differences in both recurrent amplifications (1q, 8q, and 12p) and deletions (6p, 8p, 19p, and 19q) were observed. The results of significant recurrent focal amplifications and

deletions were visualized via GISTIC 2.0 in the discovery (**Figure S5E**) and validation (**Figure S5F**) cohorts.

The tumor mutational burden (TMB) describes the number of mutations in one tumor sample and is often related to the prognosis of patients receiving clinical treatments[53]. The mutant-allele tumor heterogeneity (MATH) score reflects intratumor heterogeneity[33]. Microsatellite instability (MSI) arises from the impaired function of DNA mismatch repair proteins, and its molecular characteristics are regarded as significant genetic markers[54]. Although the TMB (**Figure 2E**), MATH score (**Figure 2F**), MSI (**Figure 2G**), and the variant allele frequency (VAF) (**Figure 2H**) were greater in the validation cohort than in the discovery cohort ($P < 0.001$), the mean values were relatively close. The 3 signatures extracted from the discovery samples presented cosine similarities of 86.7%, 93.1% and 94.4% to COSMIC signatures 6, 45, and 10a, respectively (**Figure S5G**), whereas those extracted from the validation samples presented similarities of 80.4%, 70.9%, and 77.9% to COSMIC signatures 5, 1, and 23, respectively (**Figure S5H**). Taken together, these results suggest that the genetic variation profiles of the discovery and validation PCNSL samples exhibit certain discrepancies.

The unique genomic landscape of PCNSLs in Chinese patients

To date, only single-center studies with small sample sizes have reported heterogeneity in the genomic landscape among different populations of Chinese PCNSL patients[15]. Thus, in this large, multicenter study, we further compared the genetic landscape of our cohort with previously published PCNSL/DLBCL data. The frequencies of recurrent genetic alterations in PCNSL patients were compared with those in DLBCL patients from the United States, Finland, Denmark, Israel, the United Kingdom, Hong Kong, and India[49]. Mirror bar plots (**Figure 3A**) revealed that a set of genes, such as PIM1 (62.1% vs. 16.88%), MYD88 (55% vs. 18.01%), IRF4 (15% vs. 3.98%), and EP300 (19.29% vs. 5.97%), had significantly higher mutation rates in Chinese PCNSL patients than in DLBCL patients, suggesting genetic diversity of the cancer genome between diseases. Compared with those in PCNSLs in French patients [14] (**Figure 3B**), the mutation frequencies of EP300 (19.29% vs. 3.48%) and KMT2D (48.6% vs. 22.61%) were significantly higher in the PCNSLs of Chinese patients. Compared with those

in the PCNSLs[3] of Japanese patients (**Figure 3C**), the mutation frequencies of MYD88 (55%

vs. 85.6%) and BTG2 (38.6% vs. 87.8%) were significantly lower in the PCNSLs of Chinese

patients. These data revealed a unique mutational landscape of Chinese PCNSL patients in

terms of the frequency of SNVs. Next, the frequencies of recurrent SNAs with the GISTIC2.0

program in Chinese PCNSL patients were compared with those in DLBCL patients. At the

focal level of SNAs, Chinese PCNSL samples had significant levels of 11q12.4, 18p11.21,

19q13.42, 22q11.1, and 22q11.21 deletions along with amplifications in arms 1q21.1, 4p16.3,

11q12.2, and 17q12 (**Figure 3D**). Compared with PCNSLs from Japanese patients, those

from Chinese patients also had significant levels of deletions in arms 11q12.1, 18p11.21,

19q13.42, 22q11.1, and 22q11.21 along with amplifications in arms 1q21.1, 4p16.3, 11q12.2,

and 17q12 (**Figure 3E**). These results suggest genetic diversity in the PCNSL genome

between races.

Owing to the genetic diversity of the PCNSL genome across races and the genetic diversity

between the PCNSL and DLBCL cancer genomes, the existing molecular subtyping methods

for PCNSL[14] and DLBCL[9–12,55,56] may not be applicable to Chinese PCNSL patients. We

assessed the prognostic value of these published molecular subtypes for overall survival (OS)

and progression-free survival (PFS) in our Chinese PCNSL cohort. Actionable PCNSL

classification can perfectly differentiate prognoses, thereby improving precision medicine

strategies. According to cell-of-origin (COO) molecular subtyping[55,56], there were no

significant differences in OS (*P* = 0.42) or PFS (*P* = 0.67) between the germinal center type

(GCB) and non-GCB subtypes (**Figure S6A**). Similarly, according to LymphPlex molecular

subtyping[12], no significant differences in OS (P = 0.68) or PFS (P = 0.71) were observed

between the BN2-like, MCD-like, EZB-like, N1-like, TP53, and other subtypes (**Figure S6B**).

We further categorized our data on the basis of the DLBCL subtyping (C0-C5) proposed by

M. A. Shipp et al.[10] Within our cohort, the C4 subtype was associated with shorter OS (*P* =

0.009, P = 0.013) and PFS (*P* = 0.096, P = 0.047) than the C0 and C5 subtypes, respectively

(**Figure S6C**). However, the overall efficacy of this DLBCL subtyping system in predicting

outcomes was less than satisfactory (OS, P = 0.074). The data were also analyzed via the

five subtypes (BN2, EZB, MCD, N1, and others)[9] and seven subtypes (A53, BN2, EZB, MCD,

N1, ST2, and others)[11] of DLBCL proposed by *Louis M Staudt* et al. There were no significant differences in OS ($P > 0.05$) or PFS ($P > 0.05$) between BN2, N1, and other categories within the 5-subtype model (**Figure S6D**). Overall, the seven-subtype model (**Figure S6E**) had inferior performance, with no significant differences observed in OS ($P = 0.61$) or PFS ($P = 0.47$). With respect to the four subtypes (CS1-4) of PCNSL proposed by A Alentorn [14], our data revealed that the OS ($P = 0.12$) and PFS ($P = 0.54$) of patients with the CS3 subtype were shorter than those of patients with the other three subtypes (**Figure S6F**), but the differences were not significant. Furthermore, the double-hit gene expression signature defines a distinct subgroup of germinal center B-cell-like DLBCL, which is associated with a poor prognosis[13]. However, in this study, there were no significant differences in OS ($P > 0.05$) or PFS ($P > 0.05$) between the double- or triple-hit PCNSL patients and the other PCNSL patients, as shown in **Figure S7** (A: all; B: discovery cohort; C: validation cohort). Taken together, these results suggest that the molecular subtyping of PCNSL and DLBCL, as previously published, is not applicable to Chinese PCNSL patients.

PCNSL molecular subtypes with clinical outcome implications

To establish a molecular classification system for PCNSL, we initially carried out consensus clustering on 58 patients (discovery cohort) who received immunochemotherapy (HD-MTX-based chemotherapy regimen followed by consolidation therapy) via WES data to generate cluster assignments (refer to Methods - Classification model to identify molecular types). WES data from 82 patients (validation cohort) who received the same treatment regimen were used to confirm the molecular clusters. Additionally, WES/WGS sequencing data from 36 newly diagnosed PCNSL patients at Fujian Cancer Hospital and The First Affiliated Hospital of Fujian Medical University were included as an independent external Chinese PCNSL cohort to further validate the molecular clusters[17].

The molecular classification scheme, presented in **Figure 4A**, **Table S3** and **Figure S8**, was determined independently of clinical information and finalized before the analysis of clinical data, allowing us to analyze the relationships between genetic subtypes and survival in this entire cohort. Finally, EP300, BTG1, MYD88, and IRF4 were included in the molecular subtyping of Chinese PCNSLs.

All nonsynonymous mutations in EP300, BTG1, MYD88, and IRF4 were visualized within the functional domains of the encoded proteins (**Figure S9**). As previously reported[10,57,58], MYD88 (L265P) was the most prevalent mutation, while EP300, BTG1, and IRF4 all harbored scattered mutations. We evaluated the prognostic significance of these four identified genes for OS and PFS. The survival outcomes of patients with EP300 mutations were significantly poorer than those of patients with wild-type EP300, and patients harboring mutations in either BTG1 (OS, P = 0.017; PFS, P = 0.12), MYD88 (OS, P = 0.016; PFS, P = 0.069), or IRF4 (OS, P = 0.045; PFS, P = 0.018) had favorable OS (**Figure S10A**) and PFS (**Figure S10B**) relative to individuals with their wild-type counterparts.

In the discovery cohort, we identified three subtypes of PCNSL: BMI (n = 16, 27.59%), E3 (n = 9, 15.52%), and UC (n = 33, 56.90%). The E3 subtype was defined as those with mutations in the EP300 gene. The PCNSLs that carried mutations in at least two of the three genes, BTG1, MYD88, and IRF4, were characterized as the BMI subtype. The other PCNSLs that did not fit into either of these categories were classified under the UC subtype (Unclassified). The three subtypes differed significantly in OS (*P* < 0.001) and PFS (*P* = 0.043), the BMI subtype (*P* < 0.001, *P* = 0.023) had much more favorable outcomes than the E3 and UC subtypes did, and the E3 subtype (*P* < 0.001, *P* = 0.037) had far worse outcomes than the BMI and UC subtypes did (**Figure 4B**). These differences between the three subtypes of PCNSL were detected in the validation cohort (OS, *P* = 0.020; PFS, *P* = 0.048; n = 82, tumor-only sample), and patients with E3 had significantly shorter survival times than those with BMI (OS, *P* = 0.007) or UC (OS, *P* = 0.043) (**Figure 4C**). Similar results were observed in the entire cohort (n = 140, OS, *P* < 0.0001; PFS, *P* = 0.0004; **Figure S10C**): 26 patients were classified as BMI (18.57%), 27 as E3 (19.29%), and 87 as UC (62.14%). Furthermore, the raw WES or WGS data, along with follow-up data from an independent cohort of 36 Chinese patients with PCNSL[17], were obtained. The three identified subtypes significantly differed in OS (P = 0.0004; **Figure 4D**). Specifically, the BMI subgroup (P = 0.0004, P = 0.005) had notably more favorable outcomes than the E3 and UC subgroups did. Conversely, the E3 subtype (P = 0.0004, P = 0.0231) had considerably worse outcomes than either the BMI or UC subgroup did. These results further validate the robustness of our

molecular classification.

BTK inhibitors have been shown to improve the prognosis of PCNSL patients by blocking B-cell receptor signaling pathways, which influence the growth and survival of B cells. Consequently, we further analyzed whether the use of BTK inhibitors affects the robustness of our classification. There were no significant differences in OS (P > 0.05, **Figure S11A**) or PFS (P > 0.05, **Figure S11B**) between patients who received BTK inhibitors (n = 8) and those who did not receive BTK inhibitors (n = 132). Additionally, after initial treatment options (HD-MTX combined with idarubicin (IDA), HD-MTX combined with rituximab (R), HD-MTX combined with IDA and R, or a combination of BTK inhibitors) and consolidated therapy (WBRT, stem cell transplant) were adjusted, multivariable Cox regression analysis revealed that the use of different treatment options did not affect the robustness of our classification (**Table S4**).

For the E3 subtype, the mean mutation prevalence in the PCNSLs of Asian patients[3,16,17] (16.98%) was significantly higher than that in the PCNSLs of Western patients[4,5,14,59–61] (4.53%) (**Figure 4E**). Patients with the E3 subtype had significantly poorer survival, which was not affected by the mutation site (OS, P = 0.82; PFS, P = 0.69; **Figure 4F**). Univariate Cox regression analysis revealed that PCNSL molecular subtypes are associated with clinical outcomes (**Table S5**), which was also confirmed by multivariate Cox regression hazard ratio analysis after adjusting for important confounders (age, IELSG, MSKCC, LDH levels, deep brain location, and CSF protein levels) (**Figure 4G**). These findings further confirmed that PCNSL molecular subtypes are associated with clinical outcomes. The key demographic and clinical characteristics of the three MS patients are summarized in **Table S6**.

We also constructed a Sankey diagram to visualize the correspondence between our samples and the previously described genetic subtypes of PCNSL and DLBCL (**Figure S12**). The Chinese PCNSL subtypes were significantly distinct from those previously described. Taken together, these results indicate that three unique molecular subtypes of Chinese PCNSL are associated with clinical outcomes.

Genomic landscape and tumor microenvironment across PCNSL subtypes

We further aimed to determine whether these three molecular subtypes of Chinese PCNSL

patients could affect the mutational landscape, oncogenic pathways and tumor microenvironment. Intergroup mutation distribution analysis was performed to identify unique and shared mutations. A total of 6,102 (median variants per sample = 196.5), 41,548 (median variants per sample = 1719), and 73,440 mutated genes (median variants per sample = 918) were identified in the BMI, E3, and UC subtypes, respectively, with 518 mutated genes shared among the three subtypes (**Figures 5A** and **S13A**). Key candidate cancer genes in the 140 PCNSL patients, grouped by these three subtypes, are presented in **Figure 5B**. The mutation frequencies of EP300, MYD88, IRF4, FAT4, KMT2C, KMT2A, MPEG1, PRKDC, SPEN, CARD11, RANBP2, NUMA1, NSD1, PASK, USP6, PTPN13, SLX4, ADGRA2, TET1, and NOTCH2 were significantly different ($P < 0.05$) among these three subtypes. Arm-level CNAs among the samples of the three subtypes are displayed in **Figure 5C**, showing significant differences in recurrent deletions (in 6q and 12p) but not in amplifications (**Figure 5D**). The results of significant recurrent focal amplifications and deletions were visualized by GISTIC 2.0 in the BMI, E3, and UC subtypes (**Figure S13B**). There was no significant difference ($P > 0.05$) in the MATH (**Figure 5E**) or MSI (**Figure 5F**) scores among the three subtypes, but the TMB was greater in the E3 subgroup than in the BMI ($P < 0.0001$) or UC subgroup ($P < 0.001$) (**Figure 5G**). Furthermore, the VAF was lower in the E3 subgroup than in the BMI ($P < 0.0001$) or UC subgroup ($P < 0.001$) (**Figure 5H**).

The contributions of mutational signatures across the three subtypes are illustrated in **Figure S13C**. The three signatures extracted from the E3 subtype samples displayed cosine similarities of 82.0%, 93.8%, and 95.2% to the COSMIC signatures 6, 10a, and 45, respectively (**Figure S13D**). In contrast, those extracted from BMI subtype samples presented similarities of 77.3%, 81.7%, and 58.3% to COSMIC signatures 5, 84, and 87, respectively (**Figure S13D**). The signatures from the UC subtype samples were 80.8%, 70.3%, and 91.7% and similar to the COSMIC signatures 5, 1, and 45 (**Figure S13D**), respectively.

Hematoxylin–eosin (H&E) and immunohistochemical (IHC) staining of CD2, CD5, CD10, CD19, CD20, CD79a, Ki-67, BCL2, BCL6, MUM1, c-Myc, and p53 from samples of different PCNSL subtypes revealed significant malignant progression in patients with the E3 subtype

of PCNSL, as shown in **Figure S14A** and **Table S7**. This is evidenced by the formation of poorly differentiated and aggressively growing tumors (Ki-67), which contrasts with the other two subtypes. Immunostaining revealed a greater presence of CD2+, CD5+, CD10+, CD19+, CD20+, and CD79a+ cells in the E3 subtype. Additionally, a greater presence of BCL2+, BCL6+, MUM1+, and c-Myc+ cells, which are markers associated with the diagnosis of B-cell lymphomas[62], was observed in the E3 subtype.

To explore potential therapeutic strategies for the genetic subtypes of PCNSL, we examined groups of genetic aberrations that target oncogenic signaling pathways (**Figure S14B**). Genetic events affecting NF-κB regulators that negatively regulate the stability of NF-κB-dependent mRNAs were detected in 85.2% and 84.6% of the E3 and BMI subtype samples but not in the UC subtype samples (P <0.05). These findings suggest that the E3 and BMI subtypes may be more responsive to BTK inhibitors[63]. The PI3 kinase pathway, a pathway that can indirectly activate NF-κB, is genetically altered in 66.7% of E3 patients[64]. We also noted a higher frequency of genetic alterations in chromatin histone modifiers (100%), RTK signaling (77.8%), protein homeostasis/ubiquitination (66.7%), cell cycle pathways (77.8%), genome integrity (82.5%), Wnt/B-catenin signaling (59.3%), the chromatin SWI/SNF complex (70.4%), and splicing (37%) in E3 patients. Therefore, a combination of cyclin D-Cdk4,6 and PI3 kinase inhibitors might be beneficial for patients with the E3 subtype.

Taken together, these results suggest that the three molecular subtypes of PCNSL in Chinese patients each have a unique genomic landscape, tumor microenvironment, and oncogenic pathways.

Mutational landscape and oncogenic pathways in PCNSLs with different sites of onset

Next, we aimed to determine whether the site of PCNSL onset in Chinese patients influences the mutational landscape and oncogenic pathways. **Figure 6A** shows that the areas most often affected by PCNSL were the basal ganglia, brainstem, cerebellum, corpus callosum, frontal lobe, occipital lobe, parietal lobe, temporal lobe, thalamus, and ventricles, as well as combinations of multiple sites. The most frequently involved sites of PCNSL were the frontal lobe (22.14%), temporal lobe (15.71%), and basal ganglia (8.57%).

First, we found that there was no significant difference in the proportion of different disease

sites among the three subtypes (chi-square test, corpus callosum, P = 0.887; basal ganglia,

P=0.645; thalamus, P= 0.317; brainstem, P= 1.000; ventricle, P= 0.631; front lobe, P= 0.576;

multiple, P= 0.949; temporal lobe, P= 0.214; cerebellum, P= 0.831; occipital lobe, P= 0.778;

parietal lobe, P= 0.887). BMI-related tumors were slightly more prevalent in the temporal lobe

(7 out of 26 patients vs. 15 out of 114 patients, chi-square test P = 0.082) than other subtypes

were. Conversely, E3 events were likely to be more frequent in the thalamus (2 out of 27

cases vs. 2 out of 113 cases, chi-square test P = 0.113). Larger studies are needed to confirm

this observation.

Next, we explored whether the tumor sites differed between the discovery and validation

cohorts. As shown in **Figure S15A**, there was no significant difference in the proportions of

different disease sites between the discovery and validation cohorts (chi-square test P =

0.610). Furthermore, the proportions of deep brain lesions in the discovery and validation

cohorts were not significantly different (P = 0.478).

Sixteen mutated genes were represented among all the sites (**Figure 6B**). Key candidate

cancer genes in the 140 PCNSL patients, grouped by site of disease onset, are presented in

Figure 6C. Mutations in the PIM1, CD79B, and MYD88 genes were relatively common across

all PCNSL onset sites. The frontal lobe had the highest mutation frequencies of five genes:

PIM1 (51.61%), MYD88 (51.61%), and ZFHX3 (48.39%). In the temporal lobe, the most

prevalent mutations were in PIM1 (68.18%), SC1. In the basal ganglia, the genes with the

highest mutation frequencies were PIM1 (66.67%), MYD88 (50%), ZFHX3 (41.67%), and

DTX1 (41.67%). The mutation frequencies of LRP1B and KMT2D were significantly different

(P< 0.05) among these sites. Arm-level CNAs among the samples of these sites are displayed

in **Figure 6D**, and significant differences in both recurrent amplifications (2q, 10q, and 10p)

and deletions (9p) were observed. The difference in the MATH score was significant ($P <$

0.05) (**Figure 6E**) across these sites, but there was no significant difference ($P > 0.05$) in the

TMB (**Figure 6F**) or MSI (**Figure 6G**) score between several sites.

Next, we further divided the tumor sites into deep brain and shallow brain tumor subgroups.

A total of 854 (81%) mutated genes were represented between the deep brain and shallow

brain tumor subgroups (**Figure S15B**). The mutational landscape between deep brain and

shallow brain tumors is shown in **Figure S15C**. The mutation frequencies of LRP1B, FAT1, KMT2B, CIC, PTPRB, MAP3KI, ARID1B, and CPS1 in the deep brain and shallow brain tumors were significantly different (P< 0.05). Arm-level CNAs between deep brain and shallow brain tumors are displayed in **Figure S15D**, and no significant differences (P >0.05) in either recurrent amplification or deletion were observed. There was no significant difference (P > 0.05) in the MATH (**Figure S15E**), TMB (**Figure S15F**), or MSI (**Figure S15G**) score between deep brain and shallow brain tumors.

We further conducted pathway enrichment analysis (**Figure 6H**) and revealed significant involvement of NF-κB signaling in the tumors of patients with onset in the parietal lobe and brainstem, each with an involvement rate of 100%. We also noted a significant difference (P <0.05) in the frequency of genetic alterations in transcription factors, other chromatin, immune signaling, NF-κB signaling, chromatin histone modifiers, RTK signaling, PI3K signaling, RNA abundance, the chromatin SWI/SNF complex, protein homeostasis/ubiquitination, and the cell cycle among these sites. Furthermore, we noted a significant difference (P <0.05) in the frequency of genetic alterations in the chromatin SWI/SNF complex but not in the other pathways (**Figure S15H**). Taken together, these results indicate that the mutational landscape in Chinese patients with PCNSL varies between sites of onset.

Discussion

Owing to the genetic, phenotypic, and tumor microenvironment heterogeneity of PCNSL, identifying classification and prognostic biomarkers for patients is extremely challenging, which in turn makes developing effective new therapies challenging. Here, we performed a study of 176 Chinese PCNSL patients that was adequately powered to expand the genomic landscape and explore its clinical significance utilizing an unprecedented sample size and a multicenter approach. We defined recurrent mutations, CNAs, and associated cancer candidate genes in Chinese PCNSL patients and compared these comprehensive genetic signatures with those of PCNSL patients and systemic DLBCL patients of other races, revealing that genetic heterogeneity is a defining feature of PCNSL. Our results highlight the complexity of PCNSLs, which have a median of 19.06 mutations/Mb and a median of 949.5

variants per sample.

Although the genetic variation profiles of the PCNSL discovery and validation samples were

similar, notable differences were evident in the mutation frequencies of certain genes (e.g.,

NOTCH1, NOTCH2, and KMT2C), the TMB, the MSI, and the MATH score. These

discrepancies might stem from variations in specimen types (fresh tissue versus FFPE) and

the absence of paired samples in the validation cohort[65]. Furthermore, genetic heterogeneity

between patient populations[15] and biological differences between PCNSL and DLBCL[13] may

be the main reasons why the published classifications for DLBCL and PCNSL are not

applicable to Chinese PCNSL patients. Currently, the classification frameworks for systemic

DLBCL offer valuable insights into the underlying biology based on molecular mechanisms.

However, it is important to note that biological insights do not necessarily correlate with

clinical outcomes, especially given that treatment paradigms may evolve over time.

More importantly, in our study, we identified three robust molecular subtypes of PCNSL in

Chinese patients: BMI, E3, and UC. Distinct clinical outcomes, activated cellular pathways,

and alterations in the genetic landscape distinguish the three subtypes and could be utilized

to personalize therapy for patients with different subtypes. For the E3 subtype, the mean

mutation rate was significantly higher in the PCNSLs of Chinese and Japanese patients

(16.88%) than in those of Western PCNSL patients (4.53%). This difference may be attributed

to variations in age, lifestyle, and EBV infection rates[17,41,66]. Patients with the E3 subtype had

significantly poorer survival outcomes, a finding that is consistent with reported findings in

DLBCL[67]. EP300 and CREBBP are two closely related members of the KAT3 family of histone

acetyltransferases that function similarly[68]. However, in this study, the survival patients with

EP300 mutations was significantly poorer, whereas CREBBP mutations did not contribute to

an inferior prognosis (**Figure S4**). This discrepancy might stem from the distinct roles of

EP300 and CREBBP in PCNSL. Specifically, in a recent study, EP300, but not CREBBP, was

reported to play an essential role in supporting the viability of classical Hodgkin's lymphoma

by directly modulating the expression of the oncogenic MYC/IRF4 network, surface receptor

CD30, immunoregulatory cytokine interleukin 10, and immune checkpoint protein PD-L1[69].

The molecular mechanisms through which EP300 mutations facilitate the progression of

PCNSL are still not understood. We plan to explore these mechanisms in a forthcoming study.

PCNSLs that carried mutations in at least two of three genes, namely, BTG1, MYD88, and IRF4, were characterized as the BMI subtype with favorable survival outcomes. The prognostic significance of MYD88 mutations in PCNSL patients remains inconclusive. While several studies have reported no effect on OS[70–73], only two have reported an unfavorable outcome[74,75]. In line with our findings, *Olimpia* et al.[76] and *Claudio* et al.[77] have indicated that the survival time is significantly longer in patients with PCNSLs harboring MYD88 mutations. This discrepancy may be attributed to heterogeneity in genetics, phenotype, and the tumor microenvironment across various patient populations; variances in sample size; and the adjustment of confounding variables, including age and treatment methods. In this large study, we found that Chinese PCNSL patients harboring MYD88 mutations have longer survival times. However, further research is needed to verify these findings in diverse populations. The associations of IRF4 and BTG1 mutations with clinical outcomes in PCNSL patients have not been previously reported. This results of this study revealed that mutations in both IRF4 and BTG1 were associated with favorable outcomes. IRF4 has been identified as a driver oncogene that transcriptionally regulates downstream target genes, such as MYC, and coordinates the transcriptional program with NF-κB references[78–81]. Numerous in vitro and clinical studies have highlighted the abnormal overexpression and oncogenic roles of IRF4 in various mature lymphoid neoplasms[81–84]. Therefore, we hypothesized that the reduced expression and/or decreased transcriptional activity of IRF4 resulting from such mutations may explain the association of IRF4 mutations with favorable survival outcomes in PCNSL patients. BTG1 mutation has been associated with extranodal dissemination and unfavorable outcomes in B-cell lymphoma patients[85–87]. However, in this study, BTG1 mutation was linked to favorable outcomes in PCNSL patients. In a substantial discovery cohort consisting of 533 patients with B-cell precursor acute lymphoblastic leukemia, *Blanca* and colleagues[88] reported that deletions of BTG1 alone did not affect prognosis. The complete remission rate among acute myeloid leukemia patients with low BTG1 expression was significantly greater than that among patients with high BTG1 expression[89]. The molecular mechanisms through which mutations in BTG1 decelerate the progression of PCNSL warrant further research.

This study has several limitations. First, the proportion of unclassified PCNSLs (UC subtypes) is high, possibly due to the lack of multiomics data needed for classification. However, the three-class classification based on WES proposed in this study has strong clinical practicality and translational value, as it relies solely on the four genes identified through WES. In other words, clinicians can predict patient outcomes and further perform individualized management of PCNSLs by detecting mutations in these four genes without the need for additional multiomics testing. Second, the three identified molecular subtypes can be used to predict patient outcomes. However, it remains challenging to determine whether these three molecular subtypes can still be used to accurately distinguish biological subtypes without additional RNA-seq data layers.

In summary, by performing genomic sequencing on tumor specimens from 176 Chinese PCNSL patients, we identified three molecular subtypes that can be used to predict patient outcomes. These genetic signatures associated with PCNSL outcomes illuminate the path toward individualized management of PCNSL patients and the discovery of novel treatment strategies for this challenging disease.

Data and code availability

The raw sequence data have been deposited in the Genome Sequence Archive[90] in the National Genomics Data Center[91], China National Center for Bioinformation/Beijing Institute of Genomics, Chinese Academy of Sciences (GSA-Human HRA006122), which is publicly accessible at https://ngdc.cncb.ac.cn/gsa-human. Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

The raw sequencing data of the DLBCL WES data are accessible from the TCGA database and were reanalyzed via pipelines and filter settings that were identical to those used in the present study. The raw WES data of the Japanese PCNSL patients were obtained from the Japanese Genotype–Phenotype Archive (JGA, http://trace.ddbj.nig.ac.jp/jga) and reanalyzed via pipelines and filtering settings that were identical to those used in the present study, which is hosted by DDBJ under the accession number JGAS000000000021[3]. The raw sequencing WES/WGS data of an external independent Chinese PCNSL cohort (n = 36)[17] were obtained

from the China National Center for Bioinformation/Beijing Institute of Genomics, Chinese Academy of Sciences and reanalyzed via pipelines and filtering settings that were identical to those used in the present study, under the accession number GSA-Human HRA002475.

The frequencies of recurrent genetic alterations in DLBCLs[49] and French PCNSLs[14] are accessible from the manuscript or supplemental information and were not reanalyzed via identical pipelines and filtering settings as those used in the present study.

Acknowledgment: This study was funded by the National Natural Science Foundation of China (82302582), the Shanghai Municipal Health Commission Project (20224Y0317), the Shanghai Science and Technology Commission (17430750200), the Join Breakthrough Project for New Frontier Technologies of Shanghai Hospital Development Center (SHDC12016120), and the Youth Medical Talents–Clinical Laboratory Practitioner Program (2022-65).

Competing Interests: The authors declare no potential conflicts of interest.

Author contributions: SJ. L, CX. L, DH. L, WJ. C, ZG. X, XT. Z, and Y. M conceived and designed the project. SJ. L, DH. L, WJ. C, JZ. C, J. R, and ZG. X collected the clinical samples. SJ. L, JN. W, JZ. C, LY. Z, HW. Y, and Y. S analyzed the WES data and performed bioinformatic analyses. SJ. L, DH. L, J. R, LY. Z, HW. Y, and JN. W integrated the sequencing data, drew the display items. SJ. L, DH. L, and CX. L wrote the manuscript. XT. Z and Y. M oversaw the ethical guidelines and data regulation. WJ. C and Y. M supervised the project. All of the authors contributed to the final version of the paper.

References

1. Chen, T., Liu, Y., Wang, Y., Chang, Q., Wu, J., Wang, Z., Geng, D., Yu, J.-T., Li, Y., Li, X.-Q., et al. (2022). Evidence-based expert consensus on the management of primary central nervous system lymphoma in China. J Hematol Oncol *15*, 136. https://doi.org/10.1186/s13045-022-01356-7.

2. Ferreri, A.J.M., Calimeri, T., Cwynarski, K., Dietrich, J., Grommes, C., Hoang-Xuan, K., Hu, L.S., Illerhaus, G., Nayak, L., Ponzoni, M., et al. (2023). Primary central nervous system lymphoma. Nat. Rev. Dis. Primers *9*, 29. https://doi.org/10.1038/s41572-023-00439-0.

3. Fukumura, K., Kawazu, M., Kojima, S., Ueno, T., Sai, E., Soda, M., Ueda, H., Yasuda, T., Yamaguchi, H., Lee,
J., et al. (2016). Genomic characterization of primary central nervous system lymphoma. Acta Neuropathol.
(Berl.) *131*, 865–875. https://doi.org/10.1007/s00401-016-1536-2.

4. Braggio, E., Van Wier, S., Ojha, J., McPhail, E., Asmann, Y.W., Egan, J., Da Silva, J.A., Schiff, D., Lopes, M.B.,
Decker, P.A., et al. (2015). Genome-Wide Analysis Uncovers Novel Recurrent Alterations in Primary Central
Nervous System Lymphomas. Clin. Cancer Res. *21*, 3986–3994. https://doi.org/10.1158/1078-0432.CCR-
14-2116.

5. Radke, J., Ishaque, N., Koll, R., Gu, Z., Schumann, E., Sieverling, L., Uhrig, S., Hübschmann, D., Toprak, U.H.,
López, C., et al. (2022). The genomic and transcriptional landscape of primary central nervous system
lymphoma. Nat. Commun. *13*, 2558. https://doi.org/10.1038/s41467-022-30050-y.

6. Vater, I., Montesinos-Rongen, M., Schlesner, M., Haake, A., Purschke, F., Sprute, R., Mettenmeyer, N.,
Nazzal, I., Nagel, I., Gutwein, J., et al. (2015). The mutational pattern of primary lymphoma of the central
nervous system determined by whole-exome sequencing. Leukemia *29*, 677–685.
https://doi.org/10.1038/leu.2014.264.

7. Rosenwald, A., Wright, G., Chan, W.C., Connors, J.M., Campo, E., Fisher, R.I., Gascoyne, R.D., Muller-
Hermelink, H.K., Smeland, E.B., Giltnane, J.M., et al. (2002). The use of molecular profiling to predict survival
after chemotherapy for diffuse large-B-cell lymphoma. N. Engl. J. Med. *346*, 1937–1947.
https://doi.org/10.1056/NEJMoa012914.

8. Wilson, W.H., Young, R.M., Schmitz, R., Yang, Y., Pittaluga, S., Wright, G., Lih, C.-J., Williams, P.M., Shaffer,
A.L., Gerecitano, J., et al. (2015). Targeting B cell receptor signaling with ibrutinib in diffuse large B cell
lymphoma. Nat. Med. *21*, 922–926. https://doi.org/10.1038/nm.3884.

9. Schmitz, R., Wright, G.W., Huang, D.W., Johnson, C.A., Phelan, J.D., Wang, J.Q., Roulland, S., Kasbekar, M.,
Young, R.M., Shaffer, A.L., et al. (2018). Genetics and Pathogenesis of Diffuse Large B-Cell Lymphoma. N.
Engl. J. Med. *378*, 1396–1407. https://doi.org/10.1056/NEJMoa1801445.

10. Chapuy, B., Stewart, C., Dunford, A.J., Kim, J., Kamburov, A., Redd, R.A., Lawrence, M.S., Roemer, M.G.M.,
Li, A.J., Ziepert, M., et al. (2018). Molecular subtypes of diffuse large B cell lymphoma are associated with
distinct pathogenic mechanisms and outcomes. Nat. Med. *24*, 679–690. https://doi.org/10.1038/s41591-
018-0016-8.

11. Wright, G.W., Huang, D.W., Phelan, J.D., Coulibaly, Z.A., Roulland, S., Young, R.M., Wang, J.Q., Schmitz, R.,
Morin, R.D., Tang, J., et al. (2020). A Probabilistic Classification Tool for Genetic Subtypes of Diffuse Large
B Cell Lymphoma with Therapeutic Implications. Cancer Cell *37*, 551-568.e14.
https://doi.org/10.1016/j.ccell.2020.03.015.

12. Shen, R., Fu, D., Dong, L., Zhang, M.-C., Shi, Q., Shi, Z.-Y., Cheng, S., Wang, L., Xu, P.-P., and Zhao, W.-L.
(2023). Simplified algorithm for genetic subtyping in diffuse large B-cell lymphoma. Signal Transduct.
Target. Ther. *8*, 145. https://doi.org/10.1038/s41392-023-01358-y.

13. Swerdlow, S.H., Campo, E., Pileri, S.A., Harris, N.L., Stein, H., Siebert, R., Advani, R., Ghielmini, M., Salles,

G.A., Zelenetz, A.D., et al. (2016). The 2016 revision of the World Health Organization classification of
lymphoid neoplasms. Blood *127*, 2375–2390. https://doi.org/10.1182/blood-2016-01-643569.

14. Hernández-Verdin, I., Kirasic, E., Wienand, K., Mokhtari, K., Eimer, S., Loiseau, H., Rousseau, A., Paillassa,
J., Ahle, G., Lerintiu, F., et al. (2023). Molecular and clinical diversity in primary central nervous system
lymphoma. Ann. Oncol. *34*, 186–199. https://doi.org/10.1016/j.annonc.2022.11.002.

15. Yuan, X., Yu, T., Zhao, J., Jiang, H., Hao, Y., Lei, W., Liang, Y., Li, B., and Qian, W. (2023). Analysis of the
genomic landscape of primary central nervous system lymphoma using whole-genome sequencing in
Chinese patients. Front. Med. *17*, 889–906. https://doi.org/10.1007/s11684-023-0994-x.

16. He, X., Fan, X., Shan, Y., Ji, X., Su, L., and Wang, Y. (2023). Analysis of genomic alterations in primary central
nervous system lymphoma. Medicine (Baltimore) *102*, e34931.
https://doi.org/10.1097/MD.0000000000034931.

17. Zhu, Q., Wang, J., Zhang, W., Zhu, W., Wu, Z., Chen, Y., Chen, M., Zheng, L., Tang, J., Zhang, S., et al. (2022).
Whole-Genome/Exome Sequencing Uncovers Mutations and Copy Number Variations in Primary Diffuse
Large B-Cell Lymphoma of the Central Nervous System. Front Genet *13*, 878618.
https://doi.org/10.3389/fgene.2022.878618.

18. Abrey, L.E., Batchelor, T.T., Ferreri, A.J.M., Gospodarowicz, M., Pulczynski, E.J., Zucca, E., Smith, J.R., Korfel,
A., Soussain, C., DeAngelis, L.M., et al. (2005). Report of an international workshop to standardize baseline
evaluation and response criteria for primary CNS lymphoma. J Clin Oncol *23*, 5034–5043.
https://doi.org/10.1200/JCO.2005.13.524.

19. Hoang-Xuan, K., Bessell, E., Bromberg, J., Hottinger, A.F., Preusser, M., Rudà, R., Schlegel, U., Siegal, T.,
Soussain, C., Abacioglu, U., et al. (2015). Diagnosis and treatment of primary CNS lymphoma in
immunocompetent patients: guidelines from the European Association for Neuro-Oncology. Lancet
Oncol *16*, e322-332. https://doi.org/10.1016/S1470-2045(15)00076-5.

20. Li, Q., Ma, J., Ma, Y., Lin, Z., Kang, H., and Chen, B. (2021). Improvement of outcomes of an escalated high-
dose methotrexate-based regimen for patients with newly diagnosed primary central nervous system
lymphoma: a real-world cohort study. Cancer Manag Res *13*, 6115–6122.
https://doi.org/10.2147/CMAR.S322467.

21. Freed, D., Aldana, R., Weber, J.A., and Edwards, J.S. (2017). The Sentieon Genomics Tools–A fast and
accurate solution to variant calling from next-generation sequence data. BioRxiv, 115717.

22. McLaren, W., Gil, L., Hunt, S.E., Riat, H.S., Ritchie, G.R.S., Thormann, A., Flicek, P., and Cunningham, F.
(2016). The Ensembl Variant Effect Predictor. Genome Biol *17*, 122. https://doi.org/10.1186/s13059-016-
0974-4.

23. Mayakonda, A., Lin, D.-C., Assenov, Y., Plass, C., and Koeffler, H.P. (2018). Maftools: efficient and
comprehensive analysis of somatic variants in cancer. Genome Res *28*, 1747–1756.
https://doi.org/10.1101/gr.239244.118.

24. Wang, Y., Bae, T., Thorpe, J., Sherman, M.A., Jones, A.G., Cho, S., Daily, K., Dou, Y., Ganz, J., Galor, A., et al.

(2021). Comprehensive identification of somatic nucleotide variants in human brain tissue. Genome Biol
22, 92. https://doi.org/10.1186/s13059-021-02285-3.

25. Sathirapongsasuti, J.F., Lee, H., Horst, B.A.J., Brunner, G., Cochran, A.J., Binder, S., Quackenbush, J., and
Nelson, S.F. (2011). Exome sequencing-based copy-number variation and loss of heterozygosity
detection: ExomeCNV. Bioinformatics *27*, 2648–2654. https://doi.org/10.1093/bioinformatics/btr462.

26. Favero, F., Joshi, T., Marquard, A.M., Birkbak, N.J., Krzystanek, M., Li, Q., Szallasi, Z., and Eklund, A.C. (2015).
Sequenza: allele-specific copy number and mutation profiles from tumor sequencing data. Ann Oncol *26*,
64–70. https://doi.org/10.1093/annonc/mdu479.

27. Mermel, C.H., Schumacher, S.E., Hill, B., Meyerson, M.L., Beroukhim, R., and Getz, G. (2011). GISTIC2.0
facilitates sensitive and confident localization of the targets of focal somatic copy-number alteration in
human cancers. Genome Biol *12*, R41. https://doi.org/10.1186/gb-2011-12-4-r41.

28. Chakravarty, D., Gao, J., Phillips, S.M., Kundra, R., Zhang, H., Wang, J., Rudolph, J.E., Yaeger, R., Soumerai,
T., Nissan, M.H., et al. (2017). OncoKB: A Precision Oncology Knowledge Base. JCO Precis Oncol *2017*,
PO.17.00011. https://doi.org/10.1200/PO.17.00011.

29. Vogelstein, B., Papadopoulos, N., Velculescu, V.E., Zhou, S., Diaz, L.A., and Kinzler, K.W. (2013). Cancer
genome landscapes. Science *339*, 1546–1558. https://doi.org/10.1126/science.1235122.

30. Griffith, M., Spies, N.C., Krysiak, K., McMichael, J.F., Coffman, A.C., Danos, A.M., Ainscough, B.J., Ramirez,
C.A., Rieke, D.T., Kujan, L., et al. (2017). CIViC is a community knowledgebase for expert crowdsourcing
the clinical interpretation of variants in cancer. Nat Genet *49*, 170–174. https://doi.org/10.1038/ng.3774.

31. Alexandrov, L.B., Kim, J., Haradhvala, N.J., Huang, M.N., Tian Ng, A.W., Wu, Y., Boot, A., Covington, K.R.,
Gordenin, D.A., Bergstrom, E.N., et al. (2020). The repertoire of mutational signatures in human cancer.
Nature *578*, 94–101. https://doi.org/10.1038/s41586-020-1943-3.

32. Chalmers, Z.R., Connelly, C.F., Fabrizio, D., Gay, L., Ali, S.M., Ennis, R., Schrock, A., Campbell, B., Shlien, A.,
Chmielecki, J., et al. (2017). Analysis of 100,000 human cancer genomes reveals the landscape of tumor
mutational burden. Genome Med *9*, 34. https://doi.org/10.1186/s13073-017-0424-2.

33. Mroz, E.A., and Rocco, J.W. (2013). MATH, a novel measure of intratumor genetic heterogeneity, is high
in poor-outcome classes of head and neck squamous cell carcinoma. Oral Oncol *49*, 211–215.
https://doi.org/10.1016/j.oraloncology.2012.09.007.

34. Sanchez-Vega, F., Mina, M., Armenia, J., Chatila, W.K., Luna, A., La, K.C., Dimitriadoy, S., Liu, D.L., Kantheti,
H.S., Saghafinia, S., et al. (2018). Oncogenic Signaling Pathways in The Cancer Genome Atlas. Cell *173*,
321-337.e10. https://doi.org/10.1016/j.cell.2018.03.035.

35. Garcia, M., Juhos, S., Larsson, M., Olason, P.I., Martin, M., Eisfeldt, J., DiLorenzo, S., Sandgren, J., Díaz De
Ståhl, T., Ewels, P., et al. (2020). Sarek: A portable workflow for whole-genome sequencing analysis of
germline and somatic variants. F1000Res *9*, 63. https://doi.org/10.12688/f1000research.16665.2.

36. Ewels, P.A., Peltzer, A., Fillinger, S., Patel, H., Alneberg, J., Wilm, A., Garcia, M.U., Di Tommaso, P., and

Nahnsen, S. (2020). The nf-core framework for community-curated bioinformatics pipelines. Nat
Biotechnol *38*, 276–278. https://doi.org/10.1038/s41587-020-0439-x.

37. Grüning, B., Dale, R., Sjödin, A., Chapman, B.A., Rowe, J., Tomkins-Tinch, C.H., Valieris, R., Köster, J., and
Bioconda Team (2018). Bioconda: sustainable and comprehensive software distribution for the life
sciences. Nat Methods *15*, 475–476. https://doi.org/10.1038/s41592-018-0046-7.

38. da Veiga Leprevost, F., Grüning, B.A., Alves Aflitos, S., Röst, H.L., Uszkoreit, J., Barsnes, H., Vaudel, M.,
Moreno, P., Gatto, L., Weber, J., et al. (2017). BioContainers: an open-source and community-driven
framework for software standardization. Bioinformatics *33*, 2580–2582.
https://doi.org/10.1093/bioinformatics/btx192.

39. P, D.T., M, C., Ew, F., Pp, B., E, P., and C, N. (2017). Nextflow enables reproducible computational workflows.
Nature biotechnology *35*. https://doi.org/10.1038/nbt.3820.

40. Gandhi, M.K., Hoang, T., Law, S.C., Brosda, S., O'Rourke, K., Tobin, J.W.D., Vari, F., Murigneux, V., Fink, L.,
Gunawardana, J., et al. (2021). EBV-associated primary CNS lymphoma occurring after
immunosuppression is a distinct immunobiological entity. Blood *137*, 1468–1477.
https://doi.org/10.1182/blood.2020008520.

41. Kaulen, L.D., Denisova, E., Hinz, F., Hai, L., Friedel, D., Henegariu, O., Hoffmann, D.C., Ito, J., Kourtesakis,
A., Lehnert, P., et al. (2023). Integrated genetic analyses of immunodeficiency-associated Epstein-Barr
virus- (EBV) positive primary CNS lymphomas. Acta Neuropathol *146*, 499–514.
https://doi.org/10.1007/s00401-023-02613-w.

42. Ren, W., Ye, X., Su, H., Li, W., Liu, D., Pirmoradian, M., Wang, X., Zhang, B., Zhang, Q., Chen, L., et al. (2018).
Genetic landscape of hepatitis B virus-associated diffuse large B-cell lymphoma. Blood *131*, 2670–2681.
https://doi.org/10.1182/blood-2017-11-817601.

43. Berhan, A., Bayleyegn, B., and Getaneh, Z. (2022). HIV/AIDS Associated Lymphoma: Review. Blood
Lymphat Cancer *12*, 31–45. https://doi.org/10.2147/BLCTT.S361320.

44. Pasqualucci, L., Trifonov, V., Fabbri, G., Ma, J., Rossi, D., Chiarenza, A., Wells, V.A., Grunn, A., Messina, M.,
Elliot, O., et al. (2011). Analysis of the coding genome of diffuse large B-cell lymphoma. Nat Genet *43*,
830–837. https://doi.org/10.1038/ng.892.

45. Morin, R.D., Mendez-Lago, M., Mungall, A.J., Goya, R., Mungall, K.L., Corbett, R.D., Johnson, N.A., Severson,
T.M., Chiu, R., Field, M., et al. (2011). Frequent mutation of histone-modifying genes in non-Hodgkin
lymphoma. Nature *476*, 298–303. https://doi.org/10.1038/nature10351.

46. Lohr, J.G., Stojanov, P., Lawrence, M.S., Auclair, D., Chapuy, B., Sougnez, C., Cruz-Gordillo, P., Knoechel,
B., Asmann, Y.W., Slager, S.L., et al. (2012). Discovery and prioritization of somatic mutations in diffuse
large B-cell lymphoma (DLBCL) by whole-exome sequencing. Proc Natl Acad Sci U S A *109*, 3879–3884.
https://doi.org/10.1073/pnas.1121343109.

47. Morin, R.D., Mungall, K., Pleasance, E., Mungall, A.J., Goya, R., Huff, R.D., Scott, D.W., Ding, J., Roth, A.,
Chiu, R., et al. (2013). Mutational and structural analysis of diffuse large B-cell lymphoma using whole-

genome sequencing. Blood *122*, 1256–1265. https://doi.org/10.1182/blood-2013-02-483727.

48. de Miranda, N.F.C.C., Georgiou, K., Chen, L., Wu, C., Gao, Z., Zaravinos, A., Lisboa, S., Enblad, G., Teixeira,
M.R., Zeng, Y., et al. (2014). Exome sequencing reveals novel mutation targets in diffuse large B-cell
lymphomas derived from Chinese patients. Blood *124*, 2544–2553. https://doi.org/10.1182/blood-2013-
12-546309.

49. Reddy, A., Zhang, J., Davis, N.S., Moffitt, A.B., Love, C.L., Waldrop, A., Leppa, S., Pasanen, A., Meriranta, L.,
Karjalainen-Lindsberg, M.-L., et al. (2017). Genetic and Functional Drivers of Diffuse Large B Cell
Lymphoma. Cell *171*, 481-494.e15. https://doi.org/10.1016/j.cell.2017.09.027.

50. Bruno, A., Boisselier, B., Labreche, K., Marie, Y., Polivka, M., Jouvet, A., Adam, C., Figarella-Branger, D.,
Miquel, C., Eimer, S., et al. (2014). Mutational analysis of primary central nervous system lymphoma.
Oncotarget *5*, 5065–5075. https://doi.org/10.18632/oncotarget.2080.

51. Chapuy, B., Roemer, M.G.M., Stewart, C., Tan, Y., Abo, R.P., Zhang, L., Dunford, A.J., Meredith, D.M.,
Thorner, A.R., Jordanova, E.S., et al. (2016). Targetable genetic features of primary testicular and primary
central nervous system lymphomas. Blood *127*, 869–881. https://doi.org/10.1182/blood-2015-10-
673236.

52. Alexandrov, L.B., Nik-Zainal, S., Wedge, D.C., Aparicio, S.A.J.R., Behjati, S., Biankin, A.V., Bignell, G.R., Bolli,
N., Borg, A., Børresen-Dale, A.-L., et al. (2013). Signatures of mutational processes in human cancer.
Nature *500*, 415–421. https://doi.org/10.1038/nature12477.

53. Rizvi, N.A., Hellmann, M.D., Snyder, A., Kvistborg, P., Makarov, V., Havel, J.J., Lee, W., Yuan, J., Wong, P.,
Ho, T.S., et al. (2015). Cancer immunology. Mutational landscape determines sensitivity to PD-1 blockade
in non-small cell lung cancer. Science *348*, 124–128. https://doi.org/10.1126/science.aaa1348.

54. Yamamoto, H., Watanabe, Y., Arai, H., Umemoto, K., Tateishi, K., and Sunakawa, Y. (2024). Microsatellite
instability: A 2024 update. Cancer Sci *115*, 1738–1748. https://doi.org/10.1111/cas.16160.

55. Alizadeh, A.A., Eisen, M.B., Davis, R.E., Ma, C., Lossos, I.S., Rosenwald, A., Boldrick, J.C., Sabet, H., Tran, T.,
Yu, X., et al. (2000). Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling.
Nature *403*, 503–511. https://doi.org/10.1038/35000501.

56. Hans, C.P., Weisenburger, D.D., Greiner, T.C., Gascoyne, R.D., Delabie, J., Ott, G., Müller-Hermelink, H.K.,
Campo, E., Braziel, R.M., Jaffe, E.S., et al. (2004). Confirmation of the molecular classification of diffuse
large B-cell lymphoma by immunohistochemistry using a tissue microarray. Blood *103*, 275–282.
https://doi.org/10.1182/blood-2003-05-1545.

57. Phelan, J.D., Young, R.M., Webster, D.E., Roulland, S., Wright, G.W., Kasbekar, M., Shaffer, A.L., Ceribelli,
M., Wang, J.Q., Schmitz, R., et al. (2018). A multiprotein supercomplex controlling oncogenic signalling in
lymphoma. Nature *560*, 387–391. https://doi.org/10.1038/s41586-018-0290-0.

58. Ngo, V.N., Young, R.M., Schmitz, R., Jhavar, S., Xiao, W., Lim, K.-H., Kohlhammer, H., Xu, W., Yang, Y., Zhao,
H., et al. (2011). Oncogenically active MYD88 mutations in human lymphoma. Nature *470*, 115–119.
https://doi.org/10.1038/nature09671.

59. Geng, H., Mo, S.S., Chen, L., Ballapuram, A., Tsang, M., Lu, M., Rauschecker, A.M., Wen, K.W., Devine, W.P.,
Solomon, D.A., et al. (2024). Identification of Genomic Biomarkers of Disease Progression and Survival in
Primary   CNS   Lymphoma.   Blood   Adv,   bloodadvances.2024014460.
https://doi.org/10.1182/bloodadvances.2024014460.

60. Zorofchian, S., El-Achi, H., Yan, Y., Esquenazi, Y., and Ballester, L.Y. (2018). Characterization of genomic
alterations in primary central nervous system lymphomas. J Neurooncol *140*, 509–517.
https://doi.org/10.1007/s11060-018-2990-6.

61. Barakat, M., Albitar, M., Whitney, R., and Abdulhaq, H. (2021). Diversity of genetic alterations of primary
central nervous system lymphoma in Hispanic versus non-Hispanic patients. Cancer Treat Res Commun
27, 100310. https://doi.org/10.1016/j.ctarc.2021.100310.

62. van Imhoff, G.W., Boerma, E.-J.G., van der Holt, B., Schuuring, E., Verdonck, L.F., Kluin-Nelemans, H.C.,
and Kluin, P.M. (2006). Prognostic impact of germinal center-associated proteins and chromosomal
breakpoints in poor-risk diffuse large B-cell lymphoma. J Clin Oncol *24*, 4135–4142.
https://doi.org/10.1200/JCO.2006.05.5897.

63. Iwasaki, H., Takeuchi, O., Teraguchi, S., Matsushita, K., Uehata, T., Kuniyoshi, K., Satoh, T., Saitoh, T.,
Matsushita, M., Standley, D.M., et al. (2011). The IκB kinase complex regulates the stability of cytokine-
encoding mRNA induced by TLR-IL-1R by controlling degradation of regnase-1. Nat Immunol *12*, 1167–
1175. https://doi.org/10.1038/ni.2137.

64. Kloo, B., Nagel, D., Pfeifer, M., Grau, M., Düwel, M., Vincendeau, M., Dörken, B., Lenz, P., Lenz, G., and
Krappmann, D. (2011). Critical role of PI3K signaling for NF-kappaB-dependent survival in a subset of
activated B-cell-like diffuse large B-cell lymphoma cells. Proc Natl Acad Sci U S A *108*, 272–277.
https://doi.org/10.1073/pnas.1008969108.

65. Spencer, D.H., Sehn, J.K., Abel, H.J., Watson, M.A., Pfeifer, J.D., and Duncavage, E.J. (2013). Comparison of
clinical targeted next-generation sequence data from formalin-fixed and fresh-frozen tissue specimens.
J Mol Diagn *15*, 623–633. https://doi.org/10.1016/j.jmoldx.2013.05.004.

66. Yokoyama, A., Kakiuchi, N., Yoshizato, T., Nannya, Y., Suzuki, H., Takeuchi, Y., Shiozawa, Y., Sato, Y., Aoki,
K., Kim, S.K., et al. (2019). Age-related remodelling of oesophageal epithelia by mutated cancer drivers.
Nature *565*, 312–317. https://doi.org/10.1038/s41586-018-0811-x.

67. Huang, Y.-H., Cai, K., Xu, P.-P., Wang, L., Huang, C.-X., Fang, Y., Cheng, S., Sun, X.-J., Liu, F., Huang, J.-Y.,
et al. (2021). CREBBP/EP300 mutations promoted tumor progression in diffuse large B-cell lymphoma
through altering tumor-associated macrophage polarization via FBXW7-NOTCH-CCL2/CSF1 axis. Signal
Transduct Target Ther *6*, 10. https://doi.org/10.1038/s41392-020-00437-8.

68. Meyer, S.N., Scuoppo, C., Vlasevska, S., Bal, E., Holmes, A.B., Holloman, M., Garcia-Ibanez, L., Nataraj, S.,
Duval, R., Vantrimpont, T., et al. (2019). Unique and Shared Epigenetic Programs of the CREBBP and EP300
Acetyltransferases in Germinal Center B Cells Reveal Targetable Dependencies in Lymphoma. Immunity
51, 535-547.e9. https://doi.org/10.1016/j.immuni.2019.08.006.

69. Wei, W., Song, Z., Chiba, M., Wu, W., Jeong, S., Zhang, J.-P., Kadin, M.E., Nakagawa, M., and Yang, Y.
(2023). Analysis and therapeutic targeting of the EP300 and CREBBP acetyltransferases in anaplastic large
cell lymphoma and Hodgkin lymphoma. Leukemia *37*, 396–407. https://doi.org/10.1038/s41375-022-
01774-z.

70. Nayyar, N., White, M.D., Gill, C.M., Lastrapes, M., Bertalan, M., Kaplan, A., D'Andrea, M.R., Bihun, I., Kaneb,
A., Dietrich, J., et al. (2019). MYD88 L265P mutation and CDKN2A loss are early mutational events in
primary central nervous system diffuse large B-cell lymphomas. Blood Adv *3*, 375–383.
https://doi.org/10.1182/bloodadvances.2018027672.

71. Zhou, Y., Liu, W., Xu, Z., Zhu, H., Xiao, D., Su, W., Zeng, R., Feng, Y., Duan, Y., Zhou, J., et al. (2018). Analysis
of Genomic Alteration in Primary Central Nervous System Lymphoma and the Expression of Some Related
Genes. Neoplasia *20*, 1059–1069. https://doi.org/10.1016/j.neo.2018.08.012.

72. Zheng, M., Perry, A.M., Bierman, P., Loberiza, F., Nasr, M.R., Szwajcer, D., Del Bigio, M.R., Smith, L.M.,
Zhang, W., and Greiner, T.C. (2017). Frequency of MYD88 and CD79B mutations, and MGMT methylation
in primary central nervous system diffuse large B-cell lymphoma. Neuropathology *37*, 509–516.
https://doi.org/10.1111/neup.12405.

73. Yamada, S., Ishida, Y., Matsuno, A., and Yamazaki, K. (2015). Primary diffuse large B-cell lymphomas of
central nervous system exhibit remarkably high prevalence of oncogenic MYD88 and CD79B mutations.
Leuk Lymphoma *56*, 2141–2145. https://doi.org/10.3109/10428194.2014.979413.

74. Takano, S., Hattori, K., Ishikawa, E., Narita, Y., Iwadate, Y., Yamaguchi, F., Nagane, M., Akimoto, J., Oka, H.,
Tanaka, S., et al. (2018). MyD88 Mutation in Elderly Predicts Poor Prognosis in Primary Central Nervous
System Lymphoma: Multi-Institutional Analysis. World Neurosurg *112*, e69–e73.
https://doi.org/10.1016/j.wneu.2017.12.028.

75. Hattori, K., Sakata-Yanagimoto, M., Okoshi, Y., Goshima, Y., Yanagimoto, S., Nakamoto-Matsubara, R.,
Sato, T., Noguchi, M., Takano, S., Ishikawa, E., et al. (2017). MYD88 (L265P) mutation is associated with an
unfavourable outcome of primary central nervous system lymphoma. Br J Haematol *177*, 492–494.
https://doi.org/10.1111/bjh.14080.

76. Curran, O.E., Poon, M.T.C., Gilroy, L., Torgersen, A., Smith, C., and Al-Qsous, W. (2021). MYD88 L265P
mutation in primary central nervous system lymphoma is associated with better survival: A single-center
experience. Neurooncol Adv *3*, vdab090. https://doi.org/10.1093/noajnl/vdab090.

77. Agostinelli, C., Morandi, L., Righi, S., Cirillo, L., Iommi, M., Tonon, C., Mazzatenta, D., Zoli, M., Rossi, M.,
Bagnato, G., et al. (2023). Genomic Profiling of Primary Diffuse Large B-Cell Lymphoma of the Central
Nervous System Suggests Novel Potential Therapeutic Targets. Mod Pathol *36*, 100323.
https://doi.org/10.1016/j.modpat.2023.100323.

78. Wong, R.W.J., Tan, T.K., Amanda, S., Ngoc, P.C.T., Leong, W.Z., Tan, S.H., Asamitsu, K., Hibi, Y., Ueda, R.,
Okamoto, T., et al. (2020). Feed-forward regulatory loop driven by IRF4 and NF-κB in adult T-cell
leukemia/lymphoma. Blood *135*, 934–947. https://doi.org/10.1182/blood.2019002639.

79. Tsuboi, K., Iida, S., Inagaki, H., Kato, M., Hayami, Y., Hanamura, I., Miura, K., Harada, S., Kikuchi, M., Komatsu,
H., et al. (2000). MUM1/IRF4 expression as a frequent event in mature lymphoid malignancies. Leukemia
14, 449–456. https://doi.org/10.1038/sj.leu.2401696.

80. Nakagawa, M., Shaffer, A.L., Ceribelli, M., Zhang, M., Wright, G., Huang, D.W., Xiao, W., Powell, J., Petrus,
M.N., Yang, Y., et al. (2018). Targeting the HTLV-I-Regulated BATF3/IRF4 Transcriptional Network in Adult
T Cell Leukemia/Lymphoma. Cancer Cell *34*, 286-297.e10. https://doi.org/10.1016/j.ccell.2018.06.014.

81. Weilemann, A., Grau, M., Erdmann, T., Merkel, O., Sobhiafshar, U., Anagnostopoulos, I., Hummel, M.,
Siegert, A., Hayford, C., Madle, H., et al. (2015). Essential role of IRF4 and MYC signaling for survival of
anaplastic large cell lymphoma. Blood *125*, 124–132. https://doi.org/10.1182/blood-2014-08-594507.

82. Shaffer, A.L., Emre, N.C.T., Lamy, L., Ngo, V.N., Wright, G., Xiao, W., Powell, J., Dave, S., Yu, X., Zhao, H., et
al. (2008). IRF4 addiction in multiple myeloma. Nature *454*, 226–231. https://doi.org/10.1038/nature07064.

83. Amanda, S., Tan, T.K., Ong, J.Z.L., Theardy, M.S., Wong, R.W.J., Huang, X.Z., Ali, M.Z., Li, Y., Gong, Z., Inagaki,
H., et al. (2022). IRF4 drives clonal evolution and lineage choice in a zebrafish model of T-cell lymphoma.
Nat Commun *13*, 2420. https://doi.org/10.1038/s41467-022-30053-9.

84. Yang, Y., Shaffer, A.L., Emre, N.C.T., Ceribelli, M., Zhang, M., Wright, G., Xiao, W., Powell, J., Platig, J.,
Kohlhammer, H., et al. (2012). Exploiting synthetic lethality for the therapy of ABC diffuse large B cell
lymphoma. Cancer Cell *21*, 723–737. https://doi.org/10.1016/j.ccr.2012.05.024.

85. Delage, L., Lambert, M., Bardel, É., Kundlacz, C., Chartoire, D., Conchon, A., Peugnet, A.-L., Gorka, L.,
Auberger, P., Jacquel, A., et al. (2023). BTG1 inactivation drives lymphomagenesis and promotes
lymphoma dissemination through activation of BCAR1. Blood *141*, 1209–1220.
https://doi.org/10.1182/blood.2022016943.

86. Mlynarczyk, C., Teater, M., Pae, J., Chin, C.R., Wang, L., Arulraj, T., Barisic, D., Papin, A., Hoehn, K.B., Kots,
E., et al. (2023). BTG1 mutation yields supercompetitive B cells primed for malignant transformation.
Science *379*, eabj7412. https://doi.org/10.1126/science.abj7412.

87. Tijchon, E., van Emst, L., Yuniati, L., van Ingen Schenau, D., Havinga, J., Rouault, J.-P., Hoogerbrugge, P.M.,
van Leeuwen, F.N., and Scheijen, B. (2016). Tumor suppressors BTG1 and BTG2 regulate early mouse B-
cell development. Haematologica *101*, e272-276. https://doi.org/10.3324/haematol.2015.139675.

88. Scheijen, B., Boer, J.M., Marke, R., Tijchon, E., van Ingen Schenau, D., Waanders, E., van Emst, L., van der
Meer, L.T., Pieters, R., Escherich, G., et al. (2017). Tumor suppressors BTG1 and IKZF1 cooperate during
mouse leukemia development and increase relapse risk in B-cell precursor acute lymphoblastic leukemia
patients. Haematologica *102*, 541–551. https://doi.org/10.3324/haematol.2016.153023.

89. Li, Y., Mao, X., Li, M., Li, L., Tong, X., and Huang, L. (2024). The predictive value of BTG1 for the response
of newly diagnosed acute myeloid leukemia to decitabine. Clin Epigenetics *16*, 16.
https://doi.org/10.1186/s13148-024-01627-9.

90. Chen, T., Chen, X., Zhang, S., Zhu, J., Tang, B., Wang, A., Dong, L., Zhang, Z., Yu, C., Sun, Y., et al. (2021).
The Genome Sequence Archive Family: Toward Explosive Data Growth and Diverse Data Types. Genomics

Proteomics Bioinformatics *19*, 578–583. https://doi.org/10.1016/j.gpb.2021.08.001.

91. CNCB-NGDC Members and Partners (2022). Database Resources of the National Genomics Data Center,
China National Center for Bioinformation in 2022. Nucleic Acids Res *50*, D27–D38.
https://doi.org/10.1093/nar/gkab951.

1079
1080

Figure 1. Study design and the PCNSL mutational landscape in Chinese patients

A: Study design, including workflow and data composition for the discovery cohort and validation cohort.

B: Number and frequency of recurrent mutations. Gene–sample matrix of recurrently mutated genes ranked by mutation frequency (n = 140). The total mutation density across the cohort is displayed at the top, with the variant allele fraction and trinucleotides at the bottom.

C: GISTIC2.0 results of significant recurrent focal amplifications (red) and deletions (blue). Genes affected by each focal event are annotated (n = 140). X-axis: plot of chromosomes; Y-axis: G score.

Figure 2. Comparison of mutation profiles between the discovery cohort and validation cohort of PCNSL patients

A: Principal component analysis of whole-exome sequencing (WES) data between the discovery cohort and validation cohort of PCNSL patients

B: Mutation rate of WES data between the discovery cohort and validation cohort of PCNSL patients

C: A mirror bar plot showing the frequencies of genetic alterations in the discovery cohort of PCNSL patients compared with those in the validation cohort.

D: Arm-level copy number alterations of the discovery cohort and validation cohort samples are displayed. The frequencies of amplifications and deletions were compared.

E: Comparison of the tumor mutational burden (TMB) between the discovery cohort and the validation cohort of PCNSL patients

F: The levels of mutant-allele tumor heterogeneity (MATH) in the discovery cohort and the validation cohort of PCNSL patients were compared.

G: The levels of microsatellite instability (MSI) in the discovery cohort and the validation cohort of PCNSL patients were compared.

H: The variant allele frequency (VAF) in the discovery cohort and the validation cohort of PCNSL patients were compared.

The discovery cohort included blood-paired fresh tumor tissue samples (n = 58), and the validation cohort included unpaired FFPE tumor tissue samples (n = 82). Independent-samples t tests and chi-squared tests were used. *$P < 0.05$; **$P < 0.01$; ***$P < 0.001$.

1113

Figure 3. Mutation profiles of Chinese PCNSLs, DLBCLs, and PCNSLs of other races

A: A mirror bar plot showing the frequencies of genetic alterations between Chinese PCNSL patients (n = 140) and DLBCL patients (n = 955).

B: A mirror bar plot showing the frequencies of genetic alterations between Chinese PCNSL patients (n = 140) and French PCNSL patients (n = 115).

C: A mirror bar plot showing the frequencies of genetic alterations between Chinese PCNSL patients (n = 140) and Japanese PCNSL patients (n = 41).

D: GISTIC-defined recurrent copy number focal deletions (blue) and gains (red) as mirror plots in DLBCL (TCGA, n = 46) and Chinese PCNSLs (n = 140) are shown. X-axis: plot of chromosomes; Y-axis: G score.

E: The GISTIC2.0-defined recurrent copy number focal deletions (blue) and gains (red) in Japanese PCNSL patients (n = 41) and Chinese PCNSL patients (n = 140) are shown as mirror plots. Y-axis: plot of chromosomes; X-axis: G score.

The chi-squared test was used. *P < 0.05. In the figure legend of A-C, 'Inferred' means that the mutation types were not directly provided but were calculated from the supplemental tables of the corresponding paper; thus, the results are 'Inferred', and the *in facto* mutation number will not be less than the inferred value.

Figure 4. Integrated genetic drivers reveal PCNSL molecular subtypes with clinical outcome implications

A: Schematic of the typing strategies used to identify PCNSL molecular subtypes in the discovery cohort (n = 58) and in the validation cohort (n = 82).

B: Kaplan–Meier estimates of overall survival (OS) and progression-free survival (PFS) among patients belonging to each molecular subtype in the discovery cohort.

C: Kaplan–Meier estimates of overall survival (OS) and progression-free survival (PFS) among patients belonging to each molecular subtype in the validation cohort.

D: Kaplan–Meier estimates of overall survival (OS) among patients belonging to each molecular subtype in an independent external cohort (n=36).

E: The prevalence of EP300 mutations in PCNSL patients of different races.

F: Kaplan–Meier analyses for comparisons of the overall survival (OS) and progression-free survival (PFS) of patients with mutations within and those with mutations outside the EP300 domain.

G: Hazard ratio estimates of overall survival in PCNSL patients (n = 140).

Figure 5. Mutation profiles across three molecular subtypes of PCNSL

A: Venn diagram showing unique and shared mutations among the three identified PCNSL subtypes.

B: Number and frequency of recurrent mutations and the gene–sample matrix of recurrently mutated genes among the three PCNSL subtypes. The relative abundance across the molecular subtypes is displayed on the right.

C: Arm-level copy number alterations among the three molecular subtype samples are displayed.

D: The frequencies of amplifications and deletions were compared among the three molecular subtype samples.

E: The levels of mutant-allele tumor heterogeneity (MATH) were compared among the three molecular subtype samples.

F: The levels of microsatellite instability (MSI) were compared among the three MS samples.

G: The tumor mutational burden (TMB) was compared among the three MS samples.

H: The levels of variant allele frequency (VAF) were compared among the three MS samples.

Independent-samples t tests and chi-squared tests were used. $*P < 0.05$; $**P < 0.01$; $***P < 0.001$; ns $P > 0.05$.

Figure 6. Mutational profiles across different sites of onset in PCNSL

A: The number and percentage of PCNSL patients with various molecular subtypes at different disease onset sites.

B: Venn diagram of unique and shared mutations among PCNSLs with different sites of onset.

C: The number and frequency of recurrent mutations, along with a gene–sample matrix of recurrently mutated genes in PCNSL patients at different sites of onset, are presented. The relative abundance across the molecular subtypes is displayed on the right.

D: Arm-level copy number alterations across different sites of onset are displayed. The frequencies of amplifications and deletions were compared.

E: The levels of mutant-allele tumor heterogeneity (MATH) were compared among the samples representing different sites of onset.

F: Comparisons of the tumor mutational burden (TMB) among the different sites of onset samples were performed.

G: The levels of microsatellite instability (MSI) were compared among the samples representing different sites of onset.

H: Pathways affected by oncogenes in PCNSL patients with different sites of onset.

The chi-squared test and one-way ANOVA were used. *$P < 0.05$.