The role of receptor binding and immunity in SARS-CoV-2 fitness landscape: a modeling study

Zhaojun Ding [a], Hsiang-Yu Yuan [a,b,*]

[a] Department of Biomedical Sciences, College of Biomedicine, City University of Hong Kong, Hong Kong

[b] Centre for Applied One Health Research and Policy Advice, Jockey Club College of Veterinary Medicine and Life Sciences, City University of Hong Kong, Hong Kong

*Correspondence: sean.yuan@cityu.edu.hk

SUMMARY

Despite extensive research on SARS-CoV-2 ACE2 binding and transmissibility, their

relationship concerning varying immunity remains unclear. SARS-CoV-2 RBD sequences from

Italy in GISAID, combined with multiple deep mutational scanning data, were used to calculate

ACE2 binding score and immune escape during the pandemic. We developed a COVID-19

transmission model that decomposed the effective reproduction number into three time-varying

components: viral infectiousness (representing fitness), host susceptibility, and contact rate.

After model fitting, a rugged fitness landscape, spanned by ACE2 binding score and virus-

perceived effective immunity (adjusted for viral immune escape), was observed with peaks

corresponding to individual VOCs (alpha, delta, and omicron (BA.1* and BA.2*)). Increasing

effective immunity with reduced ACE2 binding corresponded to decreasing virus fitness peaks

from alpha to delta. Among omicron sub-lineages, which exhibited enhanced immune escape,

BA.2* reached a fitness optimum while retaining slightly reduced ACE2 binding relative to

BA.1.1. The findings help explain the evolution of SARS-CoV-2.

Keywords: ACE2 binding score; Immune escape; Deep mutational scanning data; Rugged

fitness landscape; SARS-CoV-2 evolution trajectory

INTRODUCTION

The COVID-19 pandemic was characterized by repeated infection waves, driven by SARS-CoV-2 variants with new and advantageous genomic mutations. These mutations enhanced fitness in variants of concern (VOC) via a complex interplay between viral intrinsic traits and population immunity, shaped by vaccinations and prior infections [1,2]. Therefore, understanding the SARS-CoV-2 fitness landscape can provide insights into evolutionary virus dynamics, knowing whether the virus undergoes gradual or punctuated evolution, which is important for predicting and preparing for future outbreaks.

The SARS-CoV-2 fitness is associated with virus's ability to bind to ACE2 receptors and evade immune defenses[1]. Upon entering the human body, the virus uses its spike protein to bind to the ACE2 receptor on the surface of host cells. This interaction occurs primarily through the receptor-binding domain (RBD). Simultaneously, the host immune system produces neutralizing antibodies, which prevent the virus from binding to ACE2. However, SARS-CoV-2 can evade immune recognition by mutating amino acid residues within the RBD, a phenomenon known as immune escape [1,3]. However, how ACE2 binding and immune escape interact to influence virus fitness under varying immunity levels among infected hosts remains unclear.

The accumulated mutations in the RBD of SARS-CoV-2 spike protein can confer selective advantages toward a higher fitness, resulting in antigenic drift [2]. For example, the N501Y mutation in the alpha (B.1.1.7) sub-lineage may have altered spike protein binding affinity to ACE2, resulting in higher transmission rates [4,5]. This mutation appeared in the omicron sub-lineages. Such evolutionary changes help sustain the virus's presence in the population,

facilitating COVID-19's transition from a pandemic to an endemic state characterized by

periodic surges.

The fitness landscape has been developed to represent the relationships between genetic

variations and their corresponding fitness values, providing insights into the potential adaptive

trajectories of populations [2,6]. A rugged landscape occurs due to interactions between mutations

(epistasis) [6,7]. Increasing immunity can also exert high selection pressure on viruses, contributing

to antigenic drift [2,8]. When a virus evolves through fitness valleys toward a new peak, it

generates boom-and-bust patterns, likely resulting in repeated outbreaks [9]. However, when

SARS-CoV-2 evolutionary and transmission dynamics are intertwined, reconstructing the fitness

landscape becomes challenging. For example, not only virus traits, but also host immunity (e.g.

vaccination) and host behaviors (e.g. social distancing measures and public awareness) can

influence transmissibility (or effective reproduction number) [10]. Hence, quantifying viral fitness

and finding its relationship with viral traits presents a challenge.

In this study, we aimed to assess the relationship between viral fitness and traits derived from

deep mutational scanning (DMS) during the pandemic. Sequence and epidemiological data from

Italy, a country with high viral incidence rates and high vaccine coverage, were used. To quantify

and thereby separate intervention effects (i.e., vaccinations and non-pharmaceutical interventions

(NPIs)) on virus fitness, we incorporated mobility and vaccination data in an infectious disease

transmission model to calibrate fitness estimates. From simulated results, we constructed a

fitness landscape after mapping the virus's evolutionary trajectory traits (i.e., ACE2 additive sum

of binding score and immune escape) over different periods across different immunity levels.

This study was previously made available as a preprint version [11].

METHOD DETAILS

Sequence data

Italian SARS-CoV-2 spike protein sequence data (31[st] January 2020–17[th] October 2022) were

downloaded from the GISAID database [12]. To ensure data quality control, two exclusion criteria

were applied: (1) sequences with > 5% ambiguous amino acids [13] and (2) sequences < 1,263 bp

or > 1,273 bp were filtered out. Ultimately, 112,011 sequences were included for virus ACE2

receptor binding and immune escape measurements. The proportion of beta and gamma

sequences were very low, only 2.07% and 0.12%, respectively (see Supplementary Fig. S1).

Therefore, only alpha, delta, and omicron variants were reported as the main VOCs in

subsequent analyses.

Measuring virus ACE2 additive sum of binding score and immune escape

DMS data were used to calculate virus ACE2 additive sum of binding score (i.e., the sum of

individual mutational effects) by mapping binding affinities between ACE2 and SARS-CoV-2

RBD amino acid mutation sites [14]. To reduce the confounding effects of epistasis in cell binding,

we employed multiple DMS datasets generated from five variants as backgrounds, namely

Wuhan-Hu-1, Alpha, Delta, BA.1, and BA.2 [15,16]. For each viral sequence, we selected the DMS

dataset whose reference background most closely matched the sequence's ancestral lineage and

used it to compute the ACE2 additive binding score (e.g., wild-type $\rightarrow$ Wuhan-Hu-1; Alpha and

its sub-lineages → Alpha; Delta and its sub-lineages → Delta; BA.1 and its sub-lineages →

BA.1; BA.2 and its sub-lineages → BA.2).

Escape fraction data were used to determine viral immune escape by mapping reduced binding

levels between the virus and antibodies, using datasets integrated from 33 monoclonal antibodies

and polyclonal sera, as available from the SARS-CoV-2 RBD antibody escape maps

(https://jbloomlab.github.io/SARS2_RBD_Ab_escape_maps/) [17]. First, we used the Clustal

Omega tool to align all sequences [18]. Next, to calculate virus ACE2 additive sum of binding

score for each sequence, binding changes across individual amino acid mutations of RBD region

(residues from N331 to T531) were summed (see Supplementary Fig. S8) [19]. For immune escape,

we calculated the mean reduction in antibody binding for each amino acid mutation site across

33 monoclonal antibodies. Immune escape across individual amino acid mutations was summed,

like the calculation of ACE2 binding. Finally, we computed daily average values of the additive

ACE2 binding score and the additive immune escape score among all daily samples. Further

details are shown in Supplementary Methods 2.1.

Mobility data and calculating the mobility index

The mobility index, which indicated the degree of population contact, was estimated from

Google Mobility and Apple Mobility data [20,21]. Specifically, three Apple mobility data subtypes

(driving, walking, and transit movement) and four Google subtypes (grocery and pharmacy

stores, transit, retail and recreation, and workplaces) were used to calculate mobility indices

based on previous research [22].

Vaccination data

COVID-19 vaccine administration data provided by the Italian Ministry of Health were used to summarize daily vaccination numbers for first, second, and third doses [23]. During the study period (i.e., model fitting period, 31st January 2020–5th March 2022), 134,611,131 vaccine doses were recorded and sourced from five major vaccine manufacturers: Janssen, Moderna, Novavax, Pfizer, and AstraZeneca (see Supplementary Table S1). Messenger RNA (mRNA) vaccines (i.e., Pfizer and Moderna) and viral vector vaccines (i.e., Janssen and AstraZeneca) accounted for almost 99% of the Italian vaccination campaign (see Supplementary Fig. S3). Consequently, we only considered mRNA (Pfizer and Moderna) and viral vector vaccines (Janssen and AstraZeneca).

Calculating virus fitness

A modeling approach was adopted to quantify viral infectiousness, which we refer to as viral fitness throughout this study. We extended a stochastic Susceptible - Exposed - Infectious - Recovered (SEIR) model, coupled with vaccination, mobility, and immune escape data, to reproduce COVID-19 transmission dynamics and estimate virus fitness across different VOC periods (see Fig. 1A, B). The model was used to estimate the part of the transmission rate that were determined by viral ACE2 receptor binding only over time (denoted as $\beta_v(t)$) under different immunity levels, representing the change of infectiousness. The relative virus fitness was defined as $\beta_v(t)/\beta_0$, where β_0 is the average value of $\beta_v(t)$ during the first month of the pandemic, representing the wild-type.

141   The effective reproduction number (R_e) was represented as two components (see Fig. 2D): an

142   intrinsic factor (infectiousness), and extrinsic factor (susceptibility and contact rate), as below.

$$R_e = Infectiousness \times (Susceptibility \times Contact)$$

144
$$= \beta_v(t) \cdot T_g \times \left((1 - V_{e_i} \cdot e^{-c \cdot immune\_escape\_t}) \cdot \frac{S_i(t)}{N} \times M(t)\right) (1)$$

where T_g is the generation time, which is 4 days [24–26]; $V_{e_i}(t)$ refers to vaccine efficacy from

different doses (i) against the wild-type; $V_{e_i} \cdot e^{-c \cdot immune\_escape(t)}$ represents protection against

circulating strains, accounting for their immune escape. $Immune\_escape\_t$ is the daily average

immune escape of viruses; c is the scaling factor for daily immune escape; $\frac{S_i(t)}{N}$ represents the

fraction of individuals at each susceptibility level (i.e., S_0, S_1, S_2 or S_3), based on vaccine dose i

received; $M(t)$ is the daily mobility index, representing the effect of the change in contact rate

on reducing overall fitness, R_e [23]. $\beta_v(t) \times (1 - V_{e_i} \cdot e^{-c \cdot immune\_escape\_t}) \times M(t)$ represents the

transmission rate among susceptible groups with different vaccine efficacy (i.e., $\beta_{imm\_i}(t)$ in Fig.

1B). The effective reproduction number R_e was obtained after fitting the transmission model (Fig.

1B) to the daily reported cases, when daily vaccine doses were considered.

Exposed (E_i) individuals who came into contact with infected individuals will become infectious

(I_i) after $\frac{1}{\sigma}$ days. As not all cases were reported correctly and immediately, reported infectious

cases (IR) were considered based on the detection rate (d) and the confirmation delay (ρ_t) in the

transition between E_i and I_i, which was used to fit the daily reported cases. Individuals with an

infectious status would subsequently enter a removed status (i.e., R_0, R_1, R_2 or R_3) due to self-

isolation or recovery after $1/\gamma(t)$ days. Considering waning immunity among these individuals,

we assumed that recovered and vaccinated individuals lose their immunity on average after ω

days (ω = $log(0.85)/182.5$, corresponding to exponential waning with a 15% loss in protection

after 6 months), which is the same to the studies by Barnard et al. and Lau et al.'s settings [27,28]. In

addition, we also used a fast vaccine waning rate (ω = $log(0.5)/90$, 50% loss of protection after

3 months) and moderate waning rate (ω = $log(0.5)/182.5$, 50% loss of protection after 6 months)

as the sensitivity analysis to check the stability of fitness patterns under different waning

scenarios.

Population immunity and effective immunity

Population immunity (P_{imm_total}) refers to the average level of immunity to a specific infectious

agent (e.g., Wuhan-Hu-1) within a population, achieved through both natural infection and

vaccination. It is commonly expressed as the proportion of individuals in the population who are

immune, either through previous infection or through vaccination, but partial protection and

waning immunity should be considered [29,30]. These factors were incorporated into our analysis.

We further calculated the virus-perceived immunity, termed effective immunity (P_{imm_eff}) as

$P_{imm\_eff} = P_{imm\_total}(t) \cdot e^{-c \cdot immune\_escape\_t}$, which accounts for immune escape and represents

protection against currently circulating strains. More details about the parameters set and

population immunity calculations can be found in the Supplementary Methods 2.2.

Model fitting for reproduce of infection dynamics and fitness landscape constructed

The number of individuals in compartment IR (i.e., reported infectious cases), as the modeling

output, was used to fit the daily numbers of reported cases. The Particle Markov chain Monte

Carlo method [31,32], which combines particle filtering and Markov chain Monte Carlo approaches,

was used to calibrate model output using daily reported cases and then estimate model

parameters, with a total of 100,000 iterations. All model parameter estimations were obtained

using the 'mcstate' package in R [31]. A fitness landscape was fitted with virus fitness and assumed

a multivariate normal distribution, which helped visualize SARS-CoV-2 adaptive evolutionary

trajectories. More technical details about fitness landscape reconstruction can be found in the

Supplementary Methods 2.2.4.

RESULTS

The relationship between daily mobility, vaccine coverage, and daily incidence

A large variety in daily incidence and mobility was observed (Fig. 2A, C and Supplementary Fig.

S2). Despite ongoing vaccination efforts (Fig. 2B), viral outbreaks reoccurred and cumulative

cases increased rapidly during the omicron period (Fig. 2A). The effective reproduction number

was likely influenced by both virus infectiousness and susceptibility. In our model, we

considered susceptibility (determined by population immunity and immune escape) and contact

rate (determined by mobility) as extrinsic factors while virus infectiousness as an intrinsic factor

(i.e., viral fitness; see Methods for the definition).

Temporal dynamics of virus ACE2 additive sum of binding score and immune escape

We observed distinct patterns between the two viral traits across main VOCs, including alpha,

delta, and omicron (Fig. 3A). ACE2 additive binding score decreased from a high level in alpha

(B.1.1.7) to delta (B.1.617.2), with early omicron lineages (BA.1*) showing binding scores

comparable to the wild-type (Fig. 3B). Binding scores then slightly fluctuated among omicron

sub-lineages, with BA.1.1 showing an increase relative to BA.1, and BA.1# showing a decrease.

The emerging BA.2* lineages subsequently showed ACE2 binding scores between BA.1 and

BA.1.1. On the other hand, immune escape showed a gradual rise from alpha to delta, followed

by a sharp rise with the emergence of omicron, whose sub-lineages exhibited similar levels. (Fig.

3C).

SARS-CoV-2 transmission dynamics

To understand how ACE2 receptor binding might influence virus fitness, we incorporated

vaccine coverage, mobility and virus immune escape data into the model to calibrate model

fitting for daily reported cases (Fig. 1B). Our model was able to reproduce the observed

transmission dynamics, including Non-VOC and VOC-associated epidemic waves (Fig. 4A). The

relative virus fitness was increasing form late delta to omicron periods (Fig. 4B). Both vaccine-

induced immunity (Fig. 4C, green line) and total natural infection combined with vaccine-

induced immunity (Fig. 4C, red line)) increased due to the initiation of the vaccination campaign

(beginning in the late December 2020) in Italy around the same time [33]. A similar trend was

observed for effective immunity (i.e., immunity which considering the waning immunity and

immune escape of circulating VOCs) against VOCs during alpha and delta periods, but a

significant decrease was recorded in the omicron period (Fig. 4D). This decline reflects the

strong immune escape capabilities of omicron, despite the ongoing vaccination campaign.

SARS-CoV-2's fitness landscape

The phenotype-fitness landscape spanned by the additive sum of ACE2 binding score and host'

effective immunity showed rugged patterns with four major peaks, corresponding to B.1.1.7

(alpha), B.1.617.2 (delta), BA.1* (first omicron peak) and BA.2* (second omicron peak); One

minor peak, corresponding to the transition period (mixed with delta and original omicron

(B.1.529) lineages) between delta and omicron, and four valleys between these peaks. A higher

effective immunity was associated with a lower fitness peak from alpha to delta. During the

omicron period, a reduction in effective immunity from BA.1* to BA.2* together with similar

ACE2 binding scores, corresponded with increasing fitness peaks (Fig. 5A). Notably, BA.2*

maintained high immune escape scores comparable to BA.1.1, but showed slightly lower binding

scores.

To evaluate robustness of the evolutionary pattern (i.e., Fig 5A), we varied vaccine waning from

an optimistic baseline to moderate and fast scenarios (50% loss at 6 and 3 months, respectively,

see supplementary Fig. S7). The shape and key features of the phenotype–fitness landscape was

maintained, with no material shifts in variant ordering or trend direction. To help visualize the

rugged patterns, a smoothed fitness landscape was constructed after fitting the fitness data (Fig.

5B; see Methods).

The sequential change in mutations, characterized by partial replacement, appeared through the

fitness valleys toward the peaks were explored:

Alpha -> delta: During the transition between the fitness peaks at alpha and delta, the

prevalence of N501Y (ACE2 binding changes (Wuhan-Hu-1 background, denoted as Δ_{bind}): 0.96,

immune escape changes (denoted as Δ_{escape}): 0.01, see Supplementary Fig. S5), a mutation with

high ACE2 binding, began to decline after passing through the first fitness valley (see

Supplementary Fig. S5B and Fig. S6). Meanwhile, the prevalence of L452R (Δ_{bind}: 0.16,

Δ_{escape}:0.18) and T478K (Δ_{bind}: 0.04, Δescape:0.004) started to increase during this valley,

gradually replacing N501Y as the dominant mutations. Some rare mutations, like E484K (Δ_{bind}:

0.11, Δ_{escape}:0.27), observed as fluctuation during the transition between the fitness peaks at alpha

and delta. More details about the rare mutations across each VOC can be found in

Supplementary Fig S5 and S6.

Delta-> Omicron (BA.1*): After the delta variant, the virus's evolutionary pathway moved

through two fitness valleys (i.e., Valley 2 and Valley 3) with a local peak (i.e., a minor peak

during the transition period including a mixture of omicron and delta) and then reach to BA.1*,

the next peak. Meanwhile, only N501Y and T478K was present in the BA.1* without L452R

(Fig. 5C). Next, the Set 1 (G339D, S373P, S375F, K417N, S477N, E484A, Q498R and Y505H),

Set 2 mutations (S371L, G446S and G496S) and few other mutations (N440K, Q493R and

N501Y, see Supplementary Fig. S5) gradually increased as the BA.1 lineage predominated.

During this period, effective immunity increased from around 0.2 to 0.4 and then returned to the

original level, while binding values remained similar, corresponding to the suboptimal fitness.

Omicron (BA.1*) -> Omicron (BA.2*): The virus continued to evolve through Valley 4 towards

the optimal fitness peak associated with BA.2*. During this transition, the Set 1 and other

additional mutations (N440K, Q493R and N501Y) were retained while Set 2 gradually decreased

as transient mutations. In the end, the Set 3 (S371F, T376A, D405N and R408S) gradually

increased and became dominated. The mutational effects in Set 3 showed either unchanged or

reduced binding scores using BA.1 as background (Supplementary Fig. S5). As the virus

progressed toward BA.2 dominance, optimal fitness emerged, coinciding with reduced effective

immunity and a moderate change in ACE2 binding (Fig. 5B).

DISCUSSION

We introduced a mathematical framework to estimate virus fitness while reproducing the

transmission dynamics of different SARS-CoV-2 variants in Italy. For the first time, the fitness

landscape was constructed for the virus from alpha to omicron. This was achieved by

incorporating both intrinsic viral factors (e.g., infectiousness) and extrinsic factors (e.g., host

susceptibility and contact rates) into an SEIR-based model, which helps quantify virus fitness

while removing the effects from extrinsic factors. The results provide valuable insights into the

real-world evolutionary constraints of SARS-CoV-2. DMS data are used to provide a proxy of

virus traits for sequences from GISAID. While ACE2 binding scores derived from DMS were

used to characterize receptor affinity, additional binding results can be incorporated to further

refine the fitness landscape.

Rugged fitness landscapes that were observed in our research (Fig. 5B and supplementary Fig.

S7) suggest that certain factors may play important roles in the evolutionary pathway via fitness

valleys, facilitating the transition to new fitness peaks. Epistasis can play a crucial role, as

interactions between mutations are likely to lead to non-linear effects, forming rugged fitness

landscapes, as we observed [6,34,35]. Epistatic interactions remain uncertain as previous studies

have reported conflicting findings on the trend of SARS-CoV-2 ACE2 receptor binding and the

available experimental data are limited [36–38]. Hence, we adopted multiple DMS datasets, each

using a different variant or subvariant template. This potentially reduces the biases caused by

epistatic effects more as the dissociation constant for each template sequence was directly

measured rather than inferred by summing mutational effects from the wild-type [15,16].

Furthermore, the fitness valley maybe also related to deleterious effects among immune escape

mutations [39,40]. Those mutations occurred in the fitness valley may be detrimental to SARS-CoV-

2's evolution, requiring additional time to reach a fitness peak when compensatory mutations

occur. In addition, recombination can be a potential driver of evolutionary change in SARS-CoV-

2, especially when multiple variants co-circulate [41]. This event can also result in viral fitness

changes.

A possible compromise between receptor binding affinity and immune escape during RBD

evolution of omicron sub-variants from BA.2 to BA.4/5 has been shown in a recent study [42] .

The authors found that a reduction of immune escape was compromised by a high cell binding.

On the other hand, a strong immune escape mutation occurred with a reduced cell binding level.

Early omicron evolution (from BA.1* to BA.2*) showed that once a suboptimal fitness peak is

reached by strong immune escape, further adaptation occurred with fluctuating binding score.

Furthermore, the constantly changing population immunity may play an important role in

evolutionary shifts between different fitness peaks. In a previous study, the effects of immune

escape mutations on viral fitness were compared between the presence and absence of antibodies,

showing that these mutations can also affect replication [43]. This highlights that fitness changes

may also result from alterations in ACE2 binding. In our analysis, we considered immune escape

to influence both host susceptibility and effective immunity. Consequently, changes in effective

immunity modulated viral fitness through their interaction with ACE2 binding.

Implications

Firstly, continuous surveillance efforts of SARS-CoV-2 is important. Even within the same VOC,

such as omicron, virus still experienced rugged fitness landscape, leading to repeated epidemics

of varying sizes. This is exemplified by the transition from omicron BA.1* to BA.2*, where the

virus's evolutionary path traversed a fitness valley before reaching a new peak. Knowing the

change in traits will be important to understand why a new outbreak occur.

Secondly, forecasting when a large outbreak will occur is still very challenging. An optimal virus

fitness was likely to be reached until a set of mutations having strong immune escape with

required reduction in binding.

Moreover, the results indicate a complex relationship between virus fitness and host immunity,

modulated by ACE2 binding. To better understand the complex interplay between multiple viral

traits and fitness, more advanced artificial intelligence (AI) models, such as spiking neural

networks (SNNs), may provide valuable insights. Such modeling approaches together with the

phenotype-fitness landscape can help illuminate the dynamic evolutionary landscape of SARS-

CoV-2, which in turn underscores the ongoing need for regular updates to vaccines to maintain

their effectiveness against emerging variants.

Study limitations

There are certain limitations in this study. Firstly, the study only focused on RBD region in the

spike protein and we assumed the impact of each amino acid on viral traits is additive.

Nonetheless, several experimental studies which consider the epistasis effect showed similar

pattern in the order with our results of binding changes among each VOC [37,44]. Secondly, the

population mobility index was an NPI proxy as it assumed social contacts were proportional to

population mobility, which not considered the awareness-driven behavior changes on the

dynamic of mobility. There is a risk that changes in $\beta_v(t)$ may reflect changes in NPIs or

awareness, which could cause the actual detection rate to vary over time and potentially

influence the estimation of $\beta_v(t)$. Thirdly, our model categorized the population based on

administered numbers of vaccine doses, without considering age and gender heterogeneity in the

population. This simplification overlooked potential differences in vaccine responses and

exposure risks across different demographic groups. Fourthly, we acknowledge that viral fitness

may be influenced by within-host replication, survival, and other viral or host factors. While

modeling simplifies the complexity, it allows us to understand the mechanism through

quantifying and, therefore, separating individual effects. Fifthly, GISAID may contain some

noisy data. We have done initial screening to remove data with apparent errors. Lastly, but

critically, the seasonal or climatic factors were not considered. Future research should introduce

the impact of these factors on virus fitness to provide a more comprehensive understanding of

virus transmission dynamics.

Conclusions

SARS-CoV-2 mutations along with immunity influences generated a rugged fitness landscape,

which shaped viral evolutionary trajectories in Italy. Knowing the landscape can help provide a

better understanding of the complex dynamics of viral evolution and adaptation, particularly in

drug resistance and vaccine development. More importantly, host immune systems and viruses

are constantly engaged in an arms race, with each adapting to changes in the other. Timely

vaccine updates in response to shifts in the viral fitness landscape may allow for more accurate

targeting of emerging variants.

Acknowledgments

We gratefully acknowledge all researchers for contributing to, generating, and openly sharing

genome data via the Global Initiative on Sharing All Influenza Data (GISAID), as most of our

analyses were based on this invaluable resource. We thank Prof. Haogao Gu and Dr. Jingbo

Liang for helpful discussions, Ms. Fatema Khairunnasa for useful editing suggestions. We

greatly appreciate Prof. Joshua S. Weitz at University of Maryland for his meaningful

suggestions on model fitting. The authors also acknowledge support from grants funded by the

[City University of Hong Kong] under Grant [#9610416]; [General Research Fund] under Grant

[#9043514].

Author Contributions

Zhaojun Ding and Hsiang-Yu Yuan designed the study. Zhaojun Ding collected, analyzed, and

modelled the data. Zhaojun Ding and Hsiang-Yu Yuan wrote the paper. Hsiang-Yu Yuan gave

final approval for publication. All authors read and approved the final manuscript.

Data and code availability

Daily reported COVID-19 cases in Italy were collected from 'Our World in Data'

(https://github.com/owid/covid-19-data/tree/master/public/data). Deep mutational scanning data

were collected from "SARS-CoV-2 RBD DMS" (https://jbloomlab.github.io/SARS-CoV-2-

RBD_DMS/). COVID-19 vaccination data in Italy were collected from the Italian Ministry of

Health (https://github.com/italia/covid19-opendata-vaccini). Mobility data were collected from

"Google COVID-19 Community Mobility Reports" (https://www.google.com/covid19/mobility/)

and "Apple Map Mobility Trends Reports" (https://covid19.apple.com/mobility), respectively.

The codes for this study are available at https://github.com/DING219/bind_escape_code.

Declaration of interest statement

All authors declare no competing interests.

Ethics approval

This study is based on modeling using aggregate data, with no access to or analysis of

individual-level information. Therefore, the study does not involve human participants and is not

human subjects research. Ethical review and approval were not applicable.

References

1. Carabelli, A. M. *et al.* SARS-CoV-2 variant biology: immune escape, transmission and fitness. *Nat Rev Microbiol* **21**, 162–177 (2023).

2. Yewdell, J. W. Antigenic drift: Understanding COVID-19. *Immunity* **54**, 2681–2687 (2021).

3. Greaney, A. J. *et al.* Complete Mapping of Mutations to the SARS-CoV-2 Spike Receptor-Binding Domain that Escape Antibody Recognition. *Cell Host Microbe* **29**, 44-57.e9 (2021).

4. Fiorentini, S. *et al.* First detection of SARS-CoV-2 spike protein N501 mutation in Italy in August, 2020. *Lancet Infect Dis* **21**, 16 (2021).

5. Tian, F. *et al.* N501Y mutation of spike protein in SARS-CoV-2 strengthens its binding to receptor ACE2. *Elife* **10**, 69091 (2021).

6. Wright, S. The roles of mutation, inbreeding, crossbreeding and selection in evolution. *Sixth International Congress on Genetics* **1**, 356-366 (1932).

7. Kauffman, S. & Levin, S. Towards a general theory of adaptive walks on rugged landscapes. *J Theor Biol* **128**, 11-45 (1987).

8. Grenfell, B. T. *et al.* Unifying the Epidemiological and Evolutionary Dynamics of Pathogens. *Science* **303**, 327-332 (2004).

9. Koelle, K., Cobey, S., Grenfell, B. & Pascual, M. Epochal evolution shapes the phylodynamics of interpandemic influenza a (H3N2) in humans. *Science* **314**, 1898-1903 (2006).

10. Weitz, J. S., Park, S. W., Eksin, C. & Dushoff, J. Awareness-driven behavior changes can shift the shape of epidemics away from peaks and toward plateaus, shoulders, and oscillations. *Proc Natl Acad Sci USA* **117**, 32764-32771 (2020).

11. Ding, Z. & Yuan, H.-Y. Impacts of receptor binding and immunity on SARS-CoV-2 fitness landscape: a modeling study. *medRxiv* 2024.10.24.24316028 (2024) doi:10.1101/2024.10.24.24316028.

12. Elbe, S. & Buckland-Merrett, G. Data, disease and diplomacy: GISAID's innovative contribution to global health. *Global Challenges* **1**, 33–46 (2017).

13. Zhao, S., Ran, J. & Han, L. Exploring the interaction between E484K and N501Y substitutions of SARS-CoV-2 in shaping the transmission advantage of COVID-19 in Brazil: A modeling study. *Am J Trop Med Hyg* **105**, 1247–1254 (2021).

14. Starr, T. N. *et al.* Deep Mutational Scanning of SARS-CoV-2 Receptor Binding Domain Reveals Constraints on Folding and ACE2 Binding. *Cell* **182**, 1295-1310.e20 (2020).

15. Starr, T. N. *et al.* Deep mutational scans for ACE2 binding, RBD expression, and antibody escape in the SARS-CoV-2 Omicron BA.1 and BA.2 receptor-binding domains. *PLoS Pathog* **18**, e1010951 (2022).

16. Starr, T. N. *et al.* Shifting mutational constraints in the SARS-CoV-2 receptor-binding domain during viral evolution. *Science* **377**, 420-424 (2022).

17. Greaney, A. J., Starr, T. N. & Bloom, J. D. An antibody-escape estimator for mutations to the
SARS-CoV-2 receptor-binding domain. *Virus Evol* **8**, veac021 (2022).

18. Chenna, R. *et al.* Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids*
Res **31**, 3497-3500 (2003).

19. Bozdaganyan, M. E., Shaitan, K. V., Kirpichnikov, M. P., Sokolova, O. S. & Orekhov, P. S.
Computational Analysis of Mutations in the Receptor-Binding Domain of SARS-CoV-2 Spike and
Their Effects on Antibody Binding. *Viruses* **14**, 295 (2022).

20. Google COVID-19 Community Mobility Reports. https://www.google.com/covid19/mobility/
(accessed Sept 14, 2022).

21. Apple Map Mobility Trends Reports. https://covid19.apple.com/mobility (accessed Sept 11,
2022).

22. Nouvellet, P. *et al.* Reduction in mobility and COVID-19 transmission. *Nat Commun* **12**, 1090
(2021).

23. Struttura Commissariale per l'Emergenza Covid-19. Open data on COVID-19 vaccination in Italy.
https://github.com/italia/covid19-opendata-vaccini (accessed Oct 11, 2022).

24. Hart, W. S. *et al.* Generation time of the alpha and delta SARS-CoV-2 variants: an epidemiological
analysis. *Lancet Infect Dis* **22**, 603–610 (2022).

25. Xu, X. *et al.* Assessing changes in incubation period, serial interval, and generation time of SARS-
CoV-2 variants of concern: a systematic review and meta-analysis. *BMC Med* **21**, 374 (2023).

26. Arantes, I. *et al.* Comparative epidemic expansion of SARS-CoV-2 variants Delta and Omicron in
the Brazilian State of Amazonas. *Nat Commun* **14**, 2048 (2023).

27. Barnard, R. C. *et al.* Modelling the medium-term dynamics of SARS-CoV-2 transmission in England
in the Omicron era. *Nat Commun* **13**, 4879 (2022).

28. Lau, J. J. *et al.* Real-world COVID-19 vaccine effectiveness against the Omicron BA.2 variant in a
SARS-CoV-2 infection-naive population. *Nat Med* **29**, 348-357 (2023).

29. Lopman, B. A. *et al.* A framework for monitoring population immunity to SARS-CoV-2. *Ann*
Epidemiol **63**, 75-78 (2021).

30. Fontanet, A. & Cauchemez, S. COVID-19 herd immunity: where are we? *Nat Rev Immunol* **20**,
583–584 (2020).

31. Knock, E. S. *et al.* Reproducible parallel inference and simulation of stochastic state space models
using odin, dust, and mcstate. *Wellcome Open Res* **5**, 288 (2020).

32. Kucharski, A. J. *et al.* Early dynamics of transmission and control of COVID-19: a mathematical
modelling study. *Lancet Infect Dis* **20**, 553–558 (2020).

33. Ferrante, P. et al. The First 2 Years of COVID-19 in Italy: Incidence, Lethality, and Health Policies.
Front Public Health **10**, 986743 (2022).

34. Diaz-Colunga, J. *et al.* Global epistasis on fitness landscapes. *Phil. Trans. R. Soc. B* **378**, *20220053*
(2023).

35. Wang, D., Huot, M., Mohanty, V. & Shakhnovich, E. Biophysical principles predict fitness of SARS-
CoV-2 variants. *Biophys J* **121**, e2314518121 (2024).

36. Moulana, A. *et al.* Compensatory epistasis maintains ACE2 affinity in SARS-CoV-2 Omicron BA.1.
Nat Commun **13**, 7011 (2022).

37. Calvaresi, V. *et al.* Structural dynamics in the evolution of SARS-CoV-2 spike glycoprotein. *Nat*
Commun **14**, 1421 (2023).

38. Wu, L. *et al.* SARS-CoV-2 Omicron RBD shows weaker binding affinity than the currently dominant
Delta variant to human ACE2. *Signal Transduct Target Ther* **7**, 8 (2022).

39. Koelle, K. & Rasmussen, D. A. The effects of a deleterious mutation load on patterns of influenza
A/H3N2's antigenic evolution in humans. *Elife* **4**, e07361 (2015).

40. Bloom, J. D. & Neher, R. A. Fitness effects of mutations to SARS-CoV-2 proteins. *Virus Evol,* **9**
vead055.

41. Turakhia, Y. *et al.* Pandemic-scale phylogenomics reveals the SARS-CoV-2 recombination
landscape. *Nature* **609**, 994-997 (2022).

42. Li, W. *et al.* Key mechanistic features of the trade-off between antibody escape and host cell
binding in the SARS-CoV-2 Omicron variant spike proteins. *EMBO J* **43**, 1484-1498 (2024).

43. Schröder, S. *et al.* Characterization of intrinsic and effective fitness changes caused by
temporarily fixed mutations in the SARS-CoV-2 spike E484 epitope and identification of an
epistatic precondition for the evolution of E484A in variant Omicron. *Virol J* **20**, 257 (2023).

44. Han, P. *et al.* Receptor binding and complex structures of human ACE2 to spike RBD from
omicron and delta SARS-CoV-2. *Cell* **185**, 630-640.e10 (2022).

Figures

503
504

505
506
Fig. 1. Study flow and model design. (A) Schematic showing the extended SEIR
model integrating multiple data sources to estimate the SARS-CoV-2 fitness landscape.
Data inputs include mobility, immune escape, and vaccination coverage data. These
data feed into the extended SEIR model, which was calibrated using reported case data.
The model outputs include virus fitness combined with ACE2 binding data and effective

immunity, and show how these factors shape virus evolution patterns. (B) Detailed

structure of the SEIR compartment model. The model partitions the population based on

vaccination status and infection stage, highlighting transitions between susceptible (S),

exposed (E), infectious (I), reported infectious cases (IR) and recovered (R) states, with

an emphasis on the impact of vaccination doses on these dynamics. S_i, E_i, I_i, IR_i and R_i

subscripts denote compartments with different vaccine doses.

Fig. 2. Factors influencing daily COVID-19 incidences and virus fitness. (A) Daily
COVID-19 incidences in Italy. New daily COVID-19 cases in Italy are measured in log10
scales. VOC emergence is highlighted by vertical dashed lines, labeled as non-VOC,
alpha, delta, and omicron strains. (B) Vaccination coverage in Italy from the first to the

525   third vaccine doses (C) Mobility index in Italy. Population movement levels relative to

526   pre-pandemic levels (horizontal dashed line) are shown.

Fig. 3. Comprehensive analysis of SARS-CoV-2 VOC with ACE2 binding affinity and immune escape. (A) Variant prevalence over time. The bar chart shows the prevalence of different SARS-CoV-2 VOC (alpha, beta, delta, gamma, and omicron strains) from early 2020 through early 2022. (B) Additive binding scores of viral lineages under different VOC backgrounds. The value in parentheses after each VOC denotes the mean ACE2 binding scores. (C) The immune escape scores over time.

Scatter plot showing ACE2 binding and alpha, delta, and omicron variant immune escape across the timeline. BA.1* and BA.2* including BA.1, BA.2 and their sub-lineages. BA.1# means BA.1* without BA.1 and BA.1.1. BA.2# means BA.2* without BA.2.

Fig. 4. Epidemic indicator and immunological response trajectories during SARS-CoV-2 variant transition in Italy. (A) Model fitting for daily confirmed cases. The red line represents the estimated daily incidences. (B) Virus fitness and effective reproduction number. The red line shows estimated virus fitness over time, which is relative to the wild-type virus. The dashed grey line shows effective reproduction number. (C) Population immunity against wild-type. The green curve represents the average proportion of the population protected by vaccine-induced immunity, while the red curve while the red curve reflects total immunity, including vaccine and infection-acquired immunity. (D) The curve represents the immune protection in the total population against circulating VOCs (referred to as effective immunity), considering virus immune escape properties.

Fig. 5. Interplay between ACE2 additive binding score, effective immunity, and virus's fitness in SARS-CoV-2 evolution. (A) Observed relationship between ACE2 additive binding score, effective immunity and fitness. Gray dots represent variant fitness levels during a 2-week transition period following the initial outbreak of each VOC pandemic. (B) Fitness landscape of VOC. Surface topography shows the virus's fitness landscape, which reconstructed based on multivariate normal distribution. Peaks denote maximum variant fitness values in predominant periods. (C) Mutation prevalence associated with VOC transitions. Shadowed regions corresponding to valleys in panel B are marked by different colors. Some cluster mutations with similar prevalence were termed "Set1", "Set2" and "Set3".