Augmenting electronic health record data with social and environmental determinant of

health measures to understand regional factors associated with asthma exacerbations

Alana Schreibman[1], Kimberly Lactaoen[1], Jaehyun Joo[1], Patrick K. Gleeson[2], Gary E.

Weissman[1,2], Andrea J. Apter[2], Rebecca A. Hubbard[3], Blanca E. Himes[1]

[1] Department of Biostatistics, Epidemiology and Informatics, Perelman School of Medicine,

University of Pennsylvania, Philadelphia, PA 19104

[2] Division of Pulmonary, Allergy and Critical Care Medicine, Perelman School of Medicine,

University of Pennsylvania, Philadelphia, PA 19104

[3] Department of Biostatistics, School of Public Health, Brown University, Providence, RI 02912

Corresponding author:

Blanca E Himes, PhD

402 Blockley Hall

423 Guardian Drive

Philadelphia, PA 19104

Phone: (215) 573-3282

Email: bhimes@pennmedicine.upenn.edu

Abstract

Electronic health records (EHRs) provide rich data for diverse populations but often lack information on social and environmental determinants of health (SEDH) that are important for the study of complex conditions such as asthma, a chronic inflammatory lung disease. We integrated EHR data with seven SEDH datasets to conduct a retrospective cohort study of 6,656 adults with asthma. Using Penn Medicine encounter data from January 1, 2017 to December 31, 2020, we identified individual-level and spatially-varying factors associated with asthma exacerbations. Black race and prescription of an inhaled corticosteroid were strong risk factors for asthma exacerbations according to a logistic regression model of individual-level risk. A spatial generalized additive model (GAM) identified a hotspot of increased exacerbation risk (mean OR = 1.41, SD 0.14, $p < 0.001$), and inclusion of EHR-derived variables in the model attenuated the spatial variance in exacerbation odds by 34.0%, while additionally adjusting for the SEDH variables attenuated the spatial variance in exacerbation odds by 66.9%. Additional spatial GAMs adjusted one variable at a time revealed that neighborhood deprivation (OR = 1.05, 95% CI: 1.03, 1.07), Black race (OR = 1.66, 95% CI: 1.44, 1.91), and Medicaid health insurance (OR = 1.30, 95% CI: 1.15, 1.46) contributed most to the spatial variation in exacerbation odds. In spatial GAMs stratified by race, adjusting for neighborhood deprivation and health insurance type did not change the spatial distribution of exacerbation odds. Thus, while some EHR-derived and SEDH variables explained a large proportion of the spatial variance in asthma exacerbations across Philadelphia, a more detailed understanding of SEDH variables that vary by race is necessary to address asthma disparities. More broadly, our findings demonstrate how integration of information on SEDH with EHR data can improve understanding of the combination of risk factors that contribute to complex diseases.

Author summary

Electronic health records constitute an important source of data for understanding the health of large and diverse real-world populations, however, they do not routinely capture socioeconomic and environmental factors known to affect health outcomes. We show how electronic health record data can be augmented to include individual measures of air pollution exposures, neighborhood socioeconomic status, and the natural and built environment using patients' residential addresses to study asthma exacerbations, episodes of worsening disease that remain a major public health challenge in the United States. We found that on an individual patient-level, Black race and prescription of an inhaled corticosteroid were the factors most strongly associated with asthma exacerbations. In contrast, neighborhood deprivation, race, and health insurance type accounted for the most spatial variation in exacerbation risk across Philadelphia. Our findings provide insight into factors that contribute to asthma disparities in our region and present a framework for future efforts to expand the scope of electronic health record data.

Introduction

Electronic health records (EHRs) are a source of rich patient-level data for large and diverse

populations that can be used for research due to their widespread availability [1]. However, EHR

data often contain incomplete or low-quality measures of social and environmental determinants

of health (SEDH), limiting their utility for the study of complex diseases [2]. Recent efforts to

address this limitation have included developing methodologies to integrate external data on the

physical, built, and social environment via linkage with patient addresses [3–5], with high-

resolution geospatial datasets providing the closest estimate of individual exposures [6].

Integrated EHR and SEDH datasets can be used to understand both individual-level outcomes

and patterns of risk across a spatial region, thereby providing insights for both precision

medicine and precision public health efforts. Because many environmental exposures pose a

greater risk to select groups of people, integrating external information on SEDH with EHR data

is also helpful to address health disparities according to race, ethnicity, and socioeconomic status.

Asthma, a chronic disease that is characterized by inflammation and reversible narrowing

of the airways, affects over 25 million people or approximately 8% of the United States

population [7]. Racial and ethnic disparities in its morbidity and mortality are well known, and

those living in poverty are also more likely to have asthma [7–11]. The clinical goals of asthma

management are to control patients' symptoms and minimize long-term risk of lung function

decline [12]. This includes preventing asthma exacerbations, episodes of worsening disease

which require treatment with systemic steroids [13]. However, despite guideline-directed clinical

management, asthma exacerbations remain common, contributing to asthma-related morbidity

and mortality as well as higher health care costs and utilization [14,15]. Risk factors for asthma

exacerbations in adults include female sex [16], obesity [17], current or past smoking [18],

comorbid allergic rhinitis or chronic obstructive pulmonary disease (COPD) [17,19], and a

history of previous exacerbations [20]. Observational studies have also found independent

associations between asthma exacerbations and exposure to particulate matter (PM), gaseous air

pollutants, and to mixtures of pollutants such as traffic-related air pollution (TRAP) [21].

Similarly, associations between asthma exacerbations and living in poverty [22], substandard

housing conditions, such as the presence of mold and pests [23,24], and neighborhood

"greenness" [25,26] have been documented.

Because asthma exacerbations result from complex interactions among various

biological, social, and environmental factors that vary across individuals and geographically,

creating generalizable models of exacerbations remains an unachieved goal. Further, because in

the United States there is a high correlation between minoritized race or ethnicity, poverty, and

harmful environmental exposures, disentangling relationships among key variables is difficult

[9,27]. Approaches that model many asthma-related variables in specific regions may lead to the

identification of actionable strategies to reduce exacerbations locally, and using EHR data to

create these models has the advantage of providing health information for the specific catchment

region served by a given healthcare system. Few studies have linked EHRs with a diverse set of

SEDH variables to study asthma exacerbations using patient-level data [28–30], and to our

knowledge, none have used geospatial analysis techniques to understand the contribution of

these factors to the spatial distribution of exacerbation risk. Here, we show how EHR data can be

extended to include individualized measures of air pollution exposures, socioeconomic status

indices, and measures of the natural and built environment to identify local factors associated

with asthma exacerbation risk.

101
Methods

A retrospective cohort analysis was performed using de-identified EHR data from Penn

Medicine, a large health system that serves the greater Philadelphia area, from encounters dated

1/1/2017-12/31/2020. An overview of our study design is shown in Fig 1.

Ethics statement

Our study was approved by the University of Pennsylvania Institutional Review Board (IRB)

under protocol number 824789. Formal consent was not obtained, as a waiver of HIPAA

Authorization was granted for the conduct of this research.

Study population

Patient-level encounter data was obtained for adults (i.e. age ≥ 18 years) who had at least one

encounter with an International Classification of Diseases (ICD)-10 code for asthma (J45*) and

who were prescribed a short-acting β_2-agonist (SABA) (S1 Table). The most recent residential

address for each patient was obtained and geocoded using previously described methods [3].

Demographic, comorbidity, and medication data for encounters during the study period was

extracted and used to compute several variables, hereafter referred to as "EHR-derived

variables." These included: age at first encounter, sex, race, ethnicity, body mass index (BMI),

health insurance type, smoking status, chronic obstructive pulmonary disease (COPD), allergic

rhinitis, a modified Elixhauser score [31], inhaled corticosteroid (ICS) prescription, and years

followed (defined as the number of years between first and last encounter). Additional details,

including inclusion and exclusion criteria and definitions of the EHR-derived variables, are

provided in S1 Text.

Outcome

Asthma exacerbations were defined as encounters with an oral corticosteroid (OCS) prescription

(S1 Table) and either 1) a primary asthma diagnosis code (ICD-10, J45*) for encounters with

primary diagnosis codes listed or 2) a nonprimary asthma ICD-10 code for encounters without a

primary diagnosis listed but only one or two ICD-10 codes listed. A count of exacerbations

during the study period was computed for each patient.

SEDH data

Seven external datasets were integrated with our EHR dataset via linkage with patient geocodes,

creating variables that are hereafter referred to as "SEDH variables." Additional details on data

processing are reported in S1 Text, and dataset sources and spatiotemporal dimensions are

summarized in S2 Table. All processed SEDH data is available in Sensor-based Analysis of

Pollution in the Philadelphia Region with Information on Neighborhoods and the Environment

(SAPPHIRINE), a web application that integrates spatially distributed high-resolution social and

environmental data in the greater Philadelphia region to facilitate the conduct of local health

studies [32].

Air pollution exposures

Average pollutant exposures were assigned to each patient using high-resolution (~1x1 km^2)

geophysical model estimates. 2017-2019 NO_2 estimates (in parts per billion by volume, or ppbv)

and 2017-2020 PM2.5 estimates (in µg/m^3) were downloaded from resources reported in Cooper

et al. and van Donkelaar et al., respectively, temporally averaged, and linked to the study cohort

using bilinear interpolation [33,34]. Exposure to other toxic airborne chemicals from point

sources was estimated using the EPA Toxics Release Inventory (TRI) [35]. Total toxic air release

exposure by patient was computed as the sum of toxic releases (in kilograms) over the study

period within a 1-km circular buffer of each patient's residential address. Exposure to air

pollution from mobile sources was estimated as the sum of the daily vehicle distance traveled

(DVDT), a metric computed using traffic data published by the Pennsylvania Department of

Transportation, within a 300-m circular buffer [36].

Neighborhood socioeconomic environment

Socioeconomic disadvantage for each person was summarized as the Area Deprivation Index

(ADI), a validated index computed for each Census block group based on several ACS variables

[37]. A higher ADI score indicates greater disadvantage. 2018 ADI data was extracted from the

Neighborhood Atlas and was assigned to each patient by block group [38].

Built and natural environment

Asthma-related housing code violation data (i.e. pests, water damage, and indoor air

contamination) was obtained from a Philadelphia Department of Licenses and Inspections

dataset reported by OpenDataPhilly (S3 Table) [39]. For each block group, the number of

violations during the study period per 100 people (based on the 2019 ACS population estimates)

was computed and assigned to each patient. Vegetation density was summarized as the

normalized difference vegetation index (NDVI), an index with values ranging from -1.0 to 1.0

where higher positive values represent higher vegetation density. NDVI was computed in Google

Earth Engine using surface reflectance images from the Landsat 8 satellite during the study

period and assigned to each patient as the mean value within a 300-m circular buffer [40].

<u>Statistical analysis</u>

Analyses were conducted in R 4.2 [41].

Study area

To minimize bias in our geospatial analyses introduced by uneven spatial density across the Penn

Medicine catchment area, we restricted our study region to spatial areas in which the geospatial

representativeness of our EHR cohort was adequate compared to the underlying population.

Following methods described in Xie et al. [42], we computed a spatial representation ratio

(SRR), defined as the ratio between the proportion of our EHR cohort living in each census

block group and the proportion of the Philadelphia population living in that block group, as

reported by the 2019 American Community Survey (ACS). We defined our study region as

contiguous census tracts (i.e. adjacent or separated by a non-residential area such as a park) with

a mean SRR of 0.5 or greater. Only patients who resided in this study region were included in

analyses.

Modeling individual-level risk factors

Chi-squared and Kruskal-Wallis rank sum tests were used in bivariate analyses to assess

associations between patient characteristics and asthma exacerbation level (i.e. 0, 1-2, 3-4, 5+)

during the study period, and to compare the characteristics of complete cases to individuals with

missing data. To identify patient-level factors associated with asthma exacerbations, we fit

logistic regression models with asthma exacerbations as a binary case-control (0 vs >0) outcome.

This approach was chosen to match the dichotomous outcome used in spatial analyses. Logistic

regression models were initially adjusted for EHR-derived variables only, and then adjusted for

both EHR-derived and SEDH variables (EHR & SEDH-adjusted). Years followed was included

as a covariate in both models to account for variation in the length of available follow-up across

patients. Model fit was assessed using the Akaike information criterion (AIC). We checked for

multicollinearity by computing Pearson's correlation coefficients between all EHR and SEDH

variables, while selecting White race and Private health insurance type as reference levels for the

two nominal categorical variables, and we computed variance inflation factors (VIF) for all

independent variables.

Sensitivity analysis for individual-level risk factors

We conducted sensitivity analyses of individual-level risk factors by fitting negative binomial

regression models with asthma exacerbations represented as a count outcome and comparing

results to those of logistic regression models. The negative binomial regression models were

adjusted first for EHR-derived variables only and then for EHR & SEDH variables, while

including years followed as an offset in each model.

Modeling spatial risk factors

To estimate local odds of exacerbation as a function of location, spatial generalized additive

models (GAMs) were fit with a binary case-control outcome (0 vs >0 exacerbations) on a grid of

points across the study region, using the R *MapGAM* package and previously described methods

(S1 Text) [3,43]. Maps of spatial effect predictions were created for the smoothed spatial term of

each model, where the spatial odds ratio (OR) at each point represented the ratio between the

odds of exacerbation at that point and the median odds across all points. First, a univariable

model adjusted only for years followed, hereafter referred to as "unadjusted model", was fit to

identify hotspots and coldspots across the study region. Next, multivariable models adjusted for

1) only EHR-derived variables and 2) EHR & SEDH variables were compared to the unadjusted

model by computing the mean and standard deviation (SD) of the spatial ORs at points within

any overlapping hotspots, and by computing the percent difference between the variance in ORs

across the full study region in the adjusted models and the unadjusted model. To understand the

contribution of individual variables to the observed spatial effects, we fit additional models

adjusted for each EHR-derived or SEDH variable one variable at a time (all models were also

adjusted for years followed) and computed the percent difference between the variance in ORs in

these models and the unadjusted model. Model fit was assessed using the AIC.

Stratified analysis

We conducted a stratified analysis to further evaluate the association between race and asthma

exacerbations. Chi-square and Wilcoxon rank sum tests were used to assess bivariate

relationships between race and other variables. To test whether any EHR or SEDH variables that

were correlated with race influenced spatial patterns of asthma exacerbation risk independently

of race, spatial GAMs adjusted one variable at a time were fit on each race stratum using the

same approach described above. Before fitting the models, SRR selection was repeated for each

stratum to account for the uneven geographic distribution of race across the initial study region.

Results

Following selection of patients based on inclusion/exclusion criteria and spatial filtering (S1

Fig), the retrospective study cohort consisted of 6,656 asthma patients, 2,329 of whom had one

or more exacerbations (Table 1), with residence in 249 census tracts in Philadelphia (S2 Fig).

The spatial distribution of all processed SEDH datasets within the study region is shown in S3

Fig. The EHR-derived variables years followed, age, race, BMI, health insurance type, smoking

status, COPD, allergic rhinitis, Elixhauser comorbidity score, and ICS, as well as the SEDH

variable ADI were the most significantly associated with exacerbations according to bivariate

analyses ($p < 0.001$) (Table 1). Patients with more exacerbations during the study period were

followed for more years, more likely to be aged 35-54, of Black race, or class 2 or class 3 obese,

and more likely to have Medicaid health insurance, a history of smoking, COPD, allergic rhinitis,

higher Elixhauser comorbidity scores, an ICS prescription, or live in neighborhoods with higher

ADI. The proportion of patients in the study cohort who were prescribed controller medications

including ICS, leukotriene modifiers, long-acting β_2-agonists (LABA), and biologic therapies

was positively associated with exacerbation count (S4 Table). Bivariate analyses comparing the

distribution of characteristics between the study cohort and patients excluded due to missing

EHR data found statistically significant differences in years followed, age, sex, health insurance

type, COPD, allergic rhinitis, Elixhauser comorbidity score, and ICS ($p < 0.001$) (S5 Table).

Based on Pearson's correlation coefficients, there were no strong correlations between

any EHR-derived or SEDH variables, except for NO_2 and PM2.5 ($\rho = 0.77$) (S4 Fig). Moderate

correlations ($|\rho| > 0.50$) were observed between age and Medicare health insurance, Black race

and ADI, and NO_2 and NDVI. Furthermore, adjusted generalized VIFs for all variables did not

exceed 2.00, suggesting that all variables could be included in multivariable models (S6 Table).

Individual risk factors associated with asthma exacerbations

ORs for the EHR-derived and SEDH variables included in the multivariable logistic regression

model are summarized in Table 2. The EHR-derived variables years followed, age 35-54, Black

race, and ICS had the strongest positive associations with having at least one exacerbation during

the study period ($p < 0.001$) (Table 2). These effects persisted after additionally adjusting for

SEDH variables, and of the SEDH variables in the logistic regression model, only NO_2 exposure

was positively associated with exacerbations ($p = 0.0059$). Inclusion of the SEDH variables did

not improve model fit as determined by AIC (8,320 for both the EHR-adjusted and EHR &

SEDH-adjusted models). Sensitivity analyses showed that the risk factors identified by logistic

and negative binomial regression models were mostly consistent, with Black race and ICS

prescription having the strongest effects between both models ($p < 10^{-4}$), and variables such as

Medicaid health insurance and Elixhauser comorbidity score of 1-9 having statistically

significant p-values in both models though smaller in the negative binomial regression ($p <$

0.001) than the logistic regression ($p < 0.05$) (S7 Table). However, the negative binomial

regression model did not identify statistically significant associations between asthma

exacerbations and age 35-54 or NO_2 exposure as did the logistic regression, although the

directions of effect were consistent in both models. In addition, the negative binomial regression

model identified an association with the SEDH variable housing code violations ($p = 0.045$) that

was not present in the logistic regression model.

Spatial risk factors associated with exacerbations

Maps of ORs across the study region for the unadjusted, EHR-adjusted, and EHR & SEDH-

adjusted spatial GAMs are shown in Fig 2. In the unadjusted model, the global test of the null

hypothesis that asthma exacerbations were not associated with geographic location was

significant (p < 0.001) (Fig 2A). Local tests identified a statistically significant hotspot of

exacerbations (p < 0.01) in West and South Philadelphia with a mean spatial OR of 1.41 (SD

0.14). In the model adjusted for EHR-derived variables (Fig 2B) the global test statistic remained

significant (p < 0.001). Local tests again identified a hotspot (p < 0.01) in West and South

Philadelphia which had a decreased mean spatial OR of 1.27 (SD 0.055). In this EHR-adjusted

model, the variance in spatial ORs across the study region was 34.0% lower than the variance of

the unadjusted model (S5 Fig). In the model adjusted for both EHR-derived and SEDH variables,

the global test statistic remained significant (p < 0.001, Fig 2C) and local tests identified a

hotspot (p < 0.01) in West Philadelphia that overlapped geographically with the one in the other

models, although of a smaller area and with a smaller mean spatial OR of 1.24 (SD 0.042)

compared to the EHR-adjusted model. The variance in spatial ORs in the EHR & SEDH-

adjusted model was strongly attenuated (66.9% lower than that of the unadjusted model),

suggesting that these variables partially explained the spatial correlation of exacerbations (S5

Fig). The ORs for the other terms included in the spatial GAMs (i.e., the EHR-derived and

SEDH variables) are summarized in Table 3. All variables (i.e., years followed, age 35-54, Black

race, class 3 obesity, Medicaid health insurance, Elixhauser score 1-9, ICS) that were significant

in the multivariable logistic regression models (Table 2) were also significant in the spatial

GAMs (Table 3), except for NO_2 exposure, which was significant in the logistic regression

model but not in the spatial GAM.

   In the spatial GAM models adjusted one variable at a time, ADI, race, and health

insurance type most strongly attenuated the variation in spatial ORs (Fig 3), individually

accounting for 55.2%, 38.5%, and 26.5%, respectively, of the variation in the unadjusted model

(S5 Fig). In these models, each variable was positively associated with exacerbations (p <

0.001): ADI with OR = 1.05 (95% CI: 1.03, 1.07); Black race with OR = 1.66 (95% CI: 1.44,

1.91); and Medicaid health insurance with OR = 1.30 (95% CI: 1.15, 1.46) (S8 Table). The

spatial distribution of these variables followed similar patterns: high ADI and high density of

Black patients and patients with Medicaid insurance cooccurred in West, North, and South

Philadelphia (Fig 3). No other EHR (S6 Fig) or SEDH (S7 Fig) variable attenuated the hotspot

area or its effect size, however, adjustment for NO_2 levels resulted in an expansion of a coldspot

area (S7A Fig).

Stratified analysis by race

Bivariate analysis comparing the distribution of all EHR and SEDH characteristics between

patients of Black and White race found statistically significant differences for all variables except

ethnicity and ICS (Table 4). Given that ADI, race, and health insurance type had the strongest

relationship with asthma exacerbations in spatial analyses, and their spatial distributions were

similar, we performed stratified spatial analyses by race to determine whether ADI and health

insurance status remained significantly associated with asthma exacerbations within race-

stratified groups. After applying SRR inclusion criteria in strata according to race, an additional

265 Black patients were excluded from the stratified spatial analysis, resulting in a sample size of

4,363 patients (Fig 4A). The unadjusted spatial GAM for patients of Black race had a significant

global test statistic (p < 0.001) and local tests identified hotspots in West and South Philadelphia

consistent with results of the full cohort (Fig 4B). In contrast to spatial analyses in the full

cohort, adjusting for ADI and health insurance separately for Black patients did not attenuate the

spatial variance in ORs, instead increasing the variance by 4.70% and 2.79%, respectively,

compared to the unadjusted model (Fig 4C and 4D). In patients of White race, 73 additional

patients were removed after applying SRR inclusion criteria, resulting in 1,383 patients included

in the spatial model for which the global test statistic was no longer significant (p = 0.098, S8

Fig).

Discussion

Our analysis of individual-level risk factors found that Black race and ICS prescription had the

strongest positive associations with asthma exacerbations, as determined by individual-level

logistic regression and spatial GAM models. In our cohort of Penn Medicine asthma patients,

66% with no exacerbations were Black compared to 83% with 5+ exacerbations, consistent with

known racial disparities in asthma and observations in past Penn Medicine cohorts [44,45]. With

regard to the observed association between exacerbations and ICS prescription, international

asthma management guidelines underwent a major shift in 2019, recommending ICS as part of

the first-line treatment for all asthma patients [46]. This shift was not reflected in our cohort,

with only 76.0% of patients prescribed ICS. We observed exacerbations in patients in our cohort

regardless of whether they had an ICS prescription, however, the strong association between ICS

and exacerbations suggests that patients with more severe asthma were prescribed ICS more

frequently than those with milder asthma. Our sensitivity analysis found that logistic regression

identified positive associations between age 35-54 and NO_2 exposure that were not observed in a

negative binomial regression model, suggesting that these factors were associated with risk of

having at least one exacerbation compared to none, but not with having a higher count of

exacerbations.

Our spatial analysis of patient-level data revealed several important insights. First, we

observed that asthma exacerbation risk across Philadelphia was spatially correlated and

identified a hotspot with 41% higher odds of exacerbation compared to the median across the study region. This finding is consistent with the results of community-based pediatric asthma screening in Philadelphia, which have found that local asthma prevalence can vary significantly from regional or national estimates [47]. Our findings are also consistent with studies that have assessed spatial heterogeneity of pediatric and adult asthma data in other United States metropolitan areas, albeit with less granular spatial resolution. Zárate et al. observed statistically significant spatial patterning of asthma-related emergency department visits across census tracts in Central Texas [48], Grunwell et al. identified a group of contiguous census tracts in the State of Georgia with high rates of admission to the pediatric intensive care unit for asthma [49], and Harris et al. and Corburn et al. identified a statistically significant cluster of zip codes in St. Louis, Missouri and of census tracts in New York City, respectively, with elevated pediatric asthma hospitalization rates [50,51]. Our observation of a West and South Philadelphia hotspot of exacerbation risk is also consistent with analyses of past Penn Medicine cohorts [3,4] that focused on validating methods for augmenting EHR datasets rather than identifying factors associated with the hotspots.

The factors we found to be associated with asthma exacerbations according to EHR- and EHR & SEDH- adjusted spatial GAM models were largely consistent with individual-level logistic regression findings, but the spatial analysis provided an improved context to understand the individual-level results. Adjusting for all EHR and SEDH variables decreased the variance in spatial ORs by 66.9%, indicating that these variables together accounted for a large proportion of the spatial variance in exacerbation odds. By adjusting the spatial model one variable at a time, we found that ADI, race, and health insurance type most attenuated the hotspot area and effect size by reducing the variance of spatial ORs, suggesting that these variables were the most

influential in determining the spatial distribution of exacerbation risk. Our findings are consistent

with known asthma disparities by race/ethnicity and socioeconomic status [7–11], as well as past

neighborhood-level analyses of pediatric asthma: Harris et al. and Corburn et al. found that

asthma hospitalization hotspots in St. Louis and New York City had greater proportions of non-

White residents and greater rates of poverty, unemployment, high-density housing, and lack of

access to a household vehicle, although they did not test for statistical significance [50,51].

Grunwell et al. found a statistically significant difference between hotspot and non-hotspot

census tracts in Georgia for poverty, unemployment, and high-density housing, but not for race

[49].

Nearly all patient characteristics in our cohort, including health insurance type and ADI,

differed significantly between Black and White patients (Table 4). Most notably, 8.5% of White

patients had Medicaid health insurance compared to 39% of Black patients, and the median ADI

for White patients was 2.60 (IQR 1.50-4.30) compared to 8.40 for Black patients (IQR 6.50-

9.40), making it difficult to assess confounding in our spatial models. Although the VIF indicated

that all variables could be included in a multivariable model without substantially inflating

variance, collinearity between race and ADI as well as race and health insurance type (S4 Fig)

may help explain why ADI and health insurance type were statistically significant in bivariable

spatial models but not in the EHR & SEDH-adjusted spatial GAM nor in the EHR & SEDH-

adjusted logistic regression (Tables 2 and 3). We attempted to overcome some of these

limitations by stratifying our analysis by race. We found that, unlike in the full cohort, adjusting

for ADI and health insurance type for Black patients did not attenuate the variance in spatial

ORs, suggesting that in our cohort the association between these variables and asthma

exacerbations was confounded by race. Our results are consistent with previous observations that

racial disparities in asthma control persist even after accounting socioeconomic status [52], but

that it is difficult to separate out the effects of socioeconomic status from the effects of race [53].

Confounding of the asthma-socioeconomic status relationship by race has also been observed in

past neighborhood-level analyses of asthma morbidity. Zárate et al. found that spatial patterning

of asthma-related emergency department visits in Central Texas was partially explained by

socioeconomic characteristics in White patients, but not in Black or Hispanic patients [48]. More

broadly, understanding the relative contributions of many social determinants of health to asthma

is made difficult by their unequal distribution across racial/ethnic groups in the United States

[10].

Our health insurance type variable serves as a proxy for individual-level socioeconomic

status, and the relationship we measured between it and asthma exacerbation risk is potentially

mediated by several pathways including increased rates of smoking [54], psychosocial stress

[55], and obesity [56], or an unmeasured variable that varies with socioeconomic status and is

also associated with minoritized racial and ethnicity groups in the United States [57]. On the

other hand, low neighborhood-level socioeconomic status, which we measured with ADI, is

associated with differential exposure to indoor and outdoor air pollution, psychosocial stress

from neighborhood violence, and community norms surrounding health behaviors, all of which

have been linked to asthma exacerbations [27]. Due to geographic segregation by race that

resulted from structural racism and is common in many US cities, including in Philadelphia as

we observed in our study, race and neighborhood-level SES are also highly correlated [58]. Thus,

our inability to identify an association between asthma exacerbations and ADI when restricting

our analysis to Black patients may be due to a restriction of the range of people across the ADI

spectrum relative to the range observed in all patients [48]. Future work is needed to understand

what specific SEDH variables that covary with race, ADI, and health insurance type are the

primary drivers of local disparities in asthma exacerbation risk across Philadelphia.

Our results demonstrate that integration of diverse SEDH datasets with the EHR and the

use of both spatial and non-spatial modeling approaches are helpful to understand factors

contributing to complex health conditions in real world populations. In both our logistic

regression models and spatial GAMs, model fit was not improved by adjusting for SEDH

variables in addition to EHR-derived variables. However, in our spatial models, adjusting for

SEDH variables resulted in twice as much reduction of the initial variance in spatial ORs

compared to adjusting for EHR variables only. Our non-spatial and spatial models also identified

different sets of factors associated with exacerbations. For example, ICS prescription was found

to have a strong positive association with exacerbations in logistic regression models but did not

contribute to the spatial distribution of ORs; conversely, ADI attenuated the variance in spatial

ORs more strongly than all other variables tested but was not significant in either logistic

regression or negative binomial models. These findings present a framework for future efforts to

expand the scope of EHR data, which, especially as the spatial resolution of SEDH datasets

continues to increase, will allow for improved individualized exposure estimates. In the future,

integration of SEDH data into the EHR may be helpful to tailor asthma management strategies

and for health systems to create population-level interventions to improve health of their patients.

This study is strengthened by the high spatial resolution of both our SEDH data and our

analysis. Integrating the highest resolution SEDH data available during the time of the study

period allowed us to most closely approximate individual-level exposures, and analyzing EHR

data at a fine resolution allowed us to understand local health patterns that may not be visible at

the census tract or zip-code level. Additional strengths included accounting for many variables

and increasing the likelihood that Penn Medicine was patients' primary care provider by applying

SRR restriction. Our study is also subject to limitations, including some related to use of EHR

data, such as missingness, entry error, and phenotype misclassification. Additionally, the

geocoded addresses used in our study reflect residence at the time of the data pull, but they do

not account for residential mobility during the study period, nor does residence information

provide a full assessment of environmental exposures.

Conclusion

By integrating seven datasets containing information on SEDH with an EHR dataset to create

individualized exposure assessments, we identified non-spatial and spatial factors associated

with asthma exacerbations. Race and prescription of an ICS were most strongly associated with

exacerbations in individual-level models. Race also accounted for the most spatial variation in

exacerbation odds, along with ADI and health insurance type. Because these three variables had

similar spatial distributions, understanding which contributes most to disparities in asthma

exacerbations requires additional study of people living in the region identified as a hotspot. Our

findings demonstrate how integrating diverse data types and geospatial modeling approaches

with EHR data are helpful to understand complex diseases locally.

Acknowledgements

We would like to thank Sunil Thomas from the University of Pennsylvania Penn Data Store for

extracting the EHR data used for this project.

Table 1. Patient characteristics by exacerbation count levels. Shown are the characteristics of patients according to their number of exacerbations during the study period. For each exacerbation level, the number and percentage of patients are shown for categorical variables, and the Median and Interquartile Range (IQR) are shown for continuous variables.

Characteristic[a]	Number of Exacerbations				p-value[b]
	0 N = 4,327	**1-2** N = 1,810	**3-4** N = 328	**5+** N = 191	
Years followed	2.69 (1.89, 3.32)	2.81 (1.96, 3.46)	3.12 (2.48, 3.64)	3.60 (2.82, 3.84)	$<10^{-4}$
Age					$<10^{-4}$
18-34	1,498 (35%)	524 (29%)	82 (25%)	51 (27%)	
35-54	1,445 (33%)	688 (38%)	144 (44%)	81 (42%)	
55-74	1,198 (28%)	512 (28%)	87 (27%)	54 (28%)	
75+	186 (4.3%)	86 (4.8%)	15 (4.6%)	5 (2.6%)	
Sex					0.29
Male	1,020 (24%)	397 (22%)	71 (22%)	51 (27%)	
Female	3,307 (76%)	1,413 (78%)	257 (78%)	140 (73%)	
Race					$<10^{-4}$
White	1,059 (24%)	325 (18%)	52 (16%)	20 (10%)	
Black	2,860 (66%)	1,349 (75%)	260 (79%)	159 (83%)	
Unknown/Other	408 (9.4%)	136 (7.5%)	16 (4.9%)	12 (6.3%)	
Ethnicity					0.10
non-Hispanic/Latino	4,160 (96%)	1,747 (97%)	322 (98%)	188 (98%)	
Hispanic/Latino	167 (3.9%)	63 (3.5%)	6 (1.8%)	3 (1.6%)	
BMI					$<10^{-4}$
Not Overweight or Obese	907 (21%)	305 (17%)	49 (15%)	30 (16%)	
Overweight	1,003 (23%)	430 (24%)	81 (25%)	40 (21%)	
Class 1 Obesity	979 (23%)	381 (21%)	63 (19%)	37 (19%)	
Class 2 Obesity	650 (15%)	297 (16%)	40 (12%)	41 (21%)	
Class 3 Obesity	788 (18%)	397 (22%)	95 (29%)	43 (23%)	
Health insurance type					$<10^{-4}$
Private	1,976 (46%)	735 (41%)	131 (40%)	49 (26%)	
Medicaid	1,270 (29%)	625 (35%)	113 (34%)	87 (46%)	
Medicare	1,081 (25%)	450 (25%)	84 (26%)	55 (29%)	
Smoking status					$<10^{-4}$
Never Smoked	2,533 (59%)	997 (55%)	167 (51%)	84 (44%)	
Ever Smoker	1,233 (28%)	517 (29%)	110 (34%)	77 (40%)	
Current Smoker	561 (13%)	296 (16%)	51 (16%)	30 (16%)	
COPD	398 (9.2%)	219 (12%)	44 (13%)	30 (16%)	$<10^{-4}$
Allergic rhinitis	1,478 (34%)	636 (35%)	133 (41%)	104 (54%)	$<10^{-4}$
Elixhauser comorbidity score					$<10^{-4}$
<0	220 (5.1%)	85 (4.7%)	21 (6.4%)	5 (2.6%)	
0	3,035 (70%)	1,214 (67%)	193 (59%)	99 (52%)	
1-9	723 (17%)	356 (20%)	71 (22%)	60 (31%)	
10+	349 (8.1%)	155 (8.6%)	43 (13%)	27 (14%)	
ICS	3,074 (71%)	1,489 (82%)	309 (94%)	188 (98%)	$<10^{-4}$
NO$_2$ exposure	7.20 (6.86, 7.77)	7.18 (6.90, 7.70)	7.24 (6.86, 7.79)	7.14 (6.78, 7.56)	0.35
PM2.5 exposure	8.20 (7.84, 8.65)	8.16 (7.81, 8.62)	8.22 (7.86, 8.62)	8.14 (7.78, 8.58)	0.018
Toxic releases exposure	231 (5.3%)	87 (4.8%)	18 (5.5%)	10 (5.2%)	0.85
Vehicular traffic exposure					0.095

Lowest	1,079 (25%)	447 (25%)	85 (26%)	53 (28%)	
Low	1,079 (25%)	441 (24%)	96 (29%)	48 (25%)	
High	1,046 (24%)	485 (27%)	80 (24%)	53 (28%)	
Highest	1,123 (26%)	437 (24%)	67 (20%)	37 (19%)	
Area deprivation index	7.00 (3.50, 9.10)	7.80 (4.30, 9.20)	7.70 (5.30, 9.13)	7.80 (5.25, 9.30)	$<10^{-4}$
Housing violations	0.80 (0.36, 1.47)	0.83 (0.39, 1.49)	0.90 (0.45, 1.61)	0.84 (0.38, 1.28)	0.083
Normalized difference vegetation index	0.21 (0.17, 0.27)	0.21 (0.17, 0.26)	0.21 (0.16, 0.27)	0.22 (0.19, 0.26)	0.26

[a]Units are as follows: age (years), ICS (yes/no indicator of inhaled corticosteroid prescription), NO_2 (ppbv), PM2.5 ($\mu g/m^3$), toxic releases exposure (yes/no indicator of exposure), area deprivation index (unitless index scaled by dividing by 10), housing violations (housing violations per 100 people), normalized difference vegetation index (unitless index ranging from -1 to 1). See Methods for more details.

[b]Kruskal-Wallis rank sum test; Pearson's Chi-squared test

Table 2. Individual-level asthma exacerbation risk factors in multivariable logistic regression models. Shown are the adjusted odds ratios (ORs), 95% confidence intervals (CIs), and p-values for logistic regression models of asthma exacerbations as a dichotomous outcome adjusted for EHR-derived variables only and for both EHR and SEDH variables.

Characteristic[a]	EHR-adjusted			EHR & SEDH-adjusted		
	OR	95% CI	p-value	OR	95% CI	p-value
Years followed	1.16	1.09, 1.24	$<10^{-4}$	1.17	1.09, 1.24	$<10^{-4}$
Age						
18-34	—	—		—	—	
35-54	1.26	1.11, 1.44	5.2×10^{-4}	1.27	1.12, 1.45	3.6×10^{-4}
55-74	1.08	0.92, 1.28	0.35	1.09	0.92, 1.29	0.30
75+	1.29	0.96, 1.74	0.090	1.33	0.99, 1.79	0.060
Sex						
Male	—	—		—	—	
Female	0.99	0.87, 1.12	0.82	0.98	0.87, 1.12	0.79
Race						
White	—	—		—	—	
Black	1.49	1.29, 1.72	$<10^{-4}$	1.52	1.28, 1.81	$<10^{-4}$
Unknown/Other	1.01	0.79, 1.28	0.95	1.01	0.79, 1.29	0.93
Ethnicity						
Non-Hispanic/Latino	—	—		—	—	
Hispanic/Latino	0.99	0.71, 1.36	0.95	0.98	0.71, 1.35	0.92
BMI						
Not Overweight or Obese	—	—		—	—	
Overweight	1.17	0.99, 1.38	0.059	1.18	1.00, 1.39	0.054
Class 1 Obesity	0.97	0.82, 1.15	0.71	0.98	0.82, 1.16	0.80
Class 2 Obesity	1.09	0.90, 1.31	0.39	1.09	0.90, 1.31	0.39
Class 3 Obesity	1.24	1.04, 1.48	0.016	1.25	1.04, 1.49	0.015
Health insurance type						
Private	—	—		—	—	
Medicaid	1.22	1.07, 1.39	0.0034	1.19	1.04, 1.36	0.012
Medicare	0.90	0.77, 1.06	0.21	0.89	0.76, 1.04	0.15
Smoking status						
Never Smoked	—	—		—	—	
Ever Smoker	1.04	0.92, 1.18	0.51	1.04	0.92, 1.18	0.56
Current Smoker	1.17	0.99, 1.37	0.059	1.16	0.98, 1.36	0.077
COPD						
No	—	—		—	—	
Yes	1.14	0.95, 1.36	0.17	1.11	0.93, 1.34	0.24
Allergic rhinitis						
No	—	—		—	—	
Yes	1.09	0.97, 1.22	0.13	1.09	0.98, 1.22	0.11
Elixhauser comorbidity score						
<0	—	—		—	—	
0	1.20	0.94, 1.54	0.14	1.20	0.94, 1.54	0.14
1-9	1.40	1.08, 1.83	0.012	1.39	1.07, 1.82	0.015
10+	1.20	0.90, 1.62	0.22	1.19	0.89, 1.61	0.25
ICS						
No	—	—		—	—	
Yes	2.19	1.91, 2.51	$<10^{-4}$	2.20	1.92, 2.52	$<10^{-4}$
NO$_2$ exposure				1.27	1.07, 1.51	0.0059
PM2.5 exposure				0.89	0.72, 1.09	0.25

Toxic releases exposure						
No				—	—	
Yes				1.07	0.83, 1.37	0.60
Vehicular traffic exposure						
Lowest				—	—	
Low				1.01	0.87, 1.17	0.86
High				1.10	0.95, 1.28	0.19
Highest				0.95	0.82, 1.11	0.54
Area deprivation index				1.02	0.99, 1.04	0.22
Housing violations				0.96	0.90, 1.01	0.11
Normalized difference vegetation index				1.08	0.47, 2.44	0.86
AIC	8,320			8,320		

483 [a]Units are as follows: age (years), ICS (yes/no indicator of inhaled corticosteroid prescription), NO_2 (ppbv), PM2.5
484 ($\mu g/m^3$), toxic releases exposure (yes/no indicator of exposure), area deprivation index (unitless index scaled by
485 dividing by 10), housing violations (housing violations per 100 people), normalized difference vegetation index
486 (unitless index ranging from -1 to 1). See Methods for more details.

Table 3. Spatial asthma exacerbation risk factors in multivariable spatial GAMs. Shown are the adjusted odds ratios (ORs), 95% confidence intervals (CIs), and p-values for spatial GAMs of asthma exacerbations as a dichotomous outcome adjusted for EHR-derived variables only and for both EHR-derived and SEDH variables.

Characteristic[a]	EHR-adjusted			EHR & SEDH-adjusted		
	OR	95% CI	p-value	OR	95% CI	p-value
Years followed	1.17	1.10, 1.25	$<10^{-4}$	1.17	1.09, 1.25	$<10^{-4}$
Age						
18-34	—	—	—	—	—	—
35-54	1.28	1.12, 1.46	3.3×10^{-4}	1.27	1.12, 1.46	3.5×10^{-4}
55-74	1.08	0.91, 1.27	0.39	1.08	0.91, 1.27	0.38
75+	1.31	0.97, 1.76	0.078	1.32	0.98, 1.78	0.070
Sex						
Male	—	—	—	—	—	—
Female	0.99	0.87, 1.12	0.85	0.99	0.87, 1.12	0.84
Race						
White	—	—	—	—	—	—
Black	1.58	1.35, 1.84	$<10^{-4}$	1.55	1.30, 1.85	$<10^{-4}$
Unknown/Other	1.03	0.81, 1.31	0.81	1.02	0.79, 1.30	0.90
Ethnicity						
non-Hispanic/Latino	—	—	—	—	—	—
Hispanic/Latino	1.03	0.74, 1.42	0.87	1.02	0.74, 1.41	0.91
BMI						
Not Overweight or Obese	—	—	—	—	—	—
Overweight	1.18	1.01, 1.40	0.043	1.18	1.00, 1.39	0.047
Class 1 Obesity	0.98	0.83, 1.16	0.82	0.98	0.83, 1.16	0.81
Class 2 Obesity	1.09	0.90, 1.31	0.37	1.09	0.90, 1.31	0.39
Class 3 Obesity	1.26	1.05, 1.50	0.011	1.25	1.05, 1.50	0.013
Health insurance type						
Private	—	—	—	—	—	—
Medicaid	1.18	1.03, 1.35	0.014	1.18	1.03, 1.35	0.018
Medicare	0.89	0.76, 1.05	0.16	0.89	0.76, 1.04	0.15
Smoking status						
Never Smoked	—	—	—	—	—	—
Ever Smoker	1.04	0.91, 1.17	0.57	1.03	0.91, 1.17	0.59
Current Smoker	1.15	0.98, 1.35	0.094	1.15	0.98, 1.35	0.095
COPD						
No	—	—	—	—	—	—
Yes	1.10	0.92, 1.32	0.30	1.10	0.92, 1.32	0.29
Allergic rhinitis						
No	—	—	—	—	—	—
Yes	1.09	0.98, 1.22	0.12	1.10	0.98, 1.22	0.11
Elixhauser comorbidity score						
<0	—	—	—	—	—	—
0	1.20	0.94, 1.54	0.14	1.20	0.94, 1.53	0.15
1-9	1.39	1.06, 1.81	0.016	1.38	1.06, 1.80	0.018
10+	1.18	0.87, 1.59	0.28	1.17	0.87, 1.58	0.29
ICS						
No	—	—	—	—	—	—
Yes	2.22	1.93, 2.54	$<10^{-4}$	2.21	1.93, 2.54	$<10^{-4}$
NO$_2$ exposure				1.08	0.82, 1.43	0.58
PM2.5 exposure				1.08	0.81, 1.45	0.59

Toxic releases exposure						
No				—	—	—
Yes				1.06	0.82, 1.37	0.65
Vehicular traffic exposure						
Lowest				—	—	—
Low				1.02	0.88, 1.18	0.76
High				1.10	0.95, 1.27	0.22
Highest				0.97	0.83, 1.13	0.66
Area deprivation index				1.01	0.99, 1.04	0.32
Housing violations				0.97	0.91, 1.02	0.23
Normalized difference vegetation index				0.99	0.43, 2.27	0.97
AIC[b]	8,293			8,307		

492 [a]Units are as follows: age (years), ICS (yes/no indicator of inhaled corticosteroid prescription), NO_2 (ppbv), PM2.5
493 ($\mu g/m^3$), toxic releases exposure (yes/no indicator of exposure), area deprivation index (unitless index scaled by
494 dividing by 10), housing violations (housing violations per 100 people), normalized difference vegetation index
495 (unitless index ranging from -1 to 1). See Methods for more details.
496 [b]The AIC of the unadjusted model was 8,576.

Table 4. Patient characteristics by race. Shown are the number and percentage of patients in each level for categorical variables, and the Median and Interquartile Range (IQR) for continuous variables in patients of White race versus Black race.

Characteristic[a]	Race		p-value[b]
	White N = 1,456	Black N = 4,628	
Exacerbation count			$<10^{-4}$
0	1,059 (73%)	2,860 (62%)	
1-2	325 (22%)	1,349 (29%)	
3-4	52 (3.6%)	260 (5.6%)	
5+	20 (1.4%)	159 (3.4%)	
Years followed	2.62 (1.86, 3.28)	2.82 (1.98, 3.46)	$<10^{-4}$
Age			$<10^{-4}$
18-34	468 (32%)	1,458 (32%)	
35-54	481 (33%)	1,667 (36%)	
55-74	407 (28%)	1,332 (29%)	
75+	100 (6.9%)	171 (3.7%)	
Sex			$<10^{-4}$
Male	473 (32%)	905 (20%)	
Female	983 (68%)	3,723 (80%)	
Ethnicity			0.0024
non-Hispanic/Latino	1,431 (98%)	4,591 (99%)	
Hispanic/Latino	25 (1.7%)	37 (0.8%)	
BMI			$<10^{-4}$
Not Overweight or Obese	522 (36%)	633 (14%)	
Overweight	461 (32%)	936 (20%)	
Class 1 Obesity	265 (18%)	1,051 (23%)	
Class 2 Obesity	106 (7.3%)	859 (19%)	
Class 3 Obesity	102 (7.0%)	1,149 (25%)	
Health insurance type			$<10^{-4}$
Private	981 (67%)	1,621 (35%)	
Medicaid	124 (8.5%)	1,782 (39%)	
Medicare	351 (24%)	1,225 (26%)	
Smoking status			$<10^{-4}$
Never Smoked	898 (62%)	2,519 (54%)	
Ever Smoker	463 (32%)	1,328 (29%)	
Current Smoker	95 (6.5%)	781 (17%)	
COPD	115 (7.9%)	536 (12%)	$<10^{-4}$
Allergic rhinitis	571 (39%)	1,541 (33%)	$<10^{-4}$
Elixhauser comorbidity score			$<10^{-4}$
<0	32 (2.2%)	262 (5.7%)	
0	1,117 (77%)	2,986 (65%)	
1-9	222 (15%)	924 (20%)	
10+	85 (5.8%)	456 (9.9%)	
ICS	1,105 (76%)	3,510 (76%)	0.97
NO₂ exposure	7.85 (7.40, 8.18)	7.10 (6.81, 7.39)	$<10^{-4}$
PM2.5 exposure	8.72 (8.28, 8.90)	8.05 (7.77, 8.41)	$<10^{-4}$
Toxic releases exposure	115 (7.9%)	194 (4.2%)	$<10^{-4}$
Vehicular traffic exposure			$<10^{-4}$
Lowest	279 (19%)	1,283 (28%)	
Low	267 (18%)	1,244 (27%)	
High	312 (21%)	1,177 (25%)	

Highest	598 (41%)	924 (20%)	
Area deprivation index	2.60 (1.50, 4.30)	8.40 (6.50, 9.40)	$<10^{-4}$
Housing violations	0.41 (0.15, 0.74)	0.99 (0.52, 1.68)	$<10^{-4}$
Normalized difference vegetation index	0.18 (0.13, 0.25)	0.22 (0.18, 0.27)	$<10^{-4}$

501 [a]Units are as follows: age (years), ICS (yes/no indicator of inhaled corticosteroid prescription), NO_2 (ppbv), PM2.5
502 ($\mu g/m^3$), toxic releases exposure (yes/no indicator of exposure), area deprivation index (unitless index scaled by
503 dividing by 10), housing violations (housing violations per 100 people), normalized difference vegetation index
504 (unitless index ranging from -1 to 1). See Methods for more details.
505 [b]Pearson's Chi-squared test; Wilcoxon rank sum test

 Figure Legends

Figure 1. Overview of study design. Graphical overview of study design, including processing

and linkage of electronic health record (EHR) and social and environmental determinants of

health (SEDH) data, cohort selection including spatial filtering by assessing the representation of

the EHR cohort compared to the underlying population, and patient-level and geospatial analyses

on the expanded EHR dataset.

Figure 2. Spatial odds ratios (ORs) of exacerbations before and after adjusting for EHR-

derived and SEDH variables. (A) Unadjusted spatial GAM (adjusted only for years followed).

(B) Spatial GAM adjusted for EHR-derived variables only. (C) Spatial GAM adjusted for both

EHR-derived and SEDH variables. Base maps were created using the Stamen Design from

Stadia Maps.

Figure 3. Spatial distribution of individual variables that most strongly attenuated the

spatial odds ratios (ORs) of exacerbations along with corresponding spatial GAM results

adjusted for these individual variables. Spatial distribution in the study region of (A) the area

deprivation index (ADI), (B) race, and (C) health insurance type of patients. Corresponding

spatial GAMs adjusted only for years followed and (D) ADI, (E) race, or (F) health insurance

type.

Figure 4. Spatial odds ratios (ORs) of exacerbations among Black patients along with the

effects of ADI and health insurance type on this distribution. (A) SRR values for the updated

study region used in spatial GAMs for patients of Black race only (SRR = 1 indicates no

representativeness bias). (B) Unadjusted spatial GAM (adjusted only for years followed) for

patients of Black race (N = 4,363). Spatial GAMs adjusted additionally for (C) area deprivation

index (ADI) and (D) health insurance type. Base maps were created using the Stamen Design

from Stadia Maps.

References

1. Mooney SJ, Westreich DJ, El-Sayed AM. Commentary: Epidemiology in the era of big data. Epidemiology. 2015;c: 390–394. doi:10.1097/EDE.0000000000000274

2. Cook LA, Sachs J, Weiskopf NG. The quality of social determinants data in the electronic health record: a systematic review. J Am Med Inform Assoc JAMIA. 2021;29: 187–196. doi:10.1093/jamia/ocab199

3. Xie S, Greenblatt R, Levy MZ, Himes BE. Enhancing Electronic Health Record Data with Geospatial Information. AMIA Summits Transl Sci Proc. 2017;2017: 123–132.

4. Xie S, Himes BE. Approaches to link geospatially varying social, economic, and environmental factors with electronic health record data to better understand asthma exacerbations. AMIA Annu Symp Proc. 2018;2018: 1561–1570.

5. Cui Y, Eccles KM, Kwok RK, Joubert BR, Messier KP, Balshaw DM. Integrating multiscale geospatial environmental data into large population health studies: challenges and opportunities. Toxics. 2022;10: 403. doi:10.3390/toxics10070403

6. Schreibman A, Xie S, Hubbard RA, Himes BE. Linking ambient NO2 pollution measures with electronic health record data to study asthma exacerbations. AMIA Summits Transl Sci Proc. 2023;2023: 467–476.

7. CDC. Asthma: most recent national asthma data. Atlanta, GA: US Department of Health and Human Services, CDC; 2020. Available: https://www.cdc.gov/asthma/most_recent_national_asthma_data.htm

8. Leong AB, Ramsey CD, Celedón JC. The challenge of asthma in minority populations. Clin Rev Allergy Immunol. 2012;43: 156–183. doi:10.1007/s12016-011-8263-1

9. Forno E, Celedón JC. Health disparities in asthma. Am J Respir Crit Care Med. 2012;185: 1033–1035. doi:10.1164/rccm.201202-0350ED

10. Grant T, Croce E, Matsui EC. Asthma and the social determinants of health. Ann Allergy Asthma Immunol Off Publ Am Coll Allergy Asthma Immunol. 2022;128: 5–11. doi:10.1016/j.anai.2021.10.002

11. Lovinsky-Desir S, Riley IL, Bryant-Stephens T, De Keyser H, Forno E, Kozik AJ, et al. Research priorities in pediatric asthma morbidity: addressing the impacts of systemic racism on children with asthma in the United States: an official American Thoracic Society workshop report. Ann Am Thorac Soc. 2024;21: 1349–1364. doi:10.1513/AnnalsATS.202407-767ST

12. Global Initiative for Asthma (GINA). Global strategy for asthma management and prevention (2024 update). Bethesda, MD: Global Initiative for Asthma (GINA); 2024.

13. Fuhlbrigge A, Peden D, Apter AJ, Boushey HA, Camargo CA, Gern J, et al. Asthma
outcomes: exacerbations. J Allergy Clin Immunol. 2012;129: S34–S48.
doi:10.1016/j.jaci.2011.12.983

14. Krishnan V, Diette GB, Rand CS, Bilderback AL, Merriman B, Hansel NN, et al. Mortality
in patients hospitalized for asthma exacerbations in the United States. Am J Respir Crit
Care Med. 2006;174: 633–638. doi:10.1164/rccm.200601-007OC

15. Ivanova JI, Bergman R, Birnbaum HG, Colice GL, Silverman RA, McLaurin K. Effect of
asthma exacerbations on health care costs among asthmatic patients with moderate and
severe persistent asthma. J Allergy Clin Immunol. 2012;129: 1229–1235.
doi:10.1016/j.jaci.2012.01.039

16. Schatz M, Clark S, Camargo CA. Sex differences in the presentation and course of asthma
hospitalizations. Chest. 2006;129: 50–55. doi:10.1378/chest.129.1.50

17. Schatz M, Mosen DM, Kosinski M, Vollmer WM, Magid DJ, O'Connor E, et al. Predictors
of asthma control in a random sample of asthmatic patients. J Asthma. 2007;44: 341–345.
doi:10.1080/02770900701344421

18. Pedersen SE, Bateman ED, Bousquet J, Busse WW, Yoxall S, Clark TJ. Determinants of
response to fluticasone propionate and salmeterol/fluticasone propionate combination in the
Gaining Optimal Asthma controL study. J Allergy Clin Immunol. 2007;120: 1036–1042.
doi:10.1016/j.jaci.2007.07.016

19. Price D, Zhang Q, Kocevar VS, Yin DD, Thomas M. Effect of a concomitant diagnosis of
allergic rhinitis on asthma-related health care use by adults. Clin Exp Allergy. 2005;35:
282–287. doi:10.1111/j.1365-2222.2005.02182.x

20. Miller MK, Lee JH, Miller DP, Wenzel SE. Recent asthma exacerbations: A key predictor of
future exacerbations. Respir Med. 2007;101: 481–489. doi:10.1016/j.rmed.2006.07.005

21. Guarnieri M, Balmes JR. Outdoor air pollution and asthma. The Lancet. 2014;383: 1581–
1592. doi:10.1016/S0140-6736(14)60617-6

22. Weitzman M, Gortmaker S, Sobol A. Racial, social, and environmental risks for childhood
asthma. Am J Dis Child. 1990;144: 1189–1194.
doi:10.1001/archpedi.1990.02150350021016

23. Black PN, Udy AA, Brodie SM. Sensitivity to fungal allergens is a risk factor for life-
threatening asthma. Allergy. 2000;55: 501–504. doi:10.1034/j.1398-9995.2000.00293.x

24. Gelber LE, Seltzer LH, Bouzoukis JK, Pollart SM, Chapman MD, Platts-Mills TAE.
Sensitization and exposure to indoor allergens as risk factors for asthma among patients
presenting to hospital. Am Rev Respir Dis. 1993;147: 573–578.
doi:10.1164/ajrccm/147.3.573

25. Lovasi GS, O'Neil-Dunne JPM, Lu JWT, Sheehan D, Perzanowski MS, MacFaden SW, et al.
Urban tree canopy and asthma, wheeze, rhinitis, and allergic sensitization to tree pollen in a
New York City birth cohort. Environ Health Perspect. 2013;121: 494–500.
doi:10.1289/ehp.1205513

26. Lovasi GS, Quinn JW, Neckerman KM, Perzanowski MS, Rundle A. Children living in areas
with more street trees have lower prevalence of asthma. J Epidemiol Community Health.
2008;62: 647–649. doi:10.1136/jech.2007.071894

27. Gold DR, Wright R. Population disparities in asthma. Annu Rev Public Health. 2005;26: 89–
113. doi:10.1146/annurev.publhealth.26.021304.144528

28. Fecho K, Ahalt SC, Appold S, Arunachalam S, Pfaff E, Stillwell L, et al. Development and
application of an open tool for sharing and analyzing integrated clinical and environmental
exposures data: asthma use case. JMIR Form Res. 2022;6: e32357. doi:10.2196/32357

29. Beck AF, Huang B, Ryan PH, Sandel MT, Chen C, Kahn RS. Areas with high rates of police-
reported violent crime have higher rates of childhood asthma morbidity. J Pediatr.
2016;173: 175-182.e1. doi:10.1016/j.jpeds.2016.02.018

30. Rasmussen SG, Ogburn EL, McCormack M, Casey JA, Bandeen-Roche K, Mercer DG, et al.
Association between unconventional natural gas development in the Marcellus Shale and
asthma exacerbations. JAMA Intern Med. 2016;176: 1334–1343.
doi:10.1001/jamainternmed.2016.2436

31. van Walraven C, Austin PC, Jennings A, Quan H, Forster AJ. A modification of the
Elixhauser comorbidity measures into a point system for hospital death using administrative
data. Med Care. 2009;47: 626. doi:10.1097/MLR.0b013e31819432e5

32. Christie C, Xie S, Diwadkar AR, Greenblatt RE, Rizaldi A, Himes BE. Consolidated
Environmental and Social Data Facilitates Neighborhood-Level Health Studies in
Philadelphia. AMIA Annu Symp Proc. 2022;2021: 305–313.

33. Cooper MJ, Martin RV, McLinden CA, Brook JR. Inferring ground-level nitrogen dioxide
concentrations at fine spatial resolution applied to the TROPOMI satellite instrument.
Environ Res Lett. 2020;15: 104013. doi:10.1088/1748-9326/aba3a5

34. van Donkelaar A, Hammer MS, Bindle L, Brauer M, Brook JR, Garay MJ, et al. Monthly
global estimates of fine particulate matter and their uncertainty. Environ Sci Technol.
2021;55: 15287–15300. doi:10.1021/acs.est.1c05309

35. US EPA. TRI basic data files: calendar years 1987-present. 3 Mar 2013 [cited 25 Jul 2023].
Available: https://www.epa.gov/toxics-release-inventory-tri-program/tri-basic-data-files-
calendar-years-1987-present

36. RMSTRAFFIC (Traffic Volumes). 2023 [cited 25 Jul 2023]. Available: https://data-
pennshare.opendata.arcgis.com/datasets/rmstraffic-traffic-volumes/explore

37. Kind AJH, Buckingham WR. Making neighborhood-disadvantage metrics accessible — the neighborhood atlas. N Engl J Med. 2018;378: 2456–2458. doi:10.1056/NEJMp1802313

38. University of Wisconsin School of Medicine Public Health. 2018 Area Deprivation Index v3.1. Available: https://www.neighborhoodatlas.medicine.wisc.edu/. Accessed July 2022.

39. Licenses and Inspections code violations. In: OpenDataPhilly [Internet]. [cited 25 Jul 2023]. Available: https://opendataphilly.org/datasets/licenses-and-inspections-code-violations/

40. Ermida SL, Soares P, Mantas V, Göttsche F-M, Trigo IF. Google Earth Engine open-source code for land surface temperature estimation from the Landsat series. Remote Sens. 2020;12: 1471. doi:10.3390/rs12091471

41. R Core Team (2020). R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; Available: https://www.R-project.org/

42. Xie SJ, Kapos FP, Mooney SJ, Mooney S, Stephens KA, Chen C, et al. Geospatial divide in real-world EHR data: analytical workflow to assess regional biases and potential impact on health equity. AMIA Summits Transl Sci Proc. 2023;2023: 572–581.

43. Bai L, Bartell S, Bliss R, Vieira V. Mapping smoothed effect estimates from individual-level spatial data. 2022. Available: https://search.r-project.org/CRAN/refmans/MapGAM/html/MapGAM-package.html

44. Moorman JE, Akinbami LJ, Bailey CM, Zahran HS, King ME, Johnson CA, et al. National surveillance of asthma: United States, 2001-2010. Vital Health Stat 3. 2012; 1–58.

45. Greenblatt RE, Zhao EJ, Henrickson SE, Apter AJ, Hubbard RA, Himes BE. Factors associated with exacerbations among adults with asthma according to electronic health record data. Asthma Res Pract. 2019;5: 1. doi:10.1186/s40733-019-0048-y

46. Global Initiative for Asthma (GINA). Global strategy for asthma management and prevention (2019 update). Bethesda, MD: Global Initiative for Asthma (GINA); 2019.

47. Bryant-Stephens T, West C, Dirl C, Banks T, Briggs V, Rosenthal M. Asthma prevalence in philadelphia: description of two community-based methodologies to assess asthma prevalence in an inner-city population. J Asthma. 2012;49: 581–585. doi:10.3109/02770903.2012.690476

48. Zárate RebeccaA, Bhavnani D, Chambliss S, Hall EM, Zigler C, Cubbin C, et al. Neighborhood-level variability in asthma-related emergency department visits in Central Texas. J Allergy Clin Immunol. 2024;154: 933–939. doi:10.1016/j.jaci.2024.05.024

49. Grunwell JR, Opolka C, Mason C, Fitzpatrick AM. Geospatial analysis of social determinants of health identifies neighborhood hot spots associated with pediatric intensive care use for life-threatening asthma. J Allergy Clin Immunol Pract. 2022;10: 981-991.e1. doi:10.1016/j.jaip.2021.10.065

50. Harris KM. Mapping inequality: Childhood asthma and environmental injustice, a case study
of St. Louis, Missouri. Soc Sci Med. 2019;230: 91–110.
doi:10.1016/j.socscimed.2019.03.040

51. Corburn J, Osleeb J, Porter M. Urban asthma and the neighbourhood environment in New
York City. Health Place. 2006;12: 167–179. doi:10.1016/j.healthplace.2004.11.002

52. Haselkorn T, Lee JH, Mink DR, Weiss ST. Racial disparities in asthma-related health
outcomes in severe or difficult-to-treat asthma. Ann Allergy Asthma Immunol. 2008;101:
256–263. doi:10.1016/S1081-1206(10)60490-5

53. Boudreaux ED, Emond SD, Clark S, Camargo CA. Acute asthma among adults presenting to
the emergency department: the role of race/ethnicity and socioeconomic status. Chest.
2003;124: 803–812. doi:10.1378/chest.124.3.803

54. Hiscock R, Bauld L, Amos A, Fidler JA, Munafò M. Socioeconomic status and smoking: a
review. Ann N Y Acad Sci. 2012;1248: 107–123. doi:10.1111/j.1749-6632.2011.06202.x

55. Matthews KA, Gallo LC, Taylor SE. Are psychosocial factors mediators of socioeconomic
status and health connections? Ann N Y Acad Sci. 2010;1186: 146–173. doi:10.1111/j.1749-
6632.2009.05332.x

56. Wang Y, Beydoun MA. The obesity epidemic in the United States—gender, age,
socioeconomic, racial/ethnic, and geographic characteristics: a systematic review and meta-
regression analysis. Epidemiol Rev. 2007;29: 6–28. doi:10.1093/epirev/mxm007

57. Williams DR. Race, socioeconomic status, and health: the added effects of racism and
discrimination. Ann N Y Acad Sci. 1999;896: 173–188. doi:10.1111/j.1749-
6632.1999.tb08114.x

58. Charles CZ. The dynamics of racial residential segregation. Annu Rev Sociol. 2003;29: 167–
207. doi:10.1146/annurev.soc.29.010202.100002

Supporting information

S1 Text. Supplementary Methods.

S1 Figure. Flowchart of patient cohort selection. Overview of steps followed to select final patient cohort (N = 6,656) from EHR data on all Penn Medicine patients with at least one asthma ICD code (N = 86,787).

S2 Figure. Selection of study region using the spatial representation ratio (SRR). (A) SRR values, defined as the cohort population residing in a block group divided by the underlying population as reported by the 2019 American Community Survey, for all of Philadelphia and for the selected study region (inset box). SRR = 1 indicates no representativeness bias. Density plots of the study cohort (B) before and (C) after filtering for the study region. Base maps were created using the Stamen Design from Stadia Maps.

S3 Figure. Spatial distribution of SEDH datasets that were integrated with EHR data. The following maps are shown for the spatial area which comprised our study region: (A) raster of NO2 pollution levels, (B) raster of PM2.5 pollution levels, (C) point sites of toxic releases and the total summed emissions at each site, (D) line segments of roadways and the daily vehicle miles traveled (DVDT) on each, (E) housing violations per block group, normalized by the underlying 2019 American Community Survey population, (F) raster of the normalized difference vegetation index (NDVI). A map of area deprivation index (ADI), which most strongly reduced the spatial variance of odds of exacerbation risk in our spatial GAMs, is shown in Figure 3A. Base maps were created using the Stamen Design from Stadia Maps.

S4 Figure. Pairwise correlation between all EHR and SEDH variables. Measures in each box correspond to Pearson's correlation coefficients. For nominal categorical variables, reference levels are as follows: White (race), Private (health insurance type).

S5 Figure. Influence of individual EHR and SEDH variables on the odds ratio (OR) of the

unadjusted spatial GAM. Percent reduction in the variance of ORs across the study region for

one-variable-at-a-time adjusted spatial GAMs compared to the unadjusted model. OR changes

for models adjusted one variable at a time (in addition to years followed) are shown in blue.

Multivariable models (both EHR-adjusted and EHR & SEDH-adjusted) are shown in red for

comparison.

S6 Figure. Spatial GAMs adjusted one-at-a-time for the EHR-derived variables that did

not greatly reduce variance. Spatial odds ratios (ORs) of exacerbation are shown after

adjusting for years followed and one-at-a-time for the following variables whose percent

reduction in variance of ORs was less than 25: (A) age, (B) sex, (C) ethnicity, (D) BMI, (E)

smoking status, (F) COPD, (G) allergic rhinitis, (H) Elixhauser comorbidity score, (I) ICS. Base

maps were created using the Stamen Design from Stadia Maps.

S7 Figure. Spatial GAMs adjusted one-at-a-time for the SEDH variables that did not

greatly reduce variance. Spatial odds ratios (ORs) of exacerbation are shown after adjusting for

years followed and one-at-a-time for the following variables whose reduction in variance of ORs

was less than 25: (A) NO2, (B) PM2.5, (C) toxic releases exposure, (D) vehicular traffic, (E)

housing violations, (F) normalized difference vegetation index (NDVI). Base maps were created

using the Stamen Design from Stadia Maps.

S8 Figure. Spatial odds ratios (ORs) of exacerbations among White patients along with the

effects of ADI and health insurance type on this distribution. (A) SRR values for the updated

study region used in spatial GAMs for patients of White race only (SRR = 1 indicates no

representativeness bias). (B) Unadjusted spatial GAM adjusted only for years followed for

patients of White race (N = 1,383). Spatial GAMs adjusted additionally for (C) area deprivation

index (ADI) and (D) health insurance type. Base maps were created using the Stamen Design

from Stadia Maps.

S1 Table. **Generic medication names included in medication classes.** The following generic

drug names recorded in the EHR during the study period were used for asthma and exacerbation

phenotyping as well as used as independent variables in select models (i.e., ICS). Instances in

which these drugs were listed as investigational or nasal formulations were not included.

S2 Table. **Sources and spatiotemporal dimensions of geospatial datasets merged with EHR**

data.

S3 Table. **Asthma-related housing code violations extracted from the Philadelphia**

Department of Licenses and Inspections.

S4 Table. **Patient medications by exacerbation count levels.** Shown are the number and

percentage of patients receiving each of the medication types listed according to their number of

exacerbations during the study period.

S5 Table. Characteristics of complete cases and patients excluded due to missingness.

Shown are the number and percentage of patients in each level for categorical variables and the

Median and Interquartile Range (IQR) for the Years followed variable in complete cases versus

those excluded due to missingness in the sex, ethnicity, health insurance type, BMI, and smoking

status variables.

S6 Table. Adjusted Generalized Variance Inflation Factors (GVIFs) for each EHR and

SEDH variable included in the EHR & SEDH-adjusted negative binomial and logistic

regression models.

S7 Table. **Individual-level asthma exacerbation risk factors in multivariable negative**

binomial regression models. Shown are the adjusted incidence rate ratios (IRRs), 95%

confidence intervals (CIs), and p-values for negative binomial models of asthma exacerbations as

a count outcome adjusted for EHR-derived variables only and for both EHR-derived and SEDH

variables.

S8 Table. Spatial GAMs of asthma exacerbations adjusted for individual risk factors that

most changed risk. Shown are the adjusted odds ratios (ORs), 95% confidence intervals (CIs),

and p-values for spatial GAMs of asthma exacerbations as a dichotomous outcome adjusted for

years followed and one-at-a-time for ADI, race, and health insurance type, the three variables

whose percent reduction in variance of ORs was greater than 25.

Figure 1

(A) Univariable

(B) EHR-adjusted

(C) EHR & SEDH-adjusted

Figure 2

(A) ADI distribution

(D) ADI-adjusted GAM

(B) Race distribution

(E) Race-adjusted GAM

(C) Health insurance type distribution

(F) Health insurance type-adjusted GAM

Figure 3

(A) Stratified SRR selection

SRR
- ≤ 0.05
- 0.051–0.5
- 0.51–1.0
- 1.01–3.0
- 3.01–5.0
- 5.01–7.0
- 7.01+

(B) Univariable GAM

OR
- 2.0
- 1.5
- 1.0
- 0.5

(C) ADI-adjusted GAM

OR
- 2.0
- 1.5
- 1.0
- 0.5

(D) Health insurance type-adjusted GAM

OR
- 2.0
- 1.5
- 1.0
- 0.5

Figure 4