Timing of changes in Alzheimer's disease plasma biomarkers as assessed by amyloid and tau PET clocks

Marta Milà-Alomà[1,2], Duygu Tosun[1,2], Suzanne E. Schindler[3], Zachary Hausle[1,2], Yan Li[3], Kellen K. Petersen[3], Jeffrey L. Dage[4,5], Lei Du-Cuny[6], Ziad S. Saad[7], Benjamin Saef[3], Gallen Triana-Baltzer[7], David L. Raunig[8], Janaky Coomaraswamy[8], Michael Baratta[8], Emily A. Meyers[9], Yulia Mordashova[6], Carrie E. Rubel[10], Kyle Ferber[10], Hartmuth Kolb[11], Nicholas J. Ashton[12], Henrik Zetterberg[12-17], Erin G. Rosenbaugh[18], Leslie M. Shaw[19], Anthony W. Bannon[20], William Z. Potter[21], Alzheimer's Disease Neuroimaging Initiative (ADNI)**, Foundation for the National Institutes of Health (FNIH) Biomarkers Consortium Plasma Aβ and Phosphorylated Tau as Predictors of Amyloid and Tau Positivity in Alzheimer's Disease Project Team

Abstract

Plasma biomarkers for Alzheimer's disease (AD) are increasingly being used to assist in making an etiological diagnosis for cognitively impaired (CI) individuals or to identify cognitively unimpaired (CU) individuals with AD pathology who may be eligible for prevention trials. However, a better understanding of the timing of plasma biomarker changes is needed to optimize their use in clinical and research settings. The aim of this study was to evaluate the timing of change of key AD plasma biomarkers (Aβ42/Aβ40, p-tau181, p-tau217, GFAP and NfL) from six different companies, along with established AD biomarkers, using AD progression timelines based on amyloid and tau PET.

We used data from the Alzheimer's Disease Neuroimaging Initiative (ADNI), including 784 individuals with longitudinal amyloid PET and 359 individuals with longitudinal tau PET, to estimate age at amyloid and tau positivity, defined as the age at the first positive PET scan ([18F] florbetapir amyloid PET SUVR>0.78, [18F] flortaucipir tau PET SUVR>1.41, respectively). Of these, 190 individuals with an estimated age at amyloid positivity and 70 individuals with an estimated age at tau positivity had longitudinal plasma biomarker measures. Age at tau positivity was a stronger predictor of symptom onset than age at amyloid positivity in 17 individuals who progressed from CU to CI during their participation in the ADNI study

(Adj R^2 = 0.86 *vs.* Adj R^2 = 0.38), therefore was used to estimate symptom onset age for all individuals with an estimated age at tau positivity. Generalized additive mixed models (GAMMs) were used to model biomarker trajectories across years since amyloid positivity, tau positivity, and symptom onset, and to identify the earliest timepoint of biomarker abnormality when compared to a reference group of amyloid and tau PET-negative CU individuals, as well as time periods of significant change in biomarkers.

All plasma biomarkers except NfL became abnormal prior to amyloid and tau positivity. Plasma Aβ42/Aβ40 was the first biomarker to reach abnormality consistently across timelines. Plasma GFAP was the earliest biomarker to become abnormal in the tau timeline. The plasma Aβ42/Aβ40 levels reached a plateau, while plasma p-tau217, p-tau181, GFAP and NfL increased throughout disease progression. Some differences in the timing of change were observed across biomarker assays.

The primary utility of plasma Aβ42/Aβ40 may lie in early identification of individuals at high risk of AD. In contrast, p-tau217, p-tau181, GFAP and NfL increase throughout the estimated timelines, supporting their potential as biomarkers for staging and monitoring disease progression.

Author affiliations:

[1]Northern California Institute for Research and Education, San Francisco, CA, USA 94141

[2]Department of Radiology and Biomedical Imaging, University of California San Francisco, San Francisco, CA, USA 94143

[3]Department of Neurology, Washington University in St. Louis, St. Louis, MO, USA 63110

[4]Department of Neurology, Indiana University School of Medicine, 355 W 16th St Indianapolis, IN, USA 46202

[5]Stark Neurosciences Research Institute, Indiana University School of Medicine, NB Building, 320 W 15th St #414, Indianapolis, IN, USA 46202

[6]AbbVie, Knollstraße 50, 67061, Ludwigshafen am Rhein, Rheinland-Pfalz, Germany

[7]Neuroscience Biomarkers, Johnson and Johnson Innovative Medicine, 3210 Merryfield Row, San Diego, CA, USA 20892

[8]Takeda Pharmaceutical Company Ltd., 500 Kendall St., Cambridge, MA, USA 02142

[9]Alzheimer's Association, 225 N Michigan Ave., Chicago, IL, USA 60603

[10]Biogen, 225 Binney St., Cambridge, MA, USA 02142

[11]Enigma Biomedical Group, CA, USA

[12] Institute of Neuroscience and Physiology, Department of Psychiatry and Neurochemistry, The Sahlgrenska Academy at University of Gothenburg, Biskopsbogatan 27, S-431 80, Mölndal, Sweden

[13] Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Biskopsbogatan 27, S-431 80, Mölndal, Sweden

[14] UK Dementia Research Institute Fluid Biomarkers Laboratory, UK DRI at UCL, 3rd Floor, Queen Square House, Queen Square, London WC1N 3BG, United Kingdom

[15] Department of Neurodegenerative Disease, UCL Queen Square Institute of Neurology, Queen Square, London WC1N 3BG, United Kingdom

[16] Hong Kong Center for Neurodegenerative Diseases, Clear Water Bay, Units 1501-1502, 1512-1518, 15/F Building 17W, 17 Science Park W Ave, Science Park, Hong Kong, China

[17] Wisconsin Alzheimer's Disease Research Center, University of Wisconsin School of Medicine and Public Health, University of Wisconsin-Madison, 600 Highland Ave J5/1Mezzanine, Madison, WI, USA 53792

[18] The Foundation for the National Institutes of Health, 11400 Rockville Pike, Suite 600, North Bethesda, MD, USA 20852

[19] Department of Pathology and Laboratory Medicine, Perelman School of Medicine, University of Pennsylvania, 3400 Spruce St., Philadelphia, PA, USA 19104

[20] AbbVie, 1400 Sheridan Rd., North Chicago, IL, USA 60064

[21] Highly qualified expert

**Alzheimer's Disease Neuroimaging Initiative (ADNI): Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at:http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Correspondence to: Duygu Tosun, PhD

Full address: Department of Radiology and Biomedical Imaging, University of California San Francisco, San Francisco, CA, USA.

E-mail duygu.tosun@ucsf.edu

Running title: AD biomarkers by amyloid and tau clocks

Keywords: plasma biomarkers; clinical trials; longitudinal change; Estimated Years from Symptom Onset; amyloid time; tau time

Introduction

Alzheimer's disease (AD) is defined by the extracellular accumulation of amyloid plaques comprised mainly of amyloid-β (Aβ) peptide and the formation of intraneuronal neurofibrillary tangles comprised of hyperphosphorylated tau[1]. Amyloid plaques and tau tangles accumulate for many years in asymptomatic individuals during the preclinical phase of AD, but higher levels of pathology are often accompanied by neurodegeneration and inflammation culminating in the symptomatic phase which includes mild cognitive impairment (MCI) and dementia[2]. Biomarkers used to detect AD brain pathology in living individuals include Positron Emission Tomography (PET) scans that visualize the burden and distribution of amyloid and tau pathology, and measurement of proteins in the cerebrospinal fluid (CSF) or blood[3].

Blood biomarker (BBM) tests for AD are increasingly being used in research studies, clinical trials, and clinical practice[4]. Since blood collection is more feasible and accessible than PET or CSF, BBM tests are expected to expand access to AD biomarker testing, which may improve diagnostic accuracy for patients with cognitive impairment, increase access to AD-specific treatments, and accelerate the recruitment of individuals for clinical trials[5,6]. Available BBM tests include the ratio of Aβ42 to Aβ40 (Aβ42/Aβ40)[7], concentrations of tau phosphorylated at position 181 and 217 (p-tau181, p-tau217)[8], the ratio of p-tau217 to non-phosphorylated tau (%p-tau217)[9], and concentrations of neurofilament light (NfL) and glial fibrillary acidic protein (GFAP)[10,11]. BBM tests, including those for Aβ42/40 and p-tau217, start to change during the preclinical phase of AD and are used by some clinical prevention trials to identify individuals at risk of cognitive decline[12]. Additionally, some BBM levels change in response to treatments, suggesting a potential role in treatment monitoring[13,14].

Given the potential of BBMs to accelerate development of AD treatments, the Biomarker Consortium for the Foundation for the National Institutes of Health (FNIH), which includes stakeholders from academia, industry, and patient advocacy groups, coordinated measurements of plasma samples from the Alzheimer's Disease Neuroimaging Initiative (ADNI) with BBM tests from six different diagnostics companies. While the study focused on Aβ42/Aβ40, p-tau217, and p-tau181, panels from two companies additionally included GFAP and NfL. An initial cross-sectional study compared the classification accuracies and correlations of the BBM test measures with amyloid PET, tau PET, cortical thickness, and cognitive impairment[15]. A major additional aim of the FNIH project is to understand the timing

of plasma biomarker changes, including the number of years before symptom onset certain plasma biomarkers change, which may be helpful in identifying individuals who are likely to experience cognitive decline over the next several years.

The trajectories of plasma biomarkers may vary with disease stage, which might not be fully captured when assessed as a function of chronological age, given the variability in the ages at which individuals develop AD pathology and related symptoms. Estimating timelines based on disease pathophysiology and symptomatology may provide a more accurate understanding of biomarker dynamics throughout disease progression[16–18]. One approach is to use the consistent accumulation of amyloid pathology as measured by amyloid PET as a clock, which enables calculation of the age at the start of amyloid accumulation and estimation of the age at AD symptom onset for an individual[19,20]. Biomarker values can then be aligned by years from estimated symptom onset for each individual, enabling modeling of the years before symptom onset that different biomarkers change[18]. In this study, we created clocks using longitudinal amyloid PET and tau PET data, estimated the age at amyloid positivity and tau positivity, and estimated the age at AD symptom onset for individuals based on the age at tau positivity. The trajectories of plasma biomarkers, along with established AD biomarkers, were then evaluated as a function of estimated years from amyloid positivity, tau positivity, and AD symptom onset, and the relative timing of biomarker changes were determined. Notably, all datasets used by this study, including the estimated ages of positivity and symptom onset are available at adni.loni.usc.edu.

Materials and methods

1. Study participants

The current study utilized data from the ADNI database (adni.loni.usc.edu), which aims to develop and validate biomarkers for AD clinical trials[21,22]. ADNI was launched in 2003 as a public-private partnership led by Principal Investigator Michael W. Weiner, MD. The primary goal of ADNI has been to test whether serial magnetic resonance imaging (MRI), PET, other biological markers, and clinical and neuropsychological assessment can be combined to measure the progression of mild cognitive impairment (MCI) and early AD. Written informed

consent was obtained from each participant or their legally authorized representative. Race and sex were self-identified.

Amyloid and tau PET clock models were created using longitudinal [18F] florbetapir (FBP) amyloid PET (n=784) and flortaucipir (FTP) tau PET (n=359) data regardless of the availability of plasma biomarkers.

The timing of plasma biomarker change was modeled using data from the FNIH Biomarkers Consortium "trajectories" study (n=292) dataset, hereafter referred to as the "main study cohort". FNIH dataset included CU and CI ADNI participants who had plasma samples collected within six months of an amyloid PET scan for three distinct time points[15]. For individuals with more than three time points, plasma samples were selected that represented the earliest, latest, and an intermediate timepoint. For modeling the timing of change in established (non-plasma) AD biomarkers and the Clinical Dementia Rating Sum of Boxes (CDR-SB), ADNI data were not restricted to the FNIH dataset (n=757 for CDR-SB, n=743 for amyloid PET, n=627 for cortical thickness, n=569 for CSF p-tau181/Aβ42 and n=323 for tau PET). Positivity for either amyloid PET or tau PET at one timepoint was required for estimating age at amyloid and tau positivity, as explained below.

2. Plasma biomarker measurements

Plasma samples were collected, processed, and stored according to ADNI protocols[23]. As has previously been described, samples underwent analysis with C2N Diagnostics PrecivityAD2 (C2N), Fujirebio Diagnostics Lumipulse (Fujirebio), ALZpath Quanterix (ALZpath), Janssen LucentAD Quanterix (Janssen), and Roche Diagnostics NeuroToolKit (Roche) assays; a subset (84%) additionally underwent analysis with the Quanterix Neurology 4-Plex (Quanterix) assays[15].

3. CSF biomarker measurements and *APOE* genotyping

CSF Aβ42, total tau, and p-tau181 were measured using the Elecsys® β-amyloid (1–42), Elecsys Total-Tau, and Elecsys Phospho-Tau (181P) immunoassays, respectively, on a Cobas e 601 analyzer (Roche Diagnostics International Ltd, Rotkreuz, Switzerland). *APOE* genotyping was performed as part of the ADNI protocol.

4. Amyloid PET, tau PET, and MRI measures

Amyloid PET imaging was conducted at each ADNI site following standardized protocols for FBP[24]. A global standardized uptake value ratio (SUVR) was estimated across cortical summary regions, specifically the frontal, cingulate, parietal, and lateral temporal cortices, as defined by FreeSurfer v7.1. This ratio was normalized to a composite reference region as recommended in ADNI for longitudinal analyses. The composite reference region is a non-weighted average of whole cerebellum, brainstem/pons, and subcortical WM regions proposed by Koeppe[25]. Aβ PET positivity threshold was set at 0.78 SUVR for FBP, as specified for ADNI study, which corresponds roughly to 20 Centiloid units (ADNI_UCBerkeley_AmyloidPET_Methods_v2_2023-06-29.pdf).

Tau PET imaging was conducted at each ADNI site following standardized protocols for FTP. SUVRs were estimated for regions of interest (ROIs) as defined by FreeSurfer v7.1. A mesial-temporal meta-ROI was calculated that included regions with early tau PET signal, e.g., entorhinal, parahippocampus and amygdala. SUVRs were normalized to the inferior cerebellar grey matter reference region, as defined by the SUIT template[26]. A Gaussian mixture model with two components was applied to derive a threshold for early tau PET positivity using cross-sectional mesial-temporal meta-ROI data from the entire ADNI study (*n*=907), regardless of the availability of plasma biomarkers. The threshold was established at a value equal to the mean plus two standard deviations of the first component, which corresponds to the low tau burden cluster within the Gaussian mixture model. This process rendered a threshold of 1.41 SUVR for tau PET positivity (**Supplementary Fig. 1**).

Structural brain MRI data included a 3D MP-RAGE or IR-SPGR T1-weighted 3T MRI with sagittal slices. The full cortex of each subject brain was parcellated into 41 bilateral ROIs using a volumetric Desikan-Killiany-Tourville atlas, using FreeSurfer v5.1 for ADNI-GO/2 data and v6.0 for ADNI-3 data. A composite meta-ROI cortical thickness measurement was calculated that included the entorhinal, fusiform, parahippocampal, mid-temporal, inferior-temporal, and angular gyrus[27]. To address differences in image acquisition protocol and FreeSurfer version for image processing, the meta-ROI cortical thickness measure was harmonized using the ComBat-GAM method[28]. ComBAT-GAM harmonization and adjustment for normal age effects utilized all available ADNI data for unbiased modeling. Details on the harmonization of meta-ROI cortical thickness measure are provided in **Supplementary Methods**.

5. Neuropsychological and clinical variables

Participants underwent clinical assessments that included a detailed interview of a collateral source, a neurological examination of the participant, the Clinical Dementia Rating® (CDR®) and the CDR-SB[29]. Participants were classified by their clinical diagnosis as Cognitively Unimpaired (CU), Mild Cognitive Impairment (MCI) or Dementia (Dementia). For MCI or Dementia cases, clinicians assessed the potential etiology (AD or non-AD) based on clinical features[30].

6. Statistical analyses

6.1. Estimated years from amyloid and tau positivity

All available ADNI longitudinal amyloid PET and tau PET data, regardless of availability of plasma biomarker measures, were used to create amyloid and tau PET clocks by using a modified version of a previously reported approach[19]. Briefly, rates of change in amyloid (global) or tau (mesial-temporal) SUVR over time were estimated using linear mixed effect models with random intercepts and slopes. To capture non-linearity in the associations between pathology accumulation rates and pathology burden, we fitted generalized additive model (GAM) with a cubic spline to anchor rates of change as a function of the estimated pathology burden halfway through the follow-up period (midpoint SUVR).

To identify the interval over which rates of change are consistent, clock models were restricted to the SUVR range where the variance of the GAM fitted values fell below the 90th percentile of variance for all fitted values (**Supplementary Fig. 2**). The time between 0.0001 SUVR unit increases within the resulting SUVR range (0.62 to 1.11 SUVR for amyloid PET and 0.98 to 2.04 SUVR for tau PET), was calculated by integrating the inverse of modeled rates of change. In this study, amyloid time and tau time were conceptualized as the estimated time since amyloid or tau PET positivity. Therefore, amyloid or tau time were calculated by subtracting the cumulative estimated time at the positivity threshold (global cortical amyloid PET SUVR of 0.78 or mesial-temporal meta-ROI tau PET SUVR of 1.41) from the cumulative estimated time at each SUVR value. A more detailed explanation of the methods for amyloid and tau PET clock estimations is provided in **Supplementary Methods**.

Age at amyloid positivity or age at tau positivity, representing the estimated ages at which an individual first reached positivity, were estimated for those with at least one positive amyloid PET or tau PET scan available, respectively. These were calculated by subtracting the

amyloid or tau time corresponding to their positive scan from their age at the scan. For individuals with multiple scans, the age at amyloid or tau positivity was averaged across all positive scans.

Time estimates were validated in individuals who converted from a negative to a positive amyloid or tau PET scan throughout the ADNI study duration (PET converters). The actual conversion age was estimated by averaging the age at the last negative scan and at the first positive scan. The actual time since conversion in PET converters was calculated by subtracting the actual conversion age from the age at the scan. Correlations were examined for the estimated time and actual time since conversion, and for the estimated and actual time intervals between scans. Estimated ages at positivity were also validated in PET converters by testing their correlation with actual conversion age. Lastly, the estimated years from amyloid or tau positivity were calculated as the difference between the age at an outcome measurement and the age at amyloid or tau positivity, respectively.

6.2. Estimated years from symptom onset

We identified ADNI participants who progressed from CU at their first assessment to MCI or dementia due to AD by their last assessment and had available amyloid and tau PET data (clinical progressors). Individuals were not considered as clinical progressors if they were diagnosed with a non-AD etiology, classified as CU at their last assessment, or had a negative amyloid PET at the time of clinical progression. Symptom onset age was defined as the age at the first assessment with a CDR>0 and a clinical diagnosis of MCI or dementia due to AD.

Linear regression was used to model age at symptom onset as a function of age at amyloid positivity or age at tau positivity. We also evaluated for potential effects of *APOE* ε4 carrier status and sex. To validate the model, we evaluated the correlation of the actual symptom onset age with the estimated symptom onset age in clinical progressors. Because age at tau positivity was a stronger predictor of symptom onset age, it was included as only predictor in the final model for symptom onset age. This model was applied to all participants with an estimated age at tau positivity. For subsequent analyses, the estimated years from symptom onset was calculated as the age at the outcome measure minus the estimated age at symptom onset.

6.3. Timing of change in outcome measures relative to estimated years from amyloid positivity, tau positivity and symptom onset

Outcome measures were modeled as a function of each estimated timeline (estimated years from amyloid positivity, tau positivity, and symptom onset). Outcome measures examined included plasma biomarkers, CSF p-tau181/Aβ42, amyloid PET (global cortical SUVR), tau PET (mesial-temporal meta-ROI SUVR), cortical thickness meta-ROI, and CDR-SB. Each outcome was modeled using generalized additive mixture models (GAMM) with cubic spline basis and random intercepts as a function of each estimated timeline, separately. The time of abnormality for each outcome measure in relation to each estimated timeline was set at the first timepoint where the confidence intervals of the model estimates did not overlap with the confidence interval of the overall mean value of a reference group of amyloid and tau PET-negative CU (CDR=0 and clinical diagnosis of CU) individuals.

The median and 95% confidence intervals of the times of abnormality were calculated using 1000 bootstrapped samples, based on the 2.5% and 97.5% thresholds. To determine the periods of significant rate of change for each outcome measure, we identified time segments where the first derivative of the function deviated significantly from zero. Approximate derivatives were found using finite differencing with Taylor series expansions.

Within the reference group of amyloid and tau PET-negative CU participants, the association between each outcome measure and age, sex and *APOE* ε4 carrier status was assessed separately using linear regression model fits. For those outcome measures that showed significant associations with age, sex or *APOE* ε4 carrier status, their raw values in the entire sample were adjusted based on the derived effects in the reference group, and the timing of changes was analyzed using the corrected values as a sensitivity analysis.

As further sensitivity analyses, the timing of changes in plasma biomarkers was analyzed restricting the main study cohort to the subset of individuals with available plasma biomarker measures across all assays (*n*=233).

All statistical analyses and figures were created using R software (R version 4.2.2).

Data availability

Data from this study and the study methodology report may be accessed from the ADNI Laboratory of Neuro Imaging (LONI) database: adni.loni.usc.edu.

Results

1. Characteristics of the study cohorts

Data from 784 ADNI participants with longitudinal FBP PET was used for amyloid PET clock modeling (mean follow-up time of 4.7 years), and data from 359 ADNI participants with available longitudinal FTP PET (mean follow-up time of 3.0 years) was used for tau PET clock modeling (**Supplementary Table 1**).

Of the 406 participants in the main study cohort, who underwent plasma biomarker measurements as part of the FNIH project, 190 met the inclusion criteria for estimating age at amyloid positivity (i.e., had at least one positive amyloid PET scan during their participation in the ADNI study), and 70 participants satisfied the inclusion criteria for estimating age at tau positivity (i.e., had at least one positive tau PET scan during their participation in the ADNI study). 48 participants overlapped between the two cohorts. The reference group included 80 individuals who were amyloid and tau PET-negative and CU (CDR=0 and clinical diagnosis of CU) throughout the study duration and was used to establish normative distributions for AD biomarkers and cognitive measures.

The characteristics of individuals in the main study cohort with an estimated age at amyloid positivity, tau positivity, and the reference group are described at the baseline plasma sample collection (**Table 1**). In the combined cohort, mean age was 73.9 ± 7.0 years, 147 (50.3%) were women, 121 (41.4%) were *APOE* ε4 carriers, 276 (94.5%) self-identified as white, 112 (38.4%) were cognitively impaired (CDR>0) and 130 (44.5%) were amyloid PET-positive at baseline. When compared to the reference group, individuals with an estimated age at tau positivity were significantly older, while the age difference was not significant in the cohort with an estimated age at amyloid positivity. Individuals with an estimated age at amyloid or at tau positivity were more likely to carry an *APOE* ε4 allele, and those with an estimated age at amyloid positivity had significantly lower education levels than the reference group. All AD biomarkers were significantly elevated at baseline in both cohorts compared to the reference group. Notably, only 8 participants in the cohort with an estimated age at amyloid positivity and 3 participants in the cohort with an estimated age at tau positivity had a clinical diagnosis of dementia at baseline.

Baseline characteristics of individuals included in the analyses for established (non-plasma) AD biomarkers and CDR-SB in the cohorts with an estimated age at amyloid (n=635)

and tau (n=212) positivity and in the reference group (n=263), are described in **Supplementary Table 2**. Of note, in this larger sample, individuals in both cohorts were significantly older than those in the reference group.

Finally, among all individuals with both an estimated age at amyloid and tau positivity, 17 progressed from CU to MCI or dementia due to AD during their participation in the ADNI study and were included in the model used to estimate symptom onset age.

2. Creation of clocks with longitudinal ADNI amyloid PET and tau PET data

Clocks relating amyloid or tau PET SUVR to time were created and validated within the full ADNI study dataset before being applied to the main study cohort. A detailed description of the method and model validations can be found in **Supplementary Methods**.

The estimated ages at amyloid and tau positivity were highly correlated with the actual age at conversion among individuals who converted from a negative to a positive PET scan. For amyloid PET converters (n=81), the relationship was: 'age at amyloid positivity =1.01x (conversion age) – 2.39 years ± 2.47 years'; Intercept P=0.34; Spearman's ρ=0.96, $P<0.0001$. For tau PET converters (n=35), the relationship was: 'age at tau positivity =1.07 x (conversion age) – 6.7 years ± 3.33 years'; Intercept P =0.05; Spearman's ρ=0.96, $P<0.0001$; **Supplementary Fig. 3**).

In the main study cohort, the estimated ages at amyloid positivity, tau positivity, and symptom onset were 66.6 ± 10.2, 73.7 ± 9.5, and 81.0 ± 6.1 years (mean ± standard deviation), respectively.

The observed amyloid PET SUVR or tau PET SUVR values by age are displayed in **Fig. 1A** and **Fig. 1C**, respectively, demonstrating wide variability in the ages at which participants start to accumulate amyloid and tau pathology. **Fig. 1B** and **Fig. 1D** show the modeled biomarker patterns as a function of estimated time from amyloid positivity and tau positivity, respectively, and demonstrate the relatively consistent accumulation of these pathologies over time.

3. Estimation of age at symptom onset

Using the full ADNI dataset, the age at amyloid positivity for the clinical progressors (*n*=39) was predictive of the age at symptom onset (Spearman's ρ=0.63; Adj R^2=0.38; *P*<0.0001). However, the age at tau positivity was a much stronger predictor of the age at symptom onset (*n*=17; Spearman's ρ=0.93; Adj R^2=0.86; *P* < 0.0001; **Supplementary Fig. 4**). Importantly, neither *APOE ε4* status nor sex significantly influenced the estimated age at symptom onset in these models. Therefore, age at tau positivity was included as the only predictor in the final model used to estimate symptom onset. A detailed description of the method and model validations can be found in **Supplementary Methods.**

4. AD biomarker trajectories as a function of years from amyloid positivity, tau positivity and symptom onset

Trajectories of AD plasma biomarkers, established AD biomarkers, and CDR-SB are shown as a function of the three estimated AD progression timelines in **Fig. 2** and **Fig. 3.** Plasma biomarkers previously reported to be most strongly associated with amyloid PET and tau PET are shown in **Fig. 2**[15]; trajectories for the remaining plasma biomarkers are provided in **Supplementary Fig. 5** and **Supplementary Fig. 6**. Trajectories of all outcome measures as a function of age, amyloid PET, and tau PET are depicted in **Supplementary Figs. 7-9**.

5. Timing of AD biomarker changes relative to estimated years from amyloid positivity

We modeled the longitudinal trajectories of all outcome measures as a function of estimated years from amyloid positivity and estimated the time when each measure differed significantly from the reference group mean.

All plasma biomarkers, except for NfL, reached abnormal levels before the estimated time of amyloid positivity, with plasma Aβ42/Aβ40 measures becoming abnormal first **(Fig. 4A, Supplementary Table 3)**. Specifically, C2N Aβ42/Aβ40 and Roche Aβ42/Aβ40 showed abnormal levels at a median of -7.9 years before amyloid positivity, earlier than amyloid PET (-5.9 years) or CSF p-tau181/Aβ42 (-5.2 years). Notably, Fujirebio and Quanterix Aβ42/Aβ40 measures became abnormal considerably later (-5.8 and -3.4 years, respectively). C2N %p-tau217 showed abnormality -4.4 years before amyloid positivity and was closely followed by

plasma GFAP measures, Janssen and ALZpath p-tau217, Roche p-tau181 (all at median times within -3.9 to -2.9 years prior to amyloid positivity). Fujirebio p-tau217 showed abnormal levels at -1.5 years, and Quanterix p-tau181 and C2N p-tau217 did not reach abnormality until around the estimated time of amyloid positivity, along with tau PET and CDR-SB (all from -0.9 to 0.0 years before amyloid positivity). Cortical thickness reached abnormality 2.9 years after amyloid positivity, and plasma NfL was the last to show significant elevations, 9.2 or 10.5 years after amyloid positivity, for Roche or Quanterix assays, respectively.

Next, we assessed the rates of change in plasma biomarkers relative to the amyloid timeline. In **Fig.2** and **Fig.3**, thicker segments of the fitted lines indicate time periods where significant slopes were observed. All measures of p-tau217, p-tau181, GFAP and NfL consistently increased during the entire amyloid timeline ranging from -7.9 to -6.8 years before to 30.3 years after the estimated time of amyloid positivity. In contrast, plasma Aβ42/Aβ40 measures decreased early in the timeline (-7.9 to -7.1 years) but plateaued at 11.5 to 13.0 years after amyloid positivity, depending on the assay used (**Fig.2A, Supplementary Fig. 5A and 6A, Supplementary Table 4**). CSF p-tau181/Aβ42, amyloid PET, tau PET, cortical thickness and CDR-SB exhibited significant change throughout the entire amyloid timeline (**Fig. 3A, Supplementary Table 4**).

6. Timing of AD biomarker changes relative to estimated years from tau positivity

As a function of the tau timeline, plasma GFAP measures were the earliest to differ significantly from the reference group (up to a median of -11.5 years before estimated time of tau positivity), and were followed by C2N, Fujirebio and Roche Aβ42/Aβ40 measures (ranging from -9.3 to -7.1 years) and p-tau181 (-6.7 or -5.4 years for Roche and Quanterix, respectively). Amyloid PET and CSF p-tau181/Aβ42 also became abnormal within this time range (-7.8 and -7.4 years prior to tau positivity, respectively). Notably, Quanterix Aβ42/Aβ40 changed later than the rest of plasma Aβ42/Aβ40 measures, -3.3 years before tau positivity. Plasma p-tau217 measures became abnormal ranging from -5.4 to -3.9 years before tau positivity, and tau PET abnormalities were detected at -4.7 years. These were followed by abnormalities in cortical thickness (-2.8 years) and eventually plasma NfL (5.3 or 15.3 years after tau positivity for Roche and Quanterix, respectively). Of note, CDR-SB showed

abnormality at median time of -5.7 years before tau positivity (**Fig. 4B, Supplementary Table 3**).

The temporal dynamics of plasma biomarkers across the tau timeline varied depending on the assay. C2N and Roche Aβ42/Aβ40 decreased significantly during the entire tau timeline (-11.4 years before to 17.9 years after tau positivity). In contrast, Quanterix Aβ42/Aβ40 decreased from -11.4 but then plateaued 6.0 years after tau positivity, while Fujirebio Aβ42/Aβ40 decreased significantly only from -0.8 to 4.4 years. All p-tau217 measures increased significantly throughout the entire tau timeline except for C2N p-tau217, which started to significantly rise one year before tau positivity. The increases in Roche GFAP and in Roche p-tau181 were consistent the entire tau timeline, whereas Quanterix p-tau181 rise was significant until 3.6 years after tau positivity. In contrast, Quanterix plasma GFAP and NfL measures did not significantly change at any point in the tau timeline. (**Fig. 2B, Supplementary Fig. 5B and 6B, Supplementary Table 4**).

CSF p-tau181/Aβ42, amyloid PET and tau PET consistently increased during the entire tau timeline (from -12.2 or -11.9 to 19.8 or 19.4 years for CSF p-tau181/Aβ42 and amyloid PET, respectively, and from -4.7 to 21.8 for tau PET). In contrast, CDR-SB started to change at -6.3 years prior to tau positivity. Finally, cortical thickness started to decrease shortly after tau positivity (0.5 years) and persisting until the timeline upper limit (18.1 years; **Fig. 3B, Supplementary Table 4**)

7. Timing of AD biomarker changes relative to estimated years from symptom onset

Roche, C2N, and Fujirebio plasma Aβ42/Aβ40 were the earliest plasma biomarkers to become abnormal in the symptom timeline, concurrent with amyloid PET and CSF p-tau181/Aβ42 (up to -14.4 years before symptom onset). In contrast, Quanterix plasma Aβ42/Aβ40 did not significantly differ from the reference group until -8.4 years prior to symptom onset. Tau PET, plasma GFAP, p-tau217 and p-tau181 measures all reached abnormality ranging from -11.9 to -9.4 years from symptom onset. These were followed by abnormalities in CDR-SB (-6.2 years), and lastly, plasma NfL and cortical thickness (ranging from -4.3 to -2.1 years before symptom onset) (**Fig. 4C, Supplementary Table 3**).

Importantly, plasma Aβ42/Aβ40 measures did not have a significant rate of change after the estimated symptom onset. Fujirebio and Quanterix plasma Aβ42/Aβ40 significantly decreased from the lower limit of the symptom timeline (-16.7 years) but stabilized -3.9 and -0.8 years prior to symptom onset, respectively, while changes in C2N plasma Aβ42/Aβ40 were significant only in the period between -11.1 to -2.2 years. The rate of change in Roche plasma Aβ42/Aβ40 was not significant at any point as a function of the symptom timeline.

In contrast, all plasma p-tau217, as well as Quanterix p-tau181, GFAP and NfL measures steadily increased across the entire symptom timeline (-16.7 to 7.3 years from symptom onset). The rise in Roche plasma p-tau181 was significant from the timeline lower limit but stabilized 2.2 years after symptom onset (**Fig. 2C, Supplementary Fig. 5C; Supplementary Fig. 6C Supplementary Table 4**).

CSF p-tau181/Aβ42, amyloid PET, and tau PET changed consistently during the entire symptom timeline (-18.7 to 7.8 years for CSF p-tau181/Aβ42, -19.5 to 9.6 years for amyloid PET, and -13.7 to 9.8 years for tau PET). CDR-SB started to change significantly at -8.6 years and persisted throughout the timeline (10.6 years after symptom onset). Finally, the rate of change in cortical thickness became significant shortly after (-6.2 years), also persisting across the symptom timeline (9.6 years after symptom onset) (**Fig. 3C, Supplementary Table 4**).

8. Sensitivity analyses

Sex or *APOE* ε4 carrier status did not have a significant effect in any of the outcome measures in the reference group (all $P > 0.05$). In contrast, tau PET and all p-tau181, GFAP, NfL, and p-tau217 measures, except for C2N p-tau217, were significantly associated with age. For these biomarkers, the same analyses were performed using age-corrected values. The ordering of plasma biomarkers relative to the different timelines was very similar than that of non-corrected data (**Supplementary Table 5**). Most notable differences were for plasma NfL, which became abnormal significantly earlier after age adjustment (0.1 or 4.1 years in the amyloid timeline, -5.4 or -3.3 years in the tau timeline, and at -9.6 or -8.7 in the symptom timeline for Quanterix and Roche measures, respectively).

Analyses on the dynamics of age-adjusted plasma biomarkers across the three timelines rendered very similar results to those of non-corrected biomarkers (**Supplementary Table 6**). Notably, Fujirebio plasma p-tau217 did not significantly rise until -3.0 years before tau

positivity, and the rate of change in Quanterix plasma p-tau181 did not reach statistical significance at any point in the tau timeline after age adjustment.

Finally, the timing of plasma biomarker changes was assessed in the subset of individuals with available plasma biomarkers across all assays (n=149 with an age at amyloid positivity, n=49 with an age at tau positivity, n=66 in the reference group), and results were very similar to those in the full main study cohort (**Supplementary Table 7 and Supplementary Table 8**).

Discussion

The aim of this study was to evaluate the timing of plasma biomarker changes relative to AD progression timelines based on amyloid and tau PET clocks. We estimated the age at amyloid positivity and tau positivity for ADNI participants using their longitudinal PET imaging data and found that age at tau positivity was a stronger predictor of the age at symptom onset than the age at amyloid positivity. Our main findings were 1) all plasma biomarkers except NfL exhibited changes prior to the estimated time of amyloid and tau positivity, with plasma Aβ42/Aβ40 and GFAP showing the earliest alterations; 2) plasma p-tau217, p-tau181, GFAP and NfL increased consistently over the estimated AD progression timelines, while plasma Aβ42/Aβ40 only changed early in the disease; 3) the timing of plasma biomarker changes varied depending on the AD progression timeline and the assay used. These findings contribute to improving our understanding of AD pathophysiology and the dynamics of key plasma biomarkers throughout AD progression, which is crucial for optimizing their use in clinical trials and clinical practice.

Plasma Aβ42/Aβ40 biomarkers reached abnormality as early as 7.9 years before amyloid positivity, 9.3 years before tau positivity and 14.4 years before symptom onset. These were earlier or very close times to those when CSF p-tau181/Aβ42 and amyloid PET became abnormal. Our findings are consistent with studies showing changes in plasma Aβ42/Aβ40 prior to detectable amyloid accumulation in PET[31–34], and with plasma biomarkers reflecting soluble amyloid forms prior to fibrillar amyloid detectable with amyloid PET. Plasma GFAP changed shortly after Aβ42/Aβ40 in the amyloid and symptom timelines, concurrently with p-tau217 and p-tau181 measures. Interestingly, plasma GFAP showed the earliest median time of abnormality only in the tau timeline, although it was not significantly earlier than most

plasma Aβ42/Aβ40 measures, as indicated by the overlapping confidence intervals. GFAP in plasma is reported to increase as an early response to amyloid accumulation[11,35], which may explain why it reaches abnormality close to time of amyloid biomarkers. In addition, astrocyte reactivity has been shown to modulate the association of amyloid with early tau pathology[36]. As the years from tau positivity were estimated based on a mesial-temporal meta-ROI, which included areas of early tau accumulation, the early increase of plasma GFAP in the tau timeline could potentially reflect astrocyte reactivity in individuals with amyloid pathology and within the tau accumulation pathway. Plasma p-tau181 measures also changed earlier in the tau timeline, at similar times to Aβ42/Aβ40 and earlier than p-tau217. These differences, although subtle, might be reflecting disparities between p-tau isotopes and the pathophysiology they potentially reflect[37]. Of interest, tau PET became abnormal close to the time of amyloid positivity, shortly before the threshold used for tau positivity, but more than 11 years prior to symptom onset. This demonstrates the strong association of tau accumulation with AD symptoms. Finally, cortical thickness and plasma NfL were the last to change in the three timelines, as expected for markers indicating downstream neurodegeneration. Overall, the ordering of biomarker changes was in concordance with the $A/T_1/T_2/(N)$ conceptualization, in terms of amyloid biomarkers changing first and being followed by biofluid mid-region tau fragments that possibly reflect a reaction to amyloid, and later on by tau PET and neurodegeneration markers[1].

When we studied the rates of change in plasma biomarkers throughout progression timelines, plasma Aβ42/Aβ40 decrease was subtle and plateaued at 11.5 to 13.0 years after amyloid positivity. In contrast, most p-tau biomarkers and, to a lesser extent, GFAP and NfL measures, steadily increased across AD progression similarly to established AD biomarkers, reinforcing their potential for monitoring disease progression. Plasma NfL did not significantly change in the tau timeline. As a non-specific neurodegeneration biomarker, its increase may be more evident at later disease stages. Also, multiple factors, including comorbid pathologies, may contribute to plasma NfL levels and longitudinal trajectories[38–40].

Generally, results in the timing of plasma biomarker changes were consistent across assays, but some differences were observed. Quanterix Aβ42/Aβ40 reached abnormality considerably later than the rest of Aβ42/Aβ40 assays across timelines. This finding aligns with and might explain the lower association with amyloid PET observed in our initial cross-

sectional study[15]. In addition, the temporal dynamics of plasma biomarkers varied by assay, particularly for Aβ42/Aβ40 in the tau and symptom timelines. These observations may reflect variability in the assay measures and in their sensitivity to subtle changes in plasma protein levels (*e.g.,* mass-spectrometry-based or immunoassays)[7,15,41]. Differences across assays, along with differences in biomarker dynamics, are all important factors to be considered when selecting them for specific uses in trials and clinical practice.

Several previous studies have evaluated AD biomarker trajectories, including plasma biomarkers, using amyloid PET[16,42,43] or other models[31,44,45] as proxies of disease progression. However, to the best of our knowledge, only one used a time scale derived from amyloid PET[18]. Of note, the recent study by Li et al.[18] used an earlier threshold for amyloid positivity, corresponding to the "tipping point" after which individuals accumulated amyloid at a relatively consistent rate (referred to as amyloid onset). Herein, amyloid or tau time were conceptualized as time since a positive scan, as the positivity threshold is consistently used in clinical trials and research studies to define meaningful amyloid accumulation, and it is more directly related to clinically relevant outcomes. Nevertheless, the ordering of biomarker abnormality throughout AD progression was similar in both studies. The current study validated the amyloid PET clock approach in a larger multicentric sample with extensive longitudinal data from individuals at different disease stages and expanded the results with the inclusion of tau PET imaging in addition to key plasma biomarkers for AD measured with different assays, enabling cross-assay comparisons and adding robustness to our findings.

Importantly, we found that the estimated age at tau positivity based on a tau PET clock was a much better predictor of symptom onset than the age at amyloid positivity ($R^2 = 0.86$ *vs.* $R^2 = 0.38$). This result was not unexpected given the extensive evidence supporting that tau is more strongly associated with cognitive decline than amyloid alone[46–49]. However, previous studies had found amyloid onset age to be highly predictive of symptom onset[19,20]. Several factors might be accountable for this difference. Given that ADNI is a multicenter study, there is inherent heterogeneity in clinical diagnosis practices across different sites. Additionally, herein we included participants at all stages of the AD continuum. Still, the predictive value of age at amyloid positivity in our model ($R^2 = 0.38$) was very close to that of the parental age at symptom onset shown in autosomal-dominant Alzheimer's disease (ADAD) studies (R^2 ranging from 0.38 – to 0.56)[50,51]. This is noteworthy given that, unlike the expected pure AD

pathology in ADAD cases, many factors including comorbid neuropathologies significantly contribute to variability in cognitive symptoms in sporadic AD.

This study has several limitations. First, amyloid PET in ADNI is measured with FBP, which potentially influenced the accuracy of our amyloid time estimates compared to previous studies using PIB[19,52]. Second, although the effect of the main AD risk factors such as *APOE* ε4 status or sex on the age at symptom onset was evaluated, there are multiple other variables that could influence the age at symptom onset and were not examined herein (e.g. comorbidities, social determinants of health, genetic variants). Other limitations include a lower sample size for the plasma biomarker analyses compared to the non-plasma biomarkers and cognitive measures, making effects of comparable sizes less likely to be observed among plasma biomarkers. Furthermore, there is a low frequency of dementia cases in our sample, precluding the study of biomarker trajectories in advanced disease stages. Finally, although efforts are being made to increase diversity in ADNI[53], the present study focused on retrospective data and most participants were white and highly educated, underscoring the need for future studies in more diverse populations.

In summary, we evaluated timing of change in key AD plasma biomarkers across AD progression timelines using amyloid and tau PET clocks, the latter being highly predictive of symptom onset. Plasma Aβ42/Aβ40 was the earliest to reach abnormal levels consistently across the estimated AD progression timelines, but its dynamic range was limited. In contrast, p-tau217, p-tau181, GFAP and NfL showed utility for disease staging or monitoring. Additionally, our results indicate that plasma biomarker dynamics can vary depending on the timeline and the assay used, which should be considered for their use and interpretation. Importantly, subsequent studies will address the associations of rates of change in plasma biomarkers across AD progression and changes in neurodegeneration and cognitive outcomes, as well as their consistency across assays. Overall, the results of the current study contribute to optimizing the use of plasma biomarkers in clinical trials and clinical practice.

Funding

The results of the study represent results of the Foundation for the National Institutes of Health (FNIH) Biomarkers Consortium "Biomarkers Consortium, Plasma Aβ and Phosphorylated Tau as Predictors of Amyloid and Tau Positivity in Alzheimer's Disease" project. The study was made possible through the scientific and financial support of industry, academic, patient advocacy, and governmental partners. We are grateful for the contributions of the following project team members: Anthony Bannon (AbbVie), Michael Baratta (Takeda), Janaky Coomaraswamy (Takeda), Jeff Dage (Indiana University), Iwona Dobler (Takeda), Lei Du-Cuny (AbbVie), Kyle Ferber (Biogen), John Hsiao (NIA), Hartmuth Kolb (formerly with Johnson and Johnson Innovative Medicine), Emily Meyers (Alzheimer's Association), Yulia Mordashova (AbbVie), William Potter, Maria Quinton (AbbVie), Dave Raunig (Takeda), Erin Rosenbaugh (FNIH), Carrie Rubel (Biogen), Ziad Saad (Johnson and Johnson Innovative Medicine), Patricia Saletti (Alzheimer's Drug Discovery Foundation), Suzanne Schindler (Washington University in St. Louis), Leslie Shaw (University of Pennsylvania), Gallen Triana-Baltzer (Johnson and Johnson Innovative Medicine), Christopher Weber (Alzheimer's Association), Henrik Zetterberg (University of Gothenburg). Funding partners of the project include AbbVie Inc., Alzheimer's Association®, Diagnostics Accelerator at the Alzheimer's Drug Discovery Foundation, Biogen, Janssen Research & Development, LLC, and Takeda Pharmaceutical Company Limited. Private-sector funding for the study was managed by the Foundation for the National Institutes of Health.

We recognize C2N Diagnostics, Fujirebio Diagnostics with the Indiana University National Centralized Repository for Alzheimer's Disease and Related Dementias Biomarker Assay Laboratory (NCRAD-BAL), Quanterix, and Roche Diagnostics with the University of Gothenburg for performing the plasma biomarker analysis in this study. The NCRAD-BAL is supported by a cooperative agreement grant (U24 AG021886) awarded to NCRAD by the National Institute on Aging. Elecsys β-amyloid (1–42) CSF, Elecsys Phospho-Tau (181P) CSF and Elecsys Total-Tau CSF assays are approved for clinical use. COBAS and ELECSYS are trademarks of Roche. All other product names and trademarks are the property of their respective owners. The NeuroToolKit is a panel of exploratory prototype assays designed to robustly evaluate biomarkers associated with key pathologic events characteristic of AD and other neurological disorders, used for research purposes only and not approved for clinical use (Roche Diagnostics International Ltd, Rotkreuz, Switzerland).

Finally, we would like to acknowledge ADNI for the plasma samples and ADNI participant data analyzed in this study. Data collection and sharing for the Alzheimer's Disease Neuroimaging Initiative (ADNI) is funded by the National Institute on Aging (National Institutes of Health Grant U19 AG024904). The grantee organization is the Northern California Institute for Research and Education. In the past, ADNI has also received funding from the National Institute of Biomedical Imaging and Bioengineering, the Canadian Institutes of Health Research, and private sector contributions through the Foundation for the National Institutes of Health (FNIH) including generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company Limited; and Transition Therapeutics.

M.M.A receives funding from the Alzheimer's Association Research Fellowship grant program (AARF-23-1141384). Statistical analyses were supported in part by National Institute on Aging grant R01AG070941 (S.E.S.).

The Biomarkers Consortium, Plasma Aβ and Phosphorylated Tau as Predictors of Amyloid and Tau Positivity in Alzheimer's Disease Project was made possible through a public-private partnership managed by the Foundation for the National Institute of Health (FNIH) and funded by AbbVie Inc., Alzheimer's Association®, Diagnostics Accelerator at the Alzheimer's Drug Discovery Foundation, Biogen, Janssen Research & Development, LLC, and Takeda Pharmaceutical Company Limited.

ADNI Collaborators

Data used in the preparation of this article were obtained from the ADNI database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in the analysis or writing

of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Competing interests

Suzanne E. Schindler (S.E.S.) has no financial interests in any pharmaceutical companies and has not received any direct research funding from any pharmaceutical companies. S.E.S. has served on scientific advisory boards on biomarker testing and education for Eisai and Novo Nordisk and has received speaking fees for presentations on biomarker testing from Eisai, Eli Lilly, and Novo Nordisk. Henrik Zetterberg (H.K.) has no financial interests in any pharmaceutical companies and has not received any direct funding from any pharmaceutical companies. H.K. has served on scientific advisory boards and/or as a consultant for Abbvie, Acumen, Alector, Alzinova, ALZpath, Amylyx, Annexon, Apellis, Artery Therapeutics, AZTherapies, Cognito Therapeutics, CogRx, Denali, Eisai, LabCorp, Merry Life, Nervgen, Novo Nordisk, Optoceutics, Passage Bio, Pinteon Therapeutics, Prothena, Red Abbey Labs, reMYND, Roche, Samumed, Siemens Healthineers, Triplet Therapeutics, and Wave, has given lectures sponsored by Alzecure, BioArctic, Biogen, Cellectricon, Fujirebio, Lilly, Novo Nordisk, Roche, and WebMD, and is a co-founder of Brain Biomarker Solutions in Gothenburg AB (BBS), which is a part of the GU Ventures Incubator Program (outside submitted work). Janaky Coomaraswamy (J.C.), Michael Baratta (M.B.), and David L. Raunig (D.L.R.) receive salary and company stock as compensation for their employment with Takeda Pharmaceutical Company Limited. Marta Milà-Alomà (M.M.A.), Kellen K. Petersen (K.K.P.), Benjamin Saef (B.S.), Duygu Tosun (D.T.), Zachary Hausle (Z.H.), and Erin G. Rosenbaugh (E.G.R.) have nothing to disclose. Leslie M. Shaw (L.M.S.) receives funding from the NIA for ADNI4 and from NIA for the University of Pennsylvania ADRC P30 for the Biomarker Core. Jeffrey L. Dage (J.L.D.) is an inventor on patents or patent applications of Eli Lilly and Company relating to the assays, methods, reagents, and/or compositions of matter for P-tau assays and Aβ targeting therapeutics. J.L.D. has served as a consultant or on advisory boards for Eisai, Abbvie, Genotix Biotechnologies Inc, Gates Ventures, Karuna Therapeutics, AlzPath Inc., Cognito Therapeutics, Inc., and received research support from ADx Neurosciences, Fujirebio, AlzPath Inc., Roche Diagnostics and Eli Lilly and Company in the past 2 years. J.L.D. has received speaker fees from Eli Lilly and Company. J.L.D. is a founder and advisor for Monument Biosciences. J.L.D. has stock or stock options in Eli Lilly and Company, Genotix

Biotechnologies, AlzPath Inc. and Monument Biosciences. Kyle Ferber (K.F.) and Carrie E. Rubel (C.E.R.) are employees of and may own stock in Biogen. Gallen Triana-Baltzer (G.T.B.) and Ziad S. Saad (Z.S.S.) are employed by Johnson & Johnson Innovative Medicine and may receive salary and stock for their employment. Lei Du-Cuny (L.D.C.) and Yulia Mordashova (Y.M.) are employed by AbbVie Deutschland GmbH & Co. Yan Li (Y.L.) is the co-inventor of the technology "Novel Tau isoforms to predict onset of symptoms and dementia in Alzheimer's disease" which is in the process of licensing by C2N. Nicholas J. Ashton (N.J.A) has received speaking fees from Eli Lilly, Biogen, Quanterix and Alamar Biosciences. Emily A. Meyers (E.A.M.) is employed by the Alzheimer's Association. Anthony W. Bannon (A.W.B.) receives salary and company stock as compensation for his employment with AbbVie Inc. William Z. Potter (W.Z.P.) was previously employed by the National Institute of Mental Health, and he is a stockholder in Merck & Co., Inc. W.Z.P. is a Co-Chair Emeritus for the FNIH Biomarkers Consortium Neuroscience Steering Committee. Currently residing in Philadelphia, PA, W.Z.P. serves as a consultant for Karuna, Neurocrine, Neumarker, Vaaji, and receives grant support from the NIA along with stock options from Praxis Bioresearch.

References

1. Jack CR, Andrews JS, Beach TG, et al. Revised criteria for diagnosis and staging of Alzheimer's disease: Alzheimer's Association Workgroup. *Alzheimers Dement*. Published online June 27, 2024. doi:10.1002/alz.13859

2. McDade EM. Alzheimer Disease. *Continuum (Minneap Minn)*. 2022;28(3):648-675. doi:10.1212/CON.0000000000001131

3. Schindler SE, Atri A. The role of cerebrospinal fluid and other biomarker modalities in the Alzheimer's disease diagnostic revolution. *Nat Aging*. 2023;3(5):460-462. doi:10.1038/s43587-023-00400-6

4. Hansson O, Blennow K, Zetterberg H, Dage J. Blood biomarkers for Alzheimer's disease in clinical practice and trials. *Nat Aging*. 2023;3(5):506-519. doi:10.1038/s43587-023-00403-3

5. Schindler SE, Li Y, Li M, et al. Using Alzheimer's disease blood tests to accelerate clinical trial enrollment. *Alzheimers Dement*. 2023;19(4):1175-1183. doi:10.1002/alz.12754

6. Palmqvist S, Tideman P, Mattsson-Carlgren N, et al. Blood Biomarkers to Detect Alzheimer Disease in Primary Care and Secondary Care. *JAMA*. Published online July 28, 2024:e2413855. doi:10.1001/jama.2024.13855

7. Zicha S, Bateman RJ, Shaw LM, et al. Comparative analytical performance of multiple plasma Aβ42 and Aβ40 assays and their ability to predict positron emission tomography amyloid positivity. *Alzheimers Dement*. 2023;19(3):956-966. doi:10.1002/alz.12697

8. Janelidze S, Bali D, Ashton NJ, et al. Head-to-head comparison of 10 plasma phospho-tau assays in prodromal Alzheimer's disease. *Brain*. 2023;146(4):1592-1601. doi:10.1093/brain/awac333

9. Meyer MR, Kirmess KM, Eastwood S, et al. Clinical validation of the PrecivityAD2 blood test: A mass spectrometry-based test with algorithm combining %p-tau217 and Aβ42/40 ratio to identify presence of brain amyloid. *Alzheimers Dement*. 2024;20(5):3179-3192. doi:10.1002/alz.13764

10. Ashton NJ, Janelidze S, Al Khleifat A, et al. A multicentre validation study of the diagnostic value of plasma neurofilament light. *Nature communications*. 2021;12(1):3400. doi:10.1038/s41467-021-23620-z

11. Benedet AL, Milà-Alomà M, Vrillon A, et al. Differences Between Plasma and Cerebrospinal Fluid Glial Fibrillary Acidic Protein Levels Across the Alzheimer Disease Continuum. *JAMA Neurology*. 2021;78(12):1471. doi:10.1001/jamaneurol.2021.3671

12. Rissman RA, Langford O, Raman R, et al. Plasma Aβ42/Aβ40 and phospho-tau217 concentration ratios increase the accuracy of amyloid PET classification in preclinical Alzheimer's disease. *Alzheimers Dement*. 2024;20(2):1214-1224. doi:10.1002/alz.13542

13. Pontecorvo MJ, Lu M, Burnham SC, et al. Association of Donanemab Treatment With Exploratory Plasma Biomarkers in Early Symptomatic Alzheimer Disease: A Secondary

Analysis of the TRAILBLAZER-ALZ Randomized Clinical Trial. *JAMA Neurol*. 2022;79(12):1250-1259. doi:10.1001/jamaneurol.2022.3392

14. Hu Y, Cho M, Sachdev P, et al. Fluid biomarkers in the context of amyloid-targeting disease-modifying treatments in Alzheimer's disease. *Med*. 2024;0(0). doi:10.1016/j.medj.2024.08.004

15. Schindler SE, Petersen KK, Saef B, et al. Head-to-head comparison of leading blood tests for Alzheimer's disease pathology. *Alzheimers Dement*. Published online October 12, 2024. doi:10.1002/alz.14315

16. Milà-Alomà M, Ashton NJ, Shekari M, et al. Plasma p-tau231 and p-tau217 as state markers of amyloid-β pathology in preclinical Alzheimer's disease. *Nature Medicine*. 2022;28(9):1797-1801. doi:10.1038/s41591-022-01925-w

17. Janelidze S, Barthélemy NR, Salvadó G, et al. Plasma Phosphorylated Tau 217 and Aβ42/40 to Predict Early Brain Aβ Accumulation in People Without Cognitive Impairment. *JAMA Neurol*. Published online July 28, 2024:e242619. doi:10.1001/jamaneurol.2024.2619

18. Li Y, Yen D, Hendrix RD, et al. Timing of Biomarker Changes in Sporadic Alzheimer's Disease in Estimated Years from Symptom Onset. *Ann Neurol*. 2024;95(5):951-965. doi:10.1002/ana.26891

19. Schindler SE, Li Y, Buckles VD, et al. Predicting Symptom Onset in Sporadic Alzheimer Disease With Amyloid PET. *Neurology*. 2021;97(18):e1823-e1834. doi:10.1212/WNL.0000000000012775

20. Betthauser TJ, Bilgel M, Koscik RL, et al. Multi-method investigation of factors influencing amyloid onset and impairment in three cohorts. *Brain*. 2022;145(11):4065-4079. doi:10.1093/brain/awac213

21. Weiner MW, Veitch DP, Aisen PS, et al. The Alzheimer's Disease Neuroimaging Initiative 3: continued innovation for clinical trial improvement. *Alzheimers Dement*. 2017;13(5):561-571. doi:10.1016/j.jalz.2016.10.006

22. Weiner MW, Aisen PS, Jack CR, et al. The Alzheimer's disease neuroimaging initiative: progress report and future plans. *Alzheimers Dement*. 2010;6(3):202-211.e7. doi:10.1016/j.jalz.2010.03.007

23. Kang JH, Korecka M, Figurski MJ, et al. The Alzheimer's Disease Neuroimaging Initiative 2 Biomarker Core: A review of progress and plans. *Alzheimers Dement*. 2015;11(7):772-791. doi:10.1016/j.jalz.2015.05.003

24. Jagust WJ, Landau SM, Koeppe RA, et al. The Alzheimer's Disease Neuroimaging Initiative 2 PET Core: 2015. *Alzheimers Dement*. 2015;11(7):757-771. doi:10.1016/j.jalz.2015.05.001

25. Koeppe, R.A. Data analysis for amyloid PET imaging: Longitudinal studies. In: ; 2013.

26. Diedrichsen J. A spatially unbiased atlas template of the human cerebellum. *Neuroimage*. 2006;33(1):127-138. doi:10.1016/j.neuroimage.2006.05.056

27. Jack CR, Wiste HJ, Weigand SD, et al. Defining imaging biomarker cut points for brain aging and Alzheimer's disease. *Alzheimer's and Dementia*. 2017;13(3):205-216. doi:10.1016/j.jalz.2016.08.005

28. Pomponio R, Erus G, Habes M, et al. Harmonization of large MRI datasets for the analysis of brain imaging patterns throughout the lifespan. *Neuroimage*. 2020;208:116450. doi:10.1016/j.neuroimage.2019.116450

29. Morris JC, Blennow K, Ivanoiu A, et al. The Clinical Dementia Rating (CDR): current version and scoring rules. *Neurology*. 1993;43(11):2412-2414. doi:10.1212/wnl.0b013e318221ad47

30. Aisen P, Petersen R, Donohue M, et al. Clinical core of the Alzheimer's Disease Neuroimaging Initiative: progress and plans. *Alzheimers Dement*. 2010;6(3):239-246. doi:10.1016/j.jalz.2010.03.006

31. Bilgel M, An Y, Walker KA, et al. Longitudinal changes in Alzheimer's-related plasma biomarkers and brain amyloid. *Alzheimer's & Dementia*. 2023;19(10):4335-4345. doi:10.1002/alz.13157

32. Rembach A, Faux NG, Watt AD, et al. Changes in plasma amyloid beta in a longitudinal study of aging and Alzheimer's disease. *Alzheimer's & Dementia*. 2014;10(1):53-61. doi:10.1016/J.JALZ.2012.12.006

33. Pereira JB, Janelidze S, Stomrud E, et al. Plasma markers predict changes in amyloid, tau, atrophy and cognition in non-demented subjects. *Brain*. Published online June 2, 2021. doi:10.1093/brain/awab163

34. Schindler SE, Bollinger JG, Ovod V, et al. High-precision plasma β-amyloid 42/40 predicts current and future brain amyloidosis. *Neurology*. Published online August 1, 2019:10.1212/WNL.0000000000008081. doi:10.1212/WNL.0000000000008081

35. Pereira JB, Janelidze S, Smith R, et al. Plasma GFAP is an early marker of amyloid-β but not tau pathology in Alzheimer's disease. *Brain*. 2021;144(11):3505-3516. doi:10.1093/brain/awab223

36. Bellaver B, Povala G, Ferreira PCL, et al. Astrocyte reactivity influences amyloid-β effects on tau pathology in preclinical Alzheimer's disease | Nature Medicine. *Nature Medicine*. Published online May 29, 2023. doi:10.1038/s41591-023-02380-x

37. Therriault J, Vermeiren M, Servaes S, et al. Association of Phosphorylated Tau Biomarkers With Amyloid Positron Emission Tomography vs Tau Positron Emission Tomography. *JAMA Neurology*. Published online December 12, 2022. doi:10.1001/jamaneurol.2022.4485

38. Gaetani L, Blennow K, Calabresi P, Di Filippo M, Parnetti L, Zetterberg H. Neurofilament light chain as a biomarker in neurological disorders. *J Neurol Neurosurg Psychiatry*. 2019;90(8):870-881. doi:10.1136/jnnp-2018-320106

39. Khalil M, Pirpamer L, Hofer E, et al. Serum neurofilament light levels in normal aging and their association with morphologic brain changes. *Nat Commun*. 2020;11(1):812. doi:10.1038/s41467-020-14612-6

40. Mattsson N, Andreasson U, Zetterberg H, Blennow K, Alzheimer's Disease Neuroimaging Initiative. Association of Plasma Neurofilament Light With Neurodegeneration in Patients With Alzheimer Disease. *JAMA Neurol*. 2017;74(5):557-566. doi:10.1001/jamaneurol.2016.6117

41. Janelidze S, Teunissen CE, Zetterberg H, et al. Head-to-Head Comparison of 8 Plasma Amyloid-β 42/40 Assays in Alzheimer Disease. *JAMA Neurology*. Published online September 20, 2021. doi:10.1001/jamaneurol.2021.3180

42. Palmqvist S, Insel PS, Stomrud E, et al. Cerebrospinal fluid and plasma biomarker trajectories with increasing amyloid deposition in Alzheimer's disease. *EMBO Molecular Medicine*. 2019;11(12):e11170. doi:10.15252/emmm.201911170

43. Mattsson-Carlgren N, Andersson E, Janelidze S, et al. Aβ deposition is associated with increases in soluble and phosphorylated tau that precede a positive Tau PET in Alzheimer's disease. *Sci Adv*. 2020;6(16):eaaz2387. doi:10.1126/sciadv.aaz2387

44. Roe CM, Ances BM, Head D, et al. Incident cognitive impairment: longitudinal changes in molecular, structural and cognitive biomarkers. *Brain*. 2018;141(11):3233-3248. doi:10.1093/brain/awy244

45. Cogswell PM, Lundt ES, Therneau TM, et al. Modeling the temporal evolution of plasma p-tau in relation to amyloid beta and tau PET. *Alzheimers Dement*. 2024;20(2):1225-1238. doi:10.1002/alz.13539

46. Pontecorvo MJ, Devous MD, Kennedy I, et al. A multicentre longitudinal study of flortaucipir (18F) in normal ageing, mild cognitive impairment and Alzheimer's disease dementia. *Brain*. 2019;142(6):1723-1735. doi:10.1093/brain/awz090

47. Biel D, Brendel M, Rubinski A, et al. Tau-PET and in vivo Braak-staging as prognostic markers of future cognitive decline in cognitively normal to demented individuals. *Alzheimers Res Ther*. 2021;13(1):137. doi:10.1186/s13195-021-00880-x

48. Strikwerda-Brown C, Hobbs DA, Gonneaud J, et al. Association of Elevated Amyloid and Tau Positron Emission Tomography Signal With Near-Term Development of Alzheimer Disease Symptoms in Older Adults Without Cognitive Impairment. *JAMA Neurol*. 2022;79(10):975-985. doi:10.1001/jamaneurol.2022.2379

49. Ossenkoppele R, Pichet Binette A, Groot C, et al. Amyloid and tau PET-positive cognitively unimpaired individuals are at high risk for future cognitive decline. *Nature Medicine*. 2022;28(11):2381-2387. doi:10.1038/s41591-022-02049-x

50. Bateman RJ, Xiong C, Benzinger TLS, et al. Clinical and Biomarker Changes in Dominantly Inherited Alzheimer's Disease. *New England Journal of Medicine*. 2012;367(9):795-804. doi:10.1056/NEJMoa1202753

51. Ryman DC, Acosta-Baena N, Aisen PS, et al. Symptom onset in autosomal dominant Alzheimer disease: A systematic review and meta-analysis. *Neurology*. 2014;83(3):253-260. doi:10.1212/WNL.0000000000000596

52. Su Y, Flores S, Wang G, et al. Comparison of Pittsburgh compound B and florbetapir in cross-sectional and longitudinal studies. *Alzheimers Dement (Amst)*. 2019;11:180-190. doi:10.1016/j.dadm.2018.12.008

53. Weiner MW, Veitch DP, Miller MJ, et al. Increasing participant diversity in AD research: Plans for digital screening, blood testing, and a community-engaged approach in the Alzheimer's Disease Neuroimaging Initiative 4. *Alzheimers Dement*. 2023;19(1):307-317. doi:10.1002/alz.12797

Table 1. Characteristics of the study participants

	Combined (n=292)	Reference group[a] (n=80)	Cohort with an estimated age at amyloid positivity[b] (n=190)		Cohort with an estimated age at tau positivity[c] (n=70)	
			Values	P-value	Values	P-value
Demographics						
Age, years	73.9 (7.0)	72.7 (6.4)	74.2 (7.1)	0.08	75.0 (7.31)	**0.04**
Women, n (%)	147 (50.3)	36 (45.0)	100 (52.6)	0.31	37 (52.9)	0.43
APOE ε4 carriers, n (%)	121 (41.4)	17 (21.2)	97 (51.1)	**<0.0001**	38 (54.3)	**<0.0001**
Years of education	16.3 (2.7)	16.9 (2.6)	16.0 (2.6)	**0.01**	16.4 (2.8)	0.24
Race (Black/White/Other), n (%)	10 (3.4) / 276 (94.5) / 6 (2.1)	3 (3.8) / 75 (93.8) / 2 (2.4)	6 (3.2) / 182 (95.8) / 2.0 (2.0)	0.51	1 (1.4) / 66 (94.3) / 3 (4.3)	0.42
Clinical/Cognitive assessments						
CDR 0/0.5/1, n (%)	180 (61.6) / 108 (37.0) / 4 (1.4)	80 (100) / - / -	89 (46.8) / 97 (51.1) / 4 (2.1)	**<0.0001**	36 (51.4) / 34 (48.6) / 0 (0.0)	**<0.0001**
CU/MCI/Dementia, n (%)	179 (61.3) / 105 (36.0) / 8 (2.7)	80 (100.0) / - / -	87 (45.8) / 95 (50.0) / 8 (4.2)	**<0.0001**	36 (51.4) / 31 (44.3) / 3 (4.3)	**<0.0001**
CDR-SB	0.6 (1.0)	0.0 (0.0)	0.9 (1.2)	**<0.0001**	0.7 (0.9)	**<0.0001**
AD biomarkers						
CSF p-tau181/Aβ42	0.03 (0.02)	0.01 (0.00)	0.0 (0.0)	**<0.0001**	0.04 (0.02)	**<0.0001**
Amyloid PET, SUVR	0.81 (0.14)	0.7 (0.0)	0.9 (0.1)	**<0.0001**	0.8 (0.2)	**<0.0001**
Amyloid PET-positive, n (%)	130 (44.5)	0.0 (0.0)	127 (66.8)	**<0.0001**	38 (54.3)	**<0.0001**
Cortical thickness meta-ROI, mm	2.81 (0.14)	2.8 (0.1)	2.8 (0.2)	**0.018**	2.8 (0.2)	**0.038**
Tau PET, SUVR	1.30 (0.27)	1.2 (0.1)	1.4 (0.3)	**0.045**	1.6 (0.3)	**0.001**

Characteristics of the study participants with longitudinal plasma biomarker measures at their first plasma assessment. Data is shown as mean and standard deviation or percentage (%), as appropriate. Continuous variables were compared using t-test and categorical variables using Chi squared test. *P*-values refer to differences in relation to the reference group. Significant *P*-values (*P* < 0.05) are shown in bold.
[a]The reference group is comprised of individuals who were amyloid and tau PET-negative and had a CDR=0 and a clinical classification of cognitively unimpaired (CU) throughout the study duration
[b]The cohort with an estimated age at amyloid positivity comprises individuals with at least one positive amyloid PET scan throughout the study duration.
[c]The cohort with an estimated age at tau positivity comprises individuals with at least one positive tau PET scan throughout the study duration.
n=48 individuals had at least one positive amyloid scan and one positive tau scan throughout the study duration and therefore overlap between the two cohorts. Of note, the cohort with an estimated age at symptom onset is identical to the cohort with an estimated age at tau positivity (n=80).

Figure 1. Amyloid and tau PET trajectories as a function of age and estimated years from amyloid or tau positivity.

Individual amyloid (**A, B**) and tau (**C, D**) PET longitudinal trajectories, where each line connects multiple observations from the same individual. Points and line segments are color-coded according to the cognitive status assigned to each observation (Blue for CDR=0, orange for CDR=0.5, and red for CDR=>1). Triangles represent *APOE* ε4 carriers and circles represent *APOE* ε4 non-carriers. 1 value with tau PET above 4 SUVR was excluded for visualization.

Figure 2. AD plasma biomarkers as a function of AD progression timelines

Generalized Additive Mixed Models (GAMM) with cubic spline basis and random intercepts were used to model plasma biomarkers as a function of years from amyloid positivity **(A)**, years from tau positivity **(B)** and years from symptom onset **(C).** Dashed horizontal lines indicate the mean of the reference group, with shaded areas representing the 95% CI. Solid vertical line marks the time of abnormality compared to the reference group. Time periods with a significant rate of change are depicted with thicker lines. Individual trajectories are shown, where each line connects multiple observations from the same individual. Points and line segments are color-coded according to the cognitive status assigned to each observation (Blue for CDR=0, orange for CDR=0.5, and red for CDR=>1). Triangles represent *APOE* ε4 carriers and circles represent *APOE* ε4 non-carriers.

Outlier values were excluded for visualization (1 for Roche Aβ42/Aβ40, 2 for Fujirebio Aβ42/Aβ40, 1 for Janssen p-tau217, 2 for C2N p-tau217, 2 for ALZpath p-tau217, 3 for Fujirebio p-tau217, 3 for Quanterix p-tau181, 3 for Roche GFAP, 2 for Quanterix GFAP, 1 for Roche NfL and 1 for Quanterix NfL).

A. Amyloid timeline **B. Tau timeline** **C. Symptom timeline**

CDR = 0
CDR = 0.5
CDR ≥ 1

APOE ε4 non-carrier
APOE ε4 carrier

Figure 3. Established AD biomarkers and CDR-SB as a function of AD progression timelines

Generalized Additive Mixed Models (GAMM) with cubic spline basis and random intercepts were used to model established AD biomarkers and CDR-SB as a function of years from amyloid positivity **(A)**, years from tau positivity **(B)** and years from symptom onset **(C).** Dashed horizontal lines indicate the mean of the reference group, with shaded areas representing the 95% CI. Solid vertical line marks the time of abnormality compared to the reference group. Time periods with a significant rate of change are depicted with thicker lines. Individual trajectories are shown, where each line connects multiple observations from the same individual. Points and line segments are color-coded according to the cognitive status assigned to each observation (Blue for CDR=0, orange for CDR=0.5, and red for CDR=>1). Triangles represent *APOE* ε4 carriers and circles represent *APOE* ε4 non-carriers. 1 value with tau PET above 4 SUVR was excluded for visualization.

A. Amyloid timeline

Biomarker
C2N Aβ42/Aβ40
Roche Aβ42/Aβ40
Amyloid PET
Fujirebio Aβ42/Aβ40
CSF p-tau181/Aβ42
C2N %p-tau217
Roche GFAP
Quanterix GFAP
Quanterix Aβ42/Aβ40
ALZPath p-tau217
Janssen p-tau217
Roche p-tau181
Fujirebio p-tau217
Quanterix p-tau181
Tau PET
C2N p-tau217
CDR-SB
Cortical thickness meta-ROI
Roche NfL
Quanterix NfL

Estimated years from amyloid positivity

B. Tau timeline

Biomarker
Quanterix GFAP
Roche GFAP
C2N Aβ42/Aβ40
Amyloid PET
Fujirebio Aβ42/Aβ40
CSF p-tau181/Aβ42
Roche Aβ42/Aβ40
Roche p-tau181
CDR-SB
Quanterix p-tau181
ALZPath p-tau217
Janssen p-tau217
C2N %p-tau217
Tau PET
C2N p-tau217
Fujirebio p-tau217
Quanterix Aβ42/Aβ40
Cortical thickness meta-ROI
Roche NfL
Quanterix NfL

Estimated years from tau positivity

C. Symptom timeline

Biomarker
Roche Aβ42/Aβ40
Amyloid PET
C2N Aβ42/Aβ40
CSF p-tau181/Aβ42
Fujirebio Aβ42/Aβ40
Tau PET
C2N %p-tau217
Roche GFAP
Janssen p-tau217
ALZPath p-tau217
Quanterix GFAP
Fujirebio p-tau217
C2N p-tau217
Roche p-tau181
Quanterix p-tau181
Quanterix Aβ42/Aβ40
CDR-SB
Roche NfL
Quanterix NfL
Cortical thickness meta-ROI

Estimated years from symptom onset

Figure 4. Timing for AD biomarker abnormality

Estimated years from amyloid positivity **(A),** tau positivity **(B)** or symptom onset **(C)** where each outcome measure becomes abnormal compared to the reference group. Points depict the median and error bars depict the 95% CI. Dashed vertical lines at time = 0 represent the time at amyloid positivity (global cortical amyloid PET SUVR > 0.78) **(A)**, tau positivity (mesial-temporal meta-ROI SUVR > 1.41) **(B)** or symptom onset **(C)**. Established AD biomarkers and CDR-SB are shaded in grey.

Supplementary Materials

Table of Contents

Supplementary Methods

1. Estimates for the age at amyloid and tau positivity

To transform amyloid and tau PET data into a time scale, we first obtained the individual rates of change in amyloid and tau using linear mixed effect (LME) models and modelled them as a function of the pathology burden halfway of the follow-up period (midpoint SUVR). As shown in Supplementary Fig. 2A and 2D, the relationship between the rates of amyloid and tau accumulation and midpoint SUVRs was highly non-linear. Thus, Generalized Additive Models (GAM) were used to model amyloid and tau rates of change as a function of midpoint SUVR. To ensure optimal model fit and reduce the influence of data points with high variability, the variance of the GAM fitted values was assessed. A cutoff to identify high-variance values was established at the 90th percentile of the variance distribution (Supplementary Fig. 2B and 2E) and SUVR midpoint values where the variance of the GAM fitted values fell above this cutoff were excluded. As a result, the SUVR interval for amyloid clock modeling was 0.62–1.11 SUVR, and the interval for tau clock modeling was 0.98–2.04 SUVR (Supplementary Fig. 2C and 2F). Within these SUVR intervals, the time between 0.0001 SUVR unit increases was calculated by integrating the inverse of the modeled rates of change. In this study, amyloid time and tau time were conceptualized as the estimated time since amyloid or tau PET positivity. Therefore, amyloid or tau time were calculated by subtracting the cumulative estimated time at the positivity threshold (global cortical amyloid PET SUVR of 0.78 or mesial-temporal meta-ROI tau PET SUVR of 1.41) from the cumulative estimated time at each SUVR value.

To validate the amyloid and tau PET clock models, participants that converted from a negative to a positive PET status during their participation in ADNI study were identified (PET converters; n=81 for amyloid PET and n= 35 for tau PET), and their age at conversion was calculated by averaging their ages at the last negative scan and at the first positive scan. We then tested the associations between the estimated time with actual time since conversion (age at the scan minus actual age at conversion), and between estimated age at amyloid or tau positivity with actual age at conversion. Additionally, we tested the correlations between the estimated time intervals and the actual time intervals.

The resulting estimated amyloid time was strongly correlated (Spearman's $\rho = 0.91$, $P<0.0001$) with actual time since conversion in amyloid PET converters, (Supplementary Fig. 3A). Similarly, the correlation between the estimated tau time and the time since conversion in tau PET converters was Spearman's $\rho = 0.76$, $P<0.0001$(Supplementary Fig. 3D). The correlations between estimated and actual time intervals were Spearman's $\rho = 0.79$, $P<0.0001$ for amyloid time (Supplementary Fig. 3B), and Spearman's $\rho = 0.58$, $P<0.0001$ for tau time (Supplementary Fig. 3E).

Age at amyloid positivity was estimated for all individuals with at least one amyloid positive scan (n=635) as their age at the positive scan minus the corresponding amyloid time. In participants having more than one positive scan, the estimated ages for all positive scans were averaged. In the

group of amyloid PET converters, the resulting correlation between the estimated age at amyloid positivity and the actual age at conversion was Spearman's ρ = 0.96, $P<0.0001$ (Supplementary Fig. 3C) and the mean absolute error (MAE) for the estimated amyloid age was 1.63 years.

Age at tau positivity was estimated for all tau PET scans for participants with at least one positive tau scan (n=212), with a correlation between the estimated age at tau positivity and actual age at conversion in tau PET converters of Spearman's ρ = 0.96, $P<0.0001$ (Supplementary Fig. 3F), and a MAE of 2.3 years.

Additionally, the test-retest reliability of the estimates for the age at amyloid and tau positivity, as assessed by the median slope of the estimated ages across SUVR from different positive scans in individuals with more than one positive scan was -0.29 (IQR: -0.52 to 0.09) for amyloid, and -0.52 (IQR: -1.16 to -0.17) for tau.

2. Estimates for the age at symptom onset

There were 39 clinical progressors in the entire ADNI sample, which were defined as participants with a CDR = 0 and clinically classified as CU at first assessment and CDR > 0 and clinically diagnosed with an AD syndrome at last assessment, with at least one positive amyloid PET scan (*i.e.* with an estimated age at amyloid positivity). 17 of them additionally had an estimated age at tau positivity.

In this group of clinical progressors, age at symptom onset was predicted using the estimated ages at amyloid and tau positivity. Age at amyloid positivity alone was a significant predictor of the age at symptom onset (Adjusted R^2= 0.38, P<0.0001, Supplementary Fig. 4A and 4B). However, upon incorporating age at tau positivity as a predictor, the significance of age at amyloid positivity was no longer observed (*P*>0.05). Age at tau positivity alone predicted age at symptom onset with an adjusted R^2= 0.86 (Supplementary Fig. 4C), therefore, this model was applied to all participants with an estimated age at tau positivity to estimate their age at symptom onset. This approach rendered a correlation of Spearman's ρ = 0.93 between the actual symptom onset age and the estimated symptom onset age in the group of clinical progressors (Supplementary Fig. 4C and 4D). *APOE* ε4 genotype or sex did not have a significant effect on the estimated symptom onset age (*P*>0.05 for all).

Finally, to assign a corresponding timing of AD progression to each outcome measure, years from amyloid positivity, tau positivity, or symptom onset were calculated subtracting the estimated age at amyloid positivity, tau positivity or symptom onset from the age at the outcome measure.

3. Structural MRI measures

Structural brain MRI data included a 3D MP-RAGE or IR-SPGR T1-weighted 3T MRI with sagittal slices. The full cortex of each subject brain was parcellated into 41 bilateral ROIs using a volumetric Desikan-Killiany-Tourville atlas, using FreeSurfer v5.1 for ADNI-GO/2 data and v6.0 for ADNI-3 data. A composite meta-ROI cortical thickness measurement was calculated that included the entorhinal, fusiform, parahippocampal, mid-temporal, inferior-temporal, and angular gyrus[26]. To address differences in image acquisition protocol and image processing, the meta-ROI cortical thickness measure was harmonized using the ComBat-GAM method[27].

Harmonization parameters specific to each batch were estimated using data from the cognitively unimpaired individuals, while considering normal variance due to age, and sex. The age effect was modelled as a non-linear effect over the study age range. The estimated harmonization parameters were then applied to the full dataset. Finally, normal aging effects on the meta-ROI cortical thickness (harmonized) was estimated within cognitively unimpaired and amyloid-PET negative individuals and regressed out from the corresponding measures from all individuals (using age and quadratic age as predictors). The steps for ComBAT-GAM harmonization and adjusting for normal effects of age utilized all available ADNI data for unbiased modelling.

Supplementary Table 1. Baseline characteristics of the study samples used for amyloid and tau PET clock modeling

	Amyloid PET sample (n=784)	Tau PET sample* (n=359)
Age, years	73.0 (7.4)	73.4 (7.1)
Women, n (%)	371 (47.3)	177 (49.3)
APOE ε4 **carriers, n (%)**	313 (39.9)	151 (42.1)
Years of education	16.3 (2.7)	16.6 (2.5)
CDR 0/0.5/≥1, n (%)	361 (46.0) / 392 (50.0) / 31 (4.0)	222 (61.8) / 114 (31.8) / 23 (6.4)
Amyloid PET SUVR	0.8 (0.1)	0.8 (0.2)
Amyloid PET-positive, n (%)	300 (38.3)	141 (39.3)
Tau PET SUVR	1.3 (0.3)	1.3 (0.2)
Number of observations per individual	3.1 (1.2)	2.6 (0.8)
Follow-up time, years	4.7 (2.7)	3.0 (1.4)

Data is shown as mean (M) and standard deviation (SD) or percentage (%), as appropriate.

Data from the entire ADNI study was used to estimate age at amyloid and tau positivity. 784 and 359 participants had longitudinal amyloid and tau PET data and were used to estimate age at amyloid and tau positivity, respectively. 201 participants had both longitudinal amyloid and tau PET data and overlap between the two cohorts.

Supplementary Table 2. Baseline characteristics of participants included in models of established (non-plasma) AD biomarkers

	Combined (n= 1020)	Reference Amyloid and tau negative CU group (n= 263)	Cohort with an estimated age at amyloid positivity* (n= 635)		Cohort with an estimated age at tau positivity** (n=212)	
			Values	*P*-value	Values	*P*-value
Demographics						
Age, years	73.1 (6.8)	71.3 (6.4)	73.8 (6.8)	**<0.0001**	73.4 (6.6)	**0.0005**
Women, n (%)	514 (50.4)	139 (52.9)	317 (49.9)	0.47	103 (48.6)	0.41
APOE ε4 carriers, n (%)	482 (47.3)	47 (19.7)	385 (61.7)	**<0.0001**	108 (61.7)	**<0.0001**
Years of education	16.2 (2.7)	16.8 (2.5)	15.9 (2.7)	**<0.0001**	16.4 (2.45)	0.06
Clinical/Cognitive assessments						
CDR 0/0.5/≥1, n (%)	498 (48.8) / 451 (44.2) / 71 (7.0)	263 (100) / - / -	195 (30.7) / 384 (60.5) / 56 (8.9)	**<0.0001**	78 (36.8) / 118 (55.7) / 16 (7.5)	**<0.0001**
CU/MCI/ Dementia, n (%)	498 (48.8) / 394 (38.6) / 128 (12.5)	263 (100) / - / -	194 (30.6) / 338 (53.3) / 102 (16.1)	**<0.0001**	78 (36.8) / 102 (48.1) / 32 (15.1)	**<0.0001**
CDR-SB	1.2 (1.6)	0.04 (0.2)	1.7 (1.8)	**<0.0001**	1.3 (1.6)	**<0.0001**

Data is shown as mean (M) and standard deviation (SD) or percentage (%), as appropriate.

The reference group is comprised of individuals that were amyloid and tau PET-negative and had a CDR=0 and a clinical classification of cognitively unimpaired (CU) throughout the study duration. Continuous variables were compared using t-test and categorical variables using Chi squared test. *P*-values refer to differences in relation to the reference group. Significant *P*-values ($P < 0.05$) are shown in bold.

*The cohort with an estimated age at amyloid positivity comprises individuals with at least one positive amyloid PET scan throughout the study duration. Among the 635 participants, 494 had CSF ptau-181/Aβ42, 201 had tau PET, and 524 had structural MRI cortical thickness. **The cohort with an estimated age at tau positivity comprises individuals with at least one positive tau PET scan throughout the study duration. Among the 212 participants, 198 had amyloid PET, 124 had CSF ptau-181/Aβ42, and 169 had structural MRI cortical thickness. n=90 individuals had at least one positive amyloid scan and one positive tau scan throughout the study duration and therefore overlap between the two cohorts. Of note, the cohort with an estimated age at symptom onset is identical to the cohort with an estimated age at tau positivity (n=212).

Supplementary Table 3. Timing of AD biomarker abnormality

	Amyloid timeline (estimated years from amyloid positivity)	Tau timeline (estimated years from tau positivity)	Symptom timeline (estimated years from symptom onset)
C2N Aβ42/Aβ40	-7.9 [-7.9 to -6.8]	-9.3 [-9.3 to -6.8]	-14.4 [-16.7 to -11.7]
Roche Aβ42/Aβ40	-7.9 [-7.9 to -6.3]	-7.1 [-9.3 to -4.7]	-14.4 [-16.7 to -11.7]
Amyloid PET	-5.9 [-6.0 to -5.3]	-7.8 [-8.8 to -6.7]	-14.4 [-16.5 to -13.5]
Fujirebio Aβ42/Aβ40	-5.8 [-7.1 to -4.4]	-7.5 [-9.3 to -4.9]	-13.7 [-16.7 to -11.7]
CSF p-tau181/Aβ42	-5.2 [-6.3 to -4.6]	-7.4 [-8.2 to -7.1]	-14.4 [-15.5 to -13.5]
C2N %p-tau217	-4.4 [-5.0 to -3.8]	-4.9 [-5.4 to -4.2]	-11.7 [-11.7 to -11.2]
Roche GFAP	-3.9 [-5.0 to -2.3]	-11.4 [-11.4 to -6.0]	-11.3 [-11.2 to -9.8]
Quanterix GFAP	-3.5 [-4.17 to -1.52]	-11.5 [-11.5 to -7.0]	-11.3 [-12.2 to -9.8]
Quanterix Aβ42/Aβ40	-3.4 [-4.1 to -2.0]	-3.3 [-3.9 to -2.3]	-8.4 [-8.9 to -7.8]
ALZpath p-tau217	-3.2 [-3.9 to -2.3]	-5.4 [-6.0 to -4.4]	-10.2 [-11.3 to -9.6]
Janssen p-tau217	-3.0 [-3.9 to -2.3]	-5.2 [-6.0 to -4.4]	-10.2 [-11.7 to -9.9]
Roche p-tau181	-2.9 [-3.9 to -1.7]	-6.7 [-8.8 to -4.7]	-9.8 [-10.2 to -8.7]
Fujirebio p-tau217	-1.5 [-2.3 to -0.9]	-3.9 [-4.9 to -3.3]	-9.9 [-11.7 to -9.4]
Quanterix p-tau181	-0.9 [-2.2 to 0.4]	-5.4 [-7.5 to -4.2]	-9.4 [-10.2 to -8.3]
Tau PET	-0.7 [-2.5 to 0.4]	-4.7 [-4.7 to -4.3]	-11.9 [-12.7 to -11.7]
CDR-SB	0.0 [-2.4 to 1.1]	-5.7 [-6.1 to -5.2]	-6.2 [-7.3 to -5.2]
C2N p-tau217	-0.4 [-1.2 to 0.1]	-4.7 [-7.1 to -3.1]	-9.9 [-11.7 to -8.7]
Cortical thickness meta-ROI	2.9 [2.3 to 3.6]	-2.8 [-5.9 to 0.3]	-2.1 [-6.1 to -0.9]
Roche NfL	9.2 [3.8 to 12.7]	5.3 [-2.1 to 15.3]	-4.3 [-6.6 to -2.7]
Quanterix NfL	10.5 [3.8 to 15.1]	15.3 [-2.9 to 15.8]	-2.9 [-6.5 to 5.3]

Estimated years (Median [95% CI] from amyloid positivity, tau positivity and symptom onset when each outcome measure becomes abnormal compared to the reference group. Outcome measures are ordered by their times of abnormality in the amyloid timeline. The time of abnormality was defined as the first time point where the confidence interval (CI) of the model estimates did not overlap with the CI of the reference group's mean. The 95% CI of the bootstrapped times of abnormality were calculated based on the 2.5% and 97.5% thresholds using 1000 bootstrapped samples. The amyloid timeline analyses group includes individuals with at least one positive amyloid PET scan throughout the study duration (i.e. with an estimated age at amyloid positivity), and the tau and symptom timeline analyses group comprise individuals with at least one positive tau PET scan throughout the study duration (i.e. with an estimated age at tau positivity). Established AD biomarkers and CDR-SB are shaded in grey.

Supplementary Table 4. Timing of AD biomarkers significant rate of change

	Amyloid timeline (estimated years from amyloid positivity)	Tau timeline (estimated years from tau positivity)	Symptom timeline (estimated years from symptom onset)
C2N Aβ42/Aβ40	-7.9 to 12.2	-11.4 to 17.9	-11.1 to -2.2
Fujirebio Aβ42/Aβ40	-7.9 to 12.4	-0.8 to 4.4	-16.7 to -3.9
Roche Aβ42/Aβ40	-7.9 to 13.0	-11.4 to 17.9	NA
Quanterix Aβ42/Aβ40	-7.1 to 11.5	-11.4 to 6.0	-16.7 to -0.8
C2N p-tau217	-6.8 to 30.3	-1.0 to 17.9	-15.6 to 7.3
C2N %p-tau217	-7.9 to 30.3	-11.4 to 17.9	-16.7 to 7.3
Fujirebio p-tau217	-7.9 to 30.3	-11.4 to 17.9	-16.7 to 7.3
ALZpath p-tau217	-7.9 to 30.3	-11.4 to 17.9	-16.7 to 7.3
Janssen p-tau217	-7.9 to 30.3	-11.4 to 17.9	-16.7 to 7.3
Roche p-tau181	-7.9 to 30.3	-11.4 to 17.9	-16.7 to 2.2
Quanterix p-tau181	-7.1 to 29.3	-11.4 to 3.6	-16.7 to 7.3
Roche GFAP	-7.9 to 30.3	-11.4 to 17.9	-16.7 to 7.3
Quanterix GFAP	-7.1 to 29.3	NA	-16.7 to 7.3
Roche NfL	-7.9 to 30.3	NA	-16.7 to 7.3
Quanterix NfL	-7.1 to 29.3	NA	-16.7 to 7.3
CSF p-tau181/Aβ42	-9.2 to 28.4	-12.2 to 19.8	-18.7 to 7.8
Amyloid PET	-8.5 to 33.5	-11.9 to 19.4	-19.5 to 9.6
Tau PET	-3.6 to 30.3	-4.7 to 21.8	-13.7 to 9.8
Cortical thickness meta-ROI	-9.4 to 32.5	0.5 to 18.1	-6.2 to 9.6
CDR-SB	-12.0 to 34.2	-6.3 to 21.8	-8.6 to 10.6

Estimated years (Median [95% CI] from amyloid positivity, tau positivity, and symptom onset during when each outcome measure shows a significant rate of change. Periods of significant rate of change were identified as those time segments where the confidence interval (CI) of the first derivative of the function did not include zero. The amyloid timeline analyses group includes individuals with at least one positive amyloid PET scan throughout the study duration (i.e. with an estimated age at amyloid positivity), and the tau and symptom timeline analyses group comprise individuals with at least one positive tau PET scan throughout the study duration (i.e. with an estimated age at tau positivity). For values listed as not applicable (NA), the rates of change never became significant. Established AD biomarkers and CDR-SB are shaded in grey.

Supplementary Table 5. Timing of AD biomarker abnormality adjusted by age

	Amyloid timeline (estimated years from amyloid positivity)	Tau timeline (estimated years from tau positivity)	Symptom timeline (estimated years from symptom onset)
Roche GFAP	-3.9 [-5.0 to -2.3]	-11.4 [-11.4 to -6.9]	-13.7 [-15.0 to -11.7]
Quanterix GFAP	-3.6 [-5.0 to -2.0]	-11.4 [-11.4 to -6.0]	-12.2 [-5.0 to -10.2]
Roche p-tau181	-3.6 [-4.5- -2.6]	-7.1 [-9.3 to -4.9]	-11.7 [-11.7 to -10.2]
ALZpath p-tau217	-3.4 [-4.1 to -2.6]	-5.4 [-6.9 to -4.4]	-11.3 [-11.7 to -9.9]
Janssen p-tau217	-3.4 [-4.2 to -2.5]	-5.2 [-6.7 to -4.2]	-12.7 [-11.7 to -10.2]
Quanterix p-tau181	-1.7 [-3.6 to -0.3]	-7.1 [-9.3 to -4.9]	-11.7 [-13.7 to -9.8]
Fujirebio p-tau217	-1.7 [-2.5 to -1.2]	-4.7 [-6.9 to -3.3]	-11.7 [-13.7 to -11.5]
Tau PET	-0.8 [-2.5 to 0.4]	-4.7 [-4.7 to -4.3]	-12.7 [-13.7 to -11.7]
Quanterix NfL	0.1 [-4.9 to 4.1]	-5.4 [-11.4 to 13.2]	-9.6 [-15.0 to 4.8]
Roche NfL	4.1 [1.5 to 4.1]	-3.3 [-8.8 to -0.3]	-8.7 [-11.3 to -6.3]

Estimated years (Median [95% CI] from amyloid positivity, tau positivity, and symptom onset when each outcome measure becomes abnormal compared to the reference group for age-corrected outcome measures. Outcome measures are ordered by their times of abnormality in the amyloid timeline. Age correction was only applied to outcomes that showed a significant association with age. Sex and *APOE* ε4 did not have significant effects on any outcome and therefore were not used to correct outcome measures. The time of abnormality was defined as the first time point where the confidence interval (CI) of the model estimates did not overlap with the CI of the reference group's mean. The 95% CI of the bootstrapped times of abnormality were calculated based on the 2.5% and 97.5% thresholds using 1000 bootstrapped samples. The amyloid timeline analyses group includes individuals with at least one positive amyloid PET scan throughout the study duration (i.e. with an estimated age at amyloid positivity), and the tau and symptom timeline analyses group comprise individuals with at least one positive tau PET scan throughout the study duration (i.e. with an estimated age at tau positivity). Established AD biomarkers and CDR-SB are shaded in grey.

Supplementary Table 6. Timing of AD biomarkers significant rate of change adjusting by age

	Amyloid timeline (estimated years from amyloid positivity)	Tau timeline (estimated years from tau positivity)	Symptom timeline (estimated years from symptom onset)
Fujirebio p-tau217	-7.9 to 30.3	-3.0 to 17.9	-16.7 to 7.3
ALZpath p-tau217	-7.9 to 30.3	-11.4 to 17.9	-16.7 to 7.3
Janssen p-tau217	-7.9 to 30.3	-11.4 to 17.9	-16.7 to 7.3
Roche p-tau181	-7.9 to 30.3	-11.4 to 17.9	-16.7 to 0.3
Quanterix p-tau181	-7.1 to 29.3	NA	-16.7 to 7.3
Roche GFAP	-7.9 to 30.3	-11.4 to 17.9	-16.7 to 7.3
Quanterix GFAP	-7.1 to 29.3	NA	-16.7 to 7.3
Quanterix NfL	-7.1 to 29.3	NA	NA
Roche NfL	-7.9 to 30.3	NA	NA
Tau PET	-3.6 to 30.3	-4.7 to 21.8	-13.7 to 9.8

Estimated years (Median [95% CI] from amyloid positivity, tau positivity, and symptom onset during when each outcome measure shows a significant rate of change for age-corrected outcome measures. Age correction was only applied to biomarkers that showed a significant association with age. Sex and *APOE* ε4 did not have significant effects on any outcome and therefore were not used to correct outcome measures. Periods of significant rate of change were identified as those time segments where the confidence interval (CI) of the first derivative of the function did not include zero. The amyloid timeline analyses group includes individuals with at least one positive amyloid PET scan throughout the study duration (i.e. with an estimated age at amyloid positivity), and the tau and symptom timeline analyses group comprise individuals with at least one positive tau PET scan throughout the study duration (i.e. with an estimated age at tau positivity). For values listed as not applicable (NA), the rates of change never became significant. Established AD biomarkers and CDR-SB are shaded in grey.

Supplementary Table 7. Timing of plasma biomarker abnormality for the subset of individuals with available plasma biomarkers across all assays

	Amyloid timeline (estimated years from amyloid positivity)	Tau timeline (estimated years from tau positivity)	Symptom timeline (estimated years from symptom onset)
C2N Aβ42/Aβ40	-7.1 [-7.1 to -5.0]	-9.3 [-9.3 to -6.8]	-13.7 [-16.7 to -11.5]
Roche Aβ42/Aβ40	-6.8 [-7.1 to -5.0]	-7.1 [-9.30 to -4.7]	-13.7 [-16.7 to -11.3]
Amyloid PET	-5.0 [-5.8 to -4.5]	-7.1 [-7.1 to -5.4]	-11.7 [-13.7 to -11.5]
Fujirebio Aβ42/Aβ40	-5.0 [-6.4 to -4.2]	-7.1 [-9.3 to -4.9]	-11.7 [-16.7 to -10.2]
C2N %p-tau217	-4.2 [-5.0 to -3.5]	-4.9 [-6.7 to -4.7]	-11.3 [-11.7 to -10.2]
Roche GFAP	-3.9 [-5.0 to -3.0]	-11.4 [-11.4 to -6.0]	-11.8 [-12.2 to -9.7]
Quanterix GFAP	-3.9 [-4.1 to -2.0]	-9.4 [-11.4 to -5.4]	-10.2 [-11.8 to -8.9]
Quanterix Aβ42/Aβ40	-3.9 [-5.0 to -2.3]	-3.3 [-4.4 to-2.4]	-8.4 [-8.9 to -7.9]
Janssen p-tau217	-3.2 [-4.0 to -2.4]	-6.9 [-7.1 to -4.9]	-11.3 [-11.7 to -9.9]
ALZpath p-tau217	-3.0 [-3.6 to -2.3]	-5.4 [-7.1 to -4.9]	-10.2 [-11.7 to -9.8]
Roche p-tau181	-2.7 [-3.6 to -1.6]	-6.7 [-9.3 to -4.7]	-9.9 [-10.2 to -8.4]
Fujirebio p-tau217	-1.2 [-1.7 to -0.8]	-5.4 [-7.1 to -4.7]	-10.2 [-11.7 to -9.8]
C2N p-tau217	-0.35 [-1.1 to 0.2]	-6.7 [-7.1 to -4.7]	-9.9 [-11.7 to -8.4]
Quanterix p-tau181	-0.0 [-1.5 to 1.2]	-5.4 [-7.1 to -4.2]	-8.4 [-10.2 to -7.9]
Roche NfL	5.2 [3.2 to 6.4]	14.3 [-8.8 to 15.3]	-5.4 [-7.1 to -3.7]
Quanterix NfL	6.2 [3.8 to 8.5]	14.8 [-6.0 to 15.3]	-4.4 [-6.6 to -1.9]

Estimated years (Median [95% CI] from amyloid positivity, tau positivity, and symptom onset when each plasma biomarker becomes abnormal compared to the reference group in the subset of individuals with available plasma biomarkers across all assays (n=149 with an estimated age at amyloid positivity, n=49 with an estimated age at tau positivity, n=66 in the reference group. Outcome measures are ordered by their times of abnormality in the amyloid timeline. The time of abnormality was defined as the first time point where the confidence interval (CI) of the model estimates did not overlap with the CI of the reference group's mean. The 95% CI of the bootstrapped times of abnormality were calculated based on the 2.5% and 97.5% thresholds using 1000 bootstrapped samples. The amyloid timeline analyses group includes individuals with at least one positive amyloid PET scan throughout the study duration (i.e. with an estimated age at amyloid positivity), and the tau and symptom timeline analyses group comprise individuals with at least one positive tau PET scan throughout the study duration (i.e. with an estimated age at tau positivity). Amyloid PET is shaded in grey as a reference.

Supplementary Table 8. Timing of plasma biomarkers significant rate of change for the subset of individuals with available plasma biomarkers across all assays

	Amyloid timeline (estimated years from amyloid positivity)	Tau timeline (estimated years from tau positivity)	Symptom timeline (estimated years from symptom onset)
C2N Aβ42/Aβ40	-7.1 to 11.1	-11.4 to 17.9	-16.7 to 7.3
Fujirebio Aβ42/Aβ40	-7.1 to 12.2	-0.3 to 3.3	-16.7 to -2.6
Roche Aβ42/Aβ40	-7.1 to 11.1	-11.4 to 17.9	-7.4 to -3.9
Quanterix Aβ42/Aβ40	-7.1 to 11.3	-11.4 to 5.8	-16.7 to -1.0
C2N p-tau217	-6.8 to 29.2	-8.8 to 17.9	-15.6 to 7.3
C2N %p-tau217	-7.1 to 29.2	-11.4 to 17.9	-16.7 to 7.3
Fujirebio p-tau217	-7.1 to 29.2	-11.4 to 17.9	-16.7 to 7.3
ALZpath p-tau217	-7.1 to 29.2	-11.4 to 17.9	-16.7 to 7.3
Janssen p-tau217	-7.1 to 29.2	-11.4 to 17.9	-16.7 to 7.3
Roche p-tau181	-7.1 to 29.2	-11.4 to 6.1	-16.7 to 1.6
Quanterix p-tau181	-7.1 to 29.2	-11.4 to 4.1	-16.7 to 7.3
Roche GFAP	-7.1 to 29.2	NA	-16.7 to 7.3
Quanterix GFAP	-7.1 to 29.2	-11.4 to 17.9	-16.7 to 7.3
Roche NfL	-7.1 to 29.2	NA	-16.7 to 7.3
Quanterix NfL	-7.1 to 29.2	NA	-16.7 to 7.3
Amyloid PET	-7.1 to 29.2	-11.4 to 17.9	-16.7 to 7.3

Estimated years (Median [95% CI] from amyloid positivity, tau positivity, and symptom onset during when each plasma biomarker shows a significant rate of change in the subset of individuals with available plasma biomarkers across all assays. Periods of significant rate of change were identified as those time segments where the confidence interval (CI) of the first derivative of the function did not include zero. The amyloid timeline analyses group includes individuals with at least one positive amyloid PET scan throughout the study duration (i.e. with an estimated ag at amyloid positivity), and the tau and symptom timeline analyses group comprise individuals with at least one positive tau PET scan throughout the study duration (i.e. with an estimated age at tau positivity). For values listed as not applicable (NA), the biomarker's rates of change never became significant. Amyloid PET is shaded in grey as a reference.

Supplementary Fig. 1. Mixture Model Fit of the distribution of mesial-temporal meta-ROI [18F] flortaucipir SUVR in the ADNI Dataset

A Gaussian mixture model with two components was applied to derive a threshold for early tau PET positivity using cross-sectional mesial-temporal meta-ROI data from the entire ADNI study (*n*=907), regardless of the availability of plasma biomarkers. The threshold was established at a value equal to the mean plus two standard deviations of the first component, which corresponds to the low tau burden cluster within the Gaussian mixture model. This process rendered a threshold of 1.41 SUVR for tau PET positivity.

Supplementary Fig. 2. Model fits to estimate amyloid and tau PET clocks

Generalized Additive Models (GAM) with cubic spline were applied to evaluate the relationship between amyloid or tau burden halfway the follow-up period (midpoint SUVR) and the rate of change in amyloid or tau PET. A and D show the GAM model fit including the entire midpoint SUVR range for amyloid and tau PET data, respectively. B and E show the variance of the GAM fitted values across midpoint SUVR for amyloid and tau PET data, respectively, in models including the entire midpoint SUVR range (A and D). Horizontal dashed red line depicts the 90th percentile variance cutoff. C and F show the GAM model fit after excluding midpoint SUVR values above the 90th percentile variance cutoff, which resulted in an interval of 0.62-1.11 SUVR for amyloid PET, and 0.98-2.04 SUVR for tau PET. Each point depicts the value of an individual and the solid blue lines indicate the model fit. *APOE* ε4 carriers are shown in blue and *APOE* ε4 non-carriers are shown in green.

Supplementary Fig. 3. Validation of the amyloid and tau PET clock estimates

A Amyloid time = 0.41 x (time since conversion) + 0.17 years ± 0.10 years
Intercept *p* = 0.08
Adjusted R² = 0.74
Spearman's ρ = 0.91

- Aβ−negative
- Aβ−positive

B Amyloid time interval = 1.44 x (actual scan interval) − 0.31 years ± 0.11 years
Intercept *p* = 0.01
Adjusted R² = 0.61
Spearman's ρ = 0.80

C Age at amyloid positivity = 1.01 x (conversion age) − 2.39 years ± 2.47 years
Intercept *p* = 0.34
Adjusted R² = 0.93
Spearman's ρ = 0.96

D Tau time = 0.31 x (time since conversion) − 0.14 years ± 0.12 years
Intercept *p* = 0.27
Adjusted R² = 0.51
Spearman's ρ = 0.76

- Tau−negative
- Tau−positive

E Tau time interval = 1.35 x (actual scan interval) − 0.06 years ± 0.16 years
Intercept *p* = 0.72
Adjusted R² = 0.38
Spearman's ρ = 0.58

F Age at tau positivity = 1.07 x (conversion age) − 6.68 years ± 3.33 years
Intercept *p* = 0.05
Adjusted R² = 0.93
Spearman rho = 0.96

Amyloid and tau time models were validated in the group of individuals that converted from a negative to a positive amyloid or tau PET scan throughout the study duration, respectively. Actual conversion age in PET converters was calculated by averaging the ages at the last negative scan and at the first positive scan. A and D show the correlation between the estimated amyloid time (A) or tau time (D) and the actual time since conversion. B and E show the correlation between amyloid (B) or tau (E) time intervals between scans (amyloid or tau time - amyloid or tau time at first scan) and actual time intervals between scans above the positivity threshold in the whole sample. C and F show the correlations between the estimated age at amyloid positivity (C) or tau positivity (F) with the actual age at conversion in PET converters. Of note, each point at A, B, D and E depict a scan and each point at C and F depict an individual. Solid lines indicate the linear regression lines.

Supplementary Fig. 4. Model fits for the estimation of symptom onset age

Each point depicts the value of an individual and the solid line indicate the linear regression line. *APOE* ε4 carriers are shown in blue and *APOE* ε4 non-carriers are shown in green.

Linear regressions were run to estimate age at symptom onset in clinical progressors with the estimated age at amyloid positivity (A, n = 39) or tau positivity (C, n = 17) as predictors. Clinical progressors were defined as participants that had a CDR of 0 and were clinically classified as CU at their first assessment and had a CDR > 0 and a clinical diagnosis of an AD clinical syndrome (MCI or dementia due to AD) at their last assessment.

To validate the models, the correlation between the actual age at symptom onset and the estimated age at symptom onset was evaluated for the amyloid-based (B) and the tau-based model (D). As the estimated age at tau positivity was a better predictor of the age at symptom onset (Adj R^2 = 0.86 for the tau-based model vs Adj R^2=0.38 for the amyloid age-based model), we used estimated age at tau positivity to estimate age at symptom onset in all study participants with an estimated age at tau positivity available. Of note, when both age at amyloid and tau positivity were included in the model, age at amyloid positivity did not show a significant effect (P>0.05). *APOE* ε4 carriers are shown in blue and *APOE* ε4 non-carriers are shown in green.

...ere used to model ...m onset. Dashed ...e 95% CI. Solid ...ate of change are

...2 for ALZpath p-

a function of

CDR = 0
CDR = 0.5
CDR >= 1

APOE− ε4 non−carrier
APOE− ε4 carrier

Estimated years from symptom onset

basis and random intercepts were used to model ity, tau positivity, and symptom onset. Dashed shaded areas representing the 95% CI. Solid s with significant biomarker rate of change are

Aβ40, 2 for Fujirebio Aβ42/Aβ40, 3 for Roche
GFAP, 2 for Quanterix GFAP, 1 for Roche NfL and 1 for Quanterix NfL)

Supplementary Fig. 7. Biomarker trajectories as a function of age

Individual biomarker trajectories, where each line connects multiple observations from the same individual. Points and line segments are color-coded according to the cognitive status assigned to each observation (blue for individuals with a CDR = 0, orange for individuals with a CDR = 0.5, and red for individuals with a CDR >= 1). Triangles represent *APOE* ε4 carriers and circles represent *APOE* ε4 non-carriers.

Outlier values were excluded for visualization (1 for Roche Aβ42/Aβ40, 2 for Fujirebio Aβ42/Aβ40, 1 for Janssen p-tau217, 2 for C2N p-tau217, 2 for ALZpath p-tau217, 3 for Fujirebio p-tau217, 3 for Quanterix p-tau181, 3 for Roche GFAP, 2 for Quanterix GFAP, 1 for Roche NfL, 1 for Quanterix NfL and 1 for tau PET).

Supplementary Fig. 8. Biomarker trajectories as a function of amyloid PET

Individual biomarker trajectories, where each line connects multiple observations from the same individual. Points and line segments are color-coded according to the cognitive status assigned to each observation (blue for individuals with a CDR = 0, orange for individuals with a CDR = 0.5, and red for individuals with a CDR >= 1). Triangles represent *APOE* ε4 carriers and circles represent *APOE* ε4 non-carriers.

Outlier values were excluded for visualization (1 for Roche Aβ42/Aβ40, 2 for Fujirebio Aβ42/Aβ40, 1 for Janssen p-tau217, 2 for C2N p-tau217, 2 for ALZpath p-tau217, 3 for Fujirebio p-tau217, 3 for Quanterix p-tau181, 3 for Roche GFAP, 2 for Quanterix GFAP, 1 for Roche NfL, 1 for Quanterix NfL and 1 for tau PET).

Supplementary Fig. 9. Biomarker trajectories as a function of tau PET

ectories, where each line connects multiple observations from the same individual. re color-coded according to the cognitive status assigned to each observation (blue for 0, orange for individuals with a CDR = 0.5, and red for individuals with a CDR >= 1). ε4 carriers and circles represent *APOE* ε4 non-carriers. Outlier values were excluded oche Aβ42/Aβ40, 2 for Fujirebio Aβ42/Aβ40, 1 for Janssen p-tau217, 2 for C2N p-au217, 3 for Fujirebio p-tau217, 3 for Quanterix p-tau181, 3 for Roche GFAP, 2 for L, 1 for tau PET).