1 **Biobank-scale exposome-wide risk factors in CAD and T2D: observational, predictive, and**
2 **potentially causative evidence**
3

4 Sivateja Tangirala[1], Arjun K Manrai[1], John PA Ioannidis[2]*, Chirag J Patel[1]*
5
6
[1]Department of Biomedical Informatics, Harvard Medical School, Boston, Massachusetts.
[2]Stanford Prevention Research Center, Department of Medicine, Stanford University School of
Medicine, Stanford, CA.
[3]Department of Epidemiology and Population Health, Stanford University School of Medicine,
Stanford, CA
[4]Department of Biomedical Data Science, Stanford University School of Medicine, Stanford, CA
[5]Meta-Research Innovation Center at Stanford (METRICS), Stanford University, Stanford, CA
14
15 *Co-corresponding authors

16
17
18
19

Abstract

Cardiovascular disease and diabetes are intricately related and influenced by factors within the

"exposome". The exposome refers to a broad assessment of cumulative environmental

exposures, including physical-chemical, biological, behavior, ecosystem, and social influences,

experienced from conception throughout life, along with their dynamic interactions with the

genome (Vermeulen et al. 2020). Distinguishing between correlational and potentially causative

risk associations is challenging, especially at exposome scale. Here, we triangulate observational

Exposome-Wide Association Study (*ExWAS*) evidence with "randomized" evidence for the

exposome using mendelian randomization (MR) for almost 500 exposures. First, the *ExWAS*

identified 144 significant factors for coronary artery disease (CAD) and 237 for type 2 diabetes

(T2D), with 120 shared between both. These factors had modest predictive ability (variance

explained) for both phenotypes. However, a genetic-based potentially causative relationship was

deduced for only 14 factors in CAD and 16 in T2D, with seven implicated in both. Additionally,

we found strong concordance of MR-validated findings between prevalent and incident disease

associations (85.7% [12/14] for CAD and 87.5% [14/16] for T2D). Most correlational findings

pertain to lifestyle factors (particularly diet), but social educational factors are more prominently

highlighted among those with potentially causative support.

Introduction

Type 2 diabetes (T2D) and coronary artery disease (CAD) are causatively related[1] and carry a tremendous burden of disease worldwide. While genetics play a role, environmental factors – ranging from diet and pollution to social and behavioral factors – are thought to be key contributors to disease risk [2] , (Miller and Banbury Exposomic Consortium 2025)[3].The totality of these external influences, collectively referred to as the exposome [4], encompasses diverse domains, including *social* (e.g., education and income), *behavior* (dietary and lifestyle), *physical-chemical* (e.g., nutrients and chemicals), *biological* (e.g., infection), and *ecosystem* (e.g. air pollution) variables. Understanding how these exposures contribute to disease remains a major scientific challenge.

Traditionally, epidemiological and toxicological studies have focused on a few pre-selected hypotheses, associating a limited number of exposures at a time. Hypothesis-driven studies are a way to test candidate relationships and, when conducted with methodological rigor and transparency, can yield robust and reproducible findings. However, such studies may be susceptible to certain limitations, including publication bias, confirmation bias, and a focus on pre-specified exposures, which can constrain the discovery of novel or unexpected associations—particularly in complex, multi-factorial environments such as the exposome where effect sizes may be moderate.

This narrow approach has led to inconsistent findings, potential false positives, and difficulty in replicating results across populations[5]. By contrast, genetic epidemiology has revolutionized risk

factor discovery by systematically analyzing thousands of genetic variants in settings such as

genome-wide association studies (GWAS), improving reproducibility and replicability across a

number of cohorts, and enhancing triangulation[6]. However, unlike genetic variants, exposure

variables are time-varying, dynamic, and often highly correlated within themselves, making it

difficult to disentangle their individual effects [7].

Importantly, considering only a handful of factors, leads to a fragmented and irreproducible

literature of associations [5,8]. Here, inspired by the genomics paradigm, we deploy an *exposome-*

wide association study (ExWAS [9,10]) framework, systematically screening a comprehensive array

of exposures in an agnostic manner, while taking into account testing multiplicity [8,11, 12] . Second,

by integrating ExWAS with Mendelian randomization (MR), we are among the first to bridge the

gap between observational discovery and causal inference. This discovery-driven approach

contrasts with existing MR studies that often focus on pre-specified exposures, enabling the

identification of novel, unbiased relationships while maintaining the rigor of causal inference.

To "break" the correlation between exposures requires experimentation or "instrumental

variables", which serve as randomized proxies of exposure [13]. However, the degree by which

instrumental variables, or variables that can guide inference across the exposome is understudied

[7]. Instrumental variables have been important in enhancing potentially causative determinations

in other disciplines [14]. With the advent of biobank data with both measured genotypes,

exposures, and longitudinal phenotypic outcomes one can harness mendelian randomization

(MR)[1, 15, 16] as a way to scalably conduct "instrumental variable" analysis that hone in on

identifying potentially independent causative relationships among exposures and phenotypic or

disease outcomes.

Mendelian randomization (MR), which uses genetic variants as instrumental variables, offers an

important strategy to reduce confounding and reverse causation in this setting. Nonetheless, MR

can be challenged by violations of core assumptions, such as the exclusion restriction,

particularly when applied to time-varying exposures [17] (PMID: 24681576).

We combine ExWAS and MR to identify the exposomic associations with T2D and CAD

(Figure 1). First, we conducted an ExWAS systematically testing each of 495 individual factors

of the exposome using a sample of 472,240 white European participants from the UK Biobank

(UKB). and estimated its concordance with data from FinnGen, a biobank of 218,957

participants based in Finland. This approach allowed us to systematically evaluate which

exposures had the strongest genetic-based evidence for potentially causative relationships and to

distinguish between correlations and genuine risk factors.

Results

Baseline characteristics

We report the baseline characteristics of clinical and demographic variables including age, sex,

average household income, HbA1c, systolic blood pressure, diastolic blood pressure, BMI, LDL,

HDL, triglycerides, total cholesterol, family history for CAD, family history for T2D, prevalent

CAD, prevalent T2D, and smoking history of the UK Biobank (UKB) cohort participants in

Table 1. Briefly, the average age of participants was 56.76, 54.48% were female participants, and

~64% of participants had an average household income of less than 52,000 Euros. Out of the

472,240 European White individuals we utilize for our analysis, 7,426 have developed CAD and

12,050 have developed T2D (after baseline visit) while considering those who had already

developed CAD (n=10,772) or T2D (n=17,303) at baseline separately. Additionally, out of the

218,957 participants in the FinnGen sample we utilize to evaluate our findings via two sample

MR, 29,193 have developed T2D and 21,012 have developed CAD [18].

Distribution of observational association sizes in incident CAD and T2D

We associated 495 exposures (Extended Data Table 1) with CAD and T2D. With FDR<0.05, we

identified 144 significant exposures for CAD (Supplemental Table 1, Figure 2, Supplemental

Results) and 237 significant exposures for T2D (Supplemental Table 2, Supplemental Figure 1,

Supplemental Results). Association per exposomic category (as defined[4]) for each disease are in

Extended Data Table 2. For CAD, exposomic category-wise representation among significant

associations range from 2.33% (Ecosystems; exposomic factors pertaining to Ecosystems

comprise 2.33% of significant associations) to 58.9% (Lifestyle). Similarly for T2D, category-

wise representation among significant associations range from 2.59% (Ecosystems) to 52.8%

(Lifestyle) .

For CAD, the IQR of HRs for significant continuous variables (for a 1 standard deviation [SD]

change) that are protective factors is [0.89,0.94] and for risk factors it is [1.06, 1.11] while the

IQR of HRs for significant binary variables that are protective factors is [0.75, 0.83] and for risk

factors it is [1.25, 1.55]. For T2D, respectively, the IQR of HRs for significant continuous

variables (for a 1 standard deviation [SD] change) are [0.86, 0.92] and [1.07, 1.13] and the IQR

of HRs for significant binary variables are [0.64, 0.80] and [1.23, 1.53].

We found 120 significant exposures shared between T2D and CAD (119 of which are

concordant and one discordant in direction of effect). Additionally, we found 117 exposures

specific to T2D and 24 exposures specific to CAD. The Pearson correlation between the beta

coefficients of CAD and T2D is 0.56 and 0.30 among continuous and binary exposure factors,

respectively (Figure 2).

Concordance of findings between prevalent and incident disease ExWAS

Additionally, we sought to assess the concordance of findings from ExWAS-identified

associations for prevalent disease with those of incident disease. We associated 495 exposures

with prevalent CAD and T2D. With FDR<0.05, we identified 232 significant exposures for

prevalent CAD (Supplemental Table 3) and 309 significant exposures for prevalent T2D

(Supplemental Table 4, Supplemental Results).

We found 137 significant exposures shared between prevalent and incident CAD (134 of which

are concordant and 3 discordant in direction of effect). Furthermore, we found 62 exposures

specific to prevalent CAD and 24 exposures specific to incident CAD. The Pearson correlation

between the beta coefficients of prevalent and incident CAD is 0.53 and 0.38 among continuous

and binary exposure factors, respectively (Supplemental Figure 2). Additionally, considering the

prevalent disease ExWAS as our "diagnostic test", we sought to compare the concordance of the

prevalent CAD ExWAS with incident CAD ExWAS by computing the sensitivity and specificity

of the prevalent CAD ExWAS test. We estimate the sensitivity to be 84.8% and specificity to be

65.2% (Supplemental Table 5).

Similarly, we compare the concordance between exposure associations for prevalent and incident

T2D. We found 189 significant exposures shared between prevalent and incident T2D (184 of

which are concordant and 5 discordant in direction of effect). Furthermore, we found 48

exposures specific to prevalent T2D and 24 exposures specific to incident T2D. The Pearson

correlation between the beta coefficients of prevalent and incident T2D is 0.79 and 0.75 among

continuous and binary exposure factors, respectively. We estimate the sensitivity and specificity

of the prevalent T2D ExWAS test relative to incident T2D ExWAS to be 88.5% and 61.9%,

respectively (Supplemental Table 6).

MR-based assessment of observational ExWAS associations

We sought genetic-based evidence for potentially causative relationships and performed bi-

directional MR between each exposure-outcome pair for which Genome-wide Association Study

[GWAS] summary statistics were available (for 123/144 (85.4%) of CAD FDR significant

associations and 182/237 (76.8%) of T2D FDR significant associations) to test whether the

observational associations from the ExWAS are potentially causative. To enhance comparability,

we use odds ratios (ORs) of exposures computed from logistic regressions after assessing the

concordance in estimates between hazard ratios computed from Cox proportional hazard models

and odds ratios computed from logistic regressions (Supplemental Figures 3 and 4). We visualize

the concordance (and/or discordance) between ExWAS and MR ORs among these exposure-

disease pairs in Figure 3 for CAD (and Supplemental Figure 5 for T2D).

We identified 14 associations as being nominally significant (p-value less than 0.05) in forward

MR and *not* p-value significant in the reverse direction in CAD while also being found to be

concordant in direction to the corresponding ExWAS associations. The IQR of the ORs for CAD

among MR-validated associations was [0.38, 0.58] and [1.79, 6.15] for protective and risk

factors, respectively. The IQR of ORs for T2D among MR-validated associations was [0.25,

0.63] and [1.86, 2.38] for protective and risk factors, respectively.   We also formally tested the

difference in effect sizes (absolute value of beta estimates) of MR-validated factors and non-MR-

validated factors with the Mann-Whitney test. MR-validated factors had stronger effects than

non-MR-validated ones for CAD (Mann-Whitney statistic (W) = 571, p-value = 0.0044) and

possibly also for T2D (W= 935, p-value = 0.021).

The MR results for CAD and T2D are made available in Supplemental Tables 7 and 8,

respectively. Additionally, we found 12 out of the 14 total (85.7%) prevalent ExWAS-identified

associations validated by MR to be the same as the MR-validated ones for incident ExWAS for

CAD. Similarly, we found 14 out of the 16 (87.5%) total prevalent ExWAS-identified

associations validated by MR to be the same as the MR-validated ones for incident ExWAS for

T2D.

We describe the associations that were not only identified as significant from the ExWAS but

also found to be significant (p-value less than 0.05) from the MR analysis for CAD (while also

concordant in direction). The top 3 MR-validated protective factor associations (in order of

increasing MR p-value) were having the educational qualification of A levels/AS levels or

equivalent (MR OR: 0.39, MR p-value: 1.12×10^{-5}, ExWAS OR: 0.71, ExWAS FDR: 1.92×10^{-26}),

having the educational qualification of college or university degree (MR OR: 0.57, MR p-value:

3.51×10^{-5}, ExWAS OR: 0.66, ExWAS FDR: 3.57×10^{-44}), and the age participant first had sexual

intercourse (MR OR: 0.75, MR p-value: 4.08×10^{-5}, ExWAS OR: 0.82, ExWAS FDR: 3.44×10^{-45}

). The top 3 MR-validated risk factors were Townsend deprivation index at recruitment (MR

OR: 1.75, MR p-value: 5.86×10^{-3}, ExWAS OR: 1.14, ExWAS FDR: 9.23×10^{-20}), having no

educational or other professional qualifications (e.g. nursing, teaching) (MR OR: 1.83, MR p-

value: 0.0105, ExWAS OR: 1.51, ExWAS FDR: 6.22×10^{-49}), and no type of physical activity in

the last four weeks (MR OR: 10.5, MR p-value: 0.0313, ExWAS OR: 2.07, ExWAS FDR:

2.37×10^{-73}).

Additionally, MR results across the full suite of MR methods tested for are made available in

Supplemental Tables 9 and 10. Among the top ExWAS-determined hits with MR-confirmed

associations for CAD, we observed consistent effect directionality across all MR methods for

most (9/14) exposures. Furthermore, among these 9 exposures with consistently concordant

direction, statistical significance at p < 0.069 was seen in at least 3 of the 5 MR models for 2 of

them, but none of the 9 had formally statistically significant results in all 5 models. For example,

"Usual walking pace" showed directionally concordant inverse associations with CAD risk

across methods (ranging from -1.408 [simple mode] to -0.525 [IVW]) and "Age first had sexual

intercourse" showed (estimates ranging from -0.521 [simple mode] to -0.157 [MR Egger]). In

contrast, for a subset of 5 exposures, such as "Age completed full-time education" and

"Qualifications: A levels/AS levels or equivalent", and "Qualifications: A levels/AS levels or

equivalent," the majority of MR methods yielded consistent directionality, but MR-Egger results

showed discordance. These findings highlight the robustness of our primary findings while

emphasizing the need for cautious interpretation in cases where sensitivity analyses indicate

potential pleiotropy.

We also report the proportion of MR-validated exposures by exposomic category for CAD.

Among social factors, 7.87% (7/89) were MR-validated and among lifestyle factors 5.3% (7/132)

were MR-validated. More specifically, among education factors, 62.5% (5/8) were MR-validated

and among dietary factors only 4.1% (3/73) were MR-validated. None were MR-validated

among ecosystems and physical-chemical factors. Similarly, we report the proportion of MR-

validated exposures by exposomic category for T2D. Among social factors, 6.9% (6/87) were

MR-validated and among lifestyle factors 6.8% (9/132) were MR-validated. More specifically,

among education factors, 62.5% (5/8) were MR-validated and among dietary factors only 4.1%

(3/73) were MR-validated. 4.8% (1/21) were MR-validated among physical-chemical factors and

none were MR-validated among ecosystems factors.

Next, we sought to better understand the concordance of our ExWAS test with respect to MR by

computing the sensitivity and specificity of ExWAS (considering ExWAS as our "diagnostic

test"). We extended our MR analysis to compute associations for all ExWAS tested disease-

exposure pairs for which GWAS summary statistics were available [518 (52.3% of 990 total

ExWAS-tested disease-exposure pairs)]. More specifically, we estimated the degree to which

findings were both ExWAS significant and MR significant, or sensitivity (77.8% for CAD and

94.1% for T2D). Additionally, we estimated the degree to which ExWAS non-significant

associations were also found to be MR non-significant, or specificity (37.3 % for CAD and

26.3% for T2D) (Extended Data Tables 3 and 4).

We observed a difference in the sensitivity and specificity of ExWAS between the lifestyle and

social exposomic variable categories. For lifestyle factors, we estimate 70% sensitivity for CAD

and 75% for T2D and 31.3% specificity for CAD and 21.4% for T2D [Supplemental Tables 11

and 12]. For social factors, the sensitivity was 87.5% for CAD and 85.7% for T2D and

specificity was 45% for CAD and 34.6% for T2D [Supplemental Tables 13 and 14]. The

correlation between ExWAS and MR beta estimates were 0.146 and -0.0271 among binary and

continuous variables, respectively.

We identified 7 associations (5 education-related variables and 2 lifestyle factors [age first had

sexual intercourse and usual walking pace]) in both T2D and CAD (Table 2) with MR. Here, we

report two types of OR, both those obtained through ExWAS (a conventional OR) and those

from the MR. These variables included age first had sexual intercourse, educational qualification

of college or university degree, usual walking pace, having no educational or other professional

qualifications (e.g. nursing, teaching), educational qualification of A levels /AS levels or

equivalent, educational qualification of O levels/GCSEs or equivalent, age completed full time

education. For example, we identified the educational qualification of college or university

degree to be among the top associations for both T2D (MR OR: 0.407, MR p-value: 5.49×10^{-10},

ExWAS OR: 0.54, ExWAS FDR: 2.64×10^{-137}) and CAD (MR OR: 0.566, MR p-value: 3.51×10^{-5}

^{5}, ExWAS OR: 0.657, ExWAS FDR: 3.57×10^{-44}). Furthermore, we found all of the factors to be

protective (OR less than 1)   "other professional qualifications (e.g. nursing, teaching)" which we

found to be a risk conferring factor for both CAD (MR OR: 1.83, MR p-value: 0.0105, ExWAS

OR: 1.51, ExWAS FDR: 6.22×10^{-49}) and T2D (MR OR: 2.59, MR p-value: 2.68×10^{-4}, ExWAS

OR: 1.81, ExWAS FDR: 3.92×10^{-163}).

Assessing the variance of CAD and T2Ds explained by exposomic, demographic, genetic

and clinical risk factors

For incident CAD, we find that 42.3% of the variance explained by the exposome and

demographics (Extended Data Table 5) can be explained by exposomics alone (e.g., for T2D,

exposures alone explained 8% of the variance, while exposures plus demographics explained

18.8% of the variance). Additionally, 15.8% of the variance explained by the model including

MR validated exposures and demographics is explained by MR-validated exposures alone.

Moreover, we observed how the variance explained by the models change after inclusion of

T2D PRS and BMI. Exposomic variables alone account for 60.5% of the variance explained by

the full model. Similarly, MR validated exposures alone account for 18.9% of the variance

explained by the full model. We describe the T2D results in the Supplemental Results section.

Additionally, we computed the Nagelkerke R^2 from logistic regression models that were run by

serially adding exposomic factors to the previous model in order of increasing FDR-corrected p-

values in addition to the baseline demographic and clinical risk factors (Figure 4). The IQR of

the absolute value of pairwise correlations of the 196 exposures was [0.01, 0.08] (Supplemental

Figure 6). For CAD and T2D, the Nagelkerke R^2 for models ceased to increase when

incorporating beyond the top 175 exposures by FDR-corrected p-values.

Contextualization of MR-validated exposures in risk for CAD and T2D with established

clinical risk factors

Finally, we run clinical risk factor models for CAD and T2D consisting of age, sex, family

history, BMI, systolic blood pressure, diastolic blood pressure, HbA1c, LDL, HDL, triglycerides,

total cholesterol, and smoking history (inspired by prior literature [19, 20])**.** We compare the effects

of these factors with those of seven MR-validated exposures commonly implicated in CAD and

T2D (Supplemental Table 15). The magnitude of the effects of the MR-validated exposures

matches or sometimes even exceeds the magnitude of the effects of established clinical risk

factors.

Discussion

Our analysis integrates ExWAS and MR methods to characterize the exposomic architecture of

factors associated with CAD and T2D, made possible by use of biobank data, such as UK

Biobank and FinnGen. Using ExWAS, we identified 144 (out of 495 [29.1%]) and 237 factors

(out of 495 [47.9%]) for CAD and T2D, respectively. The larger number of factors identified for

T2D may reflect, at least in part, the larger power available for T2D which was more common

than CAD in the biobank sample. Overall, statistically significant ExWAS associations for these

major outcomes seem to be too numerous. This may also explain the plethora of significant

associations published in the epidemiological literature, where exposures are usually tested one

at a time or a few at a time[21].

For both CAD and T2D the magnitude of the effect sizes of ExWAS-identified factors was

modest. Furthermore, the vast majority of them showed no genetic-based evidence for a

potentially causative relationship on MR assessment. MR is an approach that uses genetic

variants as "instruments" to test for a potentially causative association between exposures and

disease [15]. From our MR analysis, the validated factors tended to have somewhat larger effect

sizes than those seen with observational correlation analyses in ExWAS. Only 14 (9.72%) and 16

(6.75%) of ExWAS findings were concordant with MR findings for the two phenotypes, CAD

and T2D, respectively. We observed higher (genetic-based) potentially causative relationship

rates among social and more specifically, education-related factors than for lifestyle factors, even

though the latter accounted for the vast majority of ExWAS associations. While ExWAS may be

sensitive when a large database like the UK Biobank is analyzed, specificity is quite low for T2D

and CAD relative to MR, if MR is seen as the standard.

The strong presence of education-related variables among the apparently potentially causative

ones is in line with other studies, including MR-based ones. For example, Tillmann et al. found

that educational attainment is potentially causative and protective (OR: 0.67 for a 1 SD increase

in higher education roughly corresponding to 3.6 years of additional schooling[22]) risk factor for

coronary artery disease in the Coronary Artery Disease Genetics Consortium

(CARDIoGRAMplusC4D), also suggesting increased levels of education have a protective effect

[22]. Additionally, we identified some factors suggestive of behavior including the age first had

sexual intercourse in association with CAD and T2D risk and this is also in accordance with

prior evidence[23]. Focused and intense "lifestyle" interventions that target weight loss for

individuals with elevated glucose have been developed that have causatively led to decreases in

T2D incidence[24]. Interventions may need to be tailored and tested targeting social and, in

particular, educational factors. Such interventions would require a broader community outlook.

Life course studies that examine education early in life may also be needed.

Our MR analyses incorporated a comprehensive suite of sensitivity methods beyond IVW,

enabling us to assess the robustness of our findings and evaluate the potential for horizontal

pleiotropy, which is a major threat to MR studies. Across the top exposures identified for CAD,

we observed directionally concordant estimates across multiple MR approaches, including MR-

Egger, weighted median, and mode-based methods. Although p-values were often attenuated in

these sensitivity approaches—likely reflecting their lower statistical power relative to IVW—the

consistent effect direction reinforces the plausibility of potentially causal relationships.

MR-Egger estimates for some exposures showed directional discordance relative to other

methods. This highlights the potential for pleiotropy, underscoring the importance of cautious

interpretation and the need for future studies with larger sample sizes and stronger instruments to

confirm these findings, and potential multi-variable approaches. Current mega multi-phenotype

studies have shown a vast potential for pleiotropy (Levin et al. 2025[25]), adding to the threat.

We also explicitly assessed potential sample overlap between the exposure and outcome GWAS

used in our analyses. To the best of our knowledge, there is no overlap between the cohorts

generating exposure GWAS summary statistics (primarily from the Neale Lab[26] and MRC IEU

OpenGWAS[27]) and the FinnGen cohort used for outcome data, minimizing the risk of inflated

type I error.

Additionally, reverse MR analyses were conducted as a sensitivity test to assess potential reverse

causative relationships, rather than to infer causative effects from disease to exposures. While

such analyses can provide insight into the directionality of associations, their interpretation is

limited by potential issues such as instrument strength and biological plausibility. Our reverse

MR findings were used to support the primary forward MR results by evaluating the likelihood

of reverse causative relationship.

Prior work have investigated potentially causative relationships for CAD and T2D through MR

analyses. For example, the study by Georgiou et al. [28] focuses on specific lifestyle factors, such

as alcohol consumption and physical activity, in relation to CAD, while the systematic review by

Yuan et al.[29] synthesizes MR findings for a range of pre-identified exposures relevant to T2D.

These studies provide valuable insights into targeted hypotheses but differ fundamentally from

our discovery-driven, agnostic approach. Specifically, prerequisites for these studies include

documented hypothesis between an exposure and an outcome. Such targeting of a specific

hypothesis may require consideration of other types of evidence, but it may succumb to

confirmation and other cognitive biases that perpetuate preferred narratives.

In contrast, our study employs an exposome-wide association study (ExWAS), which

systematically identifying potential exposure-outcome relationships across a broad range of

domains before validating these relationships with MR. The comprehensive, data-driven

methodology is analogous to GWAS in its agnostic screening of a large number of exposures,

extending the utility of MR beyond its traditional use in hypothesis-driven studies. By integrating

ExWAS with MR, our approach enables the identification of both well-characterized and novel

exposures that may have been overlooked in pre-specified analyses. For example, while some of

the MR-validated associations we identified align with previously known findings, such as the

protective role of educational attainment for CAD, our ExWAS uncovered associations with

behavioral traits, such as age at first sexual intercourse, which were not previously highlighted in

systematic MR investigations of CAD and T2D.

Our study systematically quantifies the low concordance between ExWAS findings and MR-

validated associations for CAD and T2D. This underscores and estimates the extent of false

positives in observational analyses, particularly for lifestyle and dietary factors, where we

observed minimal potentially genetic causative evidence. Notably, our results demonstrate

disproportionately high rates among education-related factors relative to other domains, a finding

consistent with prior MR studies but newly quantified across a broad exposomic landscape. This

systematic perspective reveals a disjoint between correlational and potentially causative

structures for many lifestyle factors, while providing updated evidence for social and education-

related exposures.

However, while our study highlights the potential of agnostic discovery-driven methods, it also

underscores the challenges in establishing potentially causative evidence for many exposures,

particularly lifestyle and dietary factors when using genetics as the instrument.

Among many lifestyle dietary factors, none survived MR scrutiny for potentially causative

relationships in our data. While some factors may not have been detected in MR analyses due to

low power, the overall pattern suggests a disjoint between correlational and potentially causative

structures of different types of factors associated with these two major disease phenotypes. With

rare exceptions, detected associations with lifestyle factors (most prominent among them being

dietary factors) may reflect plain correlations rather than potentially causative relationships.

This is congruent with the evidence that while millions of published articles present or discuss

observational associations with specific dietary and other related lifestyle factors, the far smaller

corpus of interventional randomized trials relevant to direct effect of these exposures yields

mostly null results[30, 31]. Moreover, other investigators using MR methods in large genetic

consortia for CAD, T2D and ischemic stroke found no potentially causative evidence for any of

the dietary factors that they tested and found only one potentially causative association (with fat)

for heart failure[32].   One alternative model to consider includes the relationships between the

wide array of exposures interrogated. For example, one hypothesis is that dietary and other

lifestyle exposures are shaped by education-related factors, but do not by themselves cause

cardiometabolic disease. Other hypotheses may be built around the correlation of lifestyle

factors with weight which is a major causative risk factor for T2D and CAD (e.g., obesity may

coexist with specific lifestyle patterns or even shape lifestyle features, but without many of these

lifestyle features causing T2D and CAD by themselves). Future studies may test more complex

potentially causative patterns from diverse directed acyclic graphs and mediation patterns among

these multiple exposures.

Overall, we provide an overview of the currently measured exposomic risk architecture shared

between T2D and CAD. Specifically, we can explain up to ~8% and ~12% in CAD and T2D

outcomes respectively with current biobank scale measurements leaving much to be ascribed to

newer measurements and instruments[4]. However, we find low concordance between MR and

ExWAS for both CAD and T2D. Restricting the variables to just MR-implicated findings, we

find that these variables explain 2 and 3% of variation, in CAD and T2D respectively. Although

a crude comparison, heritability ranges have been reported from 30-50%, leaving much to be

explained with the exposures considered in this study. The main reason for lack of concordance

includes confounding with ExWAS findings reflecting false positive associations. Additionally,

we suggest potential classes of exposures to assay for a larger number of participants and with

greater resolution in the future. For example, the physical-chemical exposome (e.g., chemical

pollutants, etc.) needs to be more comprehensively assayed at biobank scale[33].

We also note limitations of our study. First, given the small sample size of individuals of non-

White ethnic groups, we expect the results to have high uncertainty in MR approaches; therefore,

we have excluded other ethnic groups from our analysis. A systematic review of MR studies for

CAD suggests that direction of effects of modifiable risk factors tends to be similar in different

ethnic groups, but differences in magnitude of effects may not be uncommon[34]. Second, MR

methods may also have bias and flaws, and they are not a perfect gold standard by any means.

MR relies on three core assumptions: that the genetic instruments are strongly associated with

the exposure (relevance), are independent of confounders of the exposure-outcome relationship

(independence), and influence the outcome exclusively through the exposure (exclusion

restriction). Violations of these assumptions can introduce bias and invalidate causal inferences.

Furthermore, MR strictly identifies associations that are consistent with lifelong genetically

influenced variations in exposure, and thus is particularly stringent in isolating causally robust

pathways. Its assumptions—including genetic variants having a stable, lifelong impact on

exposure, no pleiotropy, and no reverse causation—make MR an excellent choice primarily for

robust causal signals with large effect sizes. However, exposures of relevance may have small or

modest effects and may act only in specific periods of the lifespan.

In this study, many of the exposures analyzed are complex, distal variables within the exposome,

rather than molecular traits that may be more proximal to genetic variation. As such, the genetic

instruments used may be more susceptible to horizontal pleiotropy, where variants influence the

outcome through alternative pathways. While we employed sensitivity analyses—including MR-

Egger regression and the weighted median estimator—to assess the robustness of our findings

and detect potential pleiotropy, these methods have limited power, particularly in settings with

weak instruments or complex traits. Moreover, limited power may lead to false-negative

potentially causative[35] findings, but some biases may actually also create larger potentially

causative effects[36]. Third, it is unclear if we have addressed confounding for all domains of

exposures as there is no consensus on which adjustment factors to use for the different domains;

future work could look at optimizing the difference between observational and MR associations

as a function of covariate selection. Results may vary depending on what analytical choices are

made[37]. Another important limitation of this study is potential under-ascertainment of T2D, as

primary care data, medication use, and laboratory measures were not incorporated. These sources

present challenges for time-to-event modeling, including unclear diagnosis timing, reverse

causative relationships, and limited data availability. To maintain consistency and ensure

temporal clarity, both prevalent and incident T2D analyses were based solely on hospital records

and self-reported diagnoses. Future studies integrating comprehensive primary care data may

improve case detection and further validate these findings. Additionally, while we adjusted for a

key socioeconomic variable, socioeconomic status is multifaceted, and residual confounding may

remain.

Finally, prevalent disease outcome data coupled with genetic variant data as instrumental

variables for MR may suffice to identify exposures, especially when working with cohorts

without sufficient longitudinal outcome follow-up data. This is evidenced by the high degree of

commonality of MR-validated exposures between prevalent and incident disease ExWAS-

identified associations (85.7% [12 out of 14 total] for CAD and 87.5% [14 out 16 total] for T2D).

However, prevalent and incident outcome data may each have its sets of distinct biases.

In conclusion, MR-validated factors for CAD and T2D are far fewer than the rich range of

factors identified by ExWAS. Potentially causative relationships also seem more commonly

documented for educational factors, while lifestyle factors have many ExWAS signals, but these

rarely have potentially causative MR support. Additional similar analyses in more cohorts with

relevant information would be useful to expand evidence on the robustness of these findings.

Online Methods

Study population

The UK Biobank cohort is a prospective cohort including over 500,000 participants of ages 40-

69 during recruitment from 2006-2010[38]. Differences between the UK Biobank cohort

individuals and the general UK population were studied by Fry et al. in order to better

understand sampling uncertainty[39]. Their study suggested that nonparticipants are more likely to

be male, younger, and live in more socioeconomically deprived areas than UK Biobank

participants [39]. Information regarding how the UK Biobank data is maintained and validated can

be found at

https://biobank.ndph.ox.ac.uk/~bbdatan/Data_cleaning_overall_doc_showcase_v1.pdf.

The National Research Ethics Service Committee North West Multi-Centre Haydock has

approved the UKB cohort research and written informed consent to participate in the study was

provided by all participants[40]. We analyzed $n = 472,240$ European White individuals (out of total

$N = 502,628$ participants in the cohort). The other top three ethnicities represented (by sample

size) in the UKB cohort (Indian, Caribbean, and African) were low in sample size (Indian n =

5,951, Caribbean n = 4,517, African n = 3,394) when integrating exposure data and therefore we

may not have adequate power for detection of associations. Given the small sample size of

individuals of non-white ethnic groups, we expect the results to have high uncertainty and thus

the correlations would be weaker; therefore, we have excluded other ethnic groups from our

analysis. Approval for the use of this data was approved by the UK Biobank (project ID: 22881).

The Harvard internal review board (IRB) deemed the research as non-human subjects research

(IRB: IRB16-2145). Formal consent was obtained by the UK Biobank (https://

biobank.ctsu.ox.ac.uk/ukb/ukb/docs/Consent.pdf).

Disease outcomes

In our study, we investigate two major cardiometabolic incident outcomes in the UKB, including

coronary artery disease (CAD) and type 2 diabetes as ascertained by using ICD-10 and self-

reported disease status information (Supplemental Table 16). We identify cases of CAD and T2D

as ones that occur after the baseline visit (incident cases) while considering individuals who have

the disease at baseline (prevalent cases) separately. Primary care data, prescription records, and

laboratory measures (e.g., HbA1c) were not included due to limitations in event timing, risk of

reverse causative relationships, and incomplete data coverage.

Categorizing ecosystems, lifestyle, social, and physical-chemical exposures

We considered 538 total variables of which we had an adequate number of complete cases to

investigate 495 "exposures" that can be categorized (as per Vermeulen et al.[4]) as ecosystems,

lifestyle, social, and physical-chemical factors throughout the paper. We utilize the data for these

exposures collected during the participants' baseline visits (2006-2010). These 495 exposures

spanned 17 UK Biobank-defined categories (e.g., education, smoking, greenspace and coastal

proximity, sun exposure, estimated nutrients yesterday) (Extended Data Table 1).

We averaged quantitative factors (e.g., infectious antigens [25 exposures]) across measurements

from multiple visits. For exposures that did not have many observations in subsequent instances,

we used only the data from the baseline visit (first instance of measurement collected during

2006–2010) (e.g., environmental factors from the estimated nutrients yesterday category). We

also performed rank-based inverse normal transformation (INT) of these factors, as was

suggested by Millard et al.[41], to address non-normality and improve the comparability of effect

estimates across exposures with differing distributions. This transformation ensures that all

continuous exposure variables are on the same scale, reduces the influence of outliers and allows

for more robust association testing in data-driven epidemiological analyses such as ExWAS . For

categorical variables (which were also collected from multiple visits of a participant to the

assessment center), we used data from the baseline visit (first instance of measurement collected

during 2006–2010) as this contained the highest number of observations. Additionally,

categorical variables with multiple levels were converted to sets of binary variables where each

such variable indicates whether a participant has a given value of this variable (as was suggested

by Millard et al.[41]). Ordinal categorical environmental factor variables were analyzed by treating

such variables as continuous variables and real-valued quantitative environmental factor

variables were scaled.

Data-driven identification of exposure-disease associations

We used Cox proportional hazard models to associate each of the 495 factors and CAD

(coronary artery disease), T2D (type 2 diabetes) [individually], while adjusting for sex, age,

assessment center, ethnicity, average total household income after tax, and 40 genetic principal

components (PCs) (computed and provided by the UK Biobank). PCs were included to account

for subtle population structure that could correlate with non-genetic exposures, minimizing

potential confounding due to population stratification.We adjust resulting p-values of

associations for multiple comparisons using the false discovery rate (FDR) approach[42]. We

report hazard ratios and FDR-adjusted *p*-values for the associations. Additionally, we run logistic

regression to enhance comparison for downstream analyses (i.e. MR).

Concordance between observational and MR-based exposure-disease associations

For each exposure-outcome pair, we conducted two-sample Mendelian randomization (MR)

analyses using the full suite of methods available in the TwoSampleMR R package [27]. The

primary method used was inverse variance weighted (IVW) MR, which assumes that all genetic

instruments are valid and provides the most precise estimates when this assumption holds. To

assess the robustness of our findings and evaluate the potential for horizontal pleiotropy, we

additionally applied complementary sensitivity analyses including MR-Egger regression,

weighted median estimator, simple mode and weighted mode estimators, and the Wald ratio

methods. These analyses enable us to assess consistency of effect direction and identify potential

violations of MR assumptions, particularly related to horizontal pleiotropy.

Furthermore, we performed MR for all associations for which there were available GWAS

summary statistics for the corresponding exposures (n = 292 exposures). We recognize that

some exposures, including air pollution, are measured on the geospatial level, and it does not

make sense to execute a MR on these[43]. We exclude such exposures and related findings from

our MR analysis. Additionally, it is important to note that educational attainment was analyzed

using different categories to align ExWAS exposures with GWAS summary data available for

the UK Biobank and public resources (e.g., Neale lab[26]). We derived these measures from the

PHESANT pipeline[41], ensuring consistency across observational and MR analyses. Moreover,

we used GWAS summary statistics for identifying instruments for each exposure from GWAS

summary statistics generated by the Neale Lab[26] and MRC IEU OpenGWAS [27] and made freely

accessible on the MR-Base platform [27]. We use the TwoSampleMR R package [27] to test the

identified instruments for each exposure for each of the two outcomes using summary statistics

of GWAS from the FinnGEN cohort (https://www.finngen.fi/en). To the best of our knowledge,

there is no overlap between cohort utilized by the Neale Lab[26] and MRC IEU OpenGWAS [27] to

derive GWAS summary statistics for a given exposure and the FinnGen cohort[18]. Summary

statistics from FinnGen were adjusted by sex, age, genotyping batch, and ten principal

components. We utilize the FinnGen sample of 218,957 participants to validate our findings via

two sample MR. Among these participants, 29,193 have developed T2D and 21,012 have

developed CAD [18]. Additionally, we used the following thresholds to ascertain genetic

instruments including p-value $< 5 \times 10^{-8}$, linkage disequilibrium (LD) R^2: 0.001, and clumping

distance of 10000 kb. We report potentially causative genetic-based estimates computed using

the inverse-variance weighting (IVW) method between each exposure-disease pair.

As a sensitivity analysis to assess potential reverse causative relationship, we performed reverse

Mendelian randomization (MR), using genetic instruments for type 2 diabetes (T2D) and

coronary artery disease (CAD) as exposures and the previously analyzed exposures as outcomes.

Reverse MR was conducted using two-sample inverse variance weighted (IVW) MR as the

primary approach. These analyses were not interpreted as evidence of causative effects but

served to assess the possibility that genetic liability for disease influenced the exposures.

Assessing the variance of CAD and T2D explained by exposomic and demographic risk

factors

We contextualize the variance (computed as Nagelkerke R^2) explained by validated exposomic

factors with demographic risk factors (Extended Data Table 5). Briefly, for each incident disease

we run two major sets of models: a) regressing incident disease status on demographic factors

and b) regressing disease status on demographic factors and exposomic factors. Additionally, we

ran sets of models where we constrained the space of exposomic factors to just MR-validated

exposures in addition to separate sets of models considering a wider array of 196 exposures

measured (albeit limited by number of complete cases), cumulatively leading to the specification

of models A-I (Extended Data Table 5).

We estimated the variance (computed as Nagelkerke R^2) explained by validated exposomic

factors with demographic risk factors (Extended Data Table 6). Briefly, for each disease we run

two major sets of logistic regression models: a) regressing disease status on demographic,

genetic risk score (for CAD and T2D) and clinical (BMI for T2D) factors and b) regressing

disease status on demographic factors and exposomic factors. Additionally, we ran sets of

models where we constrained the space of exposomic factors to just MR-validated ones in

addition to separate sets of models considering a wider array of 196 exposures measured (albeit

limited by number of complete cases). Finally, we ran logistic regression models serially adding

exposomic factors to the previous model in order of increasing FDR-corrected p-values until we

ran the final model including the full wider array of exposomic factors (in addition to the

baseline demographic and clinical risk factors). Additionally, pairwise correlations between

categorical and numeric exposure variables were computed using the "polycor" R package.

Code availability

We performed all analyses using R version 3.6.1. We made our code accessible at

(https://github.com/stejat98/CardiometabolicExWASMR).

Data availability

The data from the UK Biobank that support the findings of this study are available upon

application (https://www.ukbiobank.ac.uk/register-apply/).

Figure Legends

Figure 1. Schematic overview of analysis of exposome-wide risk architecture of T2D and CAD. This schematic diagram depicts our analytic workflow. (a) Exposure-wide association study (ExWAS) was performed to discover exposure-disease associations for T2D and CAD in the UK Biobank. (b) Two Sample bi-directional mendelian randomization (MR) was performed to replicate ExWAS-identified associations and test for potentially causative relationships. Briefly, genetic instruments for exposures were identified from the UK Biobank sample and tested on the FinnGen sample with respect to CAD and T2D

Figure 2. Exposomic architecture of CAD. This multipanel figure depicts exposome-wide findings for CAD. (a) **Volcano plot of ExWAS associations for CAD.** We visualize the FDR-corrected p-values on the negative log 10 scale versus hazard ratios of exposures. The red color indicates FDR < 0.05 significant associations and blue color indicates FDR > 0.05 associations. Top five associations are labeled. (b) **Distribution of hazard ratios of ExWAS associations for CAD and T2D stratified by significance.** The distribution of the hazard ratios (HR) computed from the absolute value of the regression beta coefficients for CAD and T2D are depicted with empirical cumulative distribution function plots and are colored by ExWAS significance. The distribution of HRs for exposures that are ExWAS FDR less than 0.05 significant are depicted in red. The distribution of HRs for exposures that are ExWAS FDR greater than 0.05 are depicted in blue. For the underlying dataset, n=472,240. (c) **Distribution of p-values of ExWAS and MR associations for CAD and T2D.** The distributions of all p-values derived from ExWAS for CAD and T2D are depicted with empirical cumulative distribution function plots. For the underlying dataset, n=472,240. (d) **Association size of CAD versus T2D for 495 exposures.**

The hazard ratios (HR) of CAD versus T2D for 495 different exposures. Exposures that are FDR

less than 5×10^{-4} significant for both CAD and T2D deviate and have HRs either less than 0.5 or

greater than 2 for either disease are labeled and shown with error bars corresponding to the 95%

confidence intervals for both CAD and T2D. The black dashed line on the scatterplot represents

a linear regression line. For the underlying dataset, n=472,240.

Figure 3. ExWAS association size versus MR causative estimate for CAD. The odds ratios

(OR) computed from ExWAS versus the OR computed from MR for 260 different exposures in

association with CAD. The blue color indicates evidence for potential reverse causative

relationship and the red color indicates no evidence for potential reverse causative relationships.

Triangles indicate associations with significant MR p-values. Horizontal and vertical black lines

demarcate OR of 1 (null association). For the underlying datasets, n=472,240 for the UK

Biobank sample and n = 218,957 for the FinnGen sample.

Figure 4. Visualizing the change in variance explained by step-wise addition of exposomic

variables to baseline model of demographic and clinical risk factors compared to exposure-

specific univariate models. We visualize the Nagelkerke R^2 of each model constructed by step-

wise addition of exposomic factors in descending order of FDR significance to the original

baseline model comprised of demographic, genetic risk score (for CAD and T2D) and clinical

(BMI for T2D) factors. Black dot indicates variance explained by the baseline model. Top

exposures (by FDR) for the multivariable analysis are labeled. We also overlay the univariate

(exposure) model results. We visualize the cumulative exposure-specific Nagelkerke R^2 as we

iterate through each univariate (exposure) model in descending order of FDR significance with

smaller point size for both CAD and T2D, separately.

Tables

Variable (units)	Label	Mean (SD) or Percentage
Age (years)	Age	56.76 (8.028)
Sex (%)	Female	54.48
Sex (%)	Male	45.52
Avg. household income (%)	< 18,000	19.07
Avg. household income (%)	18,000 to 30,999	21.74
Avg. household income (%)	31,000 to 51,999	22.43
Avg. household income (%)	52,000 to 100,000	17.55
Avg. household income (%)	> 100,000	4.65
HbA1c (mmol/mol)	HbA1c	35.96 (6.517)
Systolic pressure (mm Hg)	Systolic pressure	139.9 (19.68)
Diastolic pressure (mm Hg)	Diastolic pressure	82.16 (10.68)
BMI (kg/m2)	BMI	27.41 (4.783)
LDL (mmol/L)	LDL	3.57 (0.87)
HDL (mmol/L)	HDL	1.45 (0.383)
Triglycerides (mmol/L)	Triglycerides	1.75 (1.02)
Total cholesterol (mmol/L)	Total cholesterol	5.71 (1.14)
Family history for CAD (%)	Family history for CAD	39.8
Family history for T2D (%)	Family history for T2D	17.1
Smoking (%)	Never	53.94
Smoking (%)	Previous	35.55
Smoking (%)	Current	10.5
Prevalent CAD (%)	Prevalent CAD	2.28
Prevalent T2D (%)	Prevalent T2D	3.66

Table 1. Sample baseline characteristics.

xposure	CAD MR OR	CAD MR p-value	CAD ExWAS OR	CAD ExWAS FDR	T2D MR OR	T2D MR p-value	T2D ExWAS OR	T2D ExWAS FDR
ualifications: A vels/AS levels or uivalent	0.386	1.12E-05	0.705	1.92E-26	0.419	7.02E-04	0.627	2.97E-
ualifications: ollege or niversity degree	0.566	3.51E-05	0.657	3.57E-44	0.406	5.49E-10	0.54	2.64E-1
ge first had xual intercourse	0.749	4.08E-05	0.821	3.44E-45	0.626	1.50E-10	0.826	6.85E-
ge completed full ne education	0.598	3.59E-04	0.896	9.84E-11	0.551	1.55E-03	0.876	3.42E-
ualifications: one of the above	1.825	1.05E-02	1.51	6.22E-49	2.591	2.68E-04	1.81	3.92E-1
ualifications: O vels/GCSEs or uivalent	0.398	1.38E-02	0.797	8.78E-18	0.21	7.12E-04	0.731	2.40E-
sual walking ce	0.592	1.81E-02	0.74	7.65E-115	0.233	1.34E-06	0.625	< 1E-163

698   **Table 2. Exposomic variables significant in both MR and ExWAS and in both phenotypes**.

Table Legends

Table 1. Sample baseline characteristics. This table shows the mean (and standard deviation)

or percentage of key clinical and demographic variables for the sample of White UKB

participants that were primarily used for our analysis including age, sex, average household

income, HbA1c, systolic blood pressure, diastolic blood pressure, BMI, LDL, HDL,

triglycerides, total cholesterol, family history for CAD, family history for T2D, prevalent CAD,

prevalent T2D, and smoking history.

Table 2. Exposomic variables significant in both MR and ExWAS and in both phenotypes.

This table includes MR-derived odds ratios (ORs) and p-values and ExWAS-derived ORs and

FDR-corrected p-values for the seven exposomic variables we found significant in both MR and

ExWAS in both phenotypes.

Supplemental Table Legends

Supplemental Table 1. Exposomic variables significant in ExWAS for incident CAD. This table includes hazard ratios (HR), 95% confidence intervals (CI), FDR-corrected p-values and C-statistic from each cox proportional hazards model to systematically associate each exposure with time to event (CAD).

Supplemental Table 2. Exposomic variables significant in ExWAS for incident T2D. This table includes hazard ratios (HR), 95% confidence intervals (CI), FDR-corrected p-values and C-statistic from each cox proportional hazards model to systematically associate each exposure with time to event (T2D).

Supplemental Table 3. Exposomic variables significant in ExWAS for prevalent CAD. This table includes odds ratio (OR), 95% confidence intervals (CI), and FDR-corrected p-values from each logistic regression model to systematically associate each exposure with prevalent CAD.

Supplemental Table 4. Exposomic variables significant in ExWAS for prevalent T2D. This table includes odds ratio (OR), 95% confidence intervals (CI), and FDR-corrected p-values from each logistic regression model to systematically associate each exposure with prevalent T2D.

Supplemental Table 5. Prevalent CAD versus incident CAD ExWAS significance classification table. This table includes estimates of false positives, false negatives, true positives, true negatives when comparing prevalent CAD ExWAS (our "diagnostic test") with incident CAD ExWAS findings. True Positives were ascertained as associations that had prevalent CAD and incident CAD effects concordant in direction in addition to FDR < 0.05 significance.

Supplemental Table 6. Prevalent T2D versus incident T2D ExWAS significance classification table. This table includes estimates of false positives, false negatives, true positives, true

negatives when comparing prevalent T2D ExWAS (our "diagnostic test") with incident T2D ExWAS

findings. True Positives were ascertained as associations that had observational and MR beta

estimates concordant in direction in addition to FDR < 0.05 significance.

Supplemental Table 7. Exposomic variable associations for CAD from bi-directional two sample

MR. This table includes beta estimates and p-values for forward and reverse mendelian randomization

(MR) analysis associating each exposure (using GWAS summary statistics derived from the UK Biobank)

with CAD (using GWAS summary statistics derived from FinnGen).

752
753
754   **Supplemental Table 8. Exposomic variable associations for T2D from bi-directional two sample**

755   **MR.** This table includes beta estimates and p-values for forward and reverse mendelian randomization

756   (MR) analysis associating each exposure (using GWAS summary statistics derived from the UK Biobank)

757   with T2D (using GWAS summary statistics derived from FinnGen).

758
759   **Supplemental Table 9. Exposomic variable associations for CAD across MR methods.** This table

760   includes beta estimates and p-values for forward and reverse mendelian randomization (MR) analysis

761   associating each exposure (using GWAS summary statistics derived from the UK Biobank) with CAD

762   (using GWAS summary statistics derived from FinnGen) across the full suite of MR methods tested (MR

763   Egger, weighted median, inverse variance weighted, simple mode, and weighted mode).

764
765
766   **Supplemental Table 10. Exposomic variable associations for T2D across MR methods.** This table

767   includes beta estimates and p-values for forward and reverse mendelian randomization (MR) analysis

768   associating each exposure (using GWAS summary statistics derived from the UK Biobank) with T2D

769   (using GWAS summary statistics derived from FinnGen) across the full suite of MR methods tested (MR

770   Egger, weighted median, inverse variance weighted, simple mode, and weighted mode).

771
772
Supplemental Table 11. Observational versus MR significance classification table for lifestyle

factors in association with CAD. This table includes estimates of false positives, false negatives,

true positives, true negatives when comparing ExWAS findings with respect to gold-standard

causal estimates obtained from MR for CAD. True Positives were ascertained as associations that

had observational and MR beta estimates concordant in direction in addition to the

aforementioned observational and MR significance criteria.

779
780
Supplemental Table 12. Observational versus MR significance classification table for lifestyle

factors in association with T2D. This table includes estimates of false positives, false negatives,

true positives, true negatives when comparing ExWAS findings with respect to gold-standard

causal estimates obtained from MR for T2D. True Positives were ascertained as associations that

had observational and MR beta estimates concordant in direction in addition to the

aforementioned observational and MR significance criteria.

Supplemental Table 13. Observational versus MR significance classification table for social

factors in association with CAD. This table includes estimates of false positives, false negatives,

true positives, true negatives when comparing ExWAS findings with respect to gold-standard

causal estimates obtained from MR for CAD. True Positives were ascertained as associations that

had observational and MR beta estimates concordant in direction in addition to the

aforementioned observational and MR significance criteria.

794
795
Supplemental Table 14. Observational versus MR significance classification table for social

factors in association with T2D. This table includes estimates of false positives, false negatives,

true positives, true negatives when comparing ExWAS findings with respect to gold-standard

causal estimates obtained from MR for T2D. True Positives were ascertained as associations that

had observational and MR beta estimates concordant in direction in addition to the

aforementioned observational and MR significance criteria.

802
Supplemental Table 15. Effects of established clinical risk factors. This table reports the odds ratios

(OR) with 95% confidence intervals (CI) and p-values of established clinical risk factors from the

literature including age, sex, family history, BMI, systolic blood pressure, diastolic blood pressure,

HbA1c, LDL, HDL, triglycerides, total cholesterol, and smoking history.

Supplemental Table 16. Ascertainment of Cardiometabolic Outcomes. This table shows the ICD-10

and self-reported disease status codes used to identify coronary artery disease (CAD) and type 2 diabetes

(T2D) disease status.

Extended Data Table Legends

Extended Data Table 1. Exposure variable breakdown. This table shows the exposure

categories as defined by Vermeulen et al.[3] as well as the UKB, the number of variables within

each category, the class of the variable category (i.e. whether the category contains biomarker

(B), geographic (G) or questionnaire (Q) variables), and some example variables within each

category.

Extended Data Table 2. ExWAS-identified enrichment of exposomic categories (as defined

by Vermeulen et al.[3]) for each disease. This table shows the breakdown of the percentage of

ExWAS-identified FDR < 0.05 associations by exposure categories as defined by Vermeulen et

al for CAD and T2D.

Extended Data Table 3. Observational versus MR significance classification table for CAD.

This table includes estimates of false positives, false negatives, true positives, true negatives

when comparing ExWAS findings with respect to gold-standard causative estimates obtained

from MR for CAD. True Positives were ascertained as associations that had observational and

MR beta estimates concordant in direction in addition to the aforementioned observational and

MR significance criteria.

Extended Data Table 4. Observational versus MR significance classification table for T2D.

This table includes estimates of false positives, false negatives, true positives, true negatives

when comparing ExWAS findings with respect to gold-standard causative estimates obtained

from MR for T2D. True Positives were ascertained as associations that had observational and

MR beta estimates concordant in direction in addition to the aforementioned observational and

MR significance criteria.

Extended Data Table 5. Breakdown of variance (R2) explained by exposomic and MR-

validated exposomic factors with demographic risk factors. This table contextualizes the

variance (computed as R^2) explained by exposomic, MR-validated exposomic factors with

demographic risk factors. Delta R^2 is computed as the difference between R^2 computed from full

model and R^2 computed model only including demographic and clinical risk factor covariates.

Extended Data Table 6. Baseline demographic breakdown of samples used for models A-I.

This table shows the breakdown of the mean and shows the mean or percentage of key

demographic variables for the samples of White UKB participants that were primarily used for

our models A-I including age, sex, average household income, and assessment center.

References

1. Ross, S. *et al.* Mendelian randomization analysis supports the causal role of dysglycaemia and diabetes in the risk of coronary artery disease. *Eur. Heart J.* **36**, 1454–1462 (2015).

2. Münzel, T., Sørensen, M., Hahad, O., Nieuwenhuijsen, M. & Daiber, A. The contribution of the exposome to the burden of cardiovascular disease. *Nat. Rev. Cardiol.* **20**, 651–669 (2023).

3. Miller, G. W. & Banbury Exposomics Consortium. Integrating exposomics into biomedicine. *Science* **388**, 356–358 (2025).

4. Vermeulen, R., Schymanski, E. L., Barabási, A.-L. & Miller, G. W. The exposome and health: Where chemistry meets biology. *Science* **367**, 392–396 (2020).

5. Ioannidis, J. P. A. Why most published research findings are false. *PLoS Med* **2**, e124 (2005).

6. Munafò, M. R. & Davey Smith, G. Robust research needs many lines of evidence. *Nature* **553**, 399–401 (2018).

7. Ioannidis, J. P. A., Loy, E. Y., Poulton, R. & Chia, K. S. Researching genetic versus nongenetic determinants of disease: a comparison and proposed unification. *Sci. Transl. Med.* **1**, 7ps8 (2009).

8. Patel, C. J. & Ioannidis, J. P. A. Studying the elusive environment in large scale. *JAMA* **311**, 2173–2174 (2014).

9. Ioannidis, J. P. A. Exposure-wide epidemiology: revisiting Bradford Hill. *Stat. Med.* **35**, 1749–1762 (2016).

10. Fedak, K. M., Bernal, A., Capshaw, Z. A. & Gross, S. Applying the Bradford Hill criteria in the 21st century: how data integration has changed causal inference in molecular

epidemiology. *Emerg. Themes Epidemiol.* **12**, 14 (2015).

11. Manrai, A. K., Ioannidis, J. P. A. & Patel, C. J. Signals Among Signals: Prioritizing

Nongenetic Associations in Massive Data Sets. *Am. J. Epidemiol.* **188**, 846–850 (2019).

12. Wild, C. P. The exposome: from concept to utility. *Int. J. Epidemiol.* **41**, 24–32 (2012).

13. Katikireddi, S. V., Green, M. J., Taylor, A. E., Davey Smith, G. & Munafò, M. R.

Assessing causal relationships using genetic proxies for exposures: an introduction to

Mendelian randomization. *Addiction* **113**, 764–774 (2018).

14. DeCicca, P., Kenkel, D. & Lovenheim, M. F. The Economics of Tobacco Regulation: A

Comprehensive Review. *J Econ Lit* **60**, 883–970 (2022).

15. Davies, N. M., Holmes, M. V. & Davey Smith, G. Reading Mendelian randomisation

studies: a guide, glossary, and checklist for clinicians. *BMJ* k601 (2018).

16. Emdin, C. A., Khera, A. V. & Kathiresan, S. Mendelian Randomization. *JAMA* **318**, 1925

(2017).

17. VanderWeele, T. J., Tchetgen Tchetgen, E. J., Cornelis, M. & Kraft, P. Methodological

challenges in mendelian randomization. *Epidemiology* **25**, 427–435 (2014).

18. Kurki, M. I. *et al.* FinnGen provides genetic insights from a well-phenotyped isolated

population. *Nature* **613**, 508–518 (2023).

19. Meigs, J. B. *et al.* Genotype score in addition to common risk factors for prediction of type

2 diabetes. *N. Engl. J. Med.* **359**, 2208–2219 (2008).

20. He, Y. *et al.* Comparisons of polyexposure, polygenic, and clinical risk scores in risk

prediction of type 2 diabetes. *Diabetes Care* **44**, 935–943 (2021).

21. Janiaud, P. *et al.* Validity of observational evidence on putative risk and protective factors:

appraisal of 3744 meta-analyses on 57 topics. *BMC Med.* **19**, 157 (2021).

22. Tillmann, T. *et al.* Education and coronary heart disease: mendelian randomisation study.

BMJ **358**, j3542 (2017).

23. Mills, M. C. *et al.* Identification of 371 genetic variants for age at first sex and birth linked

to externalising behaviour. *Nat. Hum. Behav.* **5**, 1717–1730 (2021).

24. Knowler, W. C. *et al.* Reduction in the incidence of type 2 diabetes with lifestyle

intervention or metformin. *N. Engl. J. Med.* **346**, 393–403 (2002).

25. Levin, M. G. *et al.* Genome-wide assessment of pleiotropy across >1000 traits from global

biobanks. *medRxiv* (2025) doi:10.1101/2025.04.18.25326074.

26. UkBiobankGwas, G.-. N. Nealelab/UK_Biobank_GWAS: Overview of the data QC, code,

and GWAS summary output from the 2017 UK Biobank data release. *GitHub*.

27. Hemani, G. *et al.* The MR-Base platform supports systematic causal inference across the

human phenome. *Elife* **7**, (2018).

28. Georgiou, A. N. *et al.* Appraising the causal role of risk factors in coronary artery disease

and stroke: A systematic review of Mendelian randomization studies. *J. Am. Heart Assoc.*

12, e029040 (2023).

29. Yuan, S. & Larsson, S. C. An atlas on risk factors for type 2 diabetes: a wide-angled

Mendelian randomisation study. *Diabetologia* **63**, 2359–2371 (2020).

30. Trepanowski, J. F. & Ioannidis, J. P. A. Perspective: Limiting dependence on

nonrandomized studies and improving randomized trials in human nutrition research: Why

and how. *Adv. Nutr.* **9**, 367–377 (2018).

31. Young, S. S. & Karr, A. Deming, data and observational studies. *Signif. (Oxf.)* **8**, 116–120

(2011).

32. Niu, Y.-Y., Aierken, A. & Feng, L. Unraveling the link between dietary factors and

cardiovascular metabolic diseases: Insights from a two-sample Mendelian Randomization

investigation. *Heart Lung* **63**, 72–77 (2024).

33. Patel, C. J., Bhattacharya, J. & Butte, A. J. An Environment-Wide Association Study

(EWAS) on type 2 diabetes mellitus. *PLoS One* **5**, e10746 (2010).

34. Silva, S., Fatumo, S. & Nitsch, D. Mendelian randomization studies on coronary artery

disease: a systematic review and meta-analysis. *Syst. Rev.* **13**, 29 (2024).

35. Brion, M.-J. A., Shakhbazov, K. & Visscher, P. M. Calculating statistical power in

Mendelian randomization studies. *Int. J. Epidemiol.* **42**, 1497–1501 (2013).

36. Burgess, S., Thompson, S. G. & CRP CHD Genetics Collaboration. Avoiding bias from

weak instruments in Mendelian randomization studies. *Int. J. Epidemiol.* **40**, 755–764

(2011).

37. Klau, S., Hoffmann, S., Patel, C. J., Ioannidis, J. P. & Boulesteix, A.-L. Examining the

robustness of observational associations to model, measurement and sampling uncertainty

with the vibration of effects framework. *Int. J. Epidemiol.* **50**, 266–278 (2021).

38. Sudlow, C. *et al.* UK biobank: an open access resource for identifying the causes of a wide

range of complex diseases of middle and old age. *PLoS Med.* **12**, e1001779 (2015).

39. Fry, A. *et al.* Comparison of sociodemographic and health-related characteristics of UK

Biobank participants with those of the general population. *Am. J. Epidemiol.* **186**, 1026–

1034 (2017).

40. Armstrong, J. *et al.* Dynamic linkage of COVID-19 test results between Public Health

England's Second Generation Surveillance System and UK Biobank. *Microb. Genom.* **6**,

(2020).

41. Millard, L. A. C., Davies, N. M., Gaunt, T. R., Davey Smith, G. & Tilling, K. Software

Application Profile: PHESANT: a tool for performing automated phenome scans in UK

Biobank. *Int. J. Epidemiol.* **47**, 29–35 (2018).

42. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Series B Stat. Methodol.* **57**, 289–

300 (1995).

43. Au Yeung, S. L. & Gill, D. Concerns over using the Mendelian randomization design to

investigate the effect of air pollution. *Sci. Total Environ.* **917**, 170474 (2024).

Exposomic Risk for Cardiometabolic Disease

a. ExWAS

b. Causal MR

Discovery of **exposure-disease associations**

Exposure → ? → Disease

Ecosystems
Lifestyle
Physical-Chemical
Social

CAD

T2D

biobank uk

Total

ExWAS FDR < 0.05

MR p-value < 0.05
& no reverse causality

MR & ExWAS associations
concordant in direction

Replication and **causality** testing of **identified associations**

MR OR

ExWAS OR

MRBASE

biobank uk

FINNGEN