Sex difference in incidence of bipolar and anxiety disorders: findings from the Global Burden of Disease Study 2021

Xingke Zhu[1*], Qing Lv[1*]

[1] School of Pharmacy, Hubei University of Chinese Medicine, Wuhan, China.

[*]**Corresponding author:** Xingke Zhu (E-mail: krickzxk@163.com); Qing Lv (E-mail: lqql1231@163.com)

Author contributions: XZ and QL had full access to all the data in the study and will take responsibility for the integrity of the data, the accuracy of the data analysis and study concept and design. XZ contributed to the acquisition, analysis and interpretation of data. XZ and QL contributed to drafting of the manuscript. XZ contributed to the critical revision of the manuscript and supervision of the study. All authors read and approved the final manuscript.

Conflicts of interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Acknowledgments: We highly praise the collaborators of the 2021 Global Burden of Disease study for their extremely comprehensive analysis of various diseases on a global scale.

Compliance with Ethics Requirements: For Global Burden of Disease study, a waiver of informed consent was reviewed and approved by the Institutional Review Board of the University of Washington. All the information about ethical standards is available through the official website (https://www.healthdata.org/research-analysis/gbd).

Ethics approval and consent to participate: Not applicable.

Running Title: Sex difference of bipolar and anxiety disorders.

Abstract

Background: Bipolar disorder and anxiety disorders are two prominent mental disorders that represent a significant global health challenge.

Objective: The Global Burden of Disease Study 2021 (GBD 2021) was employed to evaluate sex differences in the incidence of bipolar disorder (BD) and anxiety disorder (ANX) globally by year, age, and socioeconomic status.

Method: We estimated sex-specific incidence of BD and ANX from GBD 2021 globally and in 204 countries and territories from 1990 to 2021. The sociodemographic index (SDI) was used to gauge national socioeconomic development and the Health Organization (WHO) region was used as a division of regions. Differences in age-standardized incidence rates (ASRs) by sex (absolute and relative) and risk ratios (95% confidence interval) were calculated annually and by age. Annual percent change (APC) was calculated by joinpoint regression modeling and linear regression analyses were performed to explore the socioeconomic factors associated with sex differences in incidence.

Results: The absolute and relative sex difference in ASRs of BD showed a slight declining trend during 1990 and 2021, with absolute difference decreasing from 2.50 to 1.83, and relative difference decreasing from 1.08 to 1.06; The absolute and relative sex difference in ASRs of ANX showed an increasing trend during 1990 and 2021, with absolute difference increasing from 170.02 to 208.08, and relative difference increasing from 1.35 to 1.36. Worldwide, females had a higher risk of BD and ANX than males in 1990and 2021. The highest Risk ratios of BD and ANX were observed in the European Region in 2021.The greatest relative sex difference of BD was 1.09 in the age group of 30-34. The greatest relative difference of ANX was 1.51 in the age group of 20-24. Relative sex differences of BD and ANX were significantly and positively correlated with SDI (BD, standardized $\beta = 0.27$ (95% CI, 0.22 to 0.33), $P < 0.001$; ANX, standardized $\beta = 0.80$ (95% CI, 0.47 to 1.14), $P < 0.001$).

Conclusions: Sex difference in the incidence of anxiety disorders and bipolar disorder have persisted worldwide over the past several decades, and the rates have consistently been higher among females than males. The sex difference in the global incidence of bipolar disorder has shown a slight improvement, but that in the global incidence of anxiety disorders has not been effectively mitigated. The sex difference is even more pronounced at younger ages and in more developed nations. The findings emphasize the significance of sex-specific health policies to reduce sex differences in the incidence of bipolar and anxiety disorders.

Keywords: Sex difference, Bipolar disorder, Anxiety disorder, Incidence, Trend, Age, Socioeconomic development

Introduction

Bipolar disorder and anxiety disorders are two prominent mental health conditions that impact millions of individuals across the globe [1, 2]. Anxiety disorders involve persistent and excessive worry, fear, or anxiety that interferes with an individual's ability to function effectively. Worldwide, 301.4 million people suffer from anxiety disorders [3]. Bipolar disorder, on the other hand, is characterized by extreme mood swings, including manic and depressive episodes, which can significantly impair daily functioning and quality of life. According to GBD 2019, anxiety disorders account for 39.5 million cases [3]. Both

conditions have been shown to have substantial socio-economic and health impacts. Emotional disorders, high psychosocial morbidity, and impulsive behaviors are the characteristics of bipolar disorder patients, which also increase the risk of suicide in bipolar disorder patients [4]. According to a study on the suicidal tendencies of bipolar disorder patients, there is a significantly increased risk of attempting and dying by suicide among bipolar disorder patients [5]. Similarly, deaths from non-natural causes had the highest natural mortality rate ratio among those with anxiety disorders, and anxiety disorders were linked to a significantly greater risk of death [6]. According to the total disability-adjusted life years (DALYs) attributed to mental disorders globally in 2019, anxiety disorders and bipolar disorder ranked second (22.9%) and fifth (6.8%), respectively [3].

Sex differences in the burden of mental health disorders have been a topic of significant research interest. Sex differences in the burden of BD and ANX have been an important issue of global public concern, although psychotherapy and medication are effective treatments for them. Epidemiologic and clinical research indicates that is about two times higher amongst female adolescents as amongst male adolescents, and among bipolar disorder patients, suicide attempts were 35% more common in women than in men [7, 8]. Anxiety problems also vary by gender. Women have persistently higher prevalence rates of anxiety disorders, and anxiety disorders are associated with a greater sickness burden in women than in men [9]. Biological factors such as genetics and sex hormone changes, as well as psychosocial ones such as life experiences, discrimination, and social and economic circumstances, play important roles in clarifying why men and women experience bipolar and anxiety differently [10, 11]. Thus, global patterns of sex differences in BD and ANX incidence have important implications for the development of sex-specific health policies to reduce BD and ANX.

The Global Burden of Disease Study 2021 offers an updated and comprehensive dataset to explore these sex differences in the incidence of bipolar disorder and anxiety disorders [12]. The present study investigates the sex differences in the incidence of bipolar disorder and anxiety disorders using the data provided by the Global Burden of Disease Study 2021. The findings are expected to contribute valuable insights into the epidemiology of these disorders and highlight areas where gender-specific strategies could enhance mental health outcomes globally.

Materials and methods

Data using

The Global Burden of Disease (GBD) 2021 study estimated the incidence of bipolar disorder (BD) and anxiety disorders (ANX) for both genders across 20 age groups and 204 countries and territories from 1990 onwards. This study utilizes the Bayesian meta-regression tool DisMod-MR 2.1 to integrate various data sources (including published literature, surveillance, surveys, hospital records, and clinical data) that meet specific inclusion criteria [13]. Incidence rates are age-standardized according to the GBD reference population [14]. The study employs a technique for propagating uncertainty consistent with prior methodologies, with final estimates derived from the mean of 1000 simulations, and 95% uncertainty intervals (UIs) based on the 25th and 975th percentiles. Definitions of BD and ANX adhere to DSM-IV-TR or ICD-10 criteria [15]. Data extracted include global age-standardized incidence rates (ASRs) per 100,000 population from 1990 to 2021; global and World Health Organization (WHO) regional incident cases and ASRs for 1990 and 2021;

and national ASRs for 204 countries and territories in 2021.

The sociodemographic index (SDI) is a composite measure reflecting a country's income per capita, average years of schooling, and fertility rate for females under 25 years of age, providing a comprehensive assessment of socioeconomic development [16]. The GBD study calculated the SDI as the geometric mean of these three variables using the Human Development Index methodology [17]. Quintile cutoff values for analysis were determined based on 2021 country-level SDI estimates. The SDI values range from 0 to 1, with higher values indicating more advanced socioeconomic development. In 2021, 204 countries and territories were categorized into five SDI groups: low SDI (0 < SDI < 0.466), low-middle SDI (0.466 ≤ SDI < 0.619), middle SDI (0.619 ≤ SDI < 0.712), high-middle SDI (0.712 ≤ SDI < 0.810), and high SDI (0.810 ≤ SDI < 1) [18]. Specifically, the low SDI group included 34 countries, the low-middle SDI group 42 countries, the middle SDI group 41 countries, the high-middle SDI group 48 countries, and the high SDI group 39 countries.

Statistical analyses

Global absolute (female minus male) and relative (female-to-male ratio) sex differences in incidence, along with risk ratios (RR) and their 95% confidence intervals (CIs), were calculated by year and age group. ASRs of BD and ANX were estimated using the Joinpoint Regression Model, a set of statistically linear models [19]. Annual percentage changes (APCs) and their 95% confidence intervals (CIs) were calculated. Age-standardized rates (ASRs) for males and females across 204 countries and territories were compared using the Mann–Whitney U test, accounting for multiple comparisons between socio-demographic index (SDI) groups [20]. Linear regression analyses assessed the relationship between absolute (female minus male) and relative (female-to-male ratio) sex differences in ASRs and SDI across these countries and territories. *P* values less than 0.05 were considered statistically significant. All statistical analyses were performed using R software (R Core Team, version 3.4.2, Vienna, Austria), Joinpoint software (version 4.9.1.0) from the Surveillance Research Program of the US National Cancer Institute. Zstats 1.0 (www.zstats.net) and GraphPad Prism 9.0 (San Diego, America).

Ethics Statement

All the information about ethical standards is available through the official website (https://www.healthdata.org/research-analysis/gbd).

Results

Sex difference in incidence of BD and ANX by year

Over the past 30 years, the ASRs for BD in females and males has not changed much globally: Average Annual Percent Change (AAPC) = 0.0% (95% CI, 0.0% to 0.0%), *P* < 0.001, for females, between 1990 and 2021; AAPC=0.1% (0.1% to 0.1%), *P* < 0.001, for males, between 1990 and 2021 (Fig. 1A). However, from 1990 to 2021, ASRs of BD in both sexes increased at the global level, with ASRs being 32.61 (95% UI: 27.45 to 38.60) in 1990 and 33.06 (95% UI: 27.80 to 39.60) in 2021 for females, and being 30.11 (95% UI: 25.41 to 35.57) in 1990 and 31.22 (95% UI: 26.27 to 37.34) in 2021 for males. The absolute and relative sex difference in ASRs of BD showed a slight declining trend during 1990 and 2021,

with absolute difference decreasing from 2.50 to 1.83 (Fig. 1B), and relative difference decreasing from 1.08 to 1.06 (Fig. 1C).

From 1990 to 2021, the ASRs of ANX in both females and males increased globally: AAPC = 0.7% (95% CI, 0.5% to 0.9%), $P < 0.001$, for females, between 1990 and 2021; AAPC=0.6% (0.6% to 0.7%), $P < 0.001$, for males, between 1990 and 2021 (Fig. 1E). The ASR for females was 649.32 (95% UI: 537.68 to 795.75) in 1990 and 784.53 (95% UI: 649.26 to 966.02) in 2021, while the ASR for males was 479.30 (95% UI: 402.19 to 579.75) and 576.46 in 2021 (95% UI: 486.27 to 701.84). More specifically, there was a marked upward trend in the ASR of ANX from 2019 to 2021: Annual Percent Change (APC) = 10.6% (95% CI, 8.9% to 12.4%), $P < 0.001$, for females; APC = 9.3% (95% CI, 8.9% to 9.7%), $P < 0.001$, for males. The absolute and relative sex difference in ASRs of ANX showed an increasing trend during 1990 and 2021, with absolute difference increasing from 170.02 to 208.08 (Fig. 1E), and relative difference increasing from 1.35 to 1.36 (Fig. 1F).

ASRs for BD and ANX were investigated in females and males globally and in WHO regions in 1990 and 2021. Worldwide, females had a higher risk of BD that males in 1990 (RR = 1.08; 95% CI, 1.08 to 1.09) and 2021 (RR = 1.06; 95% CI, 1.06 to 1.06). The highest RRs were observed in the European Region, being 1.19 (95% CI, 1.19 to 1.20) in 1990 and 1.20 (95% CI, 1.20 to 1.921) in 2021 (Table 1). For ANX, worldwide, females also exhibited a higher risk than males in 1990 (RR = 1.35; 95% CI, 1.34 to 1.37) and 2021 (RR = 1.36; 95% CI, 1.34 to 1.38). The highest RRs were observed in the Western Pacific Region, being 1.44 (95% CI, 1.42 to 1.47) in 1990 and in the European Region, being 1.46 ((95% CI, 1.45 to 1.50) in 2021 (Table 2).

Sex difference in incidence of BD and ANX by age

In 2021, the global incidence of BD among individuals under the age of 20 increased rapidly with age for both males and females. The highest incidence rates were 89.77 (95% UI: 65.12 to 121.90) for females and 83.75 (95% UI: 60.97 to 113.38) for males in the age group of 15-19 (Fig. 2A). The absolute sex difference in incidence rates increased rapidly with age for those under 20 years old, with the greatest absolute difference being 6.16 in the age group of 15-19 (Fig. 2B). The relative sex difference in incidence rates increased rapidly with age for those under 20 and over 80 years old, with the greatest relative difference being 1.09 in the age group of 30-34 (Fig. 2C). Supplementary Table 1 shows the incidence of BD in females and males of different ages worldwide in 2021.

In addition, the global incidence of ANX among individuals under the age of 15 increased rapidly with age for both sexes. The highest incidence rates were 1107.98 (95% UI: 846.66 to 1454.45) for females and 753.22 (95% UI: 569.77 to 998.24) for males in the age group of 10-14 (Fig. 2A). The absolute sex difference in incidence rates increased rapidly with age for those under 20 years old, with the greatest absolute difference being 354.76 in the age group of 10-14 (Fig. 2B). The relative sex difference in incidence rates decreased with age for those over 25 years old, with the greatest relative difference being 1.51 in the age group of 20-24 (Fig. 2C). Supplementary Table 2 shows the incidence of ANX in females and males of different ages worldwide in 2021.

Sex difference in incidence of BD and ANX by sociodemographic index

ASRs of BD and ANX for males and females in 204 countries and territories in 2021 were demonstrated in Fig. 3A, B, E, F. Absolute and relative sex difference in ASRs of in 204

countries and territories in 2021 were demonstrated in Fig. 3C, D, G, H.

Mann–Whitney U test showed that females had significant higher ASR of BD than males for 204 countries and territories (Z = -2.04, $P < 0.001$), with median (interquartile range) of ASRs being 39.66 (33.99 to 56.40) for females and 37.62 (31.94 to 47.43) for males. After dividing the groups by SDI, multiple comparisons revealed that females had significantly higher ASR than males in high-middle SDI group (females vs. males: 39.46 (32.28 to 58.99) vs. 33.56 (30.10 to 47.13); Z = -5.91, $P = 0.039$) and high SDI group (54.69 (37..42 to 59.43) vs. 46.43 (32.60 to 47.63); Z = -8.27, $P < 0.001$) (Fig. 4A).

Additionally, Mann–Whitney U test also showed that he ASR of ANX in females was significantly higher than that in males (Z = -246.26, $P < 0.001$): median (interquartile range) of ASRs was 848.77 (709.81 to 967.93) for females and 602.51 (544.04 to 711.66) for males. Multiple comparisons revealed that females had significantly higher ASR than males, for low SDI group (males vs. females: 694.28 (574.09 to 740.55) vs. 551.46 (488.90–612.80); Z = -142.82, $P < 0.001$), low-middle SDI group (768.53 (637.44 to 868.93) vs. 580.82 (511.16 to 669.80); Z = -187.71, P < 0.001), middle SDI group (895.81 (768.41 to 999.39) vs. 645.37 (555.47 to 719.93); Z = -250.44, P < 0.001), high-middle SDI group (918.97 (810.98 to 1009.92) vs. 660.86 (554.01 to 773.67); Z = -258.11, $P < 0.001$), and high SDI group (984.02 (856.39 to 1170.26) vs. 641.80 (546.04 to 795.65); Z = -342.23, $P < 0.001$) (Fig. 4B).

As shown in Fig. 5, absolute and relative sex differences of BD and ANX were significantly and positively correlated with SDI: Absolute sex differences of BD, standardized β = 12.46 (95% CI, 9.76 to 15.15), $P < 0.001$; Relative sex differences of BD, standardized β = 0.27 (95% CI, 0.22 to 0.33), $P < 0.001$; Absolute sex differences of ANX, standardized β = 444.65 (95% CI, 293.74 to 595.56), $P < 0.001$; Relative sex differences of ANX, standardized β = 0.80 (95% CI, 0.47 to 1.14), $P < 0.001$.

Discussion

Sex differences in the incidence of BD and ANX have persisted globally from 1990 to 2019, with rates always higher in women than in men. It is worth noting that the sex difference in BD and ANX incidence increases rapidly with age (under 20 years). In addition, countries with higher levels of socioeconomic development have greater gender differences in BD and ANX incidence.

Firstly, the results of sex difference in incidence by year showed that the incidence was consistently higher in females than in males. Previous studies have shown that the incidence of bipolar disorder and anxiety disorders is lower in men and higher in women [21, 22]. The ASRs for BD in females and males has not changed much globally although ASRs increased globally for both sexes. Over the past 30 years, gender differences in ASRs for bipolar disorder have gradually improved. In contrast, anxiety disorders have demonstrated an increasing trend in incidence and their sex differences. In terms of psychosocial factors, gender socialization puts women in a position of increased mental health vulnerability; women are allowed and encouraged to express their fears and worries, and they are more likely to seek help [23,24]. In contrast to masculine instrumentality or some male characteristics like competitiveness, femininity combined by poor self-esteem increased the development of anxiety [25, 26]. Furthermore, differences between sexes, such as brain regions involved in emotional regulation, sex hormone changes, and genetic factors, all

contribute to gender differences in disease [27-29]. More specifically, there was a marked upward trend in the ASR of ANX from 2019 to 2021. This may be related to the COVID-19 pandemic, which has had a more serious impact on women's mental health than men's, with higher levels of anxiety [30]. Therefore, the issue of gender differences in BD and the issue of sex differences in ANX remains a major global challenge in the future.

Secondly, sex differences in the incidence of BD and ANX by age revealed that the sex differences in the incidence of BD and ANX increased more significantly in younger age groups. Studies examining mental disease in various age and sex groups have shown that while there is minimal difference in the oldest age group between the sexes, females are more likely than males to experience mental illness in the youngest age group [31]. Sex hormones, the HPA axis, and the immune system interact to alter brain functioning and increase the vulnerability to mood disorders [32]. The hypothalamic-pituitary-gonadal axis stimulates the production of sex hormones differently during adolescence. Additionally, the maturation of the hypothalamic-pituitary-adrenal axis during early adolescence causes a sharp increase in testosterone levels in males and estrogen levels in females [33]. Along with the HPA axis, the immune system undergoes significant development during puberty and is greatly influenced by sex hormones. The immune system is significantly affected by sex and differences in immune responses, and it may be due to the greater concentration of androgen in males than in females, additionally, testosterone not only attenuates the activation of the HPA axis, but it also suppresses the immune response [32]. The increased immune response in females may play a significant role in the vulnerability to mental illnesses [34].

Furthermore, our results indicated that There are regional differences in the incidence of these diseases and the higher the degree of socio-demographic development, the greater the gender differences in the incidence of diseases. In countries with higher living conditions, the negative impact of emotion was greater for women than for men, and women had more mental disease, such as having more symptoms, diagnoses, or feelings of depression and remembering more nightmares [35]. In all WHO regions, especially the regions of Europe and the Americas, the incidence of BD is higher in women than in men (women have about 1.2 times the risk of men), and the incidence of ANX is higher in women than in men (women have about 1.5 times the risk of men). In countries with higher living conditions, psychosocial factors such as gender socialization may have less influence, giving women and men more freedom to pursue the values they care about, leading to greater gender differences [36]. In addition to mental illness, personality, verbal episodic memory, verbal ability, aggressive behavior, and general self-esteem also show greater gender differences in countries with higher living conditions, these factors influence people's behavior and choices[35]. This gives women and men more freedom to pursue the values they care about, not more of social-role expectations, which also contributes to greater gender disparities [35, 37]. Instead of the traditional gender division of labor based on physical attributes, greater economic prosperity will change the structure of education and careers [38]. With the emergence of new jobs that are considered appropriate for females' roles in society, more women are being socialized to fit into the careers they pursue, which brings with it mental health problems. Women who worked as housekeepers at home reported less mental health issues than those who worked in countries with southern regimes [11]. According to a representative European study, women

who were living alone—especially those who were employed—scored higher for depression

[39].

This study is constrained by GBD 2021, including data sources, statistical methods, and so forth [40]. The primary limitation of GBD analysis lies in the availability of raw data. Since country-level aggregated data are employed rather than regional data, potential bias may be introduced in geographic differences of incidence estimates. Although this study offers a global perspective on gender differences in the incidence of BD and ANX, the conclusions may not be applicable to specific regions. Considering that GBD data will be updated in the future, sex differences in the global incidence of BD and ANX over longer time periods can be further explored.

Overall, the findings highlight the importance of tailoring mental health interventions for BD and ANX to address the unique needs of different populations, particularly in regions with pronounced sex differences.

Conclusion

This study provides a comprehensive analysis of sex differences in the incidence of bipolar disorder (BD) and anxiety disorders (ANX) using data from the Global Burden of Disease Study 2021. Our study revealed that sex difference in the incidence of anxiety disorders and bipolar disorder have persisted worldwide over the past several decades, and the rates have consistently been higher among females than males. The sex difference in the global incidence of bipolar disorder has shown a slight improvement, but that in the global incidence of anxiety disorders has not been effectively mitigated. The sex difference is even more pronounced at younger ages and in more developed nations. These findings underscore the need for continued focus on sex-specific strategies in the prevention, diagnosis, and treatment of bipolar and anxiety disorders.

Availability of data and materials

The datasets analyzed during the current study are available from the publicly available website: https://www.healthdata.org/research-analysis/gbd.

Reference

[1] Gao K, Calabrese JR, 2024. Prevalence and factors associated with fatigue in patients with major depressive disorder or bipolar disorder. J Affect Disord. 362:493-501. https://doi.org/10.1016/j.jad.2024.07.016

[2] Gandy M, Woldhuis T, Wu W, et al., 2024. Cognitive behavioral interventions for depression and anxiety in adults with neurological disorders: a systematic review and meta-analysis. Psychol Med. https://doi.org/10.1017/S0033291724001995

[3] GBD 2019 Mental Disorders Collaborators, 2022. Global, regional, and national burden of 12 mental disorders in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. Lancet Psychiatry. 9(2):137-150. https://doi.org/10.1016/S2215-0366(21)00395-3

[4] Temes CM, Boccagno C, Gold AK, et al., 2024. Comorbidity of bipolar disorder and borderline personality disorder: Phenomenology, course, and treatment considerations. Bipolar Disord. https://doi.org/10.1111/bdi.13465

[5] Arnone D, Karmegam SR, Östlundh L, et al., 2024. Risk of suicidal behavior in patients with major depression and bipolar disorder - A systematic review and meta-analysis of registry-based studies. Neurosci Biobehav Rev. 159:105594. https://doi.org/10.1016/j.neubiorev.2024.105594

[6] Meier SM, Mattheisen M, Mors O, et al., 2016. Increased mortality among people with anxiety disorders: total population study. Br J Psychiatry. 209(3):216-221. https://doi.org/10.1192/bjp.bp.115.171975

[7] Mitchell RH, Metcalfe AW, et al., 2018. Sex differences in brain structure among adolescents with bipolar disorder. Bipolar Disord. https://doi.org/10.1111/bdi.12663

[8] Tondo L, Pompili M, Forte A, et al., 2016. Suicide attempts in bipolar disorders: comprehensive review of 101 reports. Acta Psychiatr Scand. 133(3):174-86. https://doi.org/10.1111/acps.12517

[9] McLean CP, Asnaani A, Litz BT, et al., 2011. Gender differences in anxiety disorders: prevalence, course of illness, comorbidity and burden of illness. J Psychiatr Res. 45(8):1027-1035. https://doi.org/10.1016/j.jpsychires.2011.03.006

[10] Zhang L, Swaab DF, 2024. Sex differences in bipolar disorder: The dorsolateral prefrontal cortex as an etiopathogenic region. Front Neuroendocrinol. 72:101115. https://doi.org/10.1016/j.yfrne.2023.101115

[11] Farhane-Medina NZ, Luque B, Tabernero C, et al., 2022. Factors associated with gender and sex differences in anxiety prevalence and comorbidity: A systematic review. Sci Prog.105(4):368504221135469. https://doi.org/10.1177/00368504221135469

[12] Zhang H, Zhou XD, Shapiro MD, et al., 2024. Global burden of metabolic diseases, 1990-2021. Metabolism. 160:155999. https://doi.org/10.1016/j.metabol.2024.155999

[13] GBD 2017 Disease and Injury Incidence and Prevalence Collaborators, 2018. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. Lancet. 392(10159):1789-1858. https://doi.org/10.1016/S0140-6736(18)32279-7

[14] GBD 2017 Mortality Collaborators, 2018. Global, regional, and national age-sex-specific mortality and life expectancy, 1950-2017: a systematic analysis for the Global Burden of Disease Study 2017. Lancet. 392(10159):1684-1735. https://doi.org/10.1016/S0140-6736(18)31891-9

[15] GBD 2021 Diseases and Injuries Collaborators, 2024. Global incidence, prevalence, years lived with disability (YLDs), disability-adjusted life-years (DALYs), and healthy life expectancy (HALE) for 371 diseases and injuries in 204 countries and territories and 811 subnational locations, 1990-2021: a systematic analysis for the Global Burden of Disease Study 2021. Lancet. 403(10440):2133-2161. https://doi.org/10.1016/S0140-6736(24)00757-8

[16] Forray AI, Coman MA, Simonescu-Colan R, et al., 2023. The Global Burden of Type 2 Diabetes Attributable to Dietary Risks: Insights from the Global Burden of Disease Study 2019. Nutrients. 15(21):4613. https://doi.org/10.3390/nu15214613

[17] Li XY, Kong XM, Yang CH, et al., 2024. Global, regional, and national burden of ischemic stroke, 1990-2021: an analysis of data from the global burden of disease study 2021. EClinicalMedicine. 75:102758. https://doi.org/10.1016/j.eclinm.2024.102758

[18] Kuang Z, Wang J, Liu K, et al. Global, regional, and national burden of tracheal, bronchus, and lung cancer and its risk factors from 1990 to 2021: findings from the global burden of disease study 2021. EClinicalMedicine. 2024;75:102804. Published 2024 Sep 5. https://doi.org/10.1016/j.eclinm.2024.102804

[19] Gao S, Yu Z, Ma X, et al., 2024. Childhood-onset systemic lupus erythematosus in China,

2016-21: a nationwide study. Lancet Child Adolesc Health. 8(10):762-772. https://doi.org/10.1016/S2352-4642(24)00172-X

[20] Nachar N, 2008. The Mann-Whitney U: a test for assessing whether two independent samples come from the same distribution. Tutor Quant Meth Psychol. https://doi.org/10.20982/ tqmp.04.1.p013

[21] Hu FH, Jia YJ, Zhao DY, et al., 2023. Gender differences in suicide among patients with bipolar disorder: A systematic review and meta-analysis. J Affect Disord. 339:601-614. https://doi.org/10.1016/j.jad.2023.07.060

[22] Asher M, Asnaani A, Aderka IM, 2017. Gender differences in social anxiety disorder: A review. Clin Psychol Rev. 56:1-12. https://doi.org/10.1016/j.cpr.2017.05.004

[23] Anyan F, Hjemdal O, 2018. Stress of home life and gender role socializations, family cohesion, and symptoms of anxiety and depression. Women Health. 58(5):548-564. https://doi.org/10.1080/03630242.2017.1316343

[24] Richmond K, Levant R, Smalley B, et al., 2015. The Femininity Ideology Scale (FIS): dimensions and its relationship to anxiety and feminine gender role stress. Women Health. 55(3):263-79. https://doi.org/10.1080/03630242.2014.996723

[25] Berke DS, Reidy D, Zeichner A, 2018. Masculinity, emotion regulation, and psychopathology: A critical review and integrated model. Clin Psychol Rev. 66:106-116. https://doi.org/10.1016/j.cpr.2018.01.004

[26] McLean CP, Anderson ER, 2009. Brave men and timid women? A review of the gender differences in fear and anxiety. Clin Psychol Rev. 29(6):496-505. https://doi.org/10.1016/j.cpr.2009.05.003

[27] Kaczkurkin AN, Raznahan A, Satterthwaite TD, 2019. Sex differences in the developing brain: insights from multimodal neuroimaging. Neuropsychopharmacology. 44(1):71-85. https://doi.org/10.1038/s41386-018-0111-z

[28] Hodes GE, Epperson CN, 2019. Sex Differences in Vulnerability and Resilience to Stress Across the Life Span. Biol Psychiatry. 86(6):421-432. https://doi.org/10.1016/j.biopsych.2019.04.028

[29] Palma-Gudiel H, Peralta V, Deuschle M, et al., 2019. Epigenetics-by-sex interaction for somatization conferred by methylation at the promoter region of SLC6A4 gene. Prog Neuropsychopharmacol Biol Psychiatry. 8;89:125-131. https://doi.org/10.1016/j.pnpbp.2018.09.002

[30] Liu S, Yang L, Zhang C, et al., 2021. Gender differences in mental health problems of healthcare workers during the coronavirus disease 2019 outbreak. J Psychiatr Res. 137:393-400. https://doi.org/10.1016/j.jpsychires.2021.03.014

[31] Höglund P, Hakelind C, Nordin S, 2020. Severity and prevalence of various types of mental ill-health in a general adult population: age and sex differences. BMC Psychiatry. 20(1):209. https://doi.org/10.1186/s12888-020-02557-5

[32] Smith KB, Murack M, Ismail N, 2023. The sex-dependent and enduring impact of pubertal stress on health and disease. Brain Res Bull. 200:110701. https://doi.org/10.1016/j.brainresbull.2023.110701

[33] Bennett DS, Ambrosini PJ, Kudes D, et al., 2005. Gender differences in adolescent depression: do symptoms differ for boys and girls? J Affect Disord. 89(1-3):35-44. https://doi.org/10.1016/j.jad.2005.05.020

[34] Kane L, Ismail N, 2019=7. Puberty as a vulnerable period to the effects of immune challenges: Focus on sex differences. Behav Brain Res. 320:374-382. https://doi.org/10.1016/j.bbr.2016.11.006

[35] Herlitz A, Hönig I, Hedebrant K, et al., 2024. A Systematic Review and New Analyses of the Gender-Equality Paradox. Perspect Psychol Sci. https://doi.org/10.1177/17456916231202685

[36] Falk A, Hermle J, 2018. Relationship of gender differences in preferences to economic development and gender equality. Science. 362(6412):eaas9899. https://doi.org/10.1126/science.aas9899

[37] Yang, K., Girgus, J.S., 2019. Are Women More Likely than Men Are to Care Excessively about Maintaining Positive Social Relationships? A Meta-Analytic Review of the Gender Difference in Sociotropy. Sex Roles 81, 157–172 (2019). https://doi.org/10.1007/s11199-018-0980-y

[38] Charles M, Bradley K, 2009. Indulging our gendered selves? Sex segregation by field of study in 44 countries. AJS. 114(4):924-76. https://doi.org/10.1086/595942

[39] Van de Velde S, Huijts T, Bracke P, et al., 2013. Macro-level gender equality and depression in men and women in Europe. Sociol Health Illn. 35(5):682-98. https://doi.org/10.1111/j.1467-9566.2012.01521.x

[40] GBD 2021 Risk Factors Collaborators, 2024. Global burden and strength of evidence for 88 risk factors in 204 countries and 811 subnational locations, 1990-2021: a systematic analysis for the Global Burden of Disease Study 2021. Lancet. 403(10440):2162-2203. https://doi.org/10.1016/S0140-6736(24)00933-4

Year and region	Female			Male			Risk ratio		
	Incidence rates	Upper	Lower	Incidence rates	Upper	Lower	Mean	Upper	Lower
1990									
Global	32.61	38.60	27.45	30.11	35.57	25.41	1.08	1.09	1.08
Eastern Mediterranean Region	44.56	54.49	36.44	40.96	49.70	33.34	1.09	1.10	1.09
European Region	47.08	56.63	39.28	39.41	47.12	32.75	1.19	1.20	1.19
Region of the Americas	56.85	66.28	49.05	52.19	60.43	45.11	1.09	1.10	1.09
South-East Asia Region	23.10	27.43	19.35	25.04	29.78	21.02	0.92	0.92	0.92
Western Pacific Region	17.66	20.81	14.91	16.21	19.02	13.73	1.09	1.09	1.09
African Region	37.82	46.22	30.91	36.50	44.42	29.82	1.04	1.04	1.04
2021									
Global	33.06	39.60	27.80	31.22	37.34	26.27	1.06	1.06	1.06
Eastern Mediterranean Region	43.43	53.06	35.41	40.32	48.91	32.68	1.08	1.08	1.08
European Region	48.44	58.29	40.24	40.45	48.36	33.61	1.20	1.21	1.20
Region of the Americas	57.07	66.57	49.12	52.57	61.29	45.34	1.09	1.09	1.08
South-East Asia Region	23.07	27.39	19.38	25.14	29.83	21.14	0.92	0.92	0.92
Western Pacific Region	17.85	21.17	15.10	16.57	19.59	14.00	1.08	1.08	1.08
African Region	37.43	45.68	30.59	36.24	44.08	29.59	1.03	1.04	1.03

Table 1. The global and WHO regional incidence of bipolar disorder for both sexes in 1990 and 2021.

WHO: World Health Organization. The confidence interval is 95%.

Incidence rates: Age-standardized incidence rates.

Table 2. The global and WHO regional incidence of anxiety disorder for both sexes in 1990 and 2021.

Year and region	Female			Male			Risk ratio		
	Incidence rates	Upper	Lower	Incidence rates	Upper	Lower	Mean	Upper	Lower
1990									
Global	649.32	795.75	537.68	479.30	579.75	402.19	1.35	1.37	1.34
Eastern Mediterranean Region	752.87	946.05	615.88	605.97	733.70	504.65	1.24	1.29	1.22
European Region	733.96	913.49	605.57	516.01	627.22	431.41	1.42	1.46	1.40
Region of the Americas	807.89	1006.83	664.27	602.69	729.42	503.77	1.34	1.38	1.32
South-East Asia Region	562.46	688.52	460.86	432.02	519.03	362.05	1.30	1.33	1.27
Western Pacific Region	620.06	750.29	516.13	429.82	511.04	364.29	1.44	1.47	1.42
African Region	580.59	725.70	476.44	477.36	591.81	393.29	1.22	1.23	1.21
2021									
Global	784.53	966.02	649.26	576.46	701.84	486.27	1.36	1.38	1.34
Eastern Mediterranean Region	885.62	1123.95	718.38	702.24	854.00	575.14	1.26	1.32	1.25
European Region	963.80	1202.67	792.66	657.98	800.01	548.42	1.46	1.50	1.45
Region of the Americas	1115.49	1407.77	909.94	802.04	987.50	660.66	1.39	1.43	1.38
South-East Asia Region	693.95	850.01	573.34	512.41	623.02	424.35	1.35	1.36	1.35
Western Pacific Region	678.82	829.63	562.44	464.04	560.07	389.17	1.46	1.48	1.45
African Region	657.70	819.67	537.75	542.65	669.98	447.05	1.21	1.22	1.20

WHO: World Health Organization. The confidence interval is 95%.

Incidence rates: Age-standardized incidence rates.

Figure Caption

Fig. 1. Sex difference in global incidence of BD and ANX from 1990 to 2021. (A) Age standardized incidence rates of BD. (B) Absolute (female minus male) sex difference in age standardized incidence rates of BD. (C) Relative (female to male ratio) sex difference in age standardized incidence rates of BD. (D) Age standardized incidence rates of ANX. (E) Absolute sex difference in age standardized incidence rates of ANX. (F) Relative sex difference in age standardized incidence rates of ANX.

Fig. 2. Sex difference in global incidence of BD and ANX in different age groups in 2021. (A) Incidence rates of BD. (B) Absolute (female minus male) sex difference in incidence rates of BD. (C) Relative (female to male ratio) sex difference in incidence rates of BD. (D) Incidence rates of ANX. (E) Absolute sex difference in incidence rates of ANX. (F) Relative sex difference in incidence rates of ANX.

Fig. 3. Sex difference in national incidence of BD and ANX in 2021. (A) Age standardized incidence rates of BD in females. (B) Age standardized incidence rates of BD in males. (C)Absolute (female minus male) sex difference in age standardized incidence rates of BD. (D) Relative (female to male ratio) sex difference in age standardized incidence rates of BD. (E) Age standardized incidence rates of ANX in females. (F) Age standardized incidence rates of ANX in males. (G)Absolute (female minus male) sex difference in age standardized incidence rates of ANX. (H) Relative (female to male ratio) sex difference in age standardized incidence rates of ANX.

Fig. 4. Sex difference in incidence of BD and ANX for SDI-based country groups. (A) Age-standardized incidence rates of BD among females and males for SDI-based country groups. (B) Age-standardized incidence rates of ANX among females and males for SDI-based country groups.

Lines inside the boxes indicate the medians, boxes the 25th and 75th percentiles, and lines outside the boxes the minimum and the maximum. * indicates $P < 0.05$; *** indicates $P < 0.001$. SDI, sociodemographic index.

Fig. 5. Sex difference in incidence of BD and ANX by SDI. (A) Linear relationship between absolute (female minus male) sex difference in age standardized incidence rates of BD and SDI. (B) Linear relationship between relative (female to male ratio) sex difference in age standardized incidence rates of BD and SDI. (C) Linear relationship between absolute (female minus male) sex difference in age standardized incidence rates of ANX and SDI. (D) Linear relationship between relative (female to male ratio) sex difference in age standardized incidence rates of ANX and SDI.

SDI, sociodemographic index.