Integrating Multidimensional Data Analytics for Precision Diagnosis of Chronic Low Back Pain

Sam Vickery[1]*, Frederick Junker[1], Rebekka Döding[1], Daniel L Belavy[1], Maia Angelova[2,3],

Chandan Karmakar[3], Luis Becker[4,5], Nima Taheri[4,5], Matthias Pumberger[4], Sandra Reitmaier[5],

Hendrik Schmidt[5]

[1] Fachbereich Pflege-, Hebammen- und Therapiewissenschaften (PHT) | Hochschule Bochum

(University of Applied Sciences) | Bochum | Germany

[2] Aston Digital Futures Institute | Aston University | Birmingham | United Kingdom

[3] School of Information Technology | Deakin University | Geelong | Australia

[4] Center for Musculoskeletal Surgery, Charité – Universitätsmedizin Berlin, Berlin, Germany

[5] Julius Wolff Institut, Berlin Institute of Health - Charité at Universitätsmedizin Berlin |

Berlin | Germany

Short title: Multimodal classification of chronic low back pain

Abstract

Low back pain (LBP) is a leading cause of disability worldwide, with up to 25% of cases become chronic (cLBP). Optimal diagnostic tools for cLBP remains unclear. Here we leveraged a comprehensive multi-dimensional data-set and machine learning-based variable importance selection to identify the most effective diagnostic tools for cLBP patient stratification. The dataset included questionnaire data, clinical and functional assessments, and spino-pelvic magnetic resonance imaging (MRI), encompassing a total of 144 parameters from 1,161 adults with (n=512) and without cLBP (n=649). Boruta and random forest were utilised for variable importance selection and cLBP classification respectively. Boruta variable selection led to pronounced variable reduction (median of all 15 datasets: 63.3%), while performing comparable to using all variables across all modality datasets. Boruta selected key variables from questionnaire, clinical, and MRI data were the most effective in distinguishing cLBP patients from controls. The most robust variables (n=9) across the whole dataset identified were psychosocial factors, neck and hip mobility, as well as lower lumbar disc herniation and degeneration. These critical variables outperformed all parameters in an unseen holdout dataset, demonstrating superior patient delineation. Paving the way for targeted diagnosis and personalized treatment strategies, ultimately enhancing clinical outcomes for cLBP patients.

Key words

Chronic low back pain; classification; data-driven; feature selection; multi-modality;
psychosocial; MRI

Introduction

In recent years, chronic low back pain (cLBP) has become one of the most prevalent and challenging conditions in clinical practice, affecting a significant portion of the global population [1,2]. Despite its widespread occurrence, diagnosing and assessing cLBP remains difficult due to the complex and multifactorial nature of the disease [3,4], which encompasses physical, psychological, and social dimensions [5]. Traditional diagnostic approaches often rely on self-reported symptoms, which can be subjective and prone to variability. Furthermore, surgical and non-surgical treatment outcomes are still inconsistent, refelected in a high rate of treatment-refactory in cLBP patients [6]. As such, there is an increasing interest in exploring more objective and comprehensive methods that incorporate multimodal data, including medical imaging, physical assessments, clinical evaluations, and patient-reported outcomes.

Machine learning algorithms provide models for identifying distinct subgroups that can help explain the occurrence and characteristics of a disease [7,8]. A previous systematic review by our team using machine learning applications in LBP [9] highlighted that a narrow range of mechanistic domains have been assessed, and sample sizes in these studies were consistently small, ranging up to only 171 participants. Consequently, using limited data and modalities limits the robustness and applicability of such models. Through gathering many data points across multiple modalities one can ascertain which variables and modalities are the most informative at distinguishing cLBP patients from asymptomatic controls. Reducing the number of variables to those that are most informative has been previously employed in predictive and classification modelling to improve accuracy [10,11]. This approach can be applied as the main outcome and not only in model preprocessing, in order to obtain a data-driven decision on the most important variables in multi-dimensional clinical data. Such a systematic data-informed investigation of back pain diagnosis in a large multi-modality sample is lacking to help inform

64   future studies in selecting which data to acquire and for clinicians in which tests to conduct.

65   Therefore, the aim of this study was to identify and compare the most informative domains and

66   variables in delineating patients with and without cLBP utilising a large multi-modality dataset.

Results

Study sample

The prospective cross-sectional study draws its data from the ongoing "Berliner Rückenstudie" ("Berlin Back Study"; https://spine.charite.de/en/spine_study/; running time: 01/01/2022 to 31/12/2025), which was registered at the German Clinical Trial Register (DRKS-ID: DRKS00027907). Recruitment procedures vary from local promotion (i.e., postal flyers, notice boards, internet approaches, and social media) at the Charité-Universitaetsmedizin Berlin, in the general public (i.e., newspapers, magazines, podcasts) to cooperation with local companies, administrative authorities, and word-of-mouth. The protocol is in accordance with the Helsinki Declaration of ethical principles [12] and has been approved by the Ethics Committee of the Charité – Universitätsmedizin Berlin (registry numbers: EA4/011/10, EA1/162/13). Written informed consent was obtained from all participants. The STROBE guideline [13] (Supplementary Table S1) and TRIPOD statement [14] (Supplementary Table S2) for prediction model development were used to report this study. Data collection started on 1st January 2022 and cut-off for inclusion in the current analysis was 5th April 2024. Data collection occurred in a research centre within a university-hospital.

Study participants were recruited through a telephone interview and excluded if they met any exclusion criteria, as well as some excluded at the testing site (Supplementary Table S3). A total of 1273 participants were included in the study at cut-off point. These participants were initially guided through self-administered questionnaires by a study coordinator. Then they continued to a clinical examination by a trained medical doctor, which included physical examinations, questions, as well as a back shape and function test. The examinations and questionnaires took a total of 90 minutes to complete. Additionally, participants were offered a

magnetic resonance imaging (MRI) within 14 days of the spino-pelvic region. During the

clinical assessment the participants were classified by the clinician as asymptomatic (no back

pain), symptomatic (cLBP), or previously suffering from cLBP. To ensure a more robust cLBP

patient classification previous symptomatic subjects were removed from the sample. The

criteria for designation as cLBP was daily back pain for more than three months. Furthermore,

participants who revoked their inclusion in the study and those who were missing demographic

data; age, sex, body mass index (BMI), and patient status were removed. This resulted in a

study sample of 1161 subjects that included 649 asymptomatic (19 – 72 years old, mean age =

40.7 ± 12.6, females = 353) and 512 cLBP (19 – 65 years old, mean age = 43.5 ± 11.7, females

= 306) participants (Table 1). This sample was sub-divided into four modalities; questionnaires

(Q), clinical physical assessment (C), back shape and function (S), and MRI (M). Each modality

was combined with demographic data (age, sex, and BMI) and then joined with all

combinations of the four modalities, resulting in 15 datasets (Figure 1A, and Supplementary

Table S4 - S18). Both complete datasets (Fig. 2) and imputed datasets were used for cLBP

classification (Fig. 1B). Imputation can lead to bias results, in particular when it comes to

variable importance selection [15]. Therefore, presentation of variable importance was only

considered for non-imputed datasets. Highly correlated ($r > 0.9$) variables were removed from

the daatsets to reduce colinearlity between dataset variables (Supplementary Figure S1 – S6).

An overview of all Berlin Back Study variables (Supplementary Table S19), those removed

during preprocessing (Supplementary Table S20), and a list of all variables (144) used for

modelling is presented in Supplementary (Supplementary Table S21).

Table 1. Demographic data of entire Berlin Back dataset

	Asymptomatic	Chronic low back pain
Sample size	649	512
Females	353	306
Age (years) mean [sd]	40.65 [12.6]	43.52 [11.67]
Body mass index mean [sd]	23.55 [2.75]	23.55 [2.84]

Pain duration mean [sd]	549.68 [532.38]
Pain Intensity mean [sd]	2.87 [1.86]

Figure 1. Modality dataset distributions and machine leaning workflow. A – Top shows the chronic low back pain (cLBP) sample size distribution across all 15 dataset modalities. Bottom presents the number of variables used for cLBP classification and variable importance selection across the 15 dataset modalities. B – Represents the machine learning workflow implemented to compare the different modalities and determine the most important variables for cLBP patient delineation using a random forest binary classification algorithm for training and testing.

Figure 2. Modality datasets age, sex, and cLBP distributions. Violin plots for non-imputed dataset
A - questionnaire; B – clinical assessment: C – back shape and function; D – MRI; E – questionnaire +
clinic; F – questionnaire + back shape and function; G – questionnaire + MRI; H – clinic + back shape
and function; I – clinic + MRI; J – back shape and function + MRI; K – questionnaire + clinic + back
shape and function; L – questionnaire + clinic + MRI; M – questionnaire + back shape and function +
MRI; N – clinic + back shape and function + MRI; O – all datasets.
130

Chronic low back pain classification

Determining modality and variable importance was conducted on 90% (n = 1045, 19 – 72 y/o,

mean age = 41.71 ± 12.28, cLBP = 469) of the whole dataset, with 10% (n = 116, 20 – 64 y/o,

mean age = 43.81 ± 12.12, cLBP = 43) used as a hold-out sample to test cLBP patient

classification performance of the most robust and important variables (Fig. 1B). Variable

selection and importance was determined using the Boruta algorithm [16] on the training data

within an iterative ten-fold train-test loop across all 15 modality datasets. Boruta is a wrapper

function selects important variables and removes unimportant variables using random random

forest (RF) [17]. A binary RF classification model was trained using the Boruta [16] selected

variables as well as all variables to classify cLBP status in the test sample across the ten-fold

train-test loop. Furthermore, imputation using missForest [18] was independently conducted on

the training and test datasets within the ten-fold loop (Supplementary Table S23). Imputed

datasets performed generally slightly worse than the full datasets. We used AUC (area under

the receiver operating characteristic (ROC) curve) as the main model performance metric. AUC

provides a good combination of sensitivity and specificity for comparing Boruta selected

variables to all variables and the different modality datasets in their performance of cLBP

patient delineation.

The best single modality for cLBP classification was Boruta selected variables from MRI (Fig.

3A) with a mean AUC of 0.645 with a 95% confidence interval (CI) of 0.618 – 0.672 and

accuracy of 0.657 (95% CI, 0.636 – 0.678). The Boruta selected questionnaire dataset modality

produced only minimally worse classification performance (mean AUC = 0.631, 95% CI =

0.610 – 0.652) than Boruta reduced and all MRI variables (mean AUC = 0.637, 95% CI = 0.599

– 0.675) using the least amount of variables (mean = 8) across all modality datasets (Fig. 3A-

C). Dual and multi modalities generally performed better than single modality models in cLBP

classification. Questionnaire, clinical physical assessment, and MRI (Q + C + M, Fig. 3C)

modality with Boruta selected variables represents the best performing modality model with a

mean AUC of 0.699 (95% CI, 0.669 – 0.729) and a mean accuracy of 0.709 (95% CI, 0.679 –

0.739). Moreover, this model showed the highest sensitivity (mean = 0.622, 95% CI – 0.568 –

0.676). The modality model showing the highest specificity was using all variables and all

modalities (Q + C + S + M, mean = 0.840, 95% CI = 0.799 – 0.881). The three best dual

modalities models, C + M (mean AUC = 0.679, 95% CI = 0.633 – 0.725), Q + M (mean AUC

= 0.674, 95% CI = 0.623 – 0.725), and Q + C (mean AUC = 0.674, 95% CI = 0.639 – 0.709)

all with Boruta selected variables (Fig. 3B), performed only slightly worse than the best model

(Boruta – Q + C + M). Overall back shape and function dataset using Boruta selected variables

produces the worst classification performance (Fig. 3A, AUC = 0.569, 95% CI = 0.538 – 0.60).

Additionally, the dual modalities continuing back shape and function data always performed

worse than those without (Fig. 3B), when using both all and Boruta selected variables.

Classification performance metrics across all models is provided in Supplementary Table S7.

Figure 3. Boruta variable reduction performance. A – C shows RF classification model performance (AUC) following the reduction of variables using Boruta and all variables in the single, dual, and multi data modalities respectively. This shows the change in performance follow variable reduction. Error bars represent 95% CI in AUC over 10-fold train-test splits. D – Shows the amount of variable reduction by using Boruta as a percentage of the total number of variables within each modality dataset.

Boruta variable importance

Boruta variable importance selection resulted in a median of 62.7% reduction in the number of

variables across all 15 datasets (Fig. 3D). This large reduction in variables from Boruta

performed comparable (eight slightly worse and seven better) compared to using all variables,

with high overlap in confidence intervals. Furthermore, three of the top five performing

modality models where those employing Boruta selected variables (Supplementary Table S22).

The greatest performance improvement following Boruta variable selection was found in the Q

+ C + M datasets with an AUC increase of 0.270 and an average reduction of 72.2 variables

(Fig. 3C). The smallest variable reduction was shown in MRI (30%) with an AUC increase of

0.008, while the largest variable reduction was found in the whole dataset (Q + C + S + M) with

a reduction of 75.7% of the variables and an AUC decrease of 0.009.

Boruta selected variables represented the best patient delineation performance in single, dual,

and multi-modality datasets (Fig. 3A-C). MRI was found to be the best single modality model

using Boruta selected variables (Fig. 3A). The most important variables (Supplementary Table

S27) were intervertebral disc (IVD) herniation L4 – L5, spinal canal width L2, IVD

degeneration L3 – L4 and L4 – L5, and spinal canal width L1, showing a mean importance

across the ten iterations of 19.49, 9.94, 9.46, 9.14, and 8.13 respectively. The best dual

modalities cLBP patient stratification model, Boruta selected C + M (Fig. 3B), showed the

second highest AUC (mean = 0.679). The MRI variable IVD herniation L4 – L5 was found to

be the most important (mean = 13.72), with clinical mobility assessments of the hip, cervical

spine, and the whole body showing high importance (Supplementary Table S31). The most

important and robust variables of the best performing model (Boruta – Q + C + M, Fig. 3C)

contained assessments from all three modalities (Supplementary Table S34). The Short-form

36 Health Status Questionnaires (SF-36) psychological well-being, SF-36 social function, and

hip pain presented a mean importance of 13.36, 13.07, and 9.71 respectively. The clinical

assessment, cervical axial rotation (left) and MRI variable IVD herniation L4 – L5 each showed

an importance of 9.47 and 9.12. These represent the top five most important variables and the

rest are provided in Table 3. The questionnaire Boruta reduced model provides a sparse model

with decent performance, meaning the modality performs comparably well with a small amount

of data. On average, the questionnaire Boruta model used 8 variables with a mean AUC

reduction of 0.068 compared to the best model (Q + C + M). The most robust and important

variables (Supplementary Table S39) were SF-36 social function (mean = 21.42), SF-36

psychological well-being (mean = 20.02), hip pain (mean =15.0), smoking in pack years (mean

= 11.8), and family history of back pain (mean = 7.27). All Boruta selected important variables

across the datasets are provided in Supplementary Table S24 – S38 and their correlation

matrices in Supplementary Figure S7 – S21.

Robust and important variables

To select the most robust and important variables for cLBP patient classification, the percentage

of selection and average importance score was calculated across the 15 datasets (Supplementary

Table S39). The variables that were selected at every opportunity (100%) across the multiple

iterations and datasets were defined as most important and solely implemented in cLBP

classification compared to all variables. This resulted in nine robust and important variables

(Fig. 4A). The variables importance score is the Z-score of the mean decrease accuracy measure

and is computed by dividing the average accuracy loss by its standard deviation. Meaning this

value shows how much the models accuracy would decrease without this variable included.

These variables represented psychosocial factors, IVD herniation and degeneration of the lower

lumbar spine, presence of hip pain, as well as mobility of the neck and general mobility of the

whole body. The performance of these nine variables were compared to all variables utilising

the hold-out dataset (Fig. 1B) in a five-fold train-test workflow to provide an unbiased

comparison. The best nine variables showed better mean accuracy (Fig. 4B, Boruta = 0.724,

95% CI = 0.593 – 0.855, All = 0.680, 95% CI = 0.625 – 0.736), AUC (Boruta = 0.664, 95% CI

= 0.514 – 0.814, All = 0.602, 95% CI = 0.538 – 0.666), and sensitivity (Boruta = 0.442, 95%

CI = 0.176 – 0.708, All = 0.298, 0.139 – 0.457), while all variables provided better specificity

(Boruta = 0.892, 95% CI = 0.795 – 0.989, All = 0.902, 95% CI = 0.803 – 1.0). Moreover, all

these nine variables showed significant univariate statistically significant differences between

cLBP patients and asymptomatic controls in the questionnaire, clinical, and MRI datasets (Fig.

4A). Utilising a Wilcoxon-Mann-Whitney test revealed reduced scores of the SF36 for social

function (u = 183751, z = -7.08, p < 0.001, effect size r = -.225) and psychological well-being

(u = 178506.5, z = -7.74, p < 0.001, r = -0.247) in people suffering from cLBP. In addition, the

occurrence of hip pain was also altered comparing people with and without cLBP (χ^2 (3, N=986)

= 40.17, p = 0.004, ω = 0.202). Comparing clinical examinations further revealed reduced

cervical axial rotation to the left (u = 221595, z = -7.98, p < 0.001, r = -0.243), reduced sit to

stand 30 second repetition (u = 226818.5, z = -6.83, p < 0.001, r = -0.208), as well as altered

general mobility ($\chi2$ (2, N=1080) = 46.81, p = 0.006, ω = 0.208) in cLBP patients. Regarding

MRI investigations, increased IVD degeneration at L2-L3 (u = 147168, z = 3.38, p = 0.011, r =

0.120) and L4-L5 (u = 147691, z = 3.43, p = 0.010, r = 0.121), as well as increased disc

herniation at L4-L5 (u = 153470.5, z = 6.02, p < 0.001, r = 0.213) were found in people suffering

from cLBP compared to asymptomatic controls. Univariate statistical results for all variable

can be found in Supplementary Table S40 – S47.

Figure 4. Most robust and important variables for chronic low back pain classification performance. A – Presents a bar plot of the nine most robust variables in order of average Boruta importance score (left). The right bar plot shows the absolute effect size (Cohen's r or ω depending on data type) comparing controls and cLBP patients of the nine robust variables. B – Column plot showing RF classification performance as mean of five-fold train-test iterations in hold-out set using Boruta selected and all variables. Column plot error bars represent 95% CI. IVD – intervertebral disc, SF-36 – short form 36 health status questionnaire, Acc – accuracy, AUC – Area under the receiver operating characteristic curve, Sens – sensitivity, Spec – specificity.

Discussion

This study employs a large multi-modal dataset and a machine learning workflow to demonstrate the importance of using data from different domains in cLBP patient delineation. Increasing the number of modalities generally lead to a model performance improvement although it seems the inclusion of back shape and motion data resulted in little to no performance improvement. Utilising Boruta in our iterative selection workflow resulted in considerable variable reduction across all datasets (median = 62.7%), while model performance remained comparable. This may reflect many variables showing little difference between patients and controls, or the underlying machanisms are more robustly captured by a small sub-set of variables. Both the best performing modality model (Q + C + M) and the most robust variables (Fig. 4), show the importance of measuring psychosocial factors, cervical axial rotation, general mobility, hip flexion, and lower lumbar spine disc degeneration and herniation ratings in cLBP patients.

The questionnaires probing the psychosocial factors, social function and psychological well-being, showed highest importance among all variables (Fig. 4A) and were the most important variables in the best sparse model (Boruta reduced questionnaire, Fig. 3A). Moreover, the quesionnaires represent the most cost effective modality to the examiner, highlighting the clinical importance of psychosocial factors in cLBP diagnosis and treatment.

Social functioning describes the ability of a person to engage in social activities, which we have shown to be an important marker for delineating cLBP patients from asymptomatic controls. Using cross-sectional data from 180 chronic low back pain patients, Ge and colleagues [19] showed that these patients reported more limitations in performing (major life tasks and) social activities as compared to subjects without cLBP, even after adjusting for influencing factors,

such as socio-demographics, lifestyle and number of diseases. Furthermore, Tagliaferri et al. [20]

were able to separate 4156 chronic back pain patients from the UK Biobank dataset into five

sub-groups based on their scores of social isolation and depressive symptoms. Interestingly,

increased social isolation was only a variable of three sub-groups, encompassing 26% of all

back pain patients (n = 1085), while the remaining subgroups showed either no changes (4.1%;

n = 776) or a reduction in social isolation scores (12%; n = 2296). This prevalence in patients

with LBP may indicate that reduced social functioning was identified by some studies, while

others did not find similar changes as compared to asymptomatic controls [21]. However, as levels

of social function (here: social participation) were found to be correlated with self-perceived

physical health status [22], a direct impact of social functioning on personal functional

impairments remains feasible.

In addition to social function, the psychological health or well-being was shown to be an

important variable in cLBP patient delineation. Using longitudinal data from the SwePain

cohort, including 9361 participants with and without chronic pain, psychological well-being

scores at baseline were able to predict pain intensity after 2 years [23]. Within this study, positive

well-being was predictive of lower pain severity in participants without and with chronic pain.

Similar conclusions were drawn from the comparison of back pain patients with different levels

of mental distress, in which patients with higher mental distress showed, among other things,

reduced psychological well-being and social function, and higher severe pain than patients with

lower mental distress [24]. Furthermore, patients suffering from chronic pain exhibit significantly

lower quality of life scores across all sub-domains, including psychological well-being [25].

Alterations in quality of life are stronger associated with changes in social functioning and

psychological well-being (via pain catastrophizing) than pain intensity it self [26], indicating the

high importance of psycho-social aspects for daily living with painful conditions such as back

pain.

Our findings highlight the potential importance of psychosocial factors in cLBP and suggest a

need for change in clinical practice. In Berlin, and more widely, the integration of psychosocial

assessment into routine cLBP care could improve treatment effectiveness. This is consistent

with previous meta-analytic evidence [27] supporting a combined biopsychosocial therapeutic

approach to optimise treatment efficacy. Early identification of psychosocial barriers such as

anxiety, depression or stress should be prioritised, with interventions such as cognitive

behavioural therapy, stress management and mental health support used alongside traditional

physical therapies.

We demonstrated that spinal herniation and degeneration observed on MRI may contribute to

pain mechanisms in the bio-psychosocial model of cLBP, consistent with previous meta-

analyses [28]. However, it is clear that MRI findings alone do not fully explain pain presence in

cLBP, highlighting the need for caution when interpreting MRI results at the individual patient

level. Interestingly, cervical spine rotation but not lumbar back motion assessments were shown

to be robust important examinations for cLBP delineation. This contradictory finding is likely

a result of two factors. First, the poor cervical rotation may be the result of neck pain that has

high comorbidity with cLBP [29] and can lead to decreased axial rotation [30]. Second, poor

psychological health has been associated with neck pain [31], which we found to also present high

importance in patient stratification and relates to the bio-physical-psychosocial interplay

present in cLBP patients [5]. A systematic review on hip mobility in LBP patients [32] showed small

to no changes in hip flexion compared to controls. As all studies had less than 110 subjects,

they were likely under powered to uncover the decreased mobility we show here and may

represent a diagnostic test for LBP. Previous studies have shown that clinical kinematic data

can effectively stratify cLBP patients into high, low and intermediate risk groups [33], suggesting

that pain correlates with reduced physical function. Persistent nociceptive input from

aggravated spinal joints/muscles may lead to reduced motor output and spinal cord excitability

[34], potentially resulting in a reduced ability to recruit specific muscles and necessitating

compensatory movement strategies. Our findings underscore that detailed movement analysis

could serve as a diagnostic biomarker for LBP, potentially rivalling medical imaging in

diagnostic accuracy and improving patient care by identifying sub-populations likely to respond

well to specific therapies or at risk of adverse outcomes.

Classification of cLBP patients has often been conducted on relatively small sample sizes (<

200) as well as utilising a single data domain [9]. Performance of such models are subject to

overfitting due to their small samples and would likely perform poorly at out-of-sample

classification in unseen external datasets [35]. Furthermore, several studies have established

classification models with high accuracy (> 0.8) at determining particular LBP symptoms [36–43],

although these lack clinical applicability in understanding the most appropriate variables and

modalities in classification of cLBP as well as the classification of the disorder in general.

Classification models created using large datasets (n > 1000) either contained psychosocial and

demographic variables, without imaging and physical variables [44,45]. On the other hand, Jin-

Heekun and colleagues [46] employed only physical variables without considering important

psychosocial factors, which have shown to be important in previous research [5,20,47,48] as well as

in our current study (Fig. 4). Our best model (Boruta – Q + C + M) performed slightly worse

than Parsaeian et al. [45] (AUC 0.693 – 0.75) and compared to Shim et al. [44] (AUC, 0.693 – 0.716),

we utilised more plentiful data points per subject in a significantly smaller sample size

(approximately 34x and 6x smaller respectively) to address the clinically relevant question of

what modalities and variables a best suited for cLBP classification.

The multimodal nature of the data utilised and the amount of subjects that have participated in

physical, imaging, and questionnaire measurements are major strengths in our current study.

Utilising the "Berlin Back Study" dataset that contains more than 500 subjects in the different

domains highlighted as lacking in a recent review by our group [9], enabling us to robustly

investigate the importance of different modalities as well as specific variables in cLBP patient

stratification. The large sample size enabled us to minimise model overfitting through cross-

validation and hold-out testing of good sample size and distributions. A more accurate

representation of a models performance is provided by out-of-sample testing that uses a new

sample population containing comparable variables. This provides a test set with minimised

sampling and dataset bias greatly improving the generalisability and applicability of the

findings.

The results of our study offer valuable insights into the potential of using a multimodal machine

learning approach for the classification of individuals with chronic low back pain (cLBP).

However, despite the promising nature of our dataset, the classification performance was

moderate. One possible explanation is the size and diversity of the dataset. Although the sample

size was substantial, it may not fully encompass the complex and multifactorial nature of cLBP,

especially when considering the broad spectrum of psychosocial, environmental, socio-

economic, and biological factors that contribute to the condition. Incorporating detailed

information regarding socio-economic status, education, and other social factors could

potentially improve model performance by capturing the broader context in which cLBP

manifests.

While our study highlights the importance of psychosocial factors in the classification of cLBP,

we acknowledge that the analysis of MRI data was limited. Several key MRI phenotypes,

including disc bulging, spondylolisthesis, osteophytes, Modic lesions, endplate abnormalities,

and high-intensity zones, were not included in the final analysis. It is also important to note that

while psychosocial factors showed strong predictive power, the MRI analysis was not

exhaustive, and a more complete MRI dataset would be necessary to make a fair comparison.

Future studies should aim to expand both the MRI and psychosocial data to provide a more

comprehensive understanding of the relative contributions of anatomical and psychosocial

factors in cLBP.

A notable limitation of this study is the absence of key social determinants of health, such as

income, education, and health insurance status, which are known to significantly influence both

the risk of developing cLBP and its persistence. While we included job-related factors, such as

posture during work, these social determinants were not available in our dataset. Incorporating

these factors into future models could enhance the accuracy of our predictions and offer a more

nuanced understanding of how socioeconomic and environmental factors intersect with clinical

and psychosocial elements in the context of cLBP. The inclusion of such variables should be a

priority in subsequent research to improve both model performance and clinical applicability.

We acknowledge the challenge of translating these findings into clinical practice. The

complexity and costs associated with collecting such comprehensive data may not be feasible

in all healthcare settings. However, the value of this research lies in its potential to inform the

development of more personalized, evidence-based treatment strategies. In future work, we aim

to explore how these multimodal data could contribute to treatment recommendations or patient

phenotyping, providing clinicians with more precise tools to tailor interventions to individual

patient needs. By advancing the field toward personalized care, we believe the clinical utility

of this approach will become more apparent. Furthermore, our study provides a cross-sectional

investigation of cLBP, whereas a prospective design would be better suited to examine the

causality of the disorder. Two different types of interviews were conducted: face-to-face

interviews (clinical examination) and electronic interviews (questionnaires). The main

difference is that body language, facial expressions and other non-verbal social cues are obvious

to the interviewer in face-to-face interviews, whereas these aspects are absent in electronic

surveys. As both surveys have advantages and disadvantages, the answers of the study

participants were weighted equally in this study. Additionally, the high number of questions

could lead to a reduction in the participants' attention and concentration.

In conclusion, while our current model shows promise, there is room for improvement.

Expanding the dataset, incorporating more detailed and diverse data sources, and exploring

alternative machine learning models are potential next steps that could enhance the predictive

power of the system. A key future application will be the use of baseline data to predict

treatment response and the transition from acute to chronic pain. This predictive capability

could enable early identification of high-risk patients, improving treatment outcomes and

advancing personalised medicine in the management of cLBP. We also believe that as more

data becomes available and our understanding of the complex interplay of factors contributing

to cLBP deepens, the clinical utility of such models will become more apparent, leading to

better patient stratification and more personalized treatment approaches.

Methods

Quantitative variables and data collection

Patients had to meet all of the following inclusion criteria: written informed consent to participate in the study, asymptomatic (no back pain) or symptomatic (cLBP) caucasian women and men aged 18–67 years, pain duration ≥12 weeks daily (cLBP only), pain localization in the lumbopelvic region (cLBP only). A telephone interview was conducted during recruiting and subjects were excluded if they met any exclusion criteria. However, some subjects came to testing that should have been excluded during the telephone interview, and were then excluded at the testing site (Supplementary Table S3). No minimal threshold for LBP intensity was defined. A list of all variables including number of missing values is shown in Supplementary Table S19.

Questionnaires

The localization, type, course, possible radiation, intensity, quality, duration, and any factors that may relieve or exacerbate the pain, as well as possible triggers or the patient's own explanations regarding the cause of the pain has been asked. The patient's medical history has been recorded, which includes any previous diseases and surgeries, and a detailed pain and general medication history as well as allergies, intolerances, and vaccination status among others. A family and social history (anamnesis) was taken (professional activity, family situation, diseases in the family, stressful situations, etc.). In addition, any past or present use of addictive substances has been asked (alcohol, nicotine, etc.).

The following questionnaires were completed within 30 minutes:

➢ Pain intensity, pain duration, and pain-related disability: von Korff et al. [49]

➢ Disability Questionnaire: Roland and Morris (RMDQ) [50]

➢   Short-form 36 Health Status Questionnaire: SF-36 [51]. The following four domains were

considered: general mental health (psychological distress and well-being), limitations in

usual role activities because of emotional problems, vitality (energy and fatigue), and

general health perceptions.

➢   International Physical Activity Questionnaire (IPAQ) [52]

➢   Self-Report Behavioural Automaticity Index (SRBAI) [53]

➢   Behavioural Regulation in Sport Questionnaire (BRSQ) [54]

➢   Tampa Scale for Kinesiophobia (TSK-GV) [55].

➢   Fear-Avoidance Belief Questionnaire (FABQ) [56].

The participants primarily answered the questionnaires in digital form using a survey program

specially developed for the study. The data were collected under similar conditions (e.g., same

room, same computer) for all subjects at the study centre.

Demographic data

During the clinical assessment, age, sex, body height, body weight, hip diameter, and waist

diameter of the subjects were recorded. BMI was chosen instead of waist hip ratio to measure

physical body size and health, as the BMI variable contained less missing values compared to

waist hip ratio (Supplementary Table S19).

Clinical examination

The clinical examination included the evaluation of organ functions (inspection, palpation,

percussion, and auscultation), the general impression, and the vital parameters of the patient

(temperature, heart rate, blood pressure, etc.). Examination was performed by an experienced

orthopaedic consultant. The neurological status was assessed by the examination of the

coordination, reflexes, sensitivity, and motor function. The evaluation of the functional

parameters, that is, the assessment of posture, shape, orientation, and movement of the lumbar

spine and pelvis, was based on current clinical standards (e.g., Ott and Schober test, 3-step

hyperextension test, passive lumbar extension test, etc.) and self-assessment by the persons

investigated. Data were documented according to their dimension using distances in cm,

degrees of angle, and number of repetitions per defined time interval or bivalent whether pain

provocation occurred. Self-assessment of functional restrictions of the back was recorded

according to a scale from 1 (best) – 10 (worst).

Back shape and function

All study participants received measurements of the back shape in the sagittal and frontal planes

during upright standing and sitting using the Idiag M360 (Idiag AG, Fehraltorf, Switzerland).

The device measures segmental angles of the thoracic and lumbar spine. In both postures, study

participants were measured upright, in flexed, extended, and in left and right lateral bending (3

repetitions, ~10 sec each). Maximum upper body flexion, extension, as well as left and right

lateral bending were performed with extended knees. During extension the arms were crossed

in front of the body. The order of performed tasks was randomised. The measurements were

performed by trained medical students. The validity and reliability were demonstrated in

previous studies [57–60].

Spino-pelvic MRI

MRIs were conducted using a 1.5 MRI scanner. Following sequences were evaluated: 1) Sag

T1 (4 mm slices), 2) Sag T2 (4 mm slices), 3) Cor STIR-T2 (4 mm slices) and 4) Axial T2

(3 mm slices). MRIs were evaluated for intervertebral disc degeneration (Pfirrmann

classification [61]), disc herniation (Kramer classification [62]), facet joint arthrosis (Fujiwara

classification [63]), osteochondrosis intervertebralis [64], spondylolisthesis (Meyerding

classification [65] and spinal canal stenosis (Schizas classification [66]) at each level of the lumbar

spine. The spino-pelvic MRI evaluation was performed blinded by two spine surgeons and a

radiologist, all of whom have many years of experience in the evaluation of spinal pathologies.

The inter-rater reliability was good-to-excellent for all measurement parameters.

Data storage

All data files electronically recorded during the study period were stored on a database server

folder (SharePoint folder) hosted by Charite-Universitaetsklinikum. A data back-up for the

database is run daily. Local study team members signed a non-disclosure agreement. They have

access to the database using a personal password and are authorized only for entries depending

on their function based on a role concept (investigator, statistician, monitor, administrator etc.).

A multilevel data validation plan was developed to guarantee the correctness and consistency

of the data. Data were entered only after a check for completeness and plausibility. Furthermore,

data were cross-checked for plausibility with previously entered data for each participant.

Questionnaires filled out on paper are stored in a lockable cabinet at the university.

Potential sources of bias and minimisation

To generally reduce the possible location and assessor bias, measurements (clinical physical

and questionnaire assessment and back shape and function) were administered by a few trained

clinicians in the same room with the same lighting. The self-administered questionnaires (von

Korff, RMDQ, SF-36, IPAQ, SRBAI, BRSQ, TSK-GV, and FABQ) were completed by the

subjects under supervision by our trained study coordinator who provided explanations for

unclear questions and mitigated possible lack of motivation to complete the questions by

assuring the subjects of the importance to complete the questionnaires. Furthermore, generic

questionnaires (SF-36 or IPAQ) were placed before specific ones (SRBAI, BRSQ) to minimize

bias from order effects.

To minimise the bias in our classification and variable selection modelling, as well as the

modality comparison, variables directly assessing back pain and questions heavily biased to

pain patients were removed. Such variables assessed back pain during particular movements or

upon physical manipulation. Furthermore, questions only back pain patients were asked for

example, pain intensity, duration, and disability, as well as pain and health biased self-

questionnaires (von Korff, RMDQ, TSK-GV, FABQ, therapies, and the SF-36 sub-categories

regarding physical function, physical role function, physical pain, health perception, and

vitality), and clinician administered questions regarding pain medication intake, previous spinal

disorder diagnosis, and participants' subjective physical health assessment were removed as

well to reduce model bias.

Outcome

The outcome target for our classification model is cLBP patients. All participants were assessed

by a clinician and diagnosed as cLBP patient, asymptomatic control, or suffered from LBP in

the past but not at present. We used the clinician diagnosis of either current cLBP patient or

asymptomatic control as our two-class target outcome. The participants with LBP in the past

were removed to enable better distinguishable groups for binary classification and variable

importance selection.

Data handling, preprocessing, cleaning and missing data

551 Total and sub-scores for the self-administered questionnaires were used after removal of pain

552 biased questionnaires (see *Potential sources of bias and minimisation*). This includes the SF-

553 36 [51], the IPAQ [52], the SRBAI [53], and the BRSQ [54]. The SF-36 was used to collect statements

554 related to the health domains 'emotional role limitation, (three items) 'social functioning' (two

items), and 'mental health' (five items). A scoring algorithm was used to convert the raw scores

into these three domains. The scores were transformed to range from zero (worst possible

health) and 100 (best possible health). The SRBAI and BRSQ collected ratings for multiple

statements on a numerical scale from 1 (strongly disagree) to 6 (strongly agree). For the SRBAI,

the total score was calculated by summing the numerical values across all 4 statements. In

contrast, sub-scores were created for the BRSQ that related to 'intrinsic motivation', 'integrated

regulation' and 'external regulation'. These scores were calculated by summing the numerical

values across two statements per sub-score. The IPAQ, recorded the average time spent per day

over the past 7 days while 'sitting', 'walking', doing 'moderate activities' (e.g., heavy lifting,

digging, aerobics, or fast bicycling), and 'vigorous activities' (e.g. carrying light loads,

bicycling at a regular pace, or doubles tennis). To estimate the energy requirements for each

activity type, the average time spent per day in minutes was multiplied by MET-score

(metabolic equivalents) of 1.5, 3.3, 4, or 8 for sitting, walking, moderate and vigorous activities,

respectively. This resulted in MET-minutes scores, describing the amount of energy in

kilocalories required for a 60 kilogram person. Finally, the MET-minutes scores were summed

across 'sitting', 'walking', 'moderate activities' and 'vigorous activities', resulting in a total

MET-minutes score per subject.

Modelling preprocessing was conducted by checking variables within each modality for very

low variance and collinearity. Variables with close to zero (variance < 1) variance were

removed. Furthermore, variables presenting a Spearman correlation greater than 0.9 were also

removed to reduce collinearity between variables. The decision on which of the correlated

variables to removed, was the variable showing less correlation to the target, cLBP patient

status. Following these cleaning steps, single data modalities were joined with demographic

data and subsequently joined with other datasets for the dual- and multi-modalities datasets.

Subjects having any missing values were removed from our main analyses and MissForest [18]

581 imputation was additionally conducted. Imputation was computed for the training and testing

582 samples within each fold independently to avoid data leakage. Following cleaning and

583 preprocessing modality dataset presented different number of subjects and variables with a

584 similar age and sex distribution across cLBP patients and asymptomatic controls (Fig. 1A).

Univariate statistics

The univariate statistics were carried out separately for continuous, ordinal and nominal data to

compare patients suffering from cLBP against asymptomatic controls using R (version 4.3.1;

www.r-project.org). As most continuous variables did not follow a normal distribution

according to the Anderson-Darling test [67], we implemented the non-parametric Wilcoxon-

Mann-Whitney test to determine significant difference between cLBP patients and

asymptomatic controls for ordinal and continuous data. Hence, u-values, z-values, r-value

(effect sizes), as well as the p-value are reported from. Nominal data were compared using the

Chi-Square test and reported with Chi^2 values, Cohen's ω-values (effect size), and p-values.

Statistical significance was determined at p≤0.05 following family wise error (FWE) [68]

correction for multiple comparisons within each modality (demographics, questionnaires,

clinical examinations, superficial spine morphology and motion, and spino-pelvic MRI).

Machine learning

Boruta variable selection

We used the Boruta method [16] for importance variable selection. Boruta utilises random forest

(RF) classification algorithm [17] with both the real variables and set of 'dummy' or 'shadow'

variables, that are created by shuffling the variable values. This creates random variables

(dummy features) that have the same distribution as the original variables, although represent

the classification accuracy of this variable randomly sampled. As these dummy variables

represent random noise, they had their possible correlation to the target (cLBP) removed. All

real and dummy variables are used to classify cLBP and variable importance is calculated.

Variable importance is calculated as the Z-score of the mean decrease in classification accuracy

following the removal of this variable from the model. The importance of the dummy variables

can be used as a reference to test the variable importance of the real variables. Through an

iterative process, real variables that have significantly greater importance than the maximum

dummy variable importance are marked as important. The variables that have a significantly

lower importance than the maximum dummy variables are deemed unimportant and removed

for the next iteration of selection. Iterations are repeated until the importance is assigned to all

variables or a user defined iteration number is reached. We used a max of 2000 iterations with

a random forest containing 1000 trees. As there are occasionally a few variables not definitively

identified as important or removed after 2000 iterations by Boruta, we only selected the

variables that have been confirmed as important in our subsequent modelling and analyses.

Random forest classification algorithm

We utilised RF [17] implemented using the ranger package [69] in R (version 4.3.1, www.r-

project.org) to classify cLBP patients and pain-free controls. As RF provides variable

importance measures and can handle categorical, ordinal, and continuous data it represents the

ideal choice to deal with the different data types present in the Berlin Back dataset. Ten-fold

cross validation was conducted during model training and hyper-parameter tuning utilising

1000 tree RF. We conducted hyper-parameter tuning using a tuning parameter search grid,

which contained number of variables to be sampled at each split (mtry) of 1 – square root of

the number of variables, and a minimum node size of 5 and 10. Therefore, using a gini split rule

a grid search was conducted with all combination of hyper-parameters to determine the best. A

ten-fold train-test loop (Fig. 1B) was conducted to determine the cLBP classification

performance within each of the 15 datasets utilising all and Boruta selected variables

independently. Model performance was calculated as follows; accuracy = (TP + TN) / (TP +

FP + TN + FN), sensitivity = TP / (TP + FN), specificity = TN / (TN + FP), and AUC (area

under the receiver operating characteristic (ROC) curve). Whereby, TP – true positive, TN –

true negative, FP – false positive, and FN – false negative. The ROC curve represents the

classification performance measured by sensitivity and specificity over a range [0, 1] of

classification thresholds.

Robust variable selection workflow

Selecting important variables and comparing the 15 different dataset modalities was conducted

on 90% (n = 1045, 19 – 72 y/o, mean age = 41.71 ± 12.28, cLBP = 469) of the preprocessed

Berlin Back study dataset, with a 10% (n = 116, 20 – 64 y/o, mean age = 43.81 ± 12.12, cLBP

= 43) hold-out set used to evaluate the most robust and important variables (Fig. 1B). The hold-

out sample comprised of participants that had data for all variables. Each 15 modality datasets

went through the iterative ten-fold train-test loop, where Boruta variable selection was

conducted on the training set. Followed by RF training using all and Boruta selected variables

implemented on the training dataset. Model performance was calculated using the test set.

Across the ten loops, the percentage a variable was selected and average importance value was

calculated across all 15 datasets. Finally, the variables that were selection within every train-

test loop and within every possible dataset were provided as the most robust and important

variables for cLBP classification. The demographic variables can be selected a maximum of 15

times, while the modality-specific variables a maximum of 8 times across the 15 dataset

modalities. These variables were then used in a five-fold train test loop on the hold-out dataset

and compared to a model trained using all variables.

Disclosures

Funding: This study is part of the Research Unit FOR 5177 funded by the German Research Foundation (DFG), Hendrik Schmidt: SCHM 2572/11-1, SCHM 2572/12-1, SCHM 2572/13-1; Sandra Reitmaeier: RE 4292/3-1, Matthias Pumberger: PU762/1-1. The analyses and contribution from the Hochschule für Gesundheit were funded, in part, by grant number 50WK2273A (to DLB) from the German AeroSpace Center (DLR).

Conflicts of interest: All authors declare no conflict of interests.

Acknowledgments

We would like to thank all patients and healthy participants for their selfless participation in this study and the participating companies for informing their employees about this study.

Data and code availability

All results in this study are provided in the (Supplementary) tables. The Berlin Back study is currently ongoing (end date 31/12/2025) and therefore the raw data used in this manuscript cannot be provided. The raw data will be openly released from the Berlin Back Study as per agreement with the funding agency following the completion of the data acquisition. A link to the raw data will be provided on the Github repository where the analysis code is located (https://github.com/viko18/BerlinBack_FeatImp/) when it is made available.

Author Contributions

CRediT Contributions:

Conceptualization – SV, HS, DLB

Methodology – SV, HS, DLB, MA, CK

Software – SV, FJ

Validation – SV

Formal Analysis – SV, FJ

Investigation – SV, LAB, NT, MP, SR

Resources – HS, DLB

Data Curation – SV FJ

Writing original draft – SV, RD, FJ, DLB, HS

Writing review & editing – All co-authors

Visualizations – SV, FJ

Supervision – DLB, HS, MA, CK

Project Administration – HS, SV

Funding Acquisition – HS, DLB

References

1. Andersson, G. B. Epidemiologic aspects on low-back pain in industry. *Spine (Phila Pa 1976)* **6**, 53–60 (1981).

2. Meucci, R. D., Fassa, A. G. & Faria, N. M. X. Prevalence of chronic low back pain: systematic review. *Rev Saude Publica* **49**, 1 (2015).

3. Pastorino, R. *et al.* Benefits and challenges of Big Data in healthcare: an overview of the European initiatives. *Eur J Public Health* **29**, 23–27 (2019).

4. Shilo, S., Rossman, H. & Segal, E. Axes of a revolution: challenges and promises of big data in healthcare. *Nat Med* **26**, 29–38 (2020).

5. Tagliaferri, S. D. *et al.* Relative contributions of the nervous system, spinal tissue and psychosocial health to non-specific low back pain: Multivariate meta-analysis. *Eur J Pain* (2021) doi:10.1002/ejp.1883.

6. Grotle, M. *et al.* Lumbar spine surgery across 15 years: trends, complications and reoperations in a longitudinal observational study from Norway. *BMJ Open* **9**, e028743 (2019).

7. Lee, W. *et al.* Identifying and Assessing Interesting Subgroups in a Heterogeneous Population. *Biomed Res Int* **2015**, 462549 (2015).

8. Lötsch, J. & Ultsch, A. Machine learning in pain research. *Pain* **159**, 623–630 (2018).

9. Tagliaferri, S. D. *et al.* Artificial intelligence to improve back pain outcomes and lessons learnt from clinical classification approaches: three systematic reviews. *npj Digital Medicine* **3**, 93 (2020).

10. Noroozi, Z., Orooji, A. & Erfannia, L. Analyzing the impact of feature selection methods on machine learning algorithms for heart disease prediction. *Sci Rep* **13**, 22588 (2023).

11. Mwangi, B., Tian, T. S. & Soares, J. C. A review of feature reduction techniques in neuroimaging. *Neuroinformatics* **12**, 229–244 (2014).

12. World Medical Association. World Medical Association Declaration of Helsinki: ethical

principles for medical research involving human subjects. *JAMA* **310**, 2191–2194 (2013).

13. von Elm, E. *et al.* The Strengthening the Reporting of Observational Studies in

Epidemiology (STROBE) Statement: Guidelines for reporting observational studies.

Preventive Medicine **45**, 247–251 (2007).

14. Collins, G. S., Reitsma, J. B., Altman, D. G. & Moons, K. G. M. Transparent Reporting of

a multivariable prediction model for Individual Prognosis or Diagnosis (TRIPOD): the

TRIPOD statement. *Ann Intern Med* **162**, 55–63 (2015).

15. Shadbahr, T. *et al.* The impact of imputation quality on machine learning classifiers for

datasets with missing values. *Commun Med* **3**, 1–15 (2023).

16. Kursa, M. B. & Rudnicki, W. R. Feature Selection with the Boruta Package. *Journal of*

Statistical Software **36**, 1–13 (2010).

17. Breiman, L. Random Forests. *Machine Learning* **45**, 5–32 (2001).

18. Stekhoven, D. J. & Bühlmann, P. MissForest—non-parametric missing value imputation

for mixed-type data. *Bioinformatics* **28**, 112–118 (2012).

19. Ge, L., Pereira, M. J., Yap, C. W. & Heng, B. H. Chronic low back pain and its impact on

physical function, mental health, and health-related quality of life: a cross-sectional study

in Singapore. *Sci Rep* **12**, 20040 (2022).

20. Tagliaferri, S. D. *et al.* Chronic back pain sub-grouped via psychosocial, brain and

physical factors using machine learning. *Sci Rep* **12**, 15194 (2022).

21. Iguti, A. M., Guimarães, M. & Barros, M. B. A. Health-related quality of life (SF-36) in

back pain: a population-based study, Campinas, São Paulo State, Brazil. *Cad Saude*

Publica **37**, e00206019 (2021).

22. Takeyachi, Y. *et al.* Correlation of low back pain with functional status, general health

perception, social participation, subjective happiness, and patient satisfaction. *Spine*

(Phila Pa 1976) **28**, 1461–1466; discussion 1467 (2003).

23. Larsson, B., Dragioti, E., Gerdle, B. & Björk, J. Positive psychological well-being

predicts lower severe pain in the general population: a 2-year follow-up study of the

SwePain cohort. *Ann Gen Psychiatry* **18**, 8 (2019).

24. Hnatešen, D. *et al.* Quality of Life and Mental Distress in Patients with Chronic Low

Back Pain: A Cross-Sectional Study. *Int J Environ Res Public Health* **19**, 10657 (2022).

25. Hadi, M. A., McHugh, G. A. & Closs, S. J. Impact of Chronic Pain on Patients' Quality of

Life: A Comparative Mixed-Methods Study. *J Patient Exp* **6**, 133–141 (2019).

26. Lamé, I. E., Peters, M. L., Vlaeyen, J. W. S., Kleef, M. v & Patijn, J. Quality of life in

chronic pain is more associated with beliefs about pain, than with pain intensity. *Eur J*

Pain **9**, 15–24 (2005).

27. Kamper, S. J. *et al.* Multidisciplinary biopsychosocial rehabilitation for chronic low back

pain: Cochrane systematic review and meta-analysis. *BMJ* **350**, h444 (2015).

28. Brinjikji, W. *et al.* MRI Findings of Disc Degeneration are More Prevalent in Adults with

Low Back Pain than in Asymptomatic Controls: A Systematic Review and Meta-

Analysis. *AJNR Am J Neuroradiol* **36**, 2394–2399 (2015).

29. von der Lippe, E. *et al.* Prevalence of back and neck pain in Germany. Results from the

BURDEN 2020 Burden of Disease Study. *J Health Monit* **6**, 2–14 (2021).

30. Rampazo, É. P. *et al.* Sensory, Motor, and Psychosocial Characteristics of Individuals

With Chronic Neck Pain: A Case Control Study. *Physical Therapy* **101**, pzab104 (2021).

31. Mansfield, M. *et al.* The association between psychosocial factors and mental health

symptoms in cervical spine pain with or without radiculopathy on health outcomes: a

systematic review. *BMC Musculoskeletal Disorders* **24**, 235 (2023).

32. Avman, M. A., Osmotherly, P. G., Snodgrass, S. & Rivett, D. A. Is there an association

between hip range of motion and nonspecific low back pain? A systematic review.

Musculoskeletal Science and Practice **42**, 38–51 (2019).

33. Abdollahi, M. *et al.* Using a Motion Sensor to Categorize Nonspecific Low Back Pain

Patients: A Machine Learning Approach. *Sensors* **20**, 3600 (2020).

34. Nijs, J. *et al.* Nociception affects motor output: a review on sensory-motor interaction

with focus on clinical implications. *Clin J Pain* **28**, 175–181 (2012).

35. Rajput, D., Wang, W.-J. & Chen, C.-C. Evaluation of a decided sample size in machine

learning applications. *BMC Bioinformatics* **24**, 48 (2023).

36. Al Imran, A., Rifat, M. R. I. & Mohammad, R. Enhancing the classification performance

of lower back pain symptoms using genetic algorithm-based feature selection. in 455–469

(Springer, 2020).

37. Abdullah, A. A., Yaakob, A. & Ibrahim, Z. Prediction of Spinal Abnormalities Using

Machine Learning Techniques. in 1–6 (IEEE, 2018).

38. Riveros, N. A. M., Espitia, B. A. C. & Pico, L. E. A. Comparison between K-means and

self-organizing maps algorithms used for diagnosis spinal column patients. *Informatics in*

Medicine Unlocked **16**, 100206 (2019).

39. Sandag, G. A., Tedry, N. E. & Lolong, S. Classification of lower back pain using K-

Nearest Neighbor algorithm. in 1–5 (IEEE, 2018).

40. Karabulut, E. M. & Ibrikci, T. Effective automated prediction of vertebral column

pathologies based on logistic model tree with SMOTE preprocessing. *Journal of Medical*

Systems **38**, 50 (2014).

41. Mathew, B., Norris, D., Hendry, D. & Waddell, G. Artificial intelligence in the diagnosis

of low-back pain and sciatica. *Spine* **13**, 168–172 (1988).

42. Vaughn, M. L., Cavill, S. J., Taylor, S. J., Foy, M. A. & Fogg, A. J. Direct explanations

for the development and use of a multi-layer perceptron network that classifies low-back-

pain patients. *International Journal of Neural Systems* **11**, 335–347 (2001).

43. Zhang, W. *et al.* Deep learning-based detection and classification of lumbar disc

herniation on magnetic resonance images. *JOR SPINE* **6**, e1276 (2023).

44. Shim, J.-G. *et al.* Machine Learning Approaches to Predict Chronic Lower Back Pain in

People Aged over 50 Years. *Medicina* **57**, 1230 (2021).

45. Parsaeian, M., Mohammad, K., Mahmoudi, M. & Zeraati, H. Comparison of logistic

regression and artificial neural network in low back pain prediction: second national

health survey. *Iranian Journal of Public Health* **41**, 86 (2012).

46. Jin-Heeku. Analysis of sitting posture using wearable sensor data and support vector

machine model. *Medico-Legal Update* **1**, 334–338 (2018).

47. Tagliaferri, S. D. *et al.* Brain structure, psychosocial, and physical health in acute and

chronic back pain: a UKBioBank study. *Pain* **163**, 1277–1290 (2022).

48. Tagliaferri, S. D. *et al.* Towards data-driven biopsychosocial classification of non-specific

chronic low back pain: a pilot study. *Sci Rep* **13**, 13112 (2023).

49. Von Korff, M., Ormel, J., Keefe, F. J. & Dworkin, S. F. Grading the severity of chronic

pain. *PAIN* **50**, 133 (1992).

50. Roland, M. & Fairbank, J. The Roland–Morris Disability Questionnaire and the Oswestry

Disability Questionnaire. *Spine* **25**, 3115 (2000).

51. Ware, J. E. & Sherbourne, C. D. The MOS 36-item short-form health survey (SF-36). I.

Conceptual framework and item selection. *Med Care* **30**, 473–483 (1992).

52. Craig, C. L. *et al.* International Physical Activity Questionnaire: 12-Country Reliability

and Validity. *Medicine & Science in Sports & Exercise* **35**, 1381 (2003).

53. Gardner, B., Abraham, C., Lally, P. & de Bruijn, G.-J. Towards parsimony in habit

measurement: Testing the convergent and predictive validity of an automaticity subscale

of the Self-Report Habit Index. *International Journal of Behavioral Nutrition and*

Physical Activity **9**, 102 (2012).

54. Lonsdale, C., Hodge, K. & Rose, E. A. The behavioral regulation in sport questionnaire

(BRSQ): Instrument development and initial validity evidence. *Journal of Sport &*

Exercise Psychology **30**, 323–355 (2008).

55. Rusu, A. C., Kreddig, N., Hallner, D., Hülsebusch, J. & Hasenbring, M. I. Fear of

movement/(Re)injury in low back pain: confirmatory validation of a German version of

the Tampa Scale for Kinesiophobia. *BMC Musculoskelet Disord* **15**, 280 (2014).

56. Waddell, G., Newton, M., Henderson, I., Somerville, D. & Main, C. J. A Fear-Avoidance

Beliefs Questionnaire (FABQ) and the role of fear-avoidance beliefs in chronic low back

pain and disability. *PAIN* **52**, 157 (1993).

57. Dreischarf, B. *et al.* Comparison of three validated systems to analyse spinal shape and

motion. *Sci Rep* **12**, 10222 (2022).

58. Guermazi, M. *et al.* [Validity and reliability of Spinal Mouse to assess lumbar flexion].

Ann Readapt Med Phys **49**, 172–177 (2006).

59. Topalidou, A., Tzagarakis, G., Souvatzis, X., Kontakis, G. & Katonis, P. Evaluation of

the reliability of a new non-invasive method for assessing the functionality and mobility

of the spine. *Acta Bioeng Biomech* **16**, 117–124 (2014).

60. Barrett, E., McCreesh, K. & Lewis, J. Reliability and validity of non-radiographic

methods of thoracic kyphosis measurement: a systematic review. *Man Ther* **19**, 10–17

(2014).

61. Pfirrmann, C. W., Metzdorf, A., Zanetti, M., Hodler, J. & Boos, N. Magnetic resonance

classification of lumbar intervertebral disc degeneration. *Spine* **26**, 1873–8 (2001).

62. Kraemer, J. Natural course and prognosis of intervertebral disc diseases. International

Society for the Study of the Lumbar Spine Seattle, Washington, June 1994. *Spine (Phila*

Pa 1976) **20**, 635–639 (1995).

63. Fujiwara, A. *et al.* The relationship between facet joint osteoarthritis and disc

degeneration of the lumbar spine: an MRI study. *Eur Spine J* **8**, 396–401 (1999).

64. Modic, M. T., Steinberg, P. M., Ross, J. S., Masaryk, T. J. & Carter, J. R. Degenerative

disk disease: assessment of changes in vertebral body marrow with MR imaging.

Radiology **166**, 193–199 (1988).

65. Meyerding, H. W. Spondyloptosis. *Surgery, Gynecology & Obstetrics* 371–377 (1932).

66. Schizas, C. *et al.* Qualitative grading of severity of lumbar spinal stenosis based on the

morphology of the dural sac on magnetic resonance images. *Spine (Phila Pa 1976)* **35**,

1919–1924 (2010).

67. Anderson, T. W. & Darling, D. A. Asymptotic Theory of Certain 'Goodness of Fit'

Criteria Based on Stochastic Processes. *The Annals of Mathematical Statistics* **23**, 193–

212 (1952).

68. Holm, S. A Simple Sequentially Rejective Multiple Test Procedure. *Scandinavian Journal*

of Statistics **6**, 65–70 (1979).

69. Wright, M. N. & Ziegler, A. ranger: A Fast Implementation of Random Forests for High

Dimensional Data in C++ and R. *Journal of Statistical Software* **77**, 1–17 (2017).