**Genomic characterization uncovers transmission dynamics of Marburg Virus in Rwanda**

**following a single zoonotic spillover event**

**Yvan Butera**[1,2, &, *], **Leon Mutesa**[1,3,4,&,*], Edyth Parker[1,5,6], Raissa Muvunyi[1,7], Esperance

Umumararungu[1,7], Alisen Ayitewala[1,4], Jean Pierre Musabyimana[1,7], Alhaji Olono[1,5], Placide

Sesonga[1,3,4], Olusola Ogunsanya[1,5], Emmanuel Kabalisa[1,7], Oluwatobi Adedokun[1,5], Nelson

Gahima[1,7], Laetitia Irankunda[1,7], Chantal Mutezemariya[1,7], Richard Niyonkuru[1,7], Arlene

Uwituze[1,7], Ithiel Uwizera[1,7], James Kagame[1,7], Arlette Umugwaneza[1,7], John Rwabuhihi[1,7],

Fidele Umwanankabandi[1,7], Valens Mbonitegeka[1,7], Edouard Ntagwabira[1,7], Etienne Kayigi[1,7],

Gerard Izuwayo[1,7], Herve Murenzi[1,7], Therese Mukankwiro[1,7], Nasson Tuyiringire[1,7], Jean Marie

Vianney Uwimana[1,7], Agnes Gasengayire[1,7], Reuben Sindayiheba[1,7], Glory-Ugochi Onyeugo[1,4],

Merawi Aragaw[1,4], Lenny Gitundu[1,4], Radjabu Bigirimana[1,4], Mosoka Fallah[1,4], Adaora

Ejikeme[1,4], Senga Sembuche[1,4], Alice Kabanda[1,4], Jean Claude Mugisha[1,3,4], Emmanuel Edwar

Siddig Francis[1,4], Pierre Gashema[1,4], Jerome Ndayisenga[1,4], Alexis Rugamba[1,3], Faustin

Kanyabwisha[1,3], Gad Murenzi[1,3], Anise Happi[5], Jean Claude Semuto Ngabonziza[1,7], Misbah

Gashegu[1,7], Ayman Ahmed[1,4,7], Noella Bigirimana[1,7], Edson Rwagasore[1,7], Muhammed

Semakula[1,2], Jean Paul Rwabihama[1,2,8], Clarisse Musanabaganwa[1,7], Eric Seruyange[1,8,9], Menelas

Nkeshimana[1,2], Theogene Twagirumugabe[1,10], David Turatsinze[1,11], Eric Remera[1,7], Noel

Gahamanyi[1,7], Sofonias Kifle Tessema[1,4], Isabelle Mukagatare[1,7], Sabin Nsanzimana[1,2],

**Christian Happi**[1,5,12, §,*], **Claude Mambo Muvunyi**[1,7,§,*]

*[&]First authors with equal contribution*

*[§]Last authors with equal contribution*

*[1]Rwanda Joint Task Force for Marburg Virus Disease Outbreak, Ministry of Health, Rwanda Biomedical Centre, Kigali, Rwanda*

*[2]Ministry of Health, Kigali, Rwanda*

*[3]Center for Human Genetics, College of Medicine and Health Sciences, University of Rwanda, Kigali, Rwanda*

*[4]Africa Centre for Disease Control and Prevention, Addis Ababa, Ethiopia*

*[5]African Center of Excellence for Genomics of Infectious Diseases, Redeemer's University, Ede, Osun State, Nigeria*

*[6]Department of Immunology and Microbiology, The Scripps Research Institute, La Jolla, CA, USA*

*[7]Rwanda Biomedical Centre, Kigali, Rwanda*

*[8]Department of Internal Medicine, College of Medicine and Health Sciences, University of Rwanda, Kigali, Rwanda*

*[9]Rwanda Military Referral and Teaching Hospital, Kigali, Rwanda*

*[10]Butare University Teaching Hospital, Huye, Rwanda*

*[11]Kigali University Teaching Hospital, Kigali, Rwanda*

*[12]Department of Immunology and Infectious Diseases, Harvard T H Chan School of Public Health, Boston, MA 02115, USA*

*Corresponding authors

Yvan Butera, MD, PhD

Ministry of Health

Kigali, Rwanda

Email: yvan.butera@moh.gov.rw

Leon Mutesa, MD, PhD

Centre for Human Genetics

College of Medicine and Health Sciences

University of Rwanda

Kigali, Rwanda

Email: l.mutesa@ur.ac.rw

Christian Happi, MSc, PhD

African Center of Excellence for Genomics of Infectious Diseases,

Redeemer's University,

Ede, Osun State, Nigeria

Email: happic@run.edu.ng

Claude Mambo Muvunyi, MD, PhD

Rwanda Biomedical Centre

Kigali, Rwanda

Email : claude.muvunyi@rbc.gov.rw

*Abstract*

The ongoing outbreak of Marburg virus disease (MVD) in Rwanda marks the third largest

historically, though it has exhibited the lowest fatality rate. Genomic analysis has identified a

lineage with limited internal diversity most closely related to a genome sequence from a sporadic

case sampled in 2014 in Uganda, though the lineages have diverged from a common ancestor that

was circulating for decades in the animal reservoir.

Notably, the data also provide evidence that the outbreak resulted from a single zoonotic

transmission event with limited human-to-human transmission, rather than multiple independent

zoonotic transmission events. The Rwandan MVD outbreak prompted a thorough investigation

that included contact tracing, clinical assessment, travel history, sequencing, and serology testing,

to trace the virus's origin. Results of investigations linked the index case to a mining cave inhabited

by *Rousettus aegyptiacus* (the Egyptian fruit bat), where three individuals tested seropositive for

IgG and IgM, further supporting the zoonotic origin of the outbreak through human-animal

interactions.

*Background*

Since its discovery in 1967, the Marburg virus disease (MVD) has emerged as a significant global

health threat, resulting in several outbreaks characterized by alarmingly high case fatality rates

ranging from 22% to 90% (1). The first reported instances of MVD occurred in simultaneous

outbreaks in Marburg and Frankfurt, Germany, as well as in Belgrade, Yugoslavia (now Serbia)

(2). To date, there have been 19 recorded outbreaks, 563 confirmed cases, and 428 deaths

worldwide (4). On September 27, 2024, Rwanda reported its first ever cases of MVD (5). The

ongoing outbreak in Rwanda has confirmed 66 cases, with a case fatality rate of approximately

23% (5).

MVD has two distinct variants: Marburg virus (MARV) and Ravn virus (RAVV), with MARV

being the more prevalent variant, responsible for the majority of sporadic MVD outbreaks

worldwide. MARV is a negative-sense single-stranded RNA virus from the Filoviridae family,

similar to Ebola and possesses inverse-complementary 3′ and 5′ termini (3). MVD is

predominantly a zoonosis, transmitting to humans from either the bat natural reservoir or through

intermediate hosts, including non-human primates. The mechanisms by which bats regulate viral

replication and maintain complex immunity in contrast to humans remain unclear due to a lack of

bat-specific research tools necessary for comparative studies, such as antibodies for flow

cytometry, genomics, and transcriptomics (1,2).  Studies confirm that MARV's natural reservoir

is the Egyptian fruit bat (*Rousettus aegyptiacus*) (6). The virus can cause viral hemorrhagic fever

in both humans and primates. However, the precise transmission dynamics between the natural

reservoir and humans or primates are insufficiently understood, though exposure to contaminated

excretions from fruit bats is likely a contributing factor (6). The virus is transmitted from human

to human, particularly through direct contact with body fluids like blood, saliva, semen, and other

body fluids from infected individuals (7).

The Marburg viral genome consists of 19,114 nucleotides encoding for seven proteins, including

glycoprotein (GP), nucleoprotein (NP) virion protein 30 (VP30), VP35, VP24, VP40 and large

viral polymerase (3).

Genomic characterization of MARV strains has revealed substantial genetic diversity across its

global phylogeny, but exhibits limited genomic variation within outbreaks (8).

Sequencing of the Marburg genome has provided deeper insights into viral evolution and identified

critical mutation hotspots during adaptation to new hosts, particularly in the VP40 protein and the

NP-VP35 intergenic region (9). These mutations can influence the virus's ability to evade the

immune system, therefore enhancing its pathogenicity.

***Statement of the specific scope of the study***

In this study, we used a near real-time genomics sequencing approach to characterize the Marburg

virus in blood samples obtained from patients in the ongoing outbreak in Rwanda. Our aim was to

use the genomic sequencing data to identify and characterize the virus and understand its evolution

and transmission dynamics during the current outbreak. This research underscores the importance

of continuous genomic surveillance which not only facilitates outbreak tracking but also informs

public health strategies for effectively controlling this deadly pathogen.

***Results***

Phylogenetic analyses support that the outbreak resulted from a single zoonotic transmission event

with limited human-to-human transmission rather than multiple independent zoonotic

transmission events (**Figure 1A**). Multiple independent zoonotic transmissions would have

resulted in a set of more diverged sequences introduced from the genetically diverse viral

population in the reservoir (10). The probable index case was a man in his 20's with an

occupational exposure to fruit bats (*Rousettus aegyptiacus*) in a mining cave environment, who

exhibited symptoms highly consisting with the classic clinical presentation of MVD. Additionally,

serological testing of three contacts of the index case exhibited MARV antibodies, further

supporting epidemiological tracing of the index case. The outbreak lineage is most closely related

to a sequence sampled in Kampala, Uganda in September 2014 from a healthcare worker (HCW),

with no secondary cases observed (KP985768). The source of the zoonotic transmission in 2014

was never identified (11). However, the Rwanda-Uganda sister lineages are significantly diverged

from one another, separated by 82 nucleotide substitutions (**Figure 1A**). In Bayesian phylogenetic

reconstructions, we estimate that the two lineages diverged from a common ancestor that circulated

in the animal reservoir in November 2008 [95% highest posterior density, or HPD: May 2007 and

June 2010] (**Supplementary Figure 1 A-B**).

The outbreak lineage in Rwanda is nested within a larger clade that includes diversity sampled

from bats in southwestern Uganda from 2007 onwards, along with two human cases identified in

Uganda in 2017 (**Figure 1A, B**). The closest bat virus (JX458855) was sampled from a juvenile

Egyptian fruit bat (*Rousettus Aegyptiacus)* in 2009 from Python Cave in Queen Elizabeth National

Park, a popular tourist attraction in south western Uganda (**Figure 1B**) (6). The lineage in Rwanda

is significantly diverged from JX458855, sharing a common ancestor that likely circulated in the

reservoir as early as June 2006 (95% HPD: December 2004 to November 2007) (**Supplementary**

**Figure 1 A-B**). *Rousettus aegyptiacus* are a cave dwelling species, with spillover events and

limited outbreaks frequently associated with mining activities. This includes outbreaks and

sporadic cases linked to Python cave and the Kitaka mine that is only 50 km away in 2007 and

2008, which both hosts *Rousettus aegyptiacus* colonies of over 50 000 bats. (6,10,12). The

divergent relationship of the Rwandan Marburg virus outbreak lineage indicates that the virus's

dispersal through host networks involved larger scale animal movement over a long period of time,

rather than a direct ancestral connection to the *Rousettus aegyptiacus* colony in southwestern

Uganda.

**Figure 1**: *A) Maximum likelihood phylogeny of the global MARV dataset. Tips are annotated by*

*country of isolation. This study's tips are relatively enlarged.  B) Map of Uganda and Rwanda.*

*The city of patients in this study (Kigali) is annotated in dark red. Python Cave and Kitaka Mine*

*in Ibanda district in southwestern Uganda are annotated in light blue and yellow, respectively.*

The index patient's close relative was admitted to Hospital 1 (H1) at the end of August 2024 but

succumbed to the illness prior to the definitive diagnosis of MVD. It is highly probable that this

patient infected healthcare workers (HCWs) at H1, where MVD was subsequently confirmed at

the end of September 2024. From 22-28 September, 26 cases were reported in HCWs at H1 and

H2, respectively. The limited genetic variation among the outbreak sequences was consistent with

a very recent common ancestor, the short sampling period, and previous observations from the

Angola MARV outbreak (**Figure 2A**) (12). A cluster of eight identical sequences was sampled,

spanning the full sampling period (C1 in **Figure 2A, B**). The majority of the C1 sequences were

isolated from healthcare workers (HCWs) at H1 from 22-28 September, with reported symptom

onset as early as mid-September (**Figure 2B**). A C1 sequence was sampled from a HCW at H2 at

the end of September, suggesting infection from an unsampled C1 case in H2 or dissemination

between the two healthcare facilities. There are recorded cases of healthcare workers moving

between healthcare settings H1 and H2. This is further supported by cluster C2, consisting of two

healthcare workers from H1 and H2 respectively that share the synonymous SNP C11070T relative

to C1. Both C2 cases report symptom onset and were sampled in the later weeks of the epidemic.

There are a further seven identical sequences (C4) sampled between H1 and H2 across the full

sampling period that are one synonymous SNP (T10031C) away from C1. C002, a healthcare

worker employed at both H1 and H2, is believed to have acquired the infection at H1. This

individual subsequently became the index case at H2, leading to a nosocomial cluster at H2 and

infections at H1. Overall, there were only three synonymous SNPs observed in the outbreak clade:

A5856G (n=1 sequence), T10031C (n=7), and C11070T (n=2). A5856G and T10031C as well as

C11070T are outside of the CDS of the GP and VP24 genes, respectively. There is therefore no

evidence that post-emergence substitutions enhanced human transmission.

**Figure 2**: *A) Maximum likelihood phylogeny of the outbreak clade, with the Kampala outgroup.*

*Sequences are colored with associated hospital. SNPs reconstructed relative to the common*

*ancestor are annotated in text, as are the clusters defined by the SNP presence. The coordinates*

*are relative to NC_001608 B) Epidemiological timeline of the sampled sequences in Figure 2A,*

*as annotated on the y-axis. The first black marker, if present, indicates the week of symptom onset*

*if distinct from week of sampling, with the second indicating the week of sampling. Sequences are*

*colored by their associated hospital (H1, H2), and partitioned by SNP presence into clusters, as*

*annotated in text. Red dashed line indicates the contact of the index case admitted to H1.*

An exhaustive investigation of all cases was conducted which led to identifying the index case.

This in-depth process involved a comprehensive collection of diverse data sources including

clinical records, travel history, antibody serology testing results, and contact tracing information.

This evidence revealed that the earliest patient was traced back to a mining cave and exhibited

symptoms highly consisting with the classic clinical presentation of MVD. Additionally, serology

testing results from three contacts linked to the index case exhibited antibodies of persons exposed

to Marburg virus. This suggests that the mining cave may have served as a potential habitat for the

MARV animal reservoir, particularly the fruit bat (*Rousettus aegyptiacus*). Following up on these

findings, the Rwanda Ministry of Health through a multisectoral collaboration launched an

investigation surveillance with the aim of screening fruit bats, mainly cave dwellers, for Marburg

virus (**Supplementary Figures 2 A, B and C**).

*Discussion*

The Rwanda MVD outbreak exhibited substantial human-to-human transmission, primarily

involving healthcare personnel, as is often observed with viral hemorrhagic fever (VHF) outbreaks

closely associated with nosocomial and occupational infections (18). However, Rwanda

effectively contained the spread from further reaching the broader community. Positive cases were

promptly isolated and treated with monoclonal antibodies and antivirals, achieving a favorable

case fatality rate of approximately 23%. This underscores the ongoing need to enhance healthcare

personnel's knowledge and attitudes regarding VHF case management.

Through evolutionary and phylogenetic analysis, we observed that the early outbreak sequences

represented very limited genomic variation, indicating that the outbreak originated from a single

zoonotic transmission event. The Rwandan lineage shared a common ancestor with sequences

originating from diversity sampled in bats in southwestern Uganda, thought the lineages had

diverged over decades in the animal reservoir. It is likely that enhanced zoonotic surveillance will

reveal many unsampled intermediates that can clarify the ecological pathways of transmission

between bat colonies in Uganda and Rwanda. Previous research indicates that bats can harbor

MARV for extended periods, with active mining areas in Rwanda providing ideal habitats for bats

and increasing the likelihood of zoonotic transmission (19).

We observed three synonymous single nucleotide polymorphisms (SNPs) outside the coding

regions of the GP and VP24 genes, respectively. Studies have shown that the Marburg virus VP24

protein interacts with the nucleoprotein (NP) and other cellular membranes, facilitating the release

of new virions from infected cells (20). Additionally, research highlights that MARV VP24

directly interacts with the human and bat Keap1 proteins, which modulate antioxidant responses

to support viral replication, a strategy likely critical for viral persistence and host adaptation (21).

Genomic sequencing of MARV in Rwanda has yielded critical insights into viral circulation

dynamics. However, as these SNPs fall outside of coding regions, it is unlikely that they

functionally contribute to enhanced viral fitness on any scale. Our results should be interpreted

within the context of our smaller sample size, which only covers the first two weeks of the

outbreak.

This research underscores the importance of continuous genomic surveillance which not only

facilitates outbreak tracking but also informs public health strategies for effectively controlling

this deadly pathogen.  In addition, previous research indicates that bats can harbor MARV for

extended periods, with active mining areas in Rwanda providing ideal habitats for bats and

increasing the likelihood of zoonotic transmission. Furthermore, Rwanda's location in a region of

consistent turbulent outbreaks, including Ebola, Rift Valley fever, COVID-19, dengue, and

Marburg, underscores the importance of regional and international collaboration to enhance

outbreak preparedness and responses for global health security.

**Reference**

1.  Mehedi M, Groseth A, Feldmann H, Ebihara H. Clinical Aspects of Marburg
Hemorrhagic Fever. Future Virol [Internet]. 2011 Sep [cited 2024 Oct 24];6(9):1091–106.
Available from: https://www.tandfonline.com/doi/full/10.2217/fvl.11.79

2.  Kilangisa LM, Max BL, Kayuni EA, Shao ER, Mashauri HL. Marburg virus disease:
lesson learned from the first outbreak encounter in Tanzania. International Journal of Surgery:
Global Health [Internet]. 2023 Jul [cited 2024 Oct 17];6(4). Available from:
https://journals.lww.com/10.1097/GH9.0000000000000186

3.  Paton, A. W., & Paton, J. C. (2018). Detection and characterization of Shiga toxigenic
*Escherichia coli* by PCR. In D. Liu (Ed.), Molecular Medical Microbiology (2nd ed., pp. 213–
239). Academic Press. https://doi.org/10.1016/B978-0-12-812026-2.00013-X

4.  Srivastava S, Sharma D, Kumar S, Sharma A, Rijal R, Asija A, et al. Emergence of Marburg
virus: a global perspective on fatal outbreaks and clinical challenges. Front Microbiol [Internet].
2023 Sep 13 [cited 2024 Oct 14];14:1239079. Available from:
https://www.frontiersin.org/articles/10.3389/fmicb.2023.1239079/full

5. Ministry of health R. Stop the Marburg virus [Internet]. 2024 Oct. Available from:
https://x.com/rwandahealth/status/1846968616278778288?s=12&t=HFAKR78RPRo-
Cmz7j7ciDQ

6.    Amman BR, Carroll SA, Reed ZD, Sealy TK, Balinandi S, Swanepoel R, et al. Seasonal

Pulses of Marburg Virus Circulation in Juvenile Rousettus aegyptiacus Bats Coincide with

Periods of Increased Risk of Human Infection. Kawaoka Y, editor. PLoS Pathog [Internet]. 2012

Oct 4 [cited 2024 Oct 18];8(10):e1002877. Available from:

https://dx.plos.org/10.1371/journal.ppat.1002877

7.    Amman BR, Schuh AJ, Albariño CG, Towner JS. Marburg Virus Persistence on Fruit as

a Plausible Route of Bat to Primate Filovirus Transmission. Viruses [Internet]. 2021 Nov 30

[cited 2024 Oct 18];13(12):2394. Available from: https://www.mdpi.com/1999-4915/13/12/2394

8.    Towner JS, Khristova ML, Sealy TK, Vincent MJ, Erickson BR, Bawiec DA, et al.

Marburgvirus Genomics and Association with a Large Hemorrhagic Fever Outbreak in Angola. J

Virol [Internet]. 2006 Jul [cited 2024 Oct 19];80(13):6497–516. Available from:

https://journals.asm.org/doi/10.1128/JVI.00069-06

9.    Wei H, Audet J, Wong G, He S, Huang X, Cutts T, et al. Deep-sequencing of Marburg

virus genome during sequential mouse passaging and cell-culture adaptation reveals extensive

changes over time. Sci Rep [Internet]. 2017 Jun 13 [cited 2024 Oct 19];7(1):3390. Available

from: https://www.nature.com/articles/s41598-017-03318-3

10.    Towner JS, Amman BR, Sealy TK, Carroll SAR, Comer JA, Kemp A, et al. Isolation of

Genetically Diverse Marburg Viruses from Egyptian Fruit Bats. Fouchier RAM, editor. PLoS

Pathog [Internet]. 2009 Jul 31 [cited 2024 Oct 25];5(7):e1000536. Available from:

https://dx.plos.org/10.1371/journal.ppat.1000536

11.     Nyakarahuka L, Ojwang J, Tumusiime A, Balinandi S, Whitmer S, Kyazze S, Kasozi S,

Wetaka M, Makumbi I, Dahlke M, Borchert J, Lutwama J, Ströher U, Rollin PE, Nichol ST,

Shoemaker TR. Isolated Case of Marburg Virus Disease, Kampala, Uganda, 2014. Emerg Infect

Dis. 2017 Jun;23(6):1001-1004. doi: 10.3201/eid2306.170047. PMID: 28518032; PMCID:

PMC5443453

12.     Carroll SA, Towner JS, Sealy TK, McMullan LK, Khristova ML, Burt FJ, et al.

Molecular Evolution of Viruses of the Family Filoviridae Based on 97 Whole-Genome

Sequences. J Virol [Internet]. 2013 Mar [cited 2024 Oct 25];87(5):2608–16. Available from:

https://journals.asm.org/doi/10.1128/JVI.03118-12

13.     Chen S. Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication

using fastp. iMeta [Internet]. 2023 May [cited 2024 Oct 24];2(2):e107. Available from:

https://onlinelibrary.wiley.com/doi/10.1002/imt2.107

14.     Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods

[Internet]. 2012 Apr [cited 2024 Oct 25];9(4):357–9. Available from:

https://www.nature.com/articles/nmeth.1923

15.    Li H. Minimap2: pairwise alignment for nucleotide sequences. Birol I, editor.

Bioinformatics [Internet]. 2018 Sep 15 [cited 2024 Oct 25];34(18):3094–100. Available from:

https://academic.oup.com/bioinformatics/article/34/18/3094/4994778

16.    Grubaugh ND, Gangavarapu K, Quick J, Matteson NL, De Jesus JG, Main BJ, et al. An

amplicon-based sequencing framework for accurately measuring intrahost virus diversity using

PrimalSeq and iVar. Genome Biol [Internet]. 2019 Dec [cited 2024 Oct 25];20(1):8. Available

from: https://genomebiology.biomedcentral.com/articles/10.1186/s13059-018-1618-7

17.    Wilm A, Aw PPK, Bertrand D, Yeo GHT, Ong SH, Wong CH, et al. LoFreq: a sequence-

quality aware, ultra-sensitive variant caller for uncovering cell-population heterogeneity from

high-throughput sequencing datasets. Nucleic Acids Research [Internet]. 2012 Dec 1 [cited 2024

Oct 25];40(22):11189–201. Available from:

https://academic.oup.com/nar/article/40/22/11189/1152727

18. Raab, M., Pfadenhauer, L.M., Millimouno, T.J. et al. Knowledge, attitudes and practices

towards viral haemorrhagic fevers amongst healthcare workers in urban and rural public healthcare

facilities in the N'zérékoré prefecture, Guinea: a cross-sectional study. BMC Public Health 20,

296 (2020). https://doi.org/10.1186/s12889-020-8433-2

19. Scarpa F, Bazzani L, Giovanetti M, Ciccozzi A, Benedetti F, Zella D, Sanna D, Casu M,

Borsetti A, Cella E, et al. Update on the Phylodynamic and Genetic Variability of Marburg Virus.

Viruses. 2023; 15(8):1721. https://doi.org/10.3390/v15081721

20.  Abir, M. H., Rahman, T., Das, A., Etu, S. N., Nafiz, I. H., Rakib, A., … Hassan, M. M. (2022). Pathogenicity and virulence of Marburg virus. Virulence, 13(1), 609–633. https://doi.org/10.1080/21505594.2022.2054760

21. Edwards MR, Johnson B, Mire CE, Xu W, Shabman RS, Speller LN, Leung DW, Geisbert TW, Amarasinghe GK, Basler CF. The Marburg virus VP24 protein interacts with Keap1 to activate the cytoprotective antioxidant response pathway. Cell Rep. 2014 Mar 27;6(6):1017-1025. doi: 10.1016/j.celrep.2014.01.043. Epub 2014 Mar 13. PMID: 24630991; PMCID: PMC3985291.

22. Kazutaka Katoh, Daron M. Standley, MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability, Molecular Biology and Evolution, Volume 30, Issue 4, April 2013, Pages 772–780, https://doi.org/10.1093/molbev/mst010

23. Bui Quang Minh, Heiko A Schmidt, Olga Chernomor, Dominik Schrempf, Michael D Woodhams, Arndt von Haeseler, Robert Lanfear, IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era, Molecular Biology and Evolution, Volume 37, Issue 5, May 2020, Pages 1530–1534, https://doi.org/10.1093/molbev/msaa015.

24. Kalyaanamoorthy, S., Minh, B., Wong, T. et al. ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods 14, 587–589 (2017). https://doi.org/10.1038/nmeth.4285.

25. Diep Thi Hoang, Olga Chernomor, Arndt von Haeseler, Bui Quang Minh, Le Sy Vinh,

UFBoot2: Improving the Ultrafast Bootstrap Approximation, Molecular Biology and Evolution,

Volume 35, Issue 2, February 2018, Pages 518–522, https://doi.org/10.1093/molbev/msx281.

26. Pavel Sagulenko, Vadim Puller, Richard A Neher, TreeTime: Maximum-likelihood

phylodynamic analysis, Virus Evolution, Volume 4, Issue 1, January 2018, vex042,

https://doi.org/10.1093/ve/vex042.

27. Carroll SA, Towner JS, Sealy TK, McMullan LK, Khristova ML, Burt FJ, Swanepoel R, Rollin

PE, Nichol ST. Molecular evolution of viruses of the family Filoviridae based on 97 whole-

genome sequences. J Virol. 2013;87:2608–2616. doi: 10.1128/JVI.03118-12.

28. Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C-H., Xie, D., Suchard, MA.,

Rambaut, A., & Drummond, A. J. (2014). BEAST 2: A Software Platform for Bayesian

Evolutionary Analysis. PLoS Computational Biology, 10(4), e1003537.

doi:10.1371/journal.pcbi.1003537.

**Acknowledgments**

We gratefully acknowledge the support of the Government of Rwanda and the Ministry of Health,

through the Rwanda Biomedical Center, for facilitating and supporting this sequencing work. We

extend our sincere appreciation to both the African Centre of Excellence for Genomics of

Infectious Diseases (ACEGID) and the Africa Centre for Disease Control and Prevention (Africa

CDC) for their invaluable contributions. This work was made possible through the collaborative

efforts of development partners and stakeholders, to whom we are deeply grateful.

**Funding:** Government of Rwanda and Development Partners.

**Author contributions**

Conceptualization: Y.B, L.M, S.N, C.H, C.M.M, Methodology: Y.B, L.M, A.A, E.U, R.M, E.P,

P.S, C.H, C.M.M Investigation: I.M, E.R, E.K, E.R, N.G Sampling: R.S, P.G, P.S, J.C.M PCR

Experiment: E.K, F.U, V.M, G.I, H.M Sequencing: Y.B, L.M, E.U, L.I, C.M, A.U, I.U, R.N, A.U,

J.K, J.P.M Bioinformatics Analysis: E.P, R.M, J.P.M, N.G, J.R, A.O, C.H Lab Enablement: S.T.K,

A.A, O.O, O.A, A.H, C.M.M Writing- Original Draft: Y.B, L.M, C.M, R.M, J.P.M, P.S Writing

Review and Editing: Y.B, L.M, E.E.S.F, E.R, C.M, M.S, M.N, T.T, D.T, E.S, M.G, J.P.R, J.N,

A.R, F.K, G.M, A.A, S.N, C.H, C.M.M.

**Diversity, equity, ethics, and inclusion [optional]:** NA

**Competing interests:** The authors declare no conflicts of interest.

**Data and code availability**

All the sequence data are available upon request on https://github.com/rbx-bfx/illumina_vsp.

*Materials and Methods*

**Ethics declaration**

The study was approved by the Rwanda National Ethics Committee (FWA Assurance No.

00001973 IRB 00001497 of IORG0001100-Protocol approval notice: N° 121/RNEC/2024).

**Patient sample collection**

We obtained whole blood samples from suspected patients presented with clinical symptoms (high

fever, severe headaches, muscle aches, fatigue, nausea, vomiting, and diarrhea) of MDV. Sample

testing was performed at Rwanda Biomedical Centre/National Reference Laboratory whereby

samples were kept in a cold chain prior to plasma separation and analysis.

**Nucleic acid extraction**

Viral RNA was extracted from 140µl plasma using the QIAamp Viral RNA Mini Kit (Qiagen,

Hilden, Germany), following the manufacturer's instructions adapted to the in-house standard

operating procedures with an elution volume of 60µl. The extracted RNA was quantified using a

Qubit fluorometer, and the samples were stored at -80°C until further analysis.

**RT-PCR and genomic sequencing**

For the detection and amplification of the target regions of the MVD, we employed the use of

RealStar® Filovirus Screen RT-PCR Kit 1.0 (Altona Diagnostics, Hamburg, Germany), which is

based on real-time PCR technology, for the qualitative detection and differentiation of Ebola and

Marburg virus specific RNA in human EDTA plasma. The RT-qPCR was conducted following

manufacturer's user guide and performed on CFX96 BIORAD machine. The amplification

followed an initial denaturation at 95°C for 5 minutes, followed by 40 cycles of denaturation at

95°C for 30 seconds, annealing at 55°C for 30 seconds, and extension at 72°C for 1 minute,

concluding with a final extension at 72°C for 5 minutes. The amplified products were analyzed via

2% agarose gel electrophoresis to confirm successful amplification and appropriate product size.

Following amplification, library construction was performed on 30 samples using the Illumina

RNA Prep with Enrichment (L) Tagmentation (IRPE) workflow and the lllumina Viral

Surveillance Panel v2 kit (VSP v2 kit). Total RNA extracts were converted to cDNA, tagmented

and thereafter amplified. Genomic region of interest was captured using a hybrid capture method.

Probes were isolated using magnetic pulldown which are selectively enriched for the desired

regions. Enriched libraries were quantified using dsDNA HS assay and Qubit fluorometer.

Libraries were thereafter denatured and normalized at a final loading concentration of 0.8pM.

Paired-end sequencing was performed using a NextSeq 550 with a 300-cycle mid-output cartridge,

with the sequencing depth aimed at a minimum coverage of 100x to ensure robust variant

detection.

**Bioinformatics Analysis**

**Genome Assembly**

We employed a reference-based genome assembly pipeline to analyze the sequencing data

generated using the Illumina Viral Surveillance Panel (VSP) v2 Kit (described above). This

workflow integrates several steps including; quality control, host genome filtering, and viral

genome assembly to accurately reconstruct viral sequences from targeted sequencing.

To ensure that only high-quality data were processed downstream, raw sequencing reads were first

processed with *fastp v. 0.23.24 (13)* for trimming adapters and filtering out low-quality bases.

Following this, human host genome sequences were filtered out by aligning the cleaned reads to

the human genome (hg38) using *Bowtie2 v.2.5.4* (14). Unmapped reads, presumed to be of viral

origin, were retained for downstream analysis. These de-hosted reads were then aligned to a

Marburg virus reference genome (NC_001608.3) using *Minimap2 v.2.28 (15)*,

The sequences had 70-97% genomic coverage. We manually masked two mutations in respective

consensus sequences that were adjacent to potential misalignment.

**Variant Calling and Consensus Generation**

Variant calling was performed using both *ivar 1.4.3 (16)* and *LoFreq v.2.1.5 (17)*. A depth

coverage threshold of 50 reads per nucleotide position was applied to ensure robust and reliable

variant detection. This threshold was also applied to the generation of consensus sequences which

was also done using *ivar 1.4.3.*,

**Phylogenetic**

We combined our eighteen higher quality sequences (coverage >= 70%) with all publicly available

MARV sequences available on Genbank (N=81). To root the tree, we included 8 Ravn virus

sequences included as an outgroup, which were pruned from the tree and excluded in all

subsequent analyses. We aligned the sequences using *Mafft v 7.52 (22)* and reconstructed a

phylogenetic tree using *IQTree 2.2.5 (23)* under modelfinder plus *(24)*, with ultrafast bootstrapping

(25). We performed the initial ancestral state reconstruction using *Treetime v0.9.3 (26)*.

**Bayesian phylogenetic reconstruction**

The current genomic data does not provide sufficient temporal signal to estimate the rate of

evolution in the outbreak clade. However, we were not interested in estimating the time to the most

recent common ancestor (tMRCA) of the outbreak clade, as determining the emergence timing of

the outbreak clade will more likely depend on the epidemiological and contact history of the index

case. We wanted to estimate tMRCA of the outbreak clade and its closest human and zoonotic

outgroups to better understand its potential zoonotic emergence pathway. We therefore

reconstructed the time-resolved phylogeny under a fixed local clock. We assumed a fixed value of

1e-3 substitutions per site per year as a plausible rate of evolution for a RNA virus in the early

stage of an outbreak for the defined outbreak clade. We allowed the remainder of the tree to evolve

under a lognormal prior centered on 5e-4. (27) We used a two-phase coalescent model: the tree

from the MRCA (Rwandan lineage) onward was modeled with an exponential growth model, with

the earlier phase modeled as a constant-population size coalescent model. We ran two independent

chains of 100 million states to ensure convergence, discarding the initial 10% of each chain as

burn-in. The chains were then combined with LogCombiner. For all subsequent analyses, we

assessed convergence using Tracer, and constructed a maximum clade credibility (MCC) tree in

TreeAnnotator 1.10 (28).

494
495

508    **Supplemental Figure 1:** *A) Time resolved maximum clade credibility tree of global MARV*

509    *dataset, with closest human (KP985768) and zoonotic (JX458855) sequence to the Rwandan*

510    *outbreak lineage annotated. B) tMRCA of the Rwandan lineage and KP985768 as well as*

511    *JX458855 respectively. C) tMRCA of root, which is truncated in A for clarity. D) Estimate clock*

512    *rate in substitutions per site per year for full tree, excluding the outbreak clade.*

517 **Supplemental Figure 2:** *A) Mining cave entrance with human activity; B) Trapping fruit bats*

518 *with net; C) Sampling bats for Marburg virus.*

# Figures

**Figure 1:** *A) Maximum likelihood phylogeny of the global MARV dataset. Tips are annotated by country of isolation. This study's tips are relatively enlarged. B) Map of Uganda and Rwanda. The city of patients in this study (Kigali) is annotated in red. Python Cave and Kitaka Mine in Ibanda district in southwest Uganda are annotated in light blue and yellow, respectively.*

**Figure 2:** *A) Maximum likelihood phylogeny of the outbreak clade, with the Kampala outgroup. Sequences are colored with associated hospital. SNPs reconstructed relative to the common ancestor are annotated in text, as are the clusters defined by the SNP presence. The coordinates are relative to NC_001608 B) Epidemiological timeline of the sampled sequences in Figure 2A, as annotated on the y-axis. The first black marker indicates the date of symptom onset, with the second indicating the date of sampling. Sequences are colored by their associated hospital (H1, H2), and partitioned by SNP presence into clusters, as annotated in text*