Sociodemographically Differential Patterns of Chronic Pain Progression Revealed by Analyzing the All of Us Research Program Data

Edwin Baldwin, BS[1], Jin Zhou, PhD[2], Wenting Luo, MS[3], W. Michael Hooten, MD[4,*], Jungwei W. Fan, PhD[5,*], Haiquan Li, PhD[1,*]

[*]Corresponding author

Email: haiquan@arizona.edu (HL), Fan.Jung-wei@mayo.edu (JWF), Hooten.William@mayo.edu (WMH)

[1]Department of Biosystems Engineering, University of Arizona, Tucson AZ USA 85721

[2]Department of Biostatistics, University of California at Los Angeles, Los Angeles, CA USA 90095

[3]Statistics and Data Science Graduate Interdisciplinary Program, University of Arizona, Tucson AZ USA 85721

[4]Division of Pain Medicine, Department of Anesthesiology. Mayo Clinic, Rochester, MN USA 55905

[5]Department of Artificial Intelligence & Informatics. Mayo Clinic, Rochester, MN 55905

Abstract

The differential progression of ten chronic overlapping pain conditions (COPC) and four comorbid mental

disorders across demographic groups have rarely been reported in the literature. To fill in this gap, we

conducted retrospective cohort analyses using All of Us Research Program data from 1970 to 2023.

Separate cohorts were created to assess the differential patterns across sex, race, and ethnicity. Logistic

regression models, controlling for demographic variables and household income level, were employed to

identify significant sociodemographic factors associated with the differential progression from one COPC

or mental condition to another. Among the 139 frequent disease pairs, we identified group-specific

patterns in 15 progression pathways. Black or African Americans with a COPC condition had a

significantly increased association in progression to other COPCs (CLBP->IBS, CLBP->MHA, or IBS->MHA,

OR$\geq$1.25, adj.p$\leq$4.0x10^{-3}) or mental disorders (CLBP->anxiety, CLBP->depression, MHA->anxiety, MHA-

>depression, OR$\geq$1.25, adj.p$\leq$1.9x10^{-2}) after developing a COPC. Females had an increased likelihood of

chronic low back pain after anxiety and depression (OR$\geq$1.12, adj.p$\leq$1.5x10^{-2}). Additionally, the lowest

income bracket was associated with an increased risk of developing another COPC from a COPC (CLBP-

>MHA, IBS->MHA, MHA->CLBP, or MHA->IBS, OR$\geq$1.44, adj.p$\leq$2.6x10^{-2}) or from a mental disorder

(depression->MHA, depression->CLBP, anxiety->CLBP, or anxiety->IBS, OR$\geq$1.50, adj.p$\leq$2.0x10^{-2}), as well

as developing a mental disorder after a COPC (CLBP->depression, CBLP->anxiety, MHA->anxiety,

OR$=$1.37, adj.p$\leq$1.6x10^{-2}). To our knowledge, this is the first study that unveils the sociodemographic

influence on COPC progression. These findings suggest the importance of considering sociodemographic

factors to achieve optimal prognostication and preemptive management of COPCs.

Author Summary

Understanding the differential progression of chronic overlapping pain conditions (COPC) can inform

patients and clinicians in optimizing the disease management. Although differential manifestations have

been extensively reported for individual COPCs, differential patterns along the progression from one

COPC to another have been rarely studied. The study addresses this gap specifically by identifying

differential COPC progression associated with sex, race, and ethnicity, providing valuable insights for

patients and clinicians to plan for their prevention or treatment strategies in the context of

sociodemographic relative risk.

Introduction

Chronic overlapping pain conditions (COPCs), defined as a subset of comorbid chronic pains that co-occur at higher-than-expected levels in the general population, are driven by shared mechanisms [1] and tend to be more common in females [1]. The National Institutes of Health Pain Consortium currently recognizes ten COPCs: chronic low back pains (CLBP), migraine (MHA), irritable bowel syndrome (IBS), temporomandibular disorders (TMD), fibromyalgia (FM), chronic fatigue syndrome (CFS), chronic tension-type headache (CTTH), endometriosis (ENDO), vulvodynia (VVD), and urologic chronic pelvic pain syndrome (UCPPS). COPCs significantly affect individuals worldwide, interfering with daily activities, mental health, and overall quality of life. Given their frequent comorbidities with mental health conditions [2], our study includes the four most frequently studied conditions—anxiety, depression, bipolar disorder, and schizophrenia [2-4].

The experience of pain is not uniform but intricately shaped by the interplay of sociodemographic factors, including sex, gender, age, race, ethnicity, and financial status (e.g., income) [5, 6], resulting in disparities known to particularly affect subpopulation [7]. Previous research reported a strong association between demographic background [5] and pain-related health outcomes [8], which implied unique pain mechanisms across racial/ethnic groups [9-11] and sexes [12-14]. Besides recognized biological differences, social determinants are shown to affect the experience of chronic pains [15]. Evidence indicates that people with financial challenges encounter greater healthcare barriers in accessing pain management [16]. Further, in urban populations, the intersection of sociodemographic factors has been shown to create unique identities that impact pain experiences [5, 17].

Although extensive research has explored disparities in individual COPC conditions, few studies have investigated sociodemographic disparities in the incremental progression among COPCs and comorbid mental disorders. To address this gap, we examined the sociodemographic disparities in COPCs using a

large-scale longitudinal cohort from the All of Us Research Program (AoURP) [18], which has enrolled

participants from a broad variety of populations.

The primary objective of the research is to discover statistically significant sociodemographic factors

associated with one of the ten COPC conditions and four mental health conditions. We sought to reveal

the differential progression patterns among these distinct COPC and mental conditions that were known

for their frequent comorbidities with COPCs [4, 19]. Validation of such differential disease manifestations

will allow for more precise prevention strategies that target the relative risk of further COPC

complications in specific subpopulations. The study represents the first of its kind by applying rigorous

statistics to identify health disparities and financial predictors in COPCs, specifically based on progression

patterns. Our results contribute to understanding the disparate, nuanced clinical experiences in COPCs,

advocating for both accessible and individualized chronic pain care that would fit each patient's unique

needs [16, 20].

Results

Of the 309,157 participants with valid electronic health records, 88,496 (28.6%) had at least one COPC

condition. Table 1 shows the demographic characteristics of the COPC cases and their comorbid mental

disorders in the AoURP version 8 dataset.

Table 1: Demographic Characteristics of the Study COPC Cases.

	All of Us Research Program (n=88,496)
Sex	
Female	62,265 (70.4%)
Male	26,231 (29.6%)

Race

American Indian or Alaska Native	934 (1.1%)
Asian	1,308 (1.5%)
Black or African American	14,641 (16.5%)
Middle Eastern or North African	405 (0.5%)
More than One Population	4,296 (4.9%)
Native Hawaiian or Other Pacific Islander	63 (0.07%)
None of these	996 (1.1%)
Unknown	12,547 (14.2%)
White	53,306 (60.2%)

Ethnicity

Hispanic	13,323 (15.1%)
Non-Hispanic	72,679 (82.1%)
Others	2,494 (2.8%)

COPC Onset

Chronic fatigue syndrome (CFS)	3,563 (4.0%)
Chronic lower back pain (CLBP)	51,480 (58.2%)
Chronic tension-type headache (CTTH)	764 (0.9%)
Endometriosis (ENDO)	1,770 (2.0%)
Fibromyalgia (FM)	6,469 (7.3%)
Irritable bowel syndrome (IBS)	8,834 (10.0%)
Migraine (MHA)	23,633 (27.7%)
Temporomandibular disorder (TMD)	2,381 (2.7%)
Urologic chronic pelvic pain syndrome (UCPPS)	947 (1.1%)

Vulvodynia (VVD)	162 (0.2%)
Mental Health Conditions	
Anxiety	30,344 (34.3%)
Bipolar disorder	4,221 (4.8%)
Depressive disorder	29,860 (33.7%)
Schizophrenia	770 (0.9%)

Note: Percentages in parentheses represent the proportion among the COPC cases only, not

the prevalence of the general population. Unknown Race included: I prefer not to answer,

none indicated, and skip. Others Ethnicity: none of these, prefer not to answer, and skip.

Intersex and transgender persons were excluded due to insufficient statistical power.

Table 2 demonstrates the elevated risk of specific COPCs in females and racial minorities, based on

multiple logistic regression studies of these variables using demographically matched cohorts.

Significance is defined as a p-value below the 0.05 threshold after the Bonferroni correction, adjusted for

other demographic factors and income levels. Except for UCPPS, all COPCs are more likely to develop in

females. For instance, fibromyalgia had an odds ratio of 9.99 in females (95% CI [8.24, 12.10], adj.p<10^{-16}

), indicating that females are about ten times more likely to be diagnosed with fibromyalgia than males.

Similarly, for migraine headaches (MHA), females exhibit an odds ratio of 3.31 (95% CI [3.09, 3.55],

adj.p<10^{-16}), suggesting females are more than tripled as likely to have MHA compared to males. Along

the race variable, Black or African American persons have an odds ratio of 1.24 (95% CI [1.16, 1.34],

adj.p<10^{-16}) in developing chronic lower back pain (CLBP) even after adjusting for income levels and

other demographic factors (e.g., sex and age).

The onset of mental disorders also showed significant demographic associations. Females present higher

odds ratios of developing anxiety and depression at 1.50 (95% CI [1.44, 1.56]) and 1.48 (95% CI [1.42,

1.54]) with adjusted p-values less than 10^{-16}. For schizophrenia, a significant odds ratio of 2.61 (95%CI

[1.93, 3.52], adj.p<10^{-16}) was observed in Black or African American persons, which corroborates the

literature [21, 22].

Table 2: Enrichment for individual pain conditions observed in females and racial minorities.

Disease	Cases	Sex		Race			
		Female	Asian	Black or African American	Middle Eastern or North African	More than One Population	American Indian or Alaska Native
CFS	3,563	1.86***	ns	0.39***	ns	ns	ns
CLBP	51,480	1.16***	0.82*	1.24***	ns	1.36***	ns
CTTH	764	2.10***	ns	ns	ns	ns	ns
FM	6,469	9.99***	0.35***	0.59***	ns	ns	ns
IBS	8,834	2.84***	0.36***	0.41***	ns	ns	ns
MHA	23,633	3.31***	0.38***	0.61***	ns	ns	ns
TMD	2,381	3.18***	ns	0.61**	ns	ns	ns
UCPPS	947	ns	ns	ns	ns	ns	ns
Anxiety	62,153	1.50***	0.37***	0.54***	0.64**	ns	0.75*
Bipolar	8,744	ns	0.31***	0.83**	0.36*	1.31*	ns
Depression	60,001	1.48***	0.41***	0.68***	0.70*	ns	ns
Schizophrenia	2,247	0.41***	ns	2.61***	ns	1.89*	ns

109 *Note*: OR: odds ratio. ns: not significant after Bonferroni correction, adj.p>0.05, *: 10^{-3}≤adj.p<0.05, **:

110 10^{-6}≤adj.p<10^{-3}, ***: adj.p<10^{-6}. The reference group for sex is male, and the reference group for race is

111 White. Since no significant racial or ethnic disparity factors have been identified for endometriosis

112 and vulvodynia—female-specific disorders—they are not shown in the table. Of note, AoU includes a

113 racial minority of Middle Eastern or North African for descendants of the region.

Fig 1. The overrepresentation of specific pairwise relations between COPC and mental conditions,

comparing females, racial minority persons, and persons with low income to the reference

sociodemographic groups. Each node represents either a specific COPC condition or a mental disorder.

For sociodemographic reference in the statistical analysis, male is the reference sex, White persons make

the reference race, non-Hispanic White is the reference ethnicity, and the highest income stratum is the

reference income level. Each arrowed edge depicts a significant temporally ordered disease pair

supported by either the reference demographics (dashed edges) or the alternative demographics (solid

edges). Edge colors indicate significant (Bonferroni-correlated p-value<0.05) association with specific

demographics (gray for White, blue for Black or African American, and purple for females), while arrow

colors indicate significant overrepresentation of the lowest household income levels in contrast to the

highest level (gray for none and blue for significant).

Fig 1 illustrates how sex and race are associated with the development of a second COPC or mental

condition following a previous condition. This network was derived from 139 frequent co-occurring

(threshold: at least 21 occurrences) pairs of conditions, where each pair of nodes must involve at least

one COPC, i.e., two COPCs or one COPC and one mental condition (Supplementary file). Each arrow

between two nodes indicates the two conditions are associated sequentially accordingly using regression

analyses using non-Hispanic White males as the reference group. Black or African American persons

show the strongest relative risk for COPC and mental disorder progression compared to White persons

(blue lines).

First, Black or African American persons are more likely to develop another COPC after the first one:

those with CLBP are more likely to develop subsequent IBS (OR=1.49, 95% CI [1.07, 2.08], adj.p=1.2x10^{-3})

and MHA (OR=1.25, 95% CI [1.03, 1.53], adj.p=4.0x10^{-3}), and those with IBS are more likely to develop

MHA (OR=1.69, 95% CI [1.02, 2.80], adj.p=1.4x10^{-2}). Second, they are more likely to develop FM after

depression (OR=1.67, 95% CI [1.20, 2.32], adj.p=2.1x10^{-6}). Third, they are more likely to develop a

mental disorder after the onset of a COPC: anxiety (OR=1.41, 95% CI [1.24, 1.61], adj.p<10^{-16}) and

depression (OR=1.32, 95% CI [1.15, 1.51], adj.p=1.2x10^{-11}) following CLBP, and anxiety (OR=1.37, 95% CI

[1.11, 1.68], adj.p=5.5X10^{-6}) and depression (OR=1.25, 95% CI [1.00, 1.55], p=1.9x10^{-2}) following

migraine. Of note, half of the results heightened complication risk of both other COPCs (e.g., MHA and

FM) and mental disorders (e.g., anxiety and depression) added to the known enrichment of CLBP in

Black or African American persons (OR=1.24, 95% CI [1.16, 1.34], adj.p<10^{-16}, Table 2), among others.

In addition, we observed a sex disparity in the progression of COPC comorbidities: females were more

likely to develop CLBP after depression (OR=1.15, 95% CI [1.04, 1.27], adj.p= 6.6x10^{-5}) or anxiety

(OR=1.12, 95% CI [1.00, 1.24], adj.p=1.5x10^{-2}), and more likely to develop schizophrenia after CLBP

(OR=1.59, 95% CI [1.01, 2.52], adj.p=2.1x10^{-2}). Note that we also accounted for income level while

examining the disparities, and the findings aligned with the literature. Specifically, we found that lower

income is associated with an elevated risk of developing another subsequent COPC or mental disorder,

compared with the highest income group ($200,000 annually). For example, persons with an annual

income below $25,000 are more likely to develop CLBP after depression (OR=1.58, 95% CI [1.17, 2.14],

adj.p=4.5x10^{-6}), anxiety (OR 1.50, 95% CI [1.12, 2.00], adj.p=4.6x10^{-5}), or MHA (OR=1.44, 95% [1.02,

2.05], adj.p=1.4x10^{-2}). Persons with the lowest household income are also more likely to develop MHA

after CLBP (OR=2.05, 95% CI [1.40, 3.01], adj.p=1.7x10^{-9}), IBS (OR=1.99, 95% CI [1.00, 3.95],

adj.p=2.6x10^{-2}), and depression (OR=2.15, 95% CI [1.42, 3.23], adj.p=2.4x10^{-9}). In addition, they were

more likely to develop IBS after MHA (OR=2.25, 95% CI [1.22, 4.16], adj.p=1.6x10^{-4}) and anxiety

(OR=1.84, 95% CI [1.01, 3.34], adj.p=2.0x10^{-2}). Finally, this group is more likely to develop anxiety

(OR=1.31, 95% CI [1.02, 1.68], adj.p=8.5x10^{-3}) and depression (OR=1.37, 95% CI [1.02, 1.83],

adj.p=9.6x10^{-3}) after CLBP, and develop anxiety after MHA (OR=1.38, 95% CI [1.01, 1.88], adj.p=1.6x10^{-2})

compared to the reference group.

Materials and Methods

Dataset and Phenotype Definition

We used the AoURP[18] version 8, released in February 2025, which involved the electronic health

records (EHR) of over 633,000 participants. Only individuals sharing EHR data were included, and those

whose data showed apparent quality or power issues were excluded, such as problematic dates, ages

(over 110 years old in any record), less than two clinical visits, and disagreement between reported sex

and gender. Variables studied include onset of each COPC and mental condition, sex at birth, age, race,

ethnicity, household income level, and region of state residence (e.g., northeast). Sex at birth, race, and

ethnicity were self-reported during the enrollment of AoU.

The determination of COPC/mental cases worked as follows:

(1) Identifying episodes of COPC and mental conditions: We cataloged COPC and mental health

conditions using the ICD-10 [23] and ICD-9 [24] codes recognized by the research community. Utilizing

terms from the literature and descriptions, we linked the conditions to the corresponding AoURP

standard concept ids, which served as the primary inclusion criteria.

(2) Establishing chronicity: A condition was deemed chronic if it was diagnosed with at least two

separate episodes, a minimum of 90 days apart, aligning with standards accepted by the pain research

community [24].

(3) Determining the earliest onset: To establish each patient's order of distinct conditions, we identified

the earliest onset for each condition based on diagnosis dates. Additionally, we required that the onset

of each condition of a patient occur at least one year (clearance period) after his/her first EHR record, to

minimize the noise from using incorrect onsets. We excluded the persons who had more than one COPC

or mental condition within the clearance period, for which the order of the condition was difficult to

establish.

Cohort Creation Process

We built a separate cohort for each demographic factor to assess its influence on developing a

COPC/mental disease or progressing to another COPC. The process of cohort selection for each analysis

proceeded as follows:

(1) Building a cohort for each demographic variable that developed a COPC/mental condition: Each

group (e.g., African American) of a demographic variable (e.g., race) was compared to a reference

group (e.g., White). We constructed a cohort of participants based on the assessed or the reference

group. The same number of participants were selected for each of the two groups, further matched

by other variables (not being assessed), such as sex at birth, birth year (categorized in decade

intervals), and race. We conducted random sampling without replacement from the group that had

more samples (usually the reference group, e.g., White), ensuring an equivalent number of

participants from the larger group for each demographic combination. Infrequently, when there

were not enough samples to achieve a complete demographic match for a particular demographic

feature combination, we opted for partial matching. The precedence for matching was sex, followed

by birth year group, and then race. We did not include gender, which is a broader social construct

concept, due to its high correlation with biological sex and insufficient data to reach the required

power.

(2) Building a cohort for each disease progression with at least 21 occurrences (per AoU policy): Starting

from any COPC/mental condition, we divided the participants into two sub-cohorts, those with and

without the start condition. We then matched the patients with the starting COPC/mental condition

against those without the condition, by sex, age interval, and race, similar to the procedure above,

but matching each case up to a sufficient number of controls to get the maximum power. The

case:control ratios ranging from 1:1 to 1:10 were explored. Then, we studied the difference in the

onset of other conditions after the start of the COPC/mental condition between the two sub-cohorts

of participants. Lastly, we assessed the differential influence of a sociodemographic group as

compared to their reference group, with regard to the disease progression patterns within this

specific cohort.

Statistical Analysis

After building the cohort for each comparison, we assessed the increased risk of developing the first or

progressing to another condition. Each COPC/mental condition other than the start condition (for the

progression study only) served as an outcome variable. We employed a logistic regression model to

assess the statistical significance of each sociodemographic group, specifically the odds ratio of

developing or progressing to the outcome condition, relative to the reference group. In the COPC onset

analysis, odds ratios were derived from the coefficient of a sociodemographic variable for risk of

outcome, while in the progression analysis, the odds ratios were from the interaction term between the

first condition and the sociodemographic variable. An odds ratio exceeding 1 with statistical significance

would suggest disparity associated with the sociodemographic factor.

The demographic variables assessed include sex, race, and ethnicity, for the known relevance in chronic

pain [7, 12-14, 25-27]. Additional covariates were incorporated including birth year, household income

level, and region of state of residence. Birth year was treated as a numerical variable, whereas sex, race,

ethnicity, and income level were handled as categorical variables. Income levels—stratified as below

$25k, $25-$50k, $50-$100k, $100-$150k, $150-$200k, and over $200k—served as a proxy for

socioeconomic status. Missing income data (Skip or Prefer not to answer in questionnaire response) was

treated as a separate category.

To address potential multiple comparison errors, we applied the Bonferroni correction for adjusting

confidence intervals (CI) for odds ratios and p-values (adj.p).

Ethics Statement

The study utilized de-identified data from the All of Us Research Program. The University of Arizona

Institutional Review Board (IRB) reviewed the study titled "Identify the heterogeneity and commonality

of chronic overlapping pain conditions (COPCs) through phenotypic and genomic perspectives" (IRB

Submission ID: STUDY00001119) and determined that it does not qualify as human research, as defined

by DHHS and FDA regulations. Consequently, IRB review and approval by this organization is not

required.

Discussion

Our results align with known disparities in developing COPC and their comorbidities, e.g., the elevated

risk for females to develop all COPCs [28, 29], except UCPPS, bipolar disorder, and schizophrenia [30].

Possible explanations may involve biological differences between sexes [13], such as hormone factors

[28], sensitization [17], and inflammatory responses [25]. Furthermore, this study provides a distinct

contribution by clarifying disparities associated with specific progression patterns in COPCs and mental

conditions [31]. For instance, our findings indicate that females are at a significantly heightened risk for

developing chronic low back pain following an anxiety or depression diagnosis. Although the impact of

psychological conditions on chronic low back pain has been widely studied [32-36], the influence and

magnitude of sex differences in the progression from a mental disorder to CLBP remain inconclusive.

These findings offer new insights into this issue.

The results indicated an increased risk of developing chronic low back pain (CLBP) in Black or African

Americans [17], who may also face a higher risk for subsequent COPCs (e.g., irritable syndrome) and

mental conditions (e.g., anxiety and depression). Although CLBP is more prevalent among African

Americans, its differential influence on other COPC pains or mental conditions has been rarely reported

in Black Americans, particularly by longitudinal and prospective studies [37]. A few studies suggested

Black population with chronic pains reported more depressive symptoms [38]. This may be related to the

racial differences in pain sensitivity and perception [26, 39-41], psychological response to chronic pain

[42], stress levels [38], and allostatic load [43, 44]. Of note, socioeconomic confounders cannot be ruled

out in the context of healthcare access and utilization [31, 45, 46], which would influence the

management and therefore containment of disease progression.

While sociodemographic disparities in the development of individual COPC conditions have been

frequently reported, studies investigating such disparities in their progression are rare. Not only are

multiple COPCs associated with more complex outcomes [47], but knowledge gained about the

differential progression should be translated into optimization of chronic pain management, cross-

specialty coordination, and creation of multi-faceted support systems customized for every

sociodemographic group, including their intersections.

Not all observed progressions were statistically significant across demographic variables (e.g., sex) for

certain pairs of COPCs. This outcome was likely due to the underpowered nature of certain cohort

groups, limiting the ability to detect subtle differences in COPC progression. The limitation in statistical

power may stem from the sample sizes for progression between some COPC pairs [48] (e.g., between

CTTH and TMD) or in certain strata, particularly when the data was segmented by multiple dimensions

such as age, sex, race, and income level. The observation may also be attributed to the modeling of

socioeconomic effects in the risk of progression since many associations between the COPC to mental

conditions could be confounded by socioeconomic factors, for instance, when a lack of sufficient medical

care exacerbates a condition [49]. Finally, sex may contribute more to the development of the first

condition, while not contributing significantly to the progression [50-52], which warrants further

verification [53]. In addition, we did not identify any disparity for the Hispanic populations, neither in

baseline nor in progression. Further research is needed to reproduce and explain the findings.

The study and results rely heavily on electronic health records, which are known to have potential biases

and noises. The All of Us Research Program is known to be enriched with underrepresented populations

and diverse health conditions, which could potentially affect the effect size of the results.

Future research should be able to build on the discovered correlations, exploring the causes and effects

of these correlations. Our progression analysis represents a promising approach to health disparity

research, especially as more data becomes available. For example, there is limited data for studying

chronic pain disparities experienced by American Indians [54] or intersex persons [13]. Such research is

crucial for developing effective pain management strategies to address the needs of specific populations

[55].

In conclusion, the study demonstrated a promising approach to systematic discovery of the differential

chronic pain progression patterns across sociodemographic groups. Our results not only reinforced

existing evidence, but also uncovered lesser-known associations, such as complications of increased

chronic low back pain in Black or African American populations. By recognizing the confluence of

sociodemographic factors, we anticipate such data-driven research will enable more contextualized

strategies in caring for our diverse populations affected by chronic pains and encourage future

prospective studies to build stronger evidence.

Acknowledgements

This work was funded by NIDCR/NIH R21DE031424. We gratefully acknowledge All of Us participants for

their contributions, without whom this research would not have been possible. We also thank the

National Institutes of Health's All of Us Research Program for making available the data and analytics

platform.

References

1.   Johnston, K.J., R. Signer, and L.M. Huckins, *Chronic overlapping pain conditions and*
nociplastic pain. Human Genetics and Genomics Advances, 2025. **6**(1).
2.   Vadivelu, N., et al., *Pain and psychology—a reciprocal relationship.* Ochsner Journal, 2017.
17(2): p. 173-180.
3.   Goesling, J., L.A. Lin, and D.J. Clauw, *Psychiatry and pain management: at the intersection*
of chronic pain and mental health. Current psychiatry reports, 2018. **20**: p. 1-8.
4.   Hooten, W.M. *Chronic pain and mental health disorders: shared neural mechanisms,*
epidemiology, and treatment. in *Mayo Clinic Proceedings.* 2016. Elsevier.
5.   Quiton, R.L., et al., *Sociodemographic patterns of pain in an urban community sample: an*
examination of intersectional effects of sex, race, age, and poverty status. Pain, 2020.
161(5): p. 1044-1051.

6.   Michael, G.E., K.A. Sporer, and G.M. Youngblood, *Women are less likely than men to receive*
312   *prehospital analgesia for isolated extremity injuries.* The American journal of emergency
313   medicine, 2007. **25**(8): p. 901-906.
314   7.   Green, C.R., et al., *The unequal burden of pain: confronting racial and ethnic disparities in*
315   *pain.* Pain medicine, 2003. **4**(3): p. 277-294.
316   8.   Hollingshead, N.A., et al., *A comparison of race-related pain stereotypes held by White and*
317   *Black individuals.* Journal of applied social psychology, 2016. **46**(12): p. 718-723.
318   9.   Todd, K.H., et al., *Ethnicity and analgesic practice.* Annals of emergency medicine, 2000.
319   **35**(1): p. 11-16.
320   10.   Edwards, R.R., et al., *Ethnic differences in pain tolerance: clinical implications in a chronic*
321   *pain population.* Psychosomatic medicine, 2001. **63**(2): p. 316-323.
322   11.   Aroke, E.N., et al., *Could epigenetics help explain racial disparities in chronic pain?* Journal
323   of pain research, 2019: p. 701-710.
324   12.   Sorge, R.E. and S.K. Totsch, *Sex differences in pain.* Journal of neuroscience research, 2017.
325   **95**(6): p. 1271-1281.
326   13.   Osborne, N.R. and K.D. Davis, *Sex and gender differences in pain*, in *International Review of*
327   *Neurobiology.* 2022, Elsevier. p. 277-307.
328   14.   Bartley, E.J. and R.B. Fillingim, *Sex differences in pain: a brief review of clinical and*
329   *experimental findings.* British journal of anaesthesia, 2013. **111**(1): p. 52-58.
330   15.   Karran, E.L., A.R. Grant, and G.L. Moseley, *Low back pain and the social determinants of*
331   *health: a systematic review and narrative synthesis.* Pain, 2020. **161**(11): p. 2476-2493.
332   16.   Thurston, K.L., et al., *A systematic review of race, sex, and socioeconomic status*
333   *differences in postoperative pain and pain management.* Journal of PeriAnesthesia Nursing,
334   2023. **38**(3): p. 504-515.
335   17.   Meints, S.M., V. Wang, and R.R. Edwards, *Sex and race differences in pain sensitization*
336   *among patients with chronic low back pain.* The Journal of Pain, 2018. **19**(12): p. 1461-1470.
337   18.   Sankar, P.L. and L.S. Parker, *The Precision Medicine Initiative's All of Us Research Program:*
338   *an agenda for research on its ethical, legal, and social issues.* Genetics in Medicine, 2017.
339   **19**(7): p. 743-750.
340   19.   Fillingim, R.B., et al., *Associations of psychologic factors with multiple chronic overlapping*
341   *pain conditions.* Journal of oral & facial pain and headache, 2020. **34**(Suppl): p. s85.
342   20.   Charleston IV, L., *Headache disparities in African-Americans in the United States: a*
343   *narrative review.* Journal of the National Medical Association, 2021. **113**(2): p. 223-229.
344   21.   Bresnahan, M., et al., *Race and risk of schizophrenia in a US birth cohort: another example*
345   *of health disparity?* International journal of epidemiology, 2007. **36**(4): p. 751-758.
346   22.   Olbert, C.M., A. Nagendra, and B. Buck, *Meta-analysis of Black vs. White racial disparity in*
347   *schizophrenia diagnosis in the United States: Do structured assessments attenuate racial*
348   *disparities?* Journal of abnormal psychology, 2018. **127**(1): p. 104.
349   23.   Schrepf, A., et al., *ICD-10 codes for the study of chronic overlapping pain conditions in*
350   *administrative databases.* The journal of pain, 2020. **21**(1-2): p. 59-70.
351   24.   Irwin, M.N. and M.A. Smith, *Validation of ICD-9 codes for identification of chronic*
352   *overlapping pain conditions.* Journal of Pain & Palliative Care Pharmacotherapy, 2022. **36**(3):
353   p. 166-177.
354   25.   Rosen, S., B. Ham, and J.S. Mogil, *Sex differences in neuroimmunity and pain.* Journal of
355   neuroscience research, 2017. **95**(1-2): p. 500-508.
356   26.   Kim, H.J., et al., *Racial and ethnic differences in experimental pain sensitivity: systematic*
357   *review and meta-analysis.* pain, 2017. **158**(2): p. 194-211.

27. Morales, M.E. and R.J. Yong, *Racial and ethnic disparities in the treatment of chronic pain*.
Pain Medicine, 2021. **22**(1): p. 75-90.
28. Gupta, S., et al., *Potential role of female sex hormones in the pathophysiology of migraine*.
Pharmacology & therapeutics, 2007. **113**(2): p. 321-340.
29. Yunus, M.B., *The role of gender in fibromyalgia syndrome*. Current rheumatology reports,
2001. **3**(2): p. 128-134.
30. Green, T., S. Flash, and A.L. Reiss, *Sex differences in psychiatric disorders: what we can*
learn from sex chromosome aneuploidies. Neuropsychopharmacology, 2019. **44**(1): p. 9-21.
31. Hamilton, T.M., J.C. Reese, and E.L. Air, *Health care disparity in pain*. Neurosurgery Clinics,
2022. **33**(3): p. 251-260.
32. Oliveira, D.S., et al., *The impact of anxiety and depression on the outcomes of chronic low*
back pain multidisciplinary pain management—a multicenter prospective cohort study in
pain clinics with one-year follow-up. Pain Medicine, 2019. **20**(4): p. 736-746.
33. Pinheiro, M.B., et al., *Symptoms of depression and risk of low back pain: a prospective co-*
twin study. The Clinical Journal of Pain, 2017. **33**(9): p. 777-785.
34. Fernandez, M., et al., *Chronic low back pain and the risk of depression or anxiety*
symptoms: insights from a longitudinal twin study. The spine journal, 2017. **17**(7): p. 905-
912.
35. Pincus, T., et al., *A systematic review of psychological factors as predictors of*
chronicity/disability in prospective cohorts of low back pain. Spine, 2002. **27**(5): p. E109-
E120.
36. Polatin, P.B., et al., *Psychiatric illness and chronic low-back pain: the mind and the spine—*
which goes first? Spine, 1993. **18**(1): p. 66-71.
37. Bazargan, M., et al., *Multi-dimensional impact of chronic low back pain among underserved*
African American and Latino older adults. International journal of environmental research
and public health, 2021. **18**(14): p. 7246.
38. Rene, C., et al., *The effect of race in older adults presenting for chronic pain management: a*
comparative study of black and white Americans. The Journal of Pain, 2003. **4**(2): p. 82-90.
39. Hollingshead, N.A., et al., *The pain experience of Hispanic Americans: a critical literature*
review and conceptual model. The Journal of Pain, 2016. **17**(5): p. 513-528.
40. Edwards, R.R., et al., *Ethnic similarities and differences in the chronic pain experience: a*
comparison of African American, Hispanic, and white patients. Pain Medicine, 2005. **6**(1): p.
88-98.
41. Riley III, J.L., et al., *Racial/ethnic differences in the experience of chronic pain*. Pain, 2002.
100(3): p. 291-298.
42. Meints, S. and R. Edwards, *Evaluating psychosocial contributions to chronic pain*
outcomes. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2018. **87**: p.
168-182.
43. Mickle, A.M., et al., *Exploring the allostatic load of pain, interference, and the buffering of*
resilience. The Journal of Pain, 2022. **23**(5): p. 30.
44. Deuster, P.A., et al., *Allostatic load and health status of African Americans and whites*.
American journal of health behavior, 2011. **35**(6): p. 641-653.
45. Foley, H.E., et al., *Association of chronic pain with comorbidities and health care utilization:*
a retrospective cohort study using health administrative data. Pain, 2021. **162**(11): p. 2737-
2749.
46. Bao, Y., R. Sturm, and T.W. Croghan, *A national study of the effect of chronic pain on the use*
of health care by depressed persons. Psychiatric services, 2003. **54**(5): p. 693-697.

47.   Logan, D.E., et al., *From one pain to many: the emergence of overlapping pains in children*
and adolescents. The Clinical Journal of Pain, 2021. **37**(6): p. 404-412.
48.   Zeng, C., et al., *Comparison of phenomic profiles in the All of Us Research Program against*
the US general population and the UK Biobank. Journal of the American Medical Informatics
Association, 2024. **31**(4): p. 846-854.
49.   Boyd, T., et al., *Differences in provider recommendations for Black/African American and*
White patients with irritable bowel syndrome. Neurogastroenterology & Motility, 2024. **36**(3):
p. e14742.
50.   Geerlings, S.W., et al., *Longitudinal relationship between pain and depression in older*
adults: sex, age and physical disability. Social Psychiatry and Psychiatric Epidemiology,
2002. **37**: p. 23-30.
51.   Parmelee, P.A., I.R. Katz, and M.P. Lawton, *The relation of pain to depression among*
institutionalized aged. Journal of Gerontology, 1991. **46**(1): p. P15-P21.
52.   Haley, W.E., J.A. Turner, and J.M. Romano, *Depression in chronic pain patients: relation to*
pain, activity, and sex differences. Pain, 1985. **23**(4): p. 337-343.
53.   Chang, M.-H., et al., *Bidirectional association between depression and fibromyalgia*
syndrome: a nationwide longitudinal study. The Journal of Pain, 2015. **16**(9): p. 895-902.
54.   Palit, S., et al., *Exploring pain processing differences in Native Americans.* Health
Psychology, 2013. **32**(11): p. 1127.
55.   Hirsh, A., et al., *The influence of patient's sex, race and depression on clinician pain*
treatment decisions. European Journal of Pain, 2013. **17**(10): p. 1569-1579.