Title: Role of the gut microbiome in frequent gut colonization with extended-spectrum β

lactamase-producing Enterobacterales among Peruvian children

Running title: Pediatric gut microbiome and ESBL-E susceptibility

Authors: Maya L. Nadimpalli[1#,] Neha Sehgal[1], Luismarcelo Rojas-Salvatierra,[2] Robert H.

Gilman[2,3], Ashlee M. Earl[4], Colin J. Worby[4], Madison Schwab,[5] Amy J. Pickering[6,7,8], Monica J.

Pajuelo[2,3#]

Author Information:□

[1]Gangarosa Department of Environmental Health, Rollins School of Public Health, Emory

University, Atlanta, GA, USA

[2]Laboratorio de Microbiología Molecular, Laboratorios de Investigación y Desarrollo, Facultad

de Ciencias e Ingeniería, Universidad Peruana Cayetano Heredia, Lima, Peru

[3]Department of International Health, Johns Hopkins Bloomberg School of Public Health,

Baltimore, Maryland, USA

[4]Infectious Disease and Microbiome Program, The Broad Institute of MIT and Harvard,

Cambridge, MA, USA

[5]Program in Population Biology, Ecology, and Evolution, Emory University, Atlanta, GA, USA

[6]Department of Civil and Environmental Engineering, University of California, Berkeley, CA,

USA

[7]Blum Center for Developing Economies, University of California, Berkeley, CA, USA

[8]Chan Zuckerberg Biohub - San Francisco, CA, USA

#Correspondence: Maya L. Nadimpalli, Claudia Nance Rollins Room 6009, 1518 Clifton Rd

NE, Atlanta GA 30322, USA. Email: **maya.l.nadimpalli@emory.edu**. ORCID:

https://orcid.org/0000-0002-6526-116X

Monica J. Pajuelo, Av. Honorio Delgado 430, San Martín de Porres 15102, Lima Peru. Email:

monica.pajuelo.t@upch.pe. ORCID: https://orcid.org/0000-0003-3662-2250

Word Count: 5172

Abstract

Gut colonization with extended-spectrum beta lactamase-producing Enterobacterales (ESBL-E)

is increasingly common among children in low and middle-income countries. Some children

nevertheless remain never or rarely colonized during early life. Understanding how this

protection is conferred could be helpful for designing future interventions to protect children's

health. Here, we investigated whether differences in gut microbiome development could

underlie differential susceptibility to ESBL-E gut colonization over time among children in peri-

urban Lima. Weekly stool and daily surveys were collected from 345 children <3 years old

during a 2016-19 study of enteric infections. A subset of children (n=12) were rarely gut-

colonized with ESBL-E from 1-16 months of age. We performed short-read metagenomic

sequencing of stool collected at 3, 6, 7, 9, 12, and 16 months from these children and a random

subset of 42 frequently colonized children, and characterized differences in their exposures and

gut microbiomes. No differences in gut taxa or functional pathways were identified over time,

though children harbored differentially abundant taxa, more unique *E. coli* strains, and a higher

abundance of bla_{CTX-M} gene copies at ESBL-E-positive versus negative timepoints. Differing

patterns of ESBL-E colonization over time among children in peri-urban Lima do not appear to

be related to differences in gut microbiome development.

Keywords: metagenomic sequencing, AMR, children's health, ESBL, microbiome

Introduction

Antimicrobial-resistant (AMR) pathogens are a public health threat, particularly in low- and middle-income (LMIC) countries like Peru where the infectious disease burden remains high. Gut colonization with extended-spectrum β-lactamase-producing Enterobacterales (ESBL-E), which are resistant to 3rd generation cephalosporins, is common among children living in LMIC cities and is a risk factor for infections, particularly in healthcare settings.[1–5] Recent studies of community-dwelling children and adults in LMICs in South America, South Asia, and Africa indicate that ESBL-E gut colonization rates vary widely, from 30-90%.[6–9]

Previous studies in low-resource settings have sought to identify risk factors for children's ESBL-E acquisition.[10–12] This is challenging to investigate in settings like peri-urban Lima, where pediatric ESBL-E gut colonization now exceeds >70%, suggesting ubiquitous environmental exposures. Some children in this setting nevertheless remain rarely or never colonized with ESBL-E early in life. Intestinal "colonization resistance" refers to the ability of gut microbiota to inhibit colonization and overgrowth of exogenous bacteria, e.g. by limiting the availability of micronutrients and carbon sources, through direct competition for adhesion receptors on the gut epithelium, and by producing toxins and antimicrobial compounds.[13] Given that ESBL-E gut colonization is a risk factor for infection and spread, especially at higher abundances,[14,15] identifying microbial taxa that inhibit children's gut colonization with ESBL-E over time, as well as early-life exposures that support the assembly and maintenance of these protective taxa, could be helpful in designing future interventions to protect children's health.

Our team recently led a prospective birth cohort study of enteric infections among 345 children living in a shantytown of peri-urban Lima. We previously used culture-based methods to characterize children's gut colonization with ESBL-producing *Escherichia coli* and ESBL-producing *Klebsiella* spp., *Enterobacter* spp., and *Citrobacter* spp. (KEC) at or near 1, 3, 4, 5, 6, 7, 9, 12, and 16 months of age for a subset of 112 children enrolled in the cohort study.[4] We noted that while most children were frequently gut-colonized with ESBL-E over time, a fraction

(n=12) were rarely or never colonized. Here, we investigated whether differences in gut

microbiome development could underlie differing patterns of susceptibility to ESBL-E gut-

colonization among this population.

Materials and methods

Stool collection

Stool samples were collected during a prospective cohort study of childhood enteric infections in

Villa El Salvador, Lima, Peru (NIH R01AI108695-01A1).[4] Pregnant women were identified

through an initial population screening in the community and through community health centers.

Full-term babies were enrolled no later than 4 weeks after birth from February 2016-May 2017

(n=345), and daily surveillance data were collected through at least two years of age. Exclusion

criteria for newborns participating in the parent cohort included low birth weight and presence of

any severe disease requiring hospitalization early in life. Starting at recruitment and weekly

thereafter, caregivers also kept a freshly soiled diaper for field worker collection; field workers

retrieved these once caretakers alerted them that soiled diapers were available for pick-up.

Upon arrival at a local lab facility, stool was aliquoted and stored at -20°C for up to 48h, then

transported to UPCH for long-term storage at -80°C.

Sample selection

As previously described,[4] Enterobacterales were cultured from weekly stool samples of all

newborns enrolled after December 2016 (112/345), in support of a secondary study focused on

enterotoxigenic *E. coli*.[16] For each sample, Enterobacterales colonies were pooled then

archived. Colonies collected at approximately 1, 3, 4, 5, 6, 7, 9, 12, and 16 months of age were

later screened for ESBL production using the double disk diffusion assay. We observed two

distinct patterns: children who were never or rarely colonized with ESBL-E (n=12), hereafter

"*rarely colonized,*" and children who were often or always colonized (n=100), hereafter

"*frequently colonized"* (**Figures S1 and S2**). Rarely colonized children first tested positive for

ESBL-E at approximately 6 months or later, and additionally were colonized at fewer than half of

all tested timepoints (up to 9 per child). Frequently colonized children were colonized at more

than half of all tested timepoints.

Exposure assessment

As previously described,[4,16] socioeconomic variables were collected at recruitment and then

once yearly. Additionally, field staff visited infants every day except Sunday to collect survey

data from the previous 24 hours. If field workers were unable to visit an infants' home (e.g., day

was Sunday), field workers collected survey data for the missing day at the next visit. At each

daily visit, caretakers were asked about children's milk consumption, antibiotic use, diarrhea

(defined as ≥3 liquid or semi-liquid stools), and symptoms of infection (e.g. stomach pain, fever,

vomiting, swelling, and loss of appetite) in the past 24 hours. Caretakers were asked to provide

medication packaging so trained field workers could confirm it was an antibiotic. Because

survey data were collected daily, this packaging was typically available. A new antibiotic course

began when caretakers reported antibiotic consumption following two days of no exposure; a

course ended when the child did not consume antibiotics for two consecutive days.

Metagenomic sequencing

We selected stool specimens collected at approximately 3, 6, 7, 9, 12, and 16 months of age for

analysis, as these were timepoints for which we had previously characterized children's ESBL-E

gut colonization, and because these timepoints span both stages of early microbiome

development as previously defined by the TEDDY study,[17] i.e., the developmental phase (3-14

months) and the transitional phase (15-30 months). We performed short-read metagenomic

sequencing on available frozen stool specimens collected at these timepoints from all rarely

colonized children (n=12 children, n=71 stool samples), and at these same timepoints from a

randomly selected subset of 42 frequently colonized children (n=238 stool samples). We used

the QIAamp PowerFecal Pro DNA Kit (Qiagen) to extract total DNA from approximately 0.25 g

of frozen feces at UPCH. Extraction blanks were included with each batch of extractions. DNA

was quantified using a Qubit 4 fluorometer (Invitrogen); DNA concentrations were below the

Qubit level of detection (<0.1 ng/μL) for all extraction blanks. DNA extracts meeting the

minimum concentration for sequencing (>10 ng/μL) were sent to the Broad Institute for short-

read, paired-end 150 bp sequencing on an Illumina Novaseq 6000 System using SP4 flow cells

to achieve 6 Gb per sample.

Processing metagenomic data

Sequencing adaptors and low quality reads were removed using bbmap.[18] Default settings were

used for adaptor removal (ktrim=r k=23 mink=11 hdist=1); we retained reads with a minimum

length of 50 bp (minlen=50) and a quality value of at least 20 (trimq=20). After tabulating the

number of raw reads per sample, we excluded samples with more or less reads than two

standard deviations from the mean from further analysis (n=11). Reads that mapped to a non-

redundant version of the Genome Reference Consortium Human Build 38 (GRCh38;

www.ncbi.nlm.nih.gov) using Bowtie2 v2.5.1 were considered "human contaminant sequences"

and discarded. Reads with human contaminant sequences removed were deposited in the

Sequence Read Archive (https://www.ncbi.nlm.nih.gov/sra) under project number

PRJNA1138246 (**Table S1**).

Profiling taxonomy, functional potential, and antibiotic resistance genes

Taxonomic assignment of short reads was performed using MetaPhlAn3 (with db v31).[19]

MetaPhlAn3 uses clade-specific marker genes to assign taxonomy at the species level; results

are then aggregated to higher taxonomic levels. Alpha diversity metrics were derived from the

MetaPhlAn3 relative abundance output using the vegan package in R, using the *diversity* and

richness functions [https://cran.r-project.org/web/packages/vegan/].

Functional annotations were performed using Humann3,[20] which uses MetaPhlAn3

output to generate organism-specific gene family and pathway abundance profiles. Briefly, for

each sample, reads were aligned to a sample-specific species-level pangenome using

Bowtie2.[21] Unclassified reads were then aligned to UniRef90, a comprehensive and non-

redundant protein database,[22] using DIAMOND.[23] Gene family abundance was annotated using

UniRef90 definitions. Gene families annotated to metabolic reactions were then further analyzed

to identify a parsimonious set of pathways explaining these observed reactions, using MinPath

and Metacyc pathway definitions.[24,25] We considered community-level (rather than organism-

specific) metabolic pathway abundance in our statistical analyses.

Antibiotic resistance genes (ARGs) were identified by mapping short reads to the

Resfinder database (v. 3.1.1) using the KMA tool.[26] Matches with >90% coverage and >95%

identity were considered true hits. We identified the number of genome equivalents in each

sample using Microbe Census.[27] To normalize for the number of bacterial genomes in each

sample, ARG abundance was calculated as fragments per kilobase per million mapped reads

(FPKM) divided by the total number of genome equivalents in that sample.[28] FPKM is a

common approach for comparing ARG abundance across samples, and is used to account for

differences in sequencing depth between samples as well as the fact that ARGs with greater

length are otherwise likely to be overrepresented, regardless of sequencing depth.[28,29] We

chose to normalize by genome equivalents, rather than total reads classified as bacteria by

MetaPhlAn3, because bacterial genome sizes can vary widely. The abundance of bla_{CTX-M}

genes (all allelic variants collectively considered as one group) and total ARGs in each sample

was determined using this approach. FPKMs were log_{10} transformed for analysis.

Characterizing *E. coli, Klebsiella* spp., and Enterobacter *spp.* strain diversity

We used the StrainGE tool[30] to characterize *E. coli, Klebsiella* spp., and Enterobacter *spp.*

strains in children's fecal metagenomes. We theorized that commensal strains could play a role

in excluding ESBL-producing *E. coli* and KEC from the gut environment through species-specific

effects. StrainGE can sensitively profile strains in a microbial community with >0.1% relative

abundance.[30] A reference database was built using complete *E. coli* genomes from NCBI's

RefSeq database (3,108 genomes downloaded March 5, 2024), and each genome was

assigned a Clermont phylotype using EzClermont v.0.70.[31] A separate reference database was

built using complete *Klebsiella* spp. and *Enterobacter* spp. genomes from NCBI's RefSeq

database (4,121 genomes downloaded June 5, 2025). *Citrobacter* spp., were relatively rare

(0.014% abundance on average) and therefore were not considered. Briefly, references in each

database were kmerized using the *straingst* kmerize command and the default k-mer size of 23.

To reduce redundancy, closely related genomes were clustered based on k-mer similarity, using

the *straingst kmersim* and *straingst cluster* commands. A single pangenome k-mer database

was then created for *E. coli* and for *Klebsiella* spp./*Enterobacter* spp., respectively, using

straingst createdb. To align sample metagenome reads to each pangenome k-mer database,

reads were first kmerized using *straingst kmerize.* We then used *straingst run* to identify

reference strains present in each sample.

<u>Data analysis</u>

We examined differences in sociodemographic characteristics, diarrhea rates, and antibiotic use

between rarely and frequently gut-colonized children using t-tests and chi square tests for all

112 children for whom we characterized ESBL-E gut colonization, and separately for the 54

children whose stool samples were sequenced.

For the subset of 54 children with fecal metagenomes, we examined whether any taxa or

metabolic pathways were differentially abundant between rarely and frequently gut-colonized

children across time using mixed effects linear regression models. An arcsine square root

transformation was applied to relative abundance values. We also used mixed effects linear

regression models to examine differences in log10 transformed species richness values,

Shannon diversity, bla_{CTX-M} FPKM and total ARG FPKM over time between groups. Transformed

relative abundance values, alpha diversity metrics, and ARG FPKM were modeled as a function

of subject (random effects term), colonization pattern, i.e., rarely versus frequently colonized

(binary fixed effects term), child age (continuous fixed effects term), and hours before diaper

retrieval (continuous fixed effects term). We also ran additionally parametrized models adjusted

for factors known to affect the gut microbiome, including child's delivery mode (binary fixed

effects term) and number of diarrhea episodes and antibiotic courses in the past 30 days

(continuous fixed effects terms), as well as socioeconomic characteristics that differed between

groups with p<0.20 (*i.e.*, maternal education and presence of a flush toilet; availability of piped

water was not considered as it was highly collinear with presence of a flush toilet). We did not

adjust for breastfeeding as there was minimal variation in this exposure over time (**Table S2**).

All taxa or metabolic pathways detected in at least 10% of samples were fitted using the nlme

package in R. Significance of associations was determined using Wald's test, and p-values

were adjusted for multiple hypothesis testing using the Benjamini-Hochberg correction.

For the subset of metagenomes for which we had matched ESBL-E gut colonization

status, we examined whether any taxa or metabolic pathways were differentially abundant

between timepoints where children were ESBL-E positive versus negative using mixed effects

logistic regression models. For any given taxa or metabolic pathway detected in at least 10% of

samples, we modeled ESBL-E positivity (yes/no) as a function of subject, child age, and hours

before diaper retrieval and the transformed relative abundance value of that taxa. We ran

additionally parametrized models that adjusted for number of recent diarrhea episodes and

antibiotic courses, maternal education, and presence of a flush toilet. We used the lme4

package in R and adjusted p-values using the Benjamini-Hochberg correction. For this same

subset of metagenomes, we examined whether log10 transformed species richness values,

Shannon diversity, bla_{CTX-M} genes or total ARGs differed between timepoints where children

were ESBL-E positive versus negative, using similar model specifications as above.

We compared the number of unique *E. coli, Klebsiella* spp., and *Enterobacter* spp.

strains between rarely and frequently colonized children using Welch two sample t-tests, which

do not assume equal variance between groups. We used Welch two sample t-tests to compare

strain richness between groups overall (pooling all timepoints together), and at specific

timepoints. Similarly, we compared strain richness at timepoints where children were ESBL-E

positive versus negative.

Statistical significance was defined by $\alpha = 0.05$ and p-values are two-sided unless

stated otherwise. All analyses were conducted in R Studio version (2024.04.1+748).

Results

Exposure profiles were similar between children who were rarely versus frequently gut-

colonized with ESBL-E

There were no significant differences in exposures that are well-known to impact the gut

microbiome, including rates of vaginal birth, antibiotic consumption rates, children's age at time

of first antibiotic consumption, or numbers of diarrhea episodes per child month between rarely

versus frequently colonized children (**Table 1)**. Rarely colonized children did consume slightly

fewer antibiotic courses overall (3.4 vs 4.5 courses/child-year, *p*=0.2), and fewer azithromycin

courses in particular (0.1 vs 0.5 courses/child-year, *p*=0.14), but differences were not

statistically significant. Only 2/112 children did not consume any antibiotics, yet both were

frequently colonized with ESBL-E.

Among children who were rarely gut-colonized with ESBL-E, there was a positive but

non-significant trend towards living in households of lower socioeconomic status (SES).

Specifically, these children were somewhat more likely to live in households that relied on a

public tap for drinking water (*p*=0.06) versus having an indoor connection, and more likely to

rely on pit latrines versus indoor flush toilets (p=0.05). Most households that relied on water taps

also used pit latrines, and vice versa. Rarely colonized children were also somewhat less likely

to be born to a mother who had completed high school (p=0.11), although this difference was

not statistically significant. Children's antibiotic use (total courses/child-year) was weakly

correlated with household reliance on public water taps (r_{pb}=0.24, p<0.05 by point-biserial

correlation) and use of a pit latrine (r_{pb}=0.25, p<0.05), but was not correlated with maternal

education attainment (r_{pb}=0.04, p=0.67).

Breastfeeding was exceptionally common among all children. On average, children

received breastmilk nearly every day in the 30 days prior to a stool sample through 16 months

of age (**Table S2**). Notably, nearly 1/3 children who were frequently colonized with ESBL-E

regularly consumed formula early in life, compared to just 1 of 12 rarely colonized children

(**Table 1**). However, this difference was not significant (p=0.29).

The distributions of these exposures were similar among the subset of 54/112 children

whose fecal metagenomes were sequenced for this study, as well as the 274/345 children

participating in the parent cohort who completed at least 16 months of data collection (**Table

S3**). Of note, both children whose ESBL status was previously ascertained (n=112) and children

whose fecal metagenomes were sequenced for this study (n=54) were more likely to be born by

Cesarean section than children who completed 16 months of data collection in the parent cohort

study (n=274) (**Table S3**).

<u>Peruvian children's gut microbiomes are dominated by *Bifidobacterium* spp. regardless of</u>

<u>ESBL-E colonization status</u>

Samples included in our final analysis included 66 fecal metagenomes from 12 rarely colonized

children and 232 fecal metagenomes from 42 frequently colonized children (n=298 total, 4-6

samples per child).

As expected given the high proportion of breastfeeding in this setting, children's gut

microbiomes were largely dominated by *Bifidobacterium* spp., comprising 58% of bacterial

reads in average at 3 months, and 34% by 16 months (**Figure 1**). *B. longum* and *B. breve*

predominated earlier in life (comprising up to 20% and 12% of bacterial reads, respectively, until

9 months of life), while *B. bifidum* consistently comprised between 10-15% of bacterial reads

(**Figure 2**). The number of unique *Bifidobacterium* spp. in children's guts ranged from 0-9

(median=4) and increased as children aged (β=0.14, SE=0.019, *p*-value<0.0001). Pooling all

timepoints together, there was no difference in the relative abundance of any *Bifidobacterium*

spp. between children who were rarely versus frequently gut-colonized with ESBL-E. Likewise,

the number of unique *Bifidobacterium* spp. did not differ between groups.

<u>No differences in diversity, relative abundance, or functional potential of gut bacteria between</u>

<u>rarely and frequently colonized children</u>

Besides *Bifidobacterium*, other dominant bacterial genera included *Blautia* (average across all

samples 3.1%), *Bacteroides* (2.1%), *Escherichia* (1.9%), *Lactobacillus* (1.1%), *Faecalibacterium*

(0.9%), and *Klebsiella* (0.8%) (**Figure 1**). Potential enteropathogens like *Klebsiella pneumoniae*

and *Enterococcus faecium* were detected among nearly all children (**Table S4**). After

adjustment for multiple comparisons, no taxa were differentially abundant between children who

were rarely versus frequently gut-colonized with ESBL-E over time in fully parameterized

models (corrected p-values all > 0.89). Results were similar for simpler models that only

adjusted for children's colonization pattern, child age, and hours before diaper retrieval (fdr-

corrected p-values all >0.66).

Like others,[17] we found the number of unique bacterial species in children's fecal

metagenomes increased as they aged, from 31 to 75, on average. The diversity of children's

fecal metagenomes also increased (0.99 Shannon diversity index on average at 3 months, to

2.31 at 16 months) (**Figure S3**). Pooling all timepoints together, we observed no differences in

diversity or richness between groups (p-values> 0.62). There were also no differences in

median richness or Shannon diversity at any specific sampling point (p>0.52 for all pairwise

differences by Wilcoxon rank sum tests).

A total of 449 metabolic pathways were identified in fecal metagenomes from at least

10% of children. After adjustment for multiple comparisons, no metabolic pathways were

differentially abundant between children who were rarely versus frequently gut-colonized with

ESBL-E over time in fully parameterized models (fdr-corrected p-values all > 0.99). Results

were similar for simpler models that only adjusted for children's colonization pattern, child age,

and hours before diaper retrieval (fdr-corrected p-values all >0.87).

Some differences in *Klebsiella* and *Enterobacter* strain richness between rarely and frequently

colonized children

E. coli strains were detected in every sample. Among the 252 unique *E. coli* strains detected

overall, most belonged to phylotypes A and B1 (**Figure 3A**). There were no differences in the

number of *E. coli* strains harbored by rarely and frequently colonized children, overall or by

timepoint (p>0.15 for all pairwise comparisons by Welch two-sample t-test) (**Figure 3A**). There

was no difference in the proportion of strains belonging to any given *E. coli* phylotype between

rarely and frequently colonized children at any time point (p>0.26 for all pairwise comparisons

by Fisher test).

Klebsiella spp. strains were detected in approximately one-third of samples (106/298).

Among the 70 unique strains detected overall, most were *K. pneumoniae* (66/70). Rarely

colonized children harbored fewer *Klebsiella* spp. strains on average (mean=1.7), pooling all

timepoints together, than frequently colonized children (mean=3.0, *p*=0.03 by Welch two-sample

t-test), (**Figure 3B**). *Enterobacter* strains were rarer, detected in only 13% of samples (39/298).

E. hormaechei (n=9), *E. asburiae* (n=4), and *E. cloacae* (n=4) comprised the 17 *Enterobacter*

strains detected. On average and across all timepoints, rarely colonized children harbored fewer

Enterobacter spp. strains (mean=0.3) than frequently colonized children (mean=1.1, *p*=0.02 by

Welch two-sample t-test) (**Figure 3C**).

<u>Some differences in ARG abundance between rarely versus frequently colonized children</u>

Rarely colonized children harbored a significantly lower abundance of bla_{CTX-M} genes (*p*<0.0001)

in their stool, on average across all time points (mean log10 bla_{CTX-M} FPKM = 4.1, SD=4.6),

compared to persistently colonized children (mean log10 bla_{CTX-M} FPKM = 4.9, SD=5.4) (**Figure**

S4). We noted significant differences at the ages of 6 (p=0.002) and 16 months (p=0.006) by

Welch two-sample test. However, there was no difference in total ARG abundance, on average

across all timepoints, between these groups (among rarely colonized children: mean log10 total

ARG FPKM=6.9, SD=1.9, among persistently colonized children: mean log10 total ARG

FPKM=7.0, SD=7.1).

<u>Differences in children's gut microbiomes and resistomes at ESBL-E positive versus negative</u>

<u>sampling points</u>

We also examined differences in children's gut microbiomes and resistomes at timepoints

where they were ESBL-E colonized versus not (ESBL-E gut colonization status available for

263/298 fecal metagenomes), regardless of whether they were rarely or frequently gut-

colonized with ESBL-E over time.

We noted several differences in children's gut microbiomes at timepoints when they

were ESBL-E colonized. First, the relative abundance of the genus *Slackia* (0.006%, fdr-

corrected p-val=0.047) in children's stool was higher at timepoints where children were not gut-

colonized with ESBL-E by culture, when controlling for child age, hours before diaper retrieval,

and repeated measurements (**Figure S5**). There were also non-significant trends towards a

higher abundance of *Erysipelotrichia* (0.47%; fdr-corrected p-val=0.18), *Proteobacteria* (0.03%,

fdr-corrected p-val=0.16) and *Bacteroidetes* (0.03%; fdr-corrected p-val=0.16) at timepoints

where were children colonized with ESBL-E by culture, and a trend towards lower abundance of

Actinobacteria at these timepoints (46%; fdr-corrected p-val=0.12). No taxa were differentially

abundant in fully parameterized models that additionally adjusted for recent diarrhea, recent

antibiotic use, maternal education, and presence of a flush toilet. Second, children harbored

more unique *E. coli* strains at sampling points where they were gut-colonized with ESBL-E

(mean=2.3 across all age groups) versus sampling points where they were not (mean=2.1,

p=0.05 by Welch two-sample t-test). However, there were no differences in the number of

Klebsiella spp. or *Enterobacter* spp. strains harbored by children at ESBL-positive versus

negative timepoints (p>0.25 for both comparisons). There were no differences in species

richness or diversity at timepoints where children were ESBL-E colonized versus not, nor any

differentially abundant metabolic pathways.

There were also significant differences in children's gut resistomes between sampling

points where children were gut-colonized with ESBL-E versus not (**Figure 4**). On average,

children harbored significantly higher bla_{CTX-M} gene FPKM (p<0.0001) at sampling points where

they were gut-colonized with ESBL-E (mean log10 bla_{CTX-M} FPKM=4.9, SD=5.4) versus

sampling points where they were not (mean log10 bla_{CTX-M} =4.4, SD=5.2). We noted significant

differences at all sampling points by Welch two-sample test (all p≤0.002). On average and

across all timepoints, children also harbored significantly higher total ARG FPKM at sampling

points (p=0.005) where they were gut-colonized with ESBL-E (mean log10 bla_{CTX-M} FPKM=7.02,

SD=7.1) versus sampling points where they were not (mean log10 bla_{CTX-M} =6.9, SD=6.8). We

noted a significant difference at 7 months of age (p=0.03) by Welch two-sample text.

Discussion

This is the first longitudinal study of the gut microbiome among South American children. As

previously reported, gut colonization with ESBL-E was prevalent among children in this peri-

urban setting and steadily increased as children aged.[4] Similar to recent studies among healthy

adults,[32–34] our findings suggest that differences in gut microbiome development do not underlie

patterns of ESBL gut colonization over time among children in peri-urban Lima, though subtle

differences may have been missed due to our small sample size. To our knowledge, this is the

first study to examine this research question among young children whose gut microbiomes are

rapidly developing.

389   Three recent Western studies and two LMIC studies have investigated associations

between ESBL-E gut colonization, gut microbial taxa, and/or the gut metabolome at single time

points.[32–36] Two case-control studies among Dutch[33] and French adults[34] found no differences in

alpha diversity, taxa abundance, or microbial functional annotations between healthy adults

who were gut-colonized with ESBL-E versus not, after controlling for age, sex, ethnicity, and

recent travel. Another study among US travelers[32] found no difference in the pre-travel gut

microbiomes of individuals who acquired ESBL-E while abroad versus those who did not,

suggesting that differences in exposures while traveling, rather than inherent differences in

intestinal colonization resistance, are more important determinants of ESBL-E acquisition. One

study of healthy Thai adults[36] reported differences in gut taxa between ESBL-E carriers and

non-carriers, but because potential confounders were not controlled for, it is challenging to

attribute differences to ESBL-E colonization status. A small study among indigenous persons in

French Guyana identified some differences in gut microbial taxa between 8 cases and 24

controls, but whether these differences preceded or were a result of ESBL-E gut colonization

could not be determined.[35] To our knowledge, no previous study has examined whether

differences in the gut microbiome over time, rather than just at a single timepoint, may underlie

ESBL-E gut colonization patterns. Our study adds to the existing literature by indicating that

patterns of rare versus frequent ESBL-E gut colonization among children who are frequently

exposed to enteropathogens and antibiotics may not primarily be a function of intestinal

colonization resistance.

Taxa that previous studies have identified to be associated with ESBL carriage or non-

carriage differed from what we observed. *Erysipelotrichia*, *Proteobacteria*, and *Bacteroidetes*,

which are commonly isolated in children's stool,[37,38] were somewhat more abundant at ESBL-

positive timepoints (*p-val*=0.16 – 0.18). *Proteobacteria* have previously been associated with gut

dysbiosis, though associations are often inconsistent among children from different world

regions.[39,40] *Slackia*, which is not well-described among children but has previously been

isolated from both gut and oral infections in adults,[41] was more abundant at ESBL-negative

timepoints though overall abundance was low (0.006%). *Actinobacteria*, which primarily

comprise *Bifidobacteria* spp. and which dominate the infant gut,[38] were also somewhat less

abundant at ESBL-negative timepoints (*p-val*=0.12). Differences between our study and

previous work[35,36] could be a result of our focus on the pediatric microbiome, differences in our

sequencing approach, differences in environmental exposures, or other factors. We note that

whether these differentially abundant taxa preceded or followed ESBL-E gut colonization cannot

be distinguished through our analytical approach.

Overall, the fact that we did not identify gut taxa or metabolic pathways that were

differentially abundant between rarely and frequently colonized children could be because such

differences do not exist, or, we may have been underpowered to detect these differences given

our modest sample size. Breastfeeding was exceptionally common in this setting and given how

strongly maternal breastmilk shapes the infant gut microbiome,[42] this may have further obscured

or minimized gut microbiome differences between groups. Given the high and increasing rates

of ESBL-E gut colonization in community settings in LMICs, we note that enrolling sufficient

children to identify small effect sizes may be challenging in future studies. Alternatively, other

gut microbiome differences that we were unable to explore may underlie differences in ESBL-E

gut colonization patterns in this setting, *e.g.*, higher bacteriocin production among commensal

strains harbored by rarely colonized children, which could favor the exclusion of ESBL-E.[43].

Interestingly, strain-level analyses using the StrainGE toolkit did identify lower richness of

Klebsiella spp. and *Enterobacter* spp. among rarely versus frequently colonized children,

suggesting differences in gut microbial dynamics could exist at the strain level between these

groups. Other studies have found that certain ESBL-E strain/gene combinations are more likely

to persist in the gut than others.[44,45] However, manipulating gut microbial strain dynamics is not

likely to be a feasible approach for protecting children from ESBL-E colonization through future

interventions.

This study had strengths and limitations. Our study represents the first longitudinal

analysis of the gut microbiome of South American children, whose fecal metagenomes are

under-represented in global databases.[28] Because daily surveys were collected, we were able

to control for time-varying exposures that are known to affect the gut microbiome, including

antibiotic use and recent diarrhea, with a low risk of recall bias. However, our study was

relatively small, and we may have been underpowered to detect differences in taxonomic

composition and functional potential between groups. Data on maternal antibiotic use during the

prenatal period, including intrapartum antibiotic prophylaxis, were not collected for the parent

study and likely had important effects on the trajectory of children's gut microbiome

development, but we could not consider those effects here. Further, as previously noted,[4] we

used culture-based methods with a relatively high lower limit of detection (~2 log10 CFU/g wet

feces) to identify ESBL-E gut colonization for this study. Rarely colonized children may have

been gut-colonized with ESBL-E at timepoints where they tested negative, just at concentrations

below the limit of detection. Since higher concentrations of multidrug-resistant Enterobacterales

in the gut are associated with elevated risks of infection,[14,15] identifying whether specific gut taxa

may suppress the growth of ESBL-E or the proliferation of bla_{CTX-M} genes remains a clinically

relevant question, especially in LMICs where multidrug-resistant infections are becoming

increasingly common.[46,47] Finally, there is no universally accepted definition of rare versus

persistent ESBL-E colonization, and studies have employed different definitions.[44,48] It is

possible that using different thresholds for classifying rare versus frequent ESBL-E colonization

would yield different results.

Conclusion

This study examined whether differences in Peruvian children's gut microbiome development

could underpin differences in early-life susceptibility to ESBL-E gut colonization. We did not

identify any significant differences in species richness, Shannon diversity, *E. coli* strain diversity,

or the relative abundance of specific taxa or metabolic pathways between rarely versus

frequently colonized children, suggesting that differences in gut microbiome development are

unlikely to explain ESBL-E gut colonization patterns in this setting. Regardless of gut

colonization patterns over time, children harbored a lower relative abundance of *Slackia* spp.,

more unique *E. coli* strains, and significantly more bla_{CTX-M} gene copies at timepoints where they

were ESBL-E positive. Overall, our findings are in line with recent studies among healthy

adults[32–34] suggesting that differences in the gut microbiome are unlikely to underlie patterns of

susceptibility to ESBL-E gut-colonization. Protecting children from ESBL-E gut colonization will

likely require that their early-life exposures to ESBL-E reservoirs are minimized.

<u>Acknowledgements</u>

We thank all field workers and the community of Villa El Salvador for their collaboration in the

parent study. A preprint of this manuscript was previously posted on medRxiv (doi:

https://doi.org/10.1101/2024.11.06.24316595).

<u>Disclosure of interest</u>

The authors report there are no competing interests to declare. This work was supported by NIH

R01AI108695-01A, NIH award KL2TR002545, and NIH award U19AI110818. MLN was

supported by Emory University and the MP3 Initiative. NS was supported by NIH award

5T32ES012870. The content is solely the responsibility of the authors and does not necessarily

represent the official views of Emory University or the MP3 Initiative.

Data Availability

Children's fecal metagenomes are available in NCBI's Sequence Read Archive

(https://www.ncbi.nlm.nih.gov/sra) under BioProject number PRJNA1138246.

Ethics approval and consent to participate

The parent cohort study was approved by the Institutional Review Boards of Asociación

Benefica PRISMA, Universidad Peruana Cayetano Heredia, and Johns Hopkins University.

Written informed consent was obtained from caretakers of the infants for both study participation

and the use of collected specimens for subsequent research. The analysis of stool specimens

for this substudy was approved by the Institutional Review Boards of Asociación Benefica

PRISMA and Universidad Peruana Cayetano Heredia (no. 201592). Metagenomic sequencing

of stool specimens was determined to be exempt by the Tufts University Health Sciences IRB

(STUDY00001328).

References

1 Cheikh A, Belefquih B, Chajai Y, Cheikhaoui Y, El Hassani A, Benouda A.
Enterobacteriaceae producing extended-spectrum β-lactamases (ESBLs) colonization as a
risk factor for developing ESBL infections in pediatric cardiac surgery patients: 'retrospective
cohort study'. *BMC Infect Dis* 2017; **17**: 237.

2 Ben-Ami R, Schwaber MJ, Navon-Venezia S, *et al.* Influx of extended-spectrum beta-
lactamase-producing enterobacteriaceae into the hospital. *Clin Infect Dis* 2006; **42**: 925–34.

3 Denis B, Lafaurie M, Donay J-L, *et al.* Prevalence, risk factors, and impact on clinical
outcome of extended-spectrum beta-lactamase-producing Escherichia coli bacteraemia: a
five-year study. *Int J Infect Dis* 2015; **39**: 1–6.

4 Nadimpalli ML, Rojas Salvatierra L, Chakraborty S, *et al.* Effects of breastfeeding on
children's gut colonization with multidrug-resistant Enterobacterales in peri-urban Lima, Peru.
Gut Microbes 2024; **16**: 2309681.

5 Nadimpalli ML, Marks SJ, Montealegre MC, *et al.* Urban informal settlements as hotspots of
antimicrobial resistance and the need to curb environmental transmission. *Nature
Microbiology* 2020; **5**: 787–95.

6 Singh SR, Mao B, Evdokimov K, *et al.* Prevalence of MDR organism (MDRO) carriage in
children and their household members in Siem Reap Province, Cambodia. *JAC-Antimicrobial
Resistance* 2020; **2**. DOI:10.1093/jacamr/dlaa097.

7 Caudell MA, Ayodo C, Ita T, *et al.* Risk Factors for Colonization With Multidrug-Resistant
Bacteria in Urban and Rural Communities in Kenya: An Antimicrobial Resistance in
Communities and Hospitals (ARCH) Study. *Clin Infect Dis* 2023; **77**: S104–10.

8 Lautenbach E, Mosepele M, Smith RM, *et al.* Risk Factors for Community Colonization With
Extended-Spectrum Cephalosporin-Resistant Enterobacterales (ESCrE) in Botswana: An
Antibiotic Resistance in Communities and Hospitals (ARCH) Study. *Clin Infect Dis* 2023; **77**:
S89–96.

9 Araos R, Smith RM, Styczynski A, *et al.* High Burden of Intestinal Colonization With
Antimicrobial-Resistant Bacteria in Chile: An Antibiotic Resistance in Communities and
Hospitals (ARCH) Study. *Clin Infect Dis* 2023; **77**: S75–81.

10 Kothari C, Gaind R, Singh LC, *et al.* Community acquisition of β-lactamase producing
Enterobacteriaceae in neonatal gut. *BMC Microbiol* 2013; **13**: 136.

11 Islam MA, Amin MB, Roy S, *et al.* Fecal colonization with multidrug-resistant E. coli among
healthy infants in rural Bangladesh. *Frontiers in Microbiology* 2019; **10**: 640.

12 Tellevik MG, Blomberg B, Kommedal Ø, Maselle SY, Langeland N, Moyo SJ. High
Prevalence of Faecal Carriage of ESBL-Producing Enterobacteriaceae among Children in
Dar es Salaam, Tanzania. *PLOS ONE* 2016; **11**: e0168024.

13 Lawley TD, Walker AW. Intestinal colonization resistance. *Immunology* 2013; **138**: 1–11.

14 Ruppé E, Lixandru B, Cojocaru R, *et al.* Relative fecal abundance of extended-spectrum-β-
lactamase-producing Escherichia coli strains and their occurrence in urinary tract infections in
women. *Antimicrob Agents Chemother* 2013; **57**: 4512–7.

15 Shimasaki T, Seekatz A, Bassis C, *et al.* Increased Relative Abundance of Klebsiella
pneumoniae Carbapenemase-producing Klebsiella pneumoniae Within the Gut Microbiota Is
Associated With Risk of Bloodstream Infection in Long-term Acute Care Hospital Patients.
Clin Infect Dis 2019; **68**: 2053–9.

16 Pajuelo MJ, Noazin S, Cabrera L, *et al.* Epidemiology of enterotoxigenic Escherichia coli and
impact on the growth of children in the first two years of life in Lima, Peru. *Front Public Health*
2024; **12**: 1332319.

17 Stewart CJ, Ajami NJ, O'Brien JL, *et al.* Temporal development of the gut microbiome in early
childhood from the TEDDY study. *Nature* 2018; **562**: 583–8.

18 Brian Bushnell. BBMap. https://sourceforge.net/projects/bbmap/.

19 Truong DT, Tett A, Pasolli E, Huttenhower C, Segata N. Microbial strain-level population
structure and genetic diversity from metagenomes. *Genome Res* 2017; **27**: 626–38.

20 Beghini F, McIver LJ, Blanco-Míguez A, *et al.* Integrating taxonomic, functional, and strain-
level profiling of diverse microbial communities with bioBakery 3. *Elife* 2021; **10**: e65088.

21 Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 2012; **9**:
357–9.

22 Suzek BE, Wang Y, Huang H, McGarvey PB, Wu CH, UniProt Consortium. UniRef clusters: a
comprehensive and scalable alternative for improving sequence similarity searches.
Bioinformatics 2015; **31**: 926–32.

23 Buchfink B, Reuter K, Drost H-G. Sensitive protein alignments at tree-of-life scale using
DIAMOND. *Nat Methods* 2021; **18**: 366–8.

24 Caspi R, Billington R, Keseler IM, *et al.* The MetaCyc database of metabolic pathways and
enzymes - a 2019 update. *Nucleic Acids Research* 2020; **48**: D445–53.

25 Ye Y, Doak TG. A Parsimony Approach to Biological Pathway Reconstruction/Inference for
Genomes and Metagenomes. *PLOS Computational Biology* 2009; **5**: e1000465.

26 Clausen PTLC, Aarestrup FM, Lund O. Rapid and precise alignment of raw reads against
redundant databases with KMA. *BMC Bioinformatics* 2018; **19**: 307.

27 Nayfach S, Pollard KS. Average genome size estimation improves comparative
metagenomics and sheds light on the functional ecology of the human microbiome. *Genome*
Biol 2015; **16**: 51.

28 Fuhrmeister ER, Harvey AP, Nadimpalli ML, *et al.* Evaluating the relationship between
community water and sanitation access and the global burden of antibiotic resistance: an
ecological study. *The Lancet Microbe* 2023; **4**: e591–600.

29 Hendriksen RS, Munk P, Njage P, *et al.* Global monitoring of antimicrobial resistance based
on metagenomics analyses of urban sewage. *Nature Communications* 2019; **10**: 1124.

30 Van Dijk LR, Walker BJ, Straub TJ, *et al.* StrainGE: a toolkit to track and characterize low-
abundance strains in complex microbial communities. *Genome Biol* 2022; **23**: 74.

31 Waters NR, Abram F, Brennan F, Holmes A, Pritchard L. Easy phylotyping of Escherichia coli
via the EzClermont web app and command-line tool. *Access Microbiol* 2020; **2**: acmi000143.

32 Worby CJ, Sridhar S, Turbett SE, *et al.* Gut microbiome perturbation, antibiotic resistance,
and Escherichia coli strain dynamics associated with international travel: a metagenomic
analysis. *The Lancet Microbe* 2023; **0**. DOI:10.1016/S2666-5247(23)00147-7.

33 Ducarmon QR, Zwittink RD, Willems RPJ, *et al.* Gut colonisation by extended-spectrum β-
lactamase-producing Escherichia coli and its association with the gut microbiome and
metabolome in Dutch adults: a matched case-control study. *Lancet Microbe* 2022; **3**: e443–
51.

34 Boyd A, El Dani M, Ajrouche R, *et al.* Gut microbiome diversity and composition in individuals
with and without extended-spectrum β-lactamase-producing *Enterobacterales* carriage: a
matched case–control study in infectious diseases department. *Clinical Microbiology and
Infection* 2024; published online March 23. DOI:10.1016/j.cmi.2024.03.016.

35 Gosalbes MJ, Vázquez-Castellanos JF, Angebault C, *et al.* Carriage of Enterobacteria
Producing Extended-Spectrum β-Lactamases and Composition of the Gut Microbiota in an
Amerindian Community. *Antimicrob Agents Chemother* 2016; **60**: 507–14.

36 Piewngam P, Quiñones M, Thirakittiwatthana W, Yungyuen T, Otto M, Kiratisin P.
Composition of the intestinal microbiota in extended-spectrum β-lactamase-producing
Enterobacteriaceae carriers and non-carriers in Thailand. *Int J Antimicrob Agents* 2019; **53**:
435–41.

37 Milani C, Duranti S, Bottacini F, *et al.* The First Microbial Colonizers of the Human Gut:
Composition, Activities, and Health Implications of the Infant Gut Microbiota. *Microbiology
and Molecular Biology Reviews* 2017; **81**: 10.1128/mmbr.00036-17.

38 Deering KE, Devine A, O'Sullivan TA, Lo J, Boyce MC, Christophersen CT. Characterizing
the Composition of the Pediatric Gut Microbiome: A Systematic Review. *Nutrients* 2020; **12**:
16.

39 Shin N-R, Whon TW, Bae J-W. Proteobacteria: microbial signature of dysbiosis in gut
microbiota. *Trends Biotechnol* 2015; **33**: 496–503.

40 Rizzatti G, Lopetuso LR, Gibiino G, Binda C, Gasbarrini A. Proteobacteria: A Common Factor
in Human Diseases. *Biomed Res Int* 2017; **2017**: 9351507.

41 Kim K-S, Rowlinson M-C, Bennion R, *et al.* Characterization of Slackia exigua isolated from
human wound infections, including abscesses of intestinal origin. *J Clin Microbiol* 2010; **48**:
1070–5.

42 Gopalakrishna KP, Hand TW. Influence of Maternal Milk on the Neonatal Intestinal
Microbiome. *Nutrients* 2020; **12**: 823.

43 Majeed H, Gillor O, Kerr B, Riley MA. Competitive interactions in Escherichia coli populations:
the role of bacteriocins. *ISME J* 2011; **5**: 71–81.

44 van Duijkeren E, Wielders CCH, Dierikx CM, *et al.* Long-term Carriage of Extended-Spectrum
β-Lactamase–Producing Escherichia coli and Klebsiella pneumoniae in the General
Population in The Netherlands. *Clinical Infectious Diseases* 2018; **66**: 1368–76.

45 Armand-Lefèvre L, Rondinaud E, Desvillechabrol D, *et al.* Dynamics of extended-spectrum
beta-lactamase-producing Enterobacterales colonization in long-term carriers following travel
abroad. *Microbial Genomics* 2021; **7**: 000576.

46 Hussein RA, AL-Kubaisy SH, Al-Ouqaili MTS. The influence of efflux pump, outer membrane
permeability and β-lactamase production on the resistance profile of multi, extensively and
pandrug resistant *Klebsiella pneumoniae*. *Journal of Infection and Public Health* 2024; **17**:
102544.

47 Antimicrobial Resistance Collaborators. Global burden of bacterial antimicrobial resistance in
2019: a systematic analysis. *Lancet* 2022; **399**: 629–55.

48 van den Bunt G, Fluit AC, Bootsma MCJ, *et al.* Dynamics of Intestinal Carriage of Extended-
Spectrum Beta-lactamase–Producing Enterobacteriaceae in the Dutch General Population,
2014–2016. *Clinical Infectious Diseases* 2020; **71**: 1847–55.

Table 1. Sociodemographic characteristics and early-life exposures among 112 Peruvian children who were rarely or frequently colonized with ESBL-E during the first 16 months of life.

	Frequently colonized (n=100)	Rarely colonized (n=12)	*p*-val
Male (n, %)	48 (48.0)	7 (58.3)	0.711
Birth mode (n, %)			0.234
Vaginal	68 (68.0)	10 (83.3)	
Cesarean section	31 (31.0)	1 (8.3)	
Missing	1 (1.0)	1 (8.3)	
Mother completed high school (n, %)	69 (69.0)	5 (41.7)	0.117
Water Source (n,%)			0.063
Other	5 (5.0)	0 (0.0)	
Public Tap	15 (15.0)	5 (41.7)	
Indoor water connection (24 hours)	80 (80.0)	7 (58.3)	
Toilet (n,%)			0.051
Other	2 (2.0)	0 (0.0)	
Pit latrine	14 (14.0)	5 (41.7)	
Indoor flush toilet	84 (84.0)	7 (58.3)	
Home has improved flooring[a]	97 (97.0)	11 (91.7)	0.906
Lives in household with more than six household members (n,%)	40 (40.0)	2 (16.7)	0.207
Received breastmilk 90% of days in first 3 months (n,%)	63 (63.0)	8 (66.7)	1
Received formula 50% of days in first 3 months (n,%)	27 (27.0)	1 (8.3)	0.29
Household owned chickens (n,%)	17 (17.0)	1 (8.3)	0.721
Total number of antibiotic courses per child-year (mean (SD))	4.45 (2.86)	3.36 (2.29)	0.207
Number of azithromycin courses per child-year (mean (SD))	0.47 (0.95)	0.06 (0.22)	0.144
Number of sulfa-trimethoprim courses per child-year (mean (SD))	1.03 (1.15)	0.98 (1.09)	0.888
Number of amoxicillin courses per child-year (mean (SD))	1.59 (1.41)	1.93 (2.36)	0.471
Number of cefalexin courses per child-year (mean (SD))	0.53 (0.90)	0.32 (0.61)	0.425
Age in months at time of first antibiotic use (mean (SD))	5.67 (4.49)	7.23 (3.75)	0.251
Number of diarrhea episodes/100-child-month (mean (SD))	6.88 (5.09)	5.28 (2.59)	0.288

[a]Improved flooring included cement, wood, tiles, and "falso piso." Unimproved flooring included dirt and sand.

639 <u>Figures</u>

Figure 1. **Proportion of bacterial genera with at least 1% average relative abundance in**

stool samples collected from Peruvian children aged 3-16 months. There were no

differences in the relative abundance of any bacterial taxa between children who were rarely

versus frequently gut-colonized with ESBL-E across all timepoints. Taxonomy profiling was

performed using MetaPhlAn3.

Figure 2. **Relative abundance of dominant *Bifidobacterium spp.* in stool samples**

collected from Peruvian children aged 3-16 months. There were no differences in the

relative abundance of any *Bifidobacterium* spp. between children who were rarely versus

frequently gut-colonized with ESBL-E across all timepoints; the number of unique

Bifidobacterium spp. also did not differ between groups. Taxonomy profiling was performed

using MetaPhlAn3.

Note: Fifteen *Bifidobacterium spp.* were detected among all children; species with at least 1%

average relative abundance are depicted here.

Figure 3. **Distribution of *E. coli*, *Klebsiella* spp, and *Enterobacter* spp. strains in stool samples from Peruvian children aged 3-16 months who were frequently vs rarely gut-colonized with ESBL-E.** Strains were identified using the StrainGE toolkit. Pooling all timepoints together, there were significantly fewer *Klebsiella* spp. and *Enterobacter* spp. strains detected among rarely versus frequently colonized children by Welch two-sample t-test.

Note: * indicates statistically significant pairwise difference in mean number of strains by Welch two-sample t-test.

670
Figure 4. **Abundance of A) total antibiotic resistance genes and B) *bla*_{CTX-M} resistance**

genes in Peruvian children's stool from 3-16 months of age. The abundance of total ARGs

and *bla*_{CTX-M} resistance genes were significantly higher when children were gut-colonized with

ESBL-producing Enterobacterales versus not, on average and across all timepoints.

Note: * denotes a statistically significant difference by Welch two sample t-test.

FPKM=Fragments per kilobase per million mapped reads. FPKM values have been log

transformed.

678