A Flexible Framework for Local-Level Estimation of the Effective Reproductive Number in Geographic Regions with Sparse Data

Md Sakhawat Hossain,[1,2*] Ravi Goyal,[3] Natasha K Martin,[3] Victor DeGruttola,[4] Mohammad Mihrab Chowdhury,[1,2] Christopher McMahan,[2,5†] Lior Rennert,[1,2*†]

1 Department of Public Health Sciences, Clemson University, Clemson, SC, USA

2 Center for Public Health Modeling and Response, Clemson University, Clemson, SC, USA

3 Division of Infectious Diseases & Global Public Health, University of California San Diego, La Jolla, CA, USA

4 Division of Biostatistics, Herbert Wertheim School of Public Health and Longevity Science, University of California San Diego, San Diego, California, USA

5 School of Mathematics and Statistics, Clemson University, Clemson, SC, USA

† Senior Authors

* liorr@clemson.edu, mdsakhh@clemson.edu

* Correspondence
Dr. Lior Rennert, Center for Public Health Modeling and Response, Department of Public Health Sciences, Clemson University, Clemson, SC 29634, USA
Email: liorr@clemson.edu

Dr. Md Sakhawat Hossain, Center for Public Health Modeling and Response, Department of Public Health Sciences, Clemson University, Clemson, SC 29634, USA
Email: mdsakhh@clemson.edu

Abstract

Background: Our research focuses on local level estimation of the effective reproductive number, which describes the transmissibility of an infectious disease and represents the average number of individuals one infectious person infects at a given time. The ability to accurately estimate the infectious disease reproductive number in geographically granular regions is critical for disaster planning and resource allocation. However, not all regions have sufficient infectious disease outcome data for estimation.

Methods: We propose a two-step approach that incorporates existing R_t estimation procedures (EpiEstim, EpiFilter, and EpiNow2) using data from geographic regions with sufficient data (step 1) into a covariate-adjusted Bayesian Integrated Nested Laplace Approximation (INLA) spatial model to predict R_t in regions with sparse or missing data (step 2). Our flexible framework effectively allows for implementing any existing estimation procedure for R_t in regions with coarse or entirely missing data. We perform external validation to evaluate predictive performance.

Results: We applied our method to estimate R_t using data from South Carolina (SC) counties and ZIP codes during the first COVID-19 wave ('Wave 1', June 16, 2020 – August 31, 2020) and the second wave ('Wave 2', December 16, 2020 – March 02, 2021). Among the three methods used in the first step, EpiNow2 yielded the highest accuracy of R_t prediction in the regions with entirely missing data. Median county-level percentage agreement (PA) was 90.9% (IQR: 89.9-92.0%) and 92.5% (IQR: 91.6-93.4%) for Wave 1 and 2, respectively. Median zip code-level PA was 95.2% (Interquartile Range, IQR: 94.4-95.7%) and 96.5% (IQR: 95.8-97.1%) for Wave 1 and 2, respectively. Using EpiEstim and EpiFilter yielded median PA ranging from 81.9%-90.0% and 87.2%-92.1% (respectively) across both waves and geographic granularities.

Conclusion: These findings demonstrate that the proposed methodology is a useful tool for small-area estimation of R_t, as our flexible framework yields high prediction accuracy for regions with entirely missing data regardless of the (step 1) estimation procedure used.

Keywords: Infectious Disease Modeling; Bayesian Statistics; Effective Reproductive Number; Small Area Estimation; Prediction

Introduction

Infectious disease outbreaks pose significant threats, impacting health, operations, and the economy. Effective outbreak management relies on the ability to estimate the intensity of local transmission and strategize appropriately, which is vital for controlling disease transmission, enhancing health outcomes, and reducing economic impacts. However, current approaches often target broad geographic areas, which may not yield actionable insights for local-level or institutional decision-making [1]. This lack of granularity can be attributed to variations in community characteristics such as age distribution, prevalence of comorbidities, and social behaviors [2,3], all of which influence disease transmission and susceptibility to severe outcomes. Moreover, policies derived from coarse data may lead to suboptimal local disease control [4,5]. Therefore, enhancing local-level transmission estimation and forecasting is essential for tailoring policies and optimizing interventions.

The effective reproductive number is a key epidemiological metric used to describe the transmissibility of infectious diseases for a specific time and setting; it is critical to understand the intensity of transmission. R_t represents the average number of people infected by one infectious individual at a given time t. An R_t value greater than 1 signifies that the disease is spreading, while an R_t less than 1 indicates that the transmission is declining [6,7]. R_t changes with the progression of an epidemic and the implementation of control measures [8–10]. It also varies spatially due to the differences in population density, mobility, social interactions, and public health policies [11,12]. Unfortunately, local-level estimation of R_t is challenging due to data sparsity or unavailability of infectious disease testing or encounter records. For example, insufficient testing and changes in case definitions or reporting can substantially impact estimates of R_t [13,14]. Due

to substantial lags in producing estimates of disease epidemiological metrics from traditional surveillance systems [15], electronic health records (EHR) obtained directly from health systems or hospitals are useful for real-time estimation of such metrics, including R_t. However, healthcare systems do not serve all populations, which inhibits the ability to forecast disease spread in communities that are not represented in the healthcare system. Additionally, estimates of R_t may be unstable for communities that are underrepresented in the healthcare system and hence offer few data points for accurate estimation.

There are several techniques for estimating the effective reproductive number; e.g., Wallinga and Tenius method [16], Cori et al. method [17], EpiFilter [6], and EpiNow2 [18]. EpiEstim is a tool integrated into widely used software platforms for estimation of the reproductive number from incidence time series, accounting for uncertainties in the serial interval distribution [17]. EpiFilter utilizes all available incidence information and does not rely on window size assumptions, making it more statistically robust, particularly during periods of low incidence [6]. The modeling approach of EpiNow2, an R software package [18], is based on the methods developed by Cori et al. and Thompson et al. [19], and it incorporates the delays from infection to reporting to estimate the effective reproductive number utilizing a range of open-source tools [20]. Although there are several methods for estimating the effective reproductive number, there is a notable gap in accurately estimating R_t for small areas with coarse or missing data. To address this gap, we develop a flexible framework that effectively allows for implementing any estimation procedure for R_t (e.g., EpiEstim, EpiFilter, EpiNow2, etc.) in regions with coarse or entirely missing data.

The motivation for this study, funded by the National Library of Medicine of the National Institutes of Health (NIH) and the Center for Forecasting and Outbreak analytics of the Centers for Disease Control and Prevention (CDC), is to inform allocation of mobile health clinics for infectious

disease interventions to medically underserved communities at high risk of infectious disease outbreaks. We address the challenge of local level R_t estimation in areas with coarse or entirely missing data by proposing a two-step estimation procedure which incorporates existing R_t estimation procedures using data from geographic regions with sufficient data (step 1) into a spatial modeling framework to predict R_t in regions with sparse or missing data (step 2). Our methodology is described in Section 2. In Section 3, we apply our method and present the results of effective reproductive number estimation of COVID-19 in South Carolina (SC) ZIP codes and counties. Our findings are discussed in Section 4.

Materials and Methods

To estimate the region-specific effective reproductive number (R_t) for an infectious disease in small areas with coarse or entirely missing data, we develop a two-step procedure. In the first step of our analyses, we estimate $R_{t,i}$, using the existing techniques EpiEstim, EpiFilter, and EpiNow2 for each geographic region i [6,17,18]. We denote the resulting estimates by $\hat{R}_{t,Estim,i}$, $\hat{R}_{t,Filter,i}$, and $\hat{R}_{t,Now,i}$, respectively. In the second step, we employ a Bayesian Integrated Laplace Approximation (INLA) model incorporating spatial and sociodemographic information. For each geographic region i with available outcome data (e.g., Covid-19 cases), we denote these estimates by $\hat{R}_{t,Estim,i}^{S}$, $\hat{R}_{t,Filter,i}^{S}$, $\hat{R}_{t,Now,i}^{S}$, which are referred to as spatially and covariates smoothed estimates. When geographic region i is not included in the spatial (and covariate-adjusted) INLA model fitting in step 2 (e.g., geographic region i has entirely missing data), we denote the prediction in this region by $\hat{R}_{t,Estim,i}^{S^{(i)}}$, $\hat{R}_{t,Filter,i}^{S^{(i)}}$, $\hat{R}_{t,Now,i}^{S^{(i)}}$, where $S^{(i)}$ indicates information from region i is not used in the model fitting in step 2. More details of initial estimation, smoothing and predictions of R_t in

regions with entirely missing data, along with their evaluations, are provided in the sections 2.2, 2.3, and 2.4 respectively.

Data Description

The number of cumulative confirmed COVID-19 cases for US counties is publicly available on the New York Times (NYT) GitHub repository [21]. The daily reported cases are calculated from the county-level cumulative data. We obtain the ZIP code level daily COVID-19 case data from Prisma Health's (SC's largest not-for-profit health care provider) COVID-19 registry. We utilize data from SC counties and ZIP codes during the first COVID-19 wave (between June 16, 2020 – August 31, 2020) and the second wave (between December 16, 2020 – March 02, 2021). Since many ZIP codes in the Prisma Health system data lack sufficient case numbers, we restrict our analysis to those ZIP codes with at least 50 cases in each period to ensure the reliability of our results (Wave 1: N = 30 ZIP codes; Wave 2: N = 45 ZIP codes). Population demographics are characterized by age, sex, race/ethnicity, employment, and insurance coverage. These data are sourced from the United States Census Bureau website [22]. The social vulnerability index (SVI) data is obtained from the Agency for Toxic Substances and Disease Registry website [23].

Initial Estimates of R_t

We utilize the techniques EpiEstim, EpiFilter, and EpiNow2 to obtain the initial estimates of the effective reproductive number. EpiEstim and EpiFilter assume the incidence I_t at time t is Poisson distributed with mean $R_t \sum_{s=1}^{t} I_{t-s} w_s$, where R_t is the effective reproductive number at time t, s is the time since infection of the case, w_s is the probability that a primary case takes time between $s - 1$ and s days to generate the secondary infection, and w_s is obtained from the serial interval distribution [6,17]. EpiEstim uses an expectation-maximization algorithm to reconstruct daily

incidence from aggregated data for estimating R_t [24]. It computes estimates of the effective reproductive number, $\hat{R}_{t,Estim,i}$, using a sliding time window; the default seven days window setting is used. EpiFilter employs a recursive Bayesian smoothing technique to estimate the effective reproductive number, $\hat{R}_{t,Filter,i}$. It unifies and extends two popular techniques EpiEstim and the Wallinga-Teunis method [6]. EpiNow2 employs a Bayesian latent variable model that incorporates a probabilistic programming language, Stan [20,25]. Detailed description of the EpiEstim, EpiFilter, and EpiNow2 methods are available in the literature [6,17,18,20]. Each of the three methods estimates R_t using different parameter or data assumptions. All three methods depend on the generation time distribution. Since infection times are often unobserved, the generation time is usually approximated using the serial interval distribution [17]. In all three methods, the parameters for both the serial interval distribution and incubation period are utilized.

From the literature, we obtain a mean of 5 days (95% CI, 4.94-5.06) and a standard deviation (SD) of 2.4 for the incubation period [26], a mean of 4.7 days and a SD of 2.9 for the serial interval distribution [27]. The 'estimate_R' function of the EpiEstim, R software package version 2.2-4 is used to obtain $\hat{R}_{t,Estim,i}$. We calculate the estimate, $\hat{R}_{t,Filter,i}$, by using the R software function 'epiSmoother' available at https://github.com/kpzoo/EpiFilter. The 'epinow' function from the R software package EpiNow2, version 1.4.0, is utilized to get $\hat{R}_{t,Now,i}$.

Proposed Covariate-Adjusted Spatial Smoothing via INLA

We propose a sociodemographic covariate adjusted, two-step Integrated Nested Laplace Approximation (INLA) model to refine the initial estimation of the effective reproductive number at the local level and to provide estimates for the regions with coarse or missing data. The use of INLA for spatial data prediction is supported by extensive research, demonstrating its effectiveness

in handling spatial dependency and missing data challenges[28,32]. INLA method is designed to derive the posterior marginal distribution for the latent Gaussian models, serves as a computationally efficient alternative to traditional Markov Chain Monte Carlo (MCMC) methods for Bayesian inference [28,29]. INLA utilizes a series of nested Laplace approximations to calculate the marginal posterior and hyperparameters of the latent Gaussian models. The INLA framework is employed to integrate the spatial dependencies of the effective reproductive number estimates into our model [30]. To fit the INLA model, we use the R software package called "R-INLA" [28,31,32]. We incorporate the effect of the spatial location (counties or ZIP codes) in the INLA function through Besag's model [33,34]. The Besag model employs spatial random effects for each area to capture the correlation among neighboring areas, adjusting the estimated R_t based on the estimates of the neighboring areas. Neighborhood structures are defined using the 'poly2nb()' function, identifying areas as neighbors if their boundary polygons share at least one vertex. Then we convert the neighborhood structure into a binary adjacency matrix using the function 'nb2mat()'. Each element of the adjacency matrix represents whether two locations are neighbors (1 if they share a boundary, 0 if they do not).

The model equation in the INLA is

$$\hat{R}_{t,m,i} = \beta_0 + \beta X_i + u_i + \varepsilon_i$$

where β_0 is the intercept of the model, β is coefficients vector for sociodemographic community-level covariates X_i, $\hat{R}_{t,m,i}$ corresponds to the estimates $\hat{R}_{t,Estim,i}$, $\hat{R}_{t,Filter,i}$, or $\hat{R}_{t,Now,i}$ at area i, the term u_i specifies the spatial random effect for location i, which depends on the neighboring location defined by the adjacency matrix, and the random Gaussian error term is ε_i.

The sociodemographic covariates used for the analyses at the county and ZIP code levels are percentage of the population for different age groups "Age 0-19", "Age 20-44", "Age 45-64", "Age 65 and over", percentage of the population for different races and ethnicity "White", "Hispanic", "Black or African American" and "others", percentage of the population "Employed" and "Unemployed", percentage of "Health Insurance Coverage", percentage of the population of "Male" and "Female", Social Vulnerability Index (SVI) and household median income. Household median income is normalized from 0 to 1, and SVI values range from 0 to 1, where higher values indicate greater social vulnerability. We consider the covariates "Age 0-19", "White", "Unemployed", "Uninsured", and "Female" population groups as the reference categories.

Assessment of Prediction of R_t for Regions with Entirely Missing Data

We employ the two-step spatial INLA model using the initial estimate, $\hat{R}_{t,m,i}$, for each method m (i.e., EpiEstim, EpiFilter, and EpiNow2) at location i, as the response variable. We validate the model using two approaches. In the first approach, we randomly select 90% of the regions to hold out for the training set and 10% for the test set. We then apply the proposed two-step spatial (covariates-adjusted) INLA model to the training set and use this model to predict the effective reproductive numbers for the geographic regions in the test set, which we denote by $\hat{R}_{t,m,i}^{S^{(i)}}$. We then compare the agreement between $\hat{R}_{t,m,i}$, and $\hat{R}_{t,m,i}^{S^{(i)}}$ for the geographic regions which were left out from the model fitting in the second step. The prediction accuracy measurement metrics are calculated using the formulas in equations 1-3. This process is repeated 200 times.

In the second validation approach, we hold out geographic regions with sufficient data for the test set – specifically, Charleston, Greenville, Horry, and Richland counties and ZIP codes "29605", "29642", "29680", and "29681". Since there is no gold standard for comparing R_t estimates, we

selected the above regions to predict R_t for test cases with entirely missing data, assuming the initial estimates from regions with sufficient data will serve as a baseline for comparison. The prediction in this approach is performed by leaving out one area at a time.

The prediction accuracy measurement metrics are calculated using the following formulas:

$$\text{Mean Absolute Error, } MAE = \frac{1}{T}\sum_{t=1}^{T}|\hat{R}_{t,m,i} - \hat{R}_{t,m,i}^{S^{(i)}}| \tag{1}$$

$$\text{Root Mean Squared Error, } RMSE = \sqrt{\frac{1}{T}\sum_{t=1}^{T}\left(\hat{R}_{t,m,i} - \hat{R}_{t,m,i}^{S^{(i)}}\right)^2} \tag{2}$$

$$\text{Percentage Agreement, } PA = \frac{1}{T}\sum_{t=1}^{T}\frac{\min\left(\hat{R}_{t,m,i}, \ \hat{R}_{t,m,i}^{S^{(i)}}\right)}{\max\left(\hat{R}_{t,m,i}, \ \hat{R}_{t,m,i}^{S^{(i)}}\right)} \tag{3}$$

where, $\hat{R}_{t,m,i}$ is the initial estimates, $\hat{R}_{t,m,i}^{S^{(i)}}$ is the prediction of effective reproductive number for each method m, at time t and location i, and T is the total number of time points (days).

Results

Initial Estimates and Spatial (Covariates-Adjusted) INLA Smoothing of R_t

For the R_t estimation in this section, we utilized the county and ZIP code level COVID-19 daily case data for the first wave (June 16, 2020, through August 31, 2020) and second wave (December 16, 2020, through March 02, 2021) of the disease in SC. **Fig 1** displays the estimates of effective reproductive number for Charleston, Greenville, Horry, and Richland counties during the second COVID-19 wave in SC. In general, proposed two-step spatial INLA estimates, $\hat{R}_{t,Estim,i}^{S}$, $\hat{R}_{t,Filter,i}^{S}$, $\hat{R}_{t,Now,i}^{S}$, demonstrated reduced peaks compared to the initial estimates, $\hat{R}_{t,Estim,i}$, $\hat{R}_{t,Filter,i}$, $\hat{R}_{t,Now,i}$.

Fig.1. Comparison of two-step spatial (covariates-adjusted) INLA smoothing with initial R_t estimates at the county level in Wave 2.

$\hat{R}^S_{t,m,i}$ (blue line) represents the two-step spatial (covariates-adjusted) INLA smoothed estimates, $\hat{R}^S_{t,Estim,i}$, $\hat{R}^S_{t,Filter,i}$, and $\hat{R}^S_{t,Now,i}$, while $\hat{R}_{t,m,i}$ (red line) represents the initial estimates, $\hat{R}_{t,Estim,i}$, $\hat{R}_{t,Filter,i}$, and $\hat{R}_{t,Now,i}$, for select Charleston, Greenville, Horry, and Richland counties during COVID 19 Wave 2 (between December 16, 2020 – March 02, 2021) in SC. The state level initial estimate of R_t is presented with dashed gray line (SC State). The plots in the right most panel present the average daily cases for the respective counties over the same period.

S1-S3 Figs in the supplementary materials (Supporting Information 1) provide a detailed

comparison of the effective reproductive number estimates, $\hat{R}_{t,Estim,i}$, $\hat{R}_{t,Filter,i}$, $\hat{R}_{t,Now,i}$ and

$\hat{R}^S_{t,Estim,i}$, $\hat{R}^S_{t,Filter,i}$, $\hat{R}^S_{t,Now,i}$ across various counties and ZIP codes. Results are presented for

Charleston, Greenville, Horry, and Richland counties and ZIP codes "29605", "29642", "29680",

and "29681", as these results serve as benchmarks when evaluating the prediction accuracy (Section 3.2). The bar graphs displaying daily COVID-19 cases revealed spikes and declines that correspond to the fluctuations in the R_t estimates. For instance, peaks in case counts align with the increases in both sets of estimates, $\hat{R}_{t,Estim,i}$, $\hat{R}_{t,Filter,i}$, $\hat{R}_{t,Now,i}$ and $\hat{R}^S_{t,Estim,i}$, $\hat{R}^S_{t,Filter,i}$, $\hat{R}^S_{t,Now,i}$. The figures show that when daily cases increase over a period, corresponding R_t estimates exceed 1.0, indicating infection growth. For example, in Fig.1, daily cases for "Charleston", "Greenville", and "Horry" counties increase between time points 10 and 25, with R_t estimates staying above 1.0. After time points 25, as cases decline almost everywhere, R_t estimates fall to 1.0 or lower, particularly with our two-step spatial INLA method (blue lines).

Prediction of R_t in Regions with Entirely Missing Data

In this section, we evaluated the accuracy of our two-step spatial INLA model in estimating $\hat{R}^{S,I}_{t,Estim}$, $\hat{R}^{S,I}_{t,Filter}$, $\hat{R}^{S,I}_{t,Now}$ for regions with entirely missing data. We utilized daily COVID-19 case data at both county and ZIP code levels for the first wave (June 16, 2020, through August 31, 2020) and second wave (December 16, 2020, through March 02, 2021) of the pandemic in SC. **Table 1** presents the prediction accuracy achieved through the two-step spatial INLA model through comparison with the one-step estimates $\hat{R}_{t,Estim,i}$, $\hat{R}_{t,Filter,i}$, and $\hat{R}_{t,Now,i}$. That is, we used the two-step spatial INLA model based on a subset of geographic regions (training set) to predict estimates of R_t for the remaining geographic region (i.e., validation set). Since there is no gold standard for comparison, we compared the predicted effective reproductive number, $\hat{R}^{S(i)}_{t,m,i}$, to the initial estimate $\hat{R}_{t,m,i}$.

At both county and ZIP code levels during the second wave of COVID-19, predicted R_t values, $\hat{R}^{S(i)}_{t,m,i}$, based on the two-step spatial INLA estimators achieved high level of accuracy (**Table 1**).

For counties, median percentage agreement was 90.0% (IQR: 89.0–90.9 %) for $\hat{R}_{t,Estim,i}^{S(i)}$, 87.2% (IQR: 86.3–88.1 %) for $\hat{R}_{t,Filter,i}^{S(i)}$, and 92.5% (IQR: 91.6–93.4 %) for $\hat{R}_{t,Now,i}^{S(i)}$. The prediction, $\hat{R}_{t,Now,i}^{S(i)}$, achieved the highest accuracy, with median PA of 92.5% at the county level. Similarly, $\hat{R}_{t,Now,i}^{S(i)}$ exhibited the highest median PA of 96.5% for SC ZIP codes. In the first wave, the accuracy was also high, particularly for $\hat{R}_{t,Now,i}^{S(i)}$, which obtained a median PA of 90.9% at the county level and 95.2% at the ZIP code level (**see Table S1**). However, $\hat{R}_{t,Estim,i}^{S(i)}$ showed the lowest accuracy, with a median PA of 81.9% at the ZIP code level, and a median PA of 85.7% at the county level. Overall, the two-step spatial (covariates-adjusted) INLA procedure showed high performance accuracy in computing $\hat{R}_{t,m,i}^{S(i)}$ for the areas with missing data when further evaluated through root mean squared error (RMSE) and mean absolute percentage error (MAPE).

Table 1. Comparison of two-step spatial (covariates-adjusted) INLA prediction, $\hat{R}_{t,m,i}^{S(i)}$, with initial estimates, $\hat{R}_{t,m,i}$, in Wave 2.

Comparison	SC County and ZIP Code Level R_t Prediction Accuracy Measurements						
	Statistic	County Level			ZIP Code Level		
		RMSE	MAPE	PA (%)	RMSE	MAPE	PA (%)
$\hat{R}_{t,Estim,i}^{S(i)}$ vs. $\hat{R}_{t,Estim,i}$	Median	0.142	0.110	90.0	0.284	0.240	83.2
	IQR	(0.13, 0.16)	(0.10, 0.13)	(89.0, 90.9)	(0.24, 0.35)	(0.18, 0.23)	(80.8, 85.1)
$\hat{R}_{t,Filter,i}^{S(i)}$ vs. $\hat{R}_{t,Filter,i}$	Median	0.187	0.147	87.2	0.110	0.086	92.1
	IQR	(0.17, 0.21)	(0.13, 0.16)	(86.3, 88.1)	(0.10, 0.12)	(0.08, 0.10)	(91.2, 92.8)
$\hat{R}_{t,Now,i}^{S(i)}$ vs. $\hat{R}_{t,Now,i}$	Median	0.104	0.082	92.5	0.044	0.036	96.5
	IQR	(0.09, 0.12)	(0.07, 0.09)	(91.6, 93.4)	(0.04, 0.06)	(0.03, 0.04)	(95.8, 97.1)

For example, the first row compares the accuracy of the predicted R_t values, $\hat{R}_{t,Estim,i}^{S^{(i)}}$, using proposed two-step spatial (covariates-adjusted) INLA model, where $S^{(i)}$ indicates information from region i is not used in the model fitting in step 2, of the counties/ZIP codes from the test set to their initial estimates, $\hat{R}_{t,Estim,i}$. Accuracy measurements were calculated, using formulas in equations 1 – 3, during CVOID 19 Wave 2 ((between December 16, 2020 – March 02, 2021) in SC. In this analysis, 90% of the areas were used as the training set, with the remaining 10% as the test set. The spatial (covariate-adjusted) INLA model was fitted to the training data, and the predictions were done for the test data set. We repeated the process 200 times and calculated the accuracy measurement metrics. The table presents the median accuracy along with the interquartile range (IQR).

In the second validation approach, we selected Charleston, Greenville, Horry, and Richland counties (county level analysis), and ZIP codes "29605", "29642", "29680", and "29681" (zip code analysis), as the test set. **Fig 2** provides a detailed comparison of the prediction $\hat{R}_{t,Estim,i}^{S^{(i)}}$, $\hat{R}_{t,Filter,i}^{S^{(i)}}$, $\hat{R}_{t,Now,i}^{S^{(i)}}$ to initial estimates $\hat{R}_{t,Estim,i}$, $\hat{R}_{t,Filter,i}$, $\hat{R}_{t,Now,i}$ across counties–Charleston, Greenville, Horry, and Richland and ZIP codes – "29605", "29642", "29680", and "29681" (also see **S4-S8 Figs**). The results comparing prediction accuracy for this validation setting are presented in **Table S2** for the first wave and **Table S3** for the second wave. During the second wave, $\hat{R}_{t,Now,i}^{S^{(i)}}$ demonstrated the highest accuracy, with a percentage agreement (PA) range of 91.1% - 94.0% across the four counties and 96.1% – 98.7 % across the four ZIP codes. $\hat{R}_{t,Filter,i}^{S^{(i)}}$ achieved the lowest PA range for counties (86.7% – 90.8%) and $\hat{R}_{t,Estim,i}^{S^{(i)}}$ achieved the lowest PA range for ZIP codes (86.9%–91.5 %). Similar results were obtained during the first wave.

Fig.2. Comparison of two-step spatial (covariates-adjusted) INLA prediction with the initial R_t estimates in Wave 2.

$\hat{R}^{S(i)}_{t,m,i}$ (blue line) represents the two-step spatial (covariates-adjusted) INLA prediction of effective reproductive number, $\hat{R}^{S(i)}_{t,Estim,i}$, $\hat{R}^{S(i)}_{t,Filter,i}$, $\hat{R}^{S(i)}_{t,Now,i}$, while $\hat{R}_{t,m,i}$ (red line) represents the initial estimates $\hat{R}_{t,Estim,i}$, $\hat{R}_{t,Filter,i}$, $\hat{R}_{t,Now,i}$, for select Charleston, Greenville, Horry, and Richland counties during COVID 19 Wave 2 (between December 16, 2020 – March 02, 2021) in SC. Here, $S^{(i)}$ *in* $\hat{R}^{S(i)}_{t,Estim,i}$ indicates that geographic region i was not included in the INLA model fitting. For example, if i=*Greenville (GVL)*, $\hat{R}^{S(GVL)}_{t,Estim,GVL}$ means Greenville County was not used in the spatial (and covariates-adjusted) INLA model fitting in step 2. The plots in the right most panel present the average daily cases for the respective counties over the same period.

Discussion

Estimation of time varying effective reproductive number R_t is necessary for understanding the transmission dynamics of infectious disease and implementing the necessary public health interventions. We developed a two-step procedure for estimating the effective reproductive number

for infectious diseases in granular geographic regions. Our approach incorporates existing R_t estimation procedures (EpiEstim, EpiFilter, and EpiNow2) using data from geographic regions with sufficient data (step 1) into a spatial modeling framework to predict R_t in regions with sparse or missing data (step 2). The choice of R_t estimation methods is not exhaustive. Our flexible framework allows for implementing any existing estimation procedure for R_t in regions with coarse or entirely missing data.

Our proposed methodology improves the estimation of the effective reproductive number R_t in small areas, especially in areas where the disease data are coarse or missing. By integrating the Bayesian spatial INLA model, this method efficiently utilizes the available data and borrows information from neighboring areas and from similar areas in SC through sociodemographic covariates, enhancing the robustness of R_t estimates. Decisions regarding the implementation, removal, or adjustment of infection control measures will be guided by a combination of epidemiological, social, and economic factors [35,36]. Effective reproductive number estimates will assist in this process by providing critical information about the potential future course of an outbreak. Through small area estimation of the effective reproductive number, we can better identify and prioritize high risk areas, which allows for optimal resource allocation [37]. For example, this current NIH and CDC-funded study is intended to better identify and prioritize medically underserved communities at high-risk of infectious disease outbreaks for priority delivery of mobile health clinics for infectious disease screening, treatment, and vaccination.

Our study has several limitations. First, our approach is focused on retroactive estimation of R_t. This is useful for forecasting outbreaks when the effective reproductive number in each region is similar between outbreaks (e.g., first two waves of Covid-19 in SC; Table S4). While this can be a reasonable assumption for endemic viruses, this is far from guaranteed for emerging viruses

where disease transmission dynamics and/or human behavior changes can substantially change with the introduction of a new variant. Therefore, a natural extension is to extend our approach for forecasting future values of R_t, particularly in granular geographic regions with sparse data.

Another limitation is that we use existing R_t estimators as a benchmark to evaluate the performance of our proposed estimator. Each method for R_t estimation has its own limitations, and there is no gold standard for comparison. Second, we evaluated model performance using three commonly used techniques (EpiEstim, EpiFilter, EpiNow2). Similar to most R_t estimation methods, EpiEstim and EpiFilter rely on the reported incidence data [6,17]. Inaccuracies or delays in reporting can lead to biased estimates. EpiNow2 makes certain assumptions about the underlying transmission dynamics, such as the generation time distribution and reporting delays. These assumptions can introduce uncertainty into the estimates, particularly during the periods of low incidence [18]. Future research may evaluate performance of other commonly used estimation procedures, including Epidemia [38], EpiInvert [39], EpiRegress [40], estimateR [41] and the Extended Kalman Filter [42].

Conclusion

This study presented a two-step spatial INLA approach to estimate the effective reproductive number in small areas with coarse or missing data. Our approach incorporated existing estimators into a Bayesian spatial model that adjusted for spatial variation by borrowing information from neighboring regions and incorporating sociodemographic covariates. Our approach provided reliable predictions of the effective reproductive number for regions with missing data. Importantly, our proposed framework is flexible, allowing for implementation of any existing estimation procedure for R_t in geographically granular regions. While our study is focused on

Covid-19, our methodology is directly translatable for estimation of the effective reproductive number for other respiratory infectious diseases.

Supporting Information 1

S1 Fig. Comparison of two-step spatial (covariates-adjusted) INLA smoothing with initial R_t estimates at the county level in Wave 1.

S2 Fig. Comparison of two-step spatial (covariates-adjusted) INLA smoothing with initial R_t estimates at the ZIP code level in Wave 1.

S3 Fig. Comparison of two-step spatial (covariates-adjusted) INLA smoothing with initial R_t estimates at the ZIP code level in Wave 2.

S1 Table. Comparison of two-step spatial (covariates-adjusted) INLA prediction, $\hat{R}_{t,m,i}^{S^{(i)}}$, with initial estimates, $\hat{R}_{t,m,i}$, in Wave 1.

S2 Table. Comparison (leave-one-out validation) of two-step spatial (covariates-adjusted) INLA prediction, $\hat{R}_{t,m,i}^{S^{(i)}}$, with initial estimates, $\hat{R}_{t,m,i}$, in Wave 1.

S3 Table. Comparison (leave-one-out validation) of two-step spatial (covariates-adjusted) INLA prediction, $\hat{R}_{t,m,i}^{S^{(i)}}$, with initial estimates, $\hat{R}_{t,m,i}$, in Wave 2.

S4 Fig. Comparison of two-step spatial (covariates-adjusted) INLA prediction with the initial R_t estimates at the county level in Wave 1.

S5 Fig. Comparison of two-step spatial (covariates-adjusted) INLA prediction with the initial R_t estimates at the ZIP code level in Wave 1.

S6 Fig. Comparison of two-step spatial (covariates-adjusted) INLA prediction with the initial R_t estimates at the ZIP code level in Wave 2.

S7 Fig. Comparison of R_t estimates of the county Greenville and its neighboring counties in Wave 2.

S8 Fig. Comparison of R_t estimates of the county Greenville and its neighboring counties in Wave 2.

S4 Table. Comparison of initial estimates and two-step INLA spatial (covariates-adjusted) estimates between Wave 1 and Wave 2.

Abbreviations

INLA	Integrated Nested Laplace Approximation
SC	South Carolina
PA	Percentage Agreement
IQR	Interquartile Range
NIH	National Institutes of Health
CDC	Center for Disease Control and Prevention
SVI	Social Vulnerability Index
NYT	New York Times
SD	Standard Deviation
CI	Confidence Interval
RMSE	Root Mean Squared Error
MAPE	Mean Absolute Percentage Error
GVL	Greenville

Acknowledgements

Thanks to Tanvir Ahammed and Dr. Jiande Wu for their assistance with processing COVID-19 case data at the ZIP code and county levels in SC.

Author Contributions

Conceptualization: MSH, RG, NKM, VDG, CM, LR

Formal Analysis: MSH

Investigation: MSH, LR

Methodology: MSH, RV, NKM, VDG, CM, LR

Project administration: LR

Software: MSH

Supervision: LR, CM

Validation: MSH

Visualization: MSH

Writing-Original Draft Preparation: MSH, LR

Writing-Review & Editing: MSH, RG, NKM, VDG, MMC, CM, LR

Funding

MSH, RG, NKM, and VDG acknowledge support from the National Library of Medicine of the National Institutes of Health (NIH) under award number R01LM014193. LR and CM acknowledge support from the National Library of Medicine of the National Institutes of Health (NIH) under award number R01LM014193 and the Center for Forecasting and Outbreak Analytics of the Centers for Disease Control and Prevention (CDC) under award number NU38FT000011. MMC recognizes support from the Center for Forecasting and Outbreak Analytics of the Centers for Disease Control and Prevention (CDC) under award number NU38FT000011. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Data Availability

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

Ethical review for this study was obtained by the Institutional Review Board of Clemson University (#2020-0150). No consent was needed for this study; retrospective data is based on medical claims and electronic health records and were de-identified to study investigators.

Consent for publication

Not applicable.

Competing Interests

The authors declare that they have no competing interests.

References

1. Ma Z, Rennert L. An epidemiological modeling framework to inform institutional-level response to infectious disease outbreaks: a Covid-19 case study. Sci Rep. 2024 Mar 27;14(1):7221.

2. Gezer F, Howard KA, Litwin AH, Martin NK, Rennert L. Identification of Factors Associated with, and Prediction of, Opioid- and HCV-Related Hospitalizations at the Zip Code Level: A Statistical Modeling Approach. SSRN Revis Invit Lancet Public Health [Internet]. 2023 Jul 13 [cited 2023 Jul 13]; Available from: https://ssrn.com/abstract=4506562

3. Borquez A, Martin NK. Fatal overdose: Predicting to prevent. Int J Drug Policy. 2022 Jun;104:103677.

4. Er S, Yang S, Zhao T. COUnty aggRegation mixup AuGmEntation (COURAGE) COVID-19 prediction. Sci Rep. 2021 Jul 12;11(1):14262.

5. CDC. Science Brief: Indicators for Monitoring COVID-19 Community Levels and Making Public Health Recommendations [Internet]. Centers for Disease Control and Prevention. 2022 [cited 2023 Jan 25]. Available from: https://www.cdc.gov/coronavirus/2019-ncov/science/science-briefs/indicators-monitoring-community-levels.html

6. Parag KV. Improved estimation of time-varying reproduction numbers at low case incidence and between epidemic waves. PLOS Comput Biol. 2021 Sep 7;17(9):e1009347.

7.   Anderson RM, May RM. Infectious Diseases of Humans: Dynamics and Control. Oxford, New York: Oxford University Press; 1992. 766 p.

8.   Andrade J, Duggan J. Inferring the effective reproductive number from deterministic and semi-deterministic compartmental models using incidence and mobility data. PLOS Comput Biol. 2022 Jun 27;18(6):e1010206.

9.   Pan A, Liu L, Wang C, Guo H, Hao X, Wang Q, et al. Association of Public Health Interventions With the Epidemiology of the COVID-19 Outbreak in Wuhan, China. JAMA. 2020 May 19;323(19):1915–23.

10. Cauchemez S, Boëlle PY, Donnelly CA, Ferguson NM, Thomas G, Leung GM, et al. Real-time estimates in early detection of SARS. Emerg Infect Dis. 2006;12(1):110.

11. Achaiah NC, Subbarajasetty SB, Shetty RM. R0 and Re of COVID-19: Can We Predict When the Pandemic Outbreak will be Contained? Indian J Crit Care Med Peer-Rev Off Publ Indian Soc Crit Care Med. 2020 Nov;24(11):1125–7.

12. Gostic KM, McGough L, Baskerville EB, Abbott S, Joshi K, Tedijanto C, et al. Practical considerations for measuring the effective reproductive number, Rt. PLOS Comput Biol. 2020 Dec 10;16(12):e1008409.

13. Sanche S, Lin YT, Xu C, Romero-Severson E, Hengartner N, Ke R. High Contagiousness and Rapid Spread of Severe Acute Respiratory Syndrome Coronavirus 2 - Volume 26, Number 7—July 2020 - Emerging Infectious Diseases journal - CDC. 2020 [cited 2020 Jul 26]; Available from: https://wwwnc.cdc.gov/eid/article/26/7/20-0282_article

14. Linka K, Peirlinck M, Kuhl E. The reproduction number of COVID-19 and its correlation with public health interventions. Comput Mech. 2020;66(4):1035–50.

15. Poirier C, Lavenu A, Bertaud V, Campillo-Gimenez B, Chazard E, Cuggia M, et al. Real Time Influenza Monitoring Using Hospital Big Data in Combination with Machine Learning Methods: Comparison Study. JMIR Public Health Surveill. 2018 Dec 21;4(4):e11361.

16. Wallinga J, Teunis P. Different Epidemic Curves for Severe Acute Respiratory Syndrome Reveal Similar Impacts of Control Measures. Am J Epidemiol. 2004 Sep 15;160(6):509–16.

17. Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to estimate time-varying reproduction numbers during epidemics. Am J Epidemiol. 2013;178(9):1505–12.

18. Abbott S, Hellewell J, Sherratt K, Gostic K, Hickson J, Badr HS, et al. EpiNow2: estimate real-time case counts and time-varying epidemiological parameters. R Package Version 01 0 [Internet]. 2020 [cited 2024 Feb 13]; Available from: https://scholar.google.com/scholar?cluster=1308097242765034406&hl=en&oi=scholarr

19. Thompson RN, Stockwin JE, van Gaalen RD, Polonsky JA, Kamvar ZN, Demarsh PA, et al. Improved inference of time-varying reproduction numbers during infectious disease outbreaks. Epidemics. 2019 Dec 1;29:100356.

20. Estimating the time-varying reproduction number of SARS-CoV-2 using … [Internet]. Wellcome Open Research | Open Access Publishing Platform. 2020 [cited 2024 Feb 8]. Available from: https://wellcomeopenresearch.org/articles/5-112

21. The New York Times. Coronavirus (Covid-19) Data in the United States [Internet]. 2021 [cited 2023 Oct 6]. Available from: https://github.com/nytimes/covid-19-data

22. U.S. Census Bureau. Census Data [Internet]. 2020. Available from: https://data.census.gov/profile

23. Agency for Toxic Substances and Disease Registry. CDC/ATSDR SVI: Data and Documentation Download [Internet]. 2022. Available from: https://www.atsdr.cdc.gov/placeandhealth/svi/data_documentation_download.html

24. Nash RK, Bhatt S, Cori A, Nouvellet P. Estimating the epidemic reproduction number from temporally aggregated incidence data: A statistical modelling approach and software tool. PLOS Comput Biol. 2023 Aug 28;19(8):e1011439.

25. Stan Development Team. "RStan: the R interface to Stan." R package version 2.32.6 [Internet]. 2024 [cited 2024 Feb 22]. Available from: https://mc-stan.org/

26. Grant R, Charmet T, Schaeffer L, Galmiche S, Madec Y, Platen CV, et al. Impact of SARS-CoV-2 Delta variant on incubation, transmission settings and vaccine effectiveness: Results from a nationwide case-control study in France. Lancet Reg Health – Eur [Internet]. 2022 Feb 1 [cited 2024 May 10];13. Available from: https://www.thelancet.com/journals/lanepe/article/PIIS2666-7762(21)00264-7/fulltext

27. Nishiura H, Linton NM, Akhmetzhanov AR. Serial interval of novel coronavirus (COVID-19) infections. Int J Infect Dis. 2020 Apr 1;93:284–6.

28. Rue H, Martino S, Chopin N. Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations. J R Stat Soc Ser B Stat Methodol. 2009;71(2):319–92.

29. Rue H, Riebler A, Sørbye SH, Illian JB, Simpson DP, Lindgren FK. Bayesian Computing with INLA: A Review. Annu Rev Stat Its Appl. 2017;4(1):395–421.

30. Schrödle B, Held L. Spatio-temporal disease mapping using INLA. Environmetrics. 2011;22(6):725–34.

31. Rue H, Martino S, Lindgren F, Simpson D, Riebler A, Krainski E. INLA: Functions Which Allow to Perform Full Bayesian Analysis of Latent Gaussian Models Using Integrated Nested Laplace Approximaxion. R package version 0.0-1389624686. 2014.

32. Bivand R, Gómez Rubio V, Rue H. Spatial Data Analysis with R-INLA with Some Extensions. J Stat Softw. 2015 Feb 1;63:1–31.

33. Besag J, Green P, Higdon D, Mengersen K. Bayesian Computation and Stochastic Systems. Stat Sci. 1995 Feb;10(1):3–41.

34. BESAG J. A candidate's formula: A curious result in Bayesian prediction. Biometrika. 1989 Mar 1;76(1):183.

35. Sebhatu A, Wennberg K, Arora-Jonsson S, Lindberg SI. Explaining the homogeneous diffusion of COVID-19 nonpharmaceutical interventions across heterogeneous countries. Proc Natl Acad Sci. 2020 Sep;117(35):21201–8.

36. Huisman JS, Scire J, Angst DC, Li J, Neher RA, Maathuis MH, et al. Estimation and worldwide monitoring of the effective reproductive number of SARS-CoV-2. Davenport MP, editor. eLife. 2022 Aug 8;11:e71345.

37. Gezer F, Howard KA, Litwin AH, Martin NK, Rennert L. Identification of factors associated with opioid-related and hepatitis C virus-related hospitalisations at the ZIP code area level in the USA: an ecological and modelling study. Lancet Public Health. 2024 Jun 1;9(6):e354–64.

38. Citation and Authors [Internet]. [cited 2024 Jun 12]. Available from: https://imperialcollegelondon.github.io/epidemia/authors.html

39. Alvarez L, Colom M, Morel JD, Morel JM. Computing the daily reproduction number of COVID-19 by inverting the renewal equation using a variational technique. Proc Natl Acad Sci. 2021 Dec 14;118(50):e2105112118.

40. Jin S, Dickens BL, Lim JT, Cook AR. EpiRegress: A Method to Estimate and Predict the Time-Varying Effective Reproduction Number. Viruses. 2022 Jul;14(7):1576.

41. Scire J, Huisman JS, Grosu A, Angst DC, Lison A, Li J, et al. estimateR: an R package to estimate and monitor the effective reproductive number. BMC Bioinformatics. 2023 Aug 11;24(1):310.

42. Hasan A, Susanto H, Tjahjono V, Kusdiantara R, Putri E, Nuraini N, et al. A new estimation method for COVID-19 time-varying reproduction number using active cases. Sci Rep. 2022 Apr 23;12(1):6675.