# Assessment of Vaccine Managers' Knowledge and Observance of Vaccination Norms and Standards in an Urban Health District of Yaounde (Cameroon) during the COVID-19 Pandemic

Fabrice Zobel Lekeumo Cheuyem[1]*, Adidja Amani[1], Brian Ngongheh Ajong[2], Raissa Katy Noa Otsali[1], Ariane Nouko[1], Edwige Omona Guissama[3], Christelle Sandrine Ngos[1], Adama Mohamadou[1], Guy Stephane Nloga[1], Michel Franck Edzamba[1], Esther Andriane Bitye Bi Mvondo[1], Larissa Linda Eyenga Ntsek[1], Christian Mouangue[4], Florence Kissougle Nkongo[5]

[1] Department of Public Health, Faculty of Medicine and Biomedical Sciences, The University of Yaounde I, Yaounde, Cameroon

[2] Health Emergencies Programme, World Health Organisation (WHO), Headquarters, Geneva, Switzerland

[3] Direction of Family Health, Ministry of Public Health, Yaounde, Cameroon

[4] Data Management Unit, Planning Department, Public Health Emergency Operations Coordination Centre, Yaounde, Cameroon

[5] Regional Delegation of Public Health for the Centre Region, Ministry of Public Health, Yaounde, Cameroon

*Corresponding author's address: Fabrice Zobel Lekeumo Cheuyem, Department of Public Health, Faculty of Medicine & Biomedical Sciences, The University of Yaoundé I, PO Box. 8526, Yaoundé, Cameroon. Tel: +237 696 57 28 07; E-mail: zobelcheuyem@gmail.com

# Abstract

**Background:** Routine childhood immunization remains a highly cost-effective public health intervention. Global immunization efforts have increased the coverage of proxy antigens in recent decades. However, annual reports indicated a progressive decline in immunization coverage in Yaounde. In addition, data management issues related to Expanded Programme of Immunization (EPI) inputs have been identified. This study aimed to identify knowledge gaps and organizational deficiencies in the implementation of routine immunization in health facilities (HFs) in the Efoulan Health District (HD) of Yaounde.

**Methods:** A descriptive cross-sectional study was conducted in Efoulan, an urban HD of Yaounde, the capital of Cameroon, in May 2022. A self-administered questionnaire was used to collect knowledge data from consenting HCWs. An observation grid was used to assess the implementation of some relevant national EPI guidelines and recommendations.

**Results:** A total of 46 HCWs providing vaccinations in the nine health areas of the HD were enrolled in the study. Almost all participants (93.5%) were unable to name all the diseases that the pentavalent vaccine protects against. More than half of the respondents (59%) were aware of the 15-month booster dose of measles-rubella (MR) vaccine included in the EPI schedule. Half of the respondents knew of the type of antigen contained in the BCG vaccine (52%). More than two-thirds of respondents (72%) had not identified the 6-month MR vaccine as recommended without risk for HIV+ children. Most of respondents were not aware of the management of opened non-lyophilized vials (50%), calculation of wastage factor (89%) and vaccine requirements (98%). Almost all respondents (87%) have poor knowledge on vaccination norms and standards. Most of the visited HFs did not follow the monthly EPI monitoring curve (87%), did not store vaccines according to standards (80%) neither have an EPI case reporting form (93%). Most of the consultants in visited HFs were not familiar with the case definitions of common vaccine preventable diseases.

**Conclusions:** The implementation of routine immunization in HFs remained poor in most of the HFs visited in Efoulan HD, due to the lack of knowledge in the EPI norms and standards of the staff. There is a need to strengthen the capacities of HCWs through ongoing training and supervision.

# Background

Routine childhood immunization remains a highly cost-effective public health intervention [1]. Immunization has contributed to significant reductions in morbidity and mortality from vaccine-preventable diseases (VPDs) worldwide [2]. Global immunization efforts have saved about 154 million lives over the past 50 years, or about 6 lives every minute of every year. Most of these lives saved were infants (101 million) [2,3]. Global immunization efforts have increased coverage of the third dose of the diphtheria-tetanus-pertussis (DTP3) vaccine from near zero in 1974 to an impressive 84% today. This progress, which reached a high of 86% in 2019 before the COVID-19 pandemic, demonstrates the effectiveness of global vaccine programs [4].

Vaccination activities for children aged 0-11 months and pregnant women, led by the Expanded Programme on Immunization (EPI) of the Ministry of Public Health (MINSANTE), are a priority objective in Cameroon's health strategy [5]. The Efoulan Health District (HD) of Yaounde, with its 54 vaccination centers spread across all health sectors, regularly carries out routine and mass vaccination campaigns to meet the targets set by the health authorities [6] .

However, in recent years, annual reports indicated progressive decline in vaccination coverage in the Efoulan HD. This was characterized by a decrease in pentavalent vaccine (penta) 3 coverage from 85.8% to 72% between 2020 and 2021 (a decrease of 15.4%) and an increase in penta 1-3 specific drop-out rates from -1.5% to 1.8% [7,8].

In addition, in resource-limited countries such as Cameroon, the emergence of the coronavirus (COVID-19) had a significant impact on the health system and disrupted the implementation of EPI activities by diverting attention and health resources to the COVID-19 response [9,10].

The reports of the coordination meetings with EPI focal points, HD officers and health facility (HF) representatives highlighted several challenges, including COVID-19-related constraints leading to lower uptake of immunization services by parents, failure of technical and organizational components, quality and management issues related to human resources in health facilities, especially in health facilities identified as having problems in the HD.

In addition, data management issues related to EPI inputs were identified, motivating the conduction of an operational research study to understand the root causes and promote evidence-based decision-making. This study was designed to identify knowledge gaps among immunization stakeholders and describe the organizational pattern in the implementation of routine vaccination in problematic health facilities in the Efoulan HD of Yaounde.

# Methods

**Study Type:** An institutional-based cross-sectional study was conducted in the Efoulan HD

**Study Period:** The study was conducted throughout the month of May 2022.

**Study Site:** The Efoulan HD is one of the 32 districts of the Central Region. It comprises nine health areas (eight urban and one rural). The population in 2021 was estimated at 453,047 inhabitants covering an area of over 67 km² (density of 6,761 inhabitants per km²). The HD counts 117 health facilities, including seven public, three denominational and 107 private facilities (Figure 1).

**Fig. 1** Localization of Efoulan among Yaounde Health Districts [11]

**Study population**: Health workers (HCWs) vaccinating in the Efoulan HD and problematic health facilities of the HD.

**Study Participants:** HCWs in a HF who vaccinates in the Efoulan HD and who provided informed consent were included.

**Sample Size and Sampling Technique:** To ensure the representativeness of all nine health areas, HCWs were enrolled using a purposive sampling technique. For observational analysis, 15 HFs identified as problematic were visited.

**Data Collection Tool and Procedures**: Participants were contacted during coordination meetings and field visits. A pre-tested and self-administered questionnaire of 28 questions was administered, covering socio-demographic information, knowledge of priority targets, vaccination schedule, technique and input management. An observation grid of 20 pre-registered questions on Google Forms collected observational data from 15 sampled health facilities.

**Data Processing and Analysis**: Each participant was assigned a score based on the knowledge assessment (16 questions, 1 point each). The results were then classified into three categories: good (≥75%), average (50-74%), and poor (<50%). [12]. Data was entered into Google Forms, extracted into Microsoft Office Excel 2016 and analyzed using IBM SPSS version 26. Charts were generated using Microsoft Office Excel 2016. In cross-tabulations, proportions were compared using chi-squared or Fisher's exact tests. A p-value<0.05 was considered statistically significant.

**Operational Definition**: Problematic HFs were described as those with poor performance in implementing immunization programs (low vaccination coverage for the penta 3 [<50%] and poor data management in terms of timeliness, completeness and coherence of data recorded in DHIS2) during the year 2021 and who had difficulty mitigating challenges after HD supervision.

# Results

## Study participant's profile

A total of 46 HCWs from the HFs providing vaccinations in the nine health areas of the HD were enrolled in the study. They were aged 27-53 years, with half of the participants aged between 27 and 36 years. Most of participants were female (84.8%) and had higher educational level (63%) (Table 1).

**Table 1** Socio-demographic characteristics of participants, Efoulan Health District, May 2022 (*n*=46)

| Characteristics | Count (n) | Frequency (%) |
|---|---|---|
| **Health area** | | |
| Afanoyoa | 5 | 10.9 |
| Ahala 1 | 3 | 6.5 |
| Ahala 2 | 4 | 8.7 |
| Efoulan | 4 | 8.7 |
| Ngoa-ekelle | 1 | 2.2 |
| Nsimeyong 1 | 8 | 17.4 |
| Nsimeyong 2 | 9 | 19.6 |
| Nsimeyong 3 | 9 | 19.6 |
| Obili | 3 | 6.5 |
| **Age** | | |
| 18-24 | 3 | 6.5 |
| 25-34 | 26 | 56.5 |
| 35-44 | 11 | 23.9 |
| 45-49 | 3 | 6.5 |
| 50+ | 3 | 6.5 |
| **Gender** | | |
| Female | 39 | 84.8 |
| Male | 7 | 15.2 |
| **Educational level** | | |
| Primary | 1 | 2.2 |
| Secondary | 16 | 34.8 |
| Higher | 29 | 63.0 |

Half of the respondents were nurses (58.7%) and vaccinators (54.3%). Half of the participants had already spent more than 2 years in the immunization service and in the health centers where they worked (Table 2).

**Table 2** Socio-professional characteristics of participants, Efoulan Health District, May 2022 (*n*=46)

| Characteristics | Count (n) | Frequency (%) |
|---|---|---|
| **Grade** | | |
| Assistant nurse | 13 | 28.3 |
| Assistant laboratory technician | 1 | 2.2 |
| Nurse | 27 | 58.7 |
| Midwife | 3 | 6.5 |
| Senior laboratory technician | 2 | 4.3 |
| **Position in the HF** | | |
| Head of health center | 5 | 10.9 |
| Health area manager | 1 | 2.2 |
| Healthcare worker | 14 | 30.4 |
| VPDs focal point | 1 | 2.2 |
| Head of vaccination unit | 25 | 54.3 |

*HF: Health facility; VPD: Vaccine Preventable Disease*

Half of the participants (50%) had already received training on the EPI, and for almost two-thirds of them (65.2%), this training dated back to more than a year. Just over half of the participants reported that they had been supervised as part of their EPI activities, and for the most of them (70.4%) this supervision was less than a year old.

**Knowledge on immunization**

Less than one-third of participants (26%) identified active artificial immunity as the type of immunity provided by vaccination. More than one-third of participants (40%) had no knowledge on this topic (Figure 2).

**Fig. 2** Knowledge of the type of immunity provided by vaccination among healthcare workers, Efoulan Health District, May 2022 (*n*=46)

Almost all participants (93.5%) were unable to name all the diseases the pentavalent vaccine protects against. *Haemophilus influenzae* type b was the antigen they knew least about. It should be noted that almost a third of respondents (30.1%) had no knowledge of the antigens contained in this flagship EPI vaccine (Figure 3).

**Fig. 3** Knowledge of diseases prevented by the pentavalent vaccine among healthcare workers in Efoulan Health District, May 2022 (*n*=46)

More than half of the respondents (59%) were aware of the 15-month booster dose of the measles-rubella (MR)2 vaccine included in the EPI schedule (Figure 4).

**Fig. 4** Healthcare workers' Knowledge of measles-rubella booster at 15 months in the Cameroon EPI schedule, Efoulan Health District, May 2022 (*n*=46)

Half of respondents had Knowledge of the type of antigen contained in the BCG vaccine (52%) (Figure 5).

**Fig. 5** Knowledge of the type of antigen contained in the BCG vaccine among healthcare workers, Efoulan Health District, May 2022

## Knowledge of priority target populations

A low proportion of participants (6.5%) were aware of the current proportion of surviving children aged 0-11 months expected in the Efoulan HDs, which is 2.5%. Slightly more than half of the respondents (59%) had no knowledge of this value, two-thirds of whom (66.7%) had never received EPI training ($p$=0.020). In addition, more than half the respondents had no knowledge of the expected proportion of pregnant women, which is 2.7% of the total HD population (Table 3).

**Table 3** Knowledge of priority target proportions among healthcare workers, Efoulan Health District, May 2022 (*n=*46)

| Expected proportion (%) | EPI training received n (%) | | Total (100%) | *p*-value |
|---|---|---|---|---|
| | **No** *n*=13 | **Yes** *n*=13 | | |
| **Surviving children** | | | | |
| 2.5 (Yes) | 1 (33) | 2 (67) | 3 | |
| 2.7 | 3 (43) | 4 (57) | 7 | |
| 3.4 | 0 | 6 (100) | 6 | 0.020 |
| 4.5 | 1 (33) | 2 (67) | 3 | |
| I don't know | 18 (67) | 9 (33) | 27 | |
| **Pregnant women** | | | | |
| 2.5 | 1 (20) | 4 (80) | 5 | |
| 2.7 (Yes) | 1 (20) | 4 (80) | 5 | 0.289 |
| 3.4 | 1 (50) | 1 (50) | 2 | |
| 4.5 | 2 (50) | 2 (50) | 4 | |
| I don't know | 18 (60) | 12 (40) | 30 | |

*EPI: Expanded Programme of Immunization*

Most of participants (87%) did not identify all the vaccines (PCV, pentavalent and RR) that could be given to an unvaccinated one year or more. The most cited vaccine was the Measles-Rubella vaccine (Figure 6).

**Fig. 6** Knowledge of Vaccines receivable by an unvaccinated child of one year old or more, Efoulan Health District, May 2022 (*n*=46)

More than two-thirds of respondents (72%) had not identified the 6-month MR vaccine as recommended without risk for HIV+ children (Figure 7).

*MR: Measles-Rubella vaccine*

**Fig. 7** Knowledge of vaccines recommended without risk for HIV+ children among healthcare workers, Efoulan Health District, May 2022 (*n*=46)

Less than half of respondents (44%) had identified the optimal 28-day interval between two doses of EPI vaccine (Figure 8).

**Fig. 8** Knowledge of the optimal period between two EPI vaccine doses, Efoulan HD, May 2022 (*n*=46)

Less than half of the respondents were aware of the correct angle for intramuscular injection during the vaccination process (43%), with more than a quarter of participants (26%) having no idea about this standard (Figure 9).

**Fig. 9** Knowledge of the ideal angle for intramuscular vaccination, Efoulan HD, May 2022 (*n*=46)

Less than a third of HCWs know the distance of 5 km from which the advanced strategy must be implemented. In contrast, more than half of respondents (52%) did not know the correct distance (Figure 10).

**Fig. 10** Knowledge of distance requiring organization of advanced vaccination strategy, Efoulan HD, May 2022 (*n=46*)

## Logistic Component of Vaccination

When asked what was the EPI tracer antigen, more than half of the participants correctly identified (41%) penta 3 and more than a quarter (28%) had no idea (Figure 11).

*MR: Measles-Rubella, Penta: Pentavalent vaccine*

**Fig.11** Knowledge of EPI tracer antigen, Efoulan HD, May 2022 (*n*=46)

Most of respondents were not aware of the various components of EPI input management, particularly managing opened non-lyophilized vials (50%), calculating the wastage factor (89%), calculating vaccine requirements (98%). Participants who reported to have received training were significantly more knowledgeable about input management than untrained health workers (Table 4).

**Table 4** Knowledge of EPI input management, Efoulan HD, May 2022 (*n=46*)

| Variables | EPI training received n (%) | | Total (100%) | *p*-value |
|---|---|---|---|---|
| | **No** *n*=13 | **Yes** *n*=13 | | |
| **Management of opened, non-lyophilized vials** | | | | |
| Should be discarded after vaccination session | 3 (30) | 7 (70) | 10 | |
| Should not be used after 28 days | 11 (48) | 12 (52) | 23 | 0.320 |
| Could be used after 28 days | 2 (67) | 1 (33) | 3 | |
| Did not know how | 7 (70) | 3 (30) | 10 | |
| **Loss factor** | | | | |
| No | 1 (11) | 8 (89) | 9 | |
| Yes | 1 (20) | 4 (80) | 5 | **0.006** |
| Did not know how to calculate | 21 (66) | 11 (34) | 32 | |
| **Calculation of the requirement dose** | | | | |
| No | 2 (13) | 13 (87) | 15 | |
| Yes | 0 (0) | 1 (100) | 1 | **0.001** |
| Did not know how to calculate | 21 (70) | 9 (30) | 30 | |

*EPI: Expanded Programme of Immunization*

## Global Score

Of the sixteen questions assessed, the global score ranged from 1/16 (6%) to 11/16 (69%). Half of the participants scored below 25%. The modal score was 3/16 (19%). Almost all respondents (87%) have poor Knowledge on vaccination norms and standards (Table 5).

**Table 5** Score of respondents by previous participation in EPI training, Efoulan Health District, May 2022 (*n=46*)

| Knowledge score | EPI training received n (%) | | Total (100%) | *p*-value |
|---|---|---|---|---|
| | **No** *n*=13 | **Yes** *n*=13 | | |
| Poor | 21 (53) | 19 (47) | 40 | 0.665 |
| Average | 2 (33) | 4 (67) | 6 | |

*EPI: Expanded Program of Immunization*

## Observance of EPI Norms and Standards

Observation of the effective implementation of EPI norms and standards was conducted in both public and private HFs of the Efoulan HD identified as problematic (Table 6).

**Table 6** Health facilities visited in the health areas of the Efoulan Health District, May 2022

| Health area | Count (n) | Frequency (%) |
|---|---|---|
| Afanoyoa | 2 | 13.3 |
| Ahala 1 | 1 | 6.7 |
| Ahala 2 | 2 | 13.3 |
| Efoulan | 1 | 6.7 |
| Nsimeyong 1 | 1 | 6.7 |
| Nsimeyong 2 | 2 | 13.3 |
| Nsimeyong 3 | 2 | 13.3 |
| Obili | 4 | 26.7 |
| **Total** | **15** | **100** |

The HFs vaccinating in the Efoulan HD showed several deficiencies in the application of Cameroon's EPI norms and standards. Most of the assessed HD did not follow the monthly EPI monitoring curve (87%), did not store vaccines according to standards (80%) neither have an EPI case reporting form (93%).

Regarding surveillance, most of the consultants in the HFs visited were not familiar with the cases definitions of common VPDs (poliomyelitis, yellow fever, measles) (Table 7).

**Table 7** Health facility compliance with some relevant national EPI guidelines, May 2022 (n= 15)

| Guideline | Frequency *n* (%) | |
|---|---|---|
| | No | Yes |
| **Organization** | | |
| Availability of an immunization manager | 0 | 15 (100) |
| Availability of a VPD focal point | 1 (7) | 14 (93) |
| **Implementation and monitoring of the program** | | |
| EPI calendar posted | 1 (7) | 14 (93) |
| EPI monitoring curve available | 7 (47) | 8 (53) |
| Monitoring curve plotted and up to date | **13 (87)** | 2 (13) |
| Vaccination registers available and correctly filled in | 5 (33) | 10 (67) |
| Lost to follow-up register available and completed | **10 (67)** | 5 (33) |
| **Cold chain** | | |
| Refrigerator available and functional | 1 (7) | 14 (93) |
| Contingency plan posted near the refrigerator | 9 (60) | 6 (40) |
| Fridge temperature record available | 5 (33) | 10 (67) |
| Fridge temperature record updated | **11 (73)** | 4 (27) |
| Fridge temperature measuring device operational | 6 (40) | 9 (60) |
| All EPI antigens available | 9 (60) | 6 (40) |
| Vaccines stored according to standards in the cold chain | **12 (80)** | 3 (20) |
| Tainted vaccines in cold chain | 9 (60) | 6 (40) |
| **Surveillance** | | |
| Notification form of VPDs available | **14 (93)** | 1 (7) |
| Definition of measles case known to consultant | 10 (67) | 5 (33) |
| Definition of AFP cases known to the consultant | **12 (80)** | 3 (20) |
| Yellow fever case definition known to the consultant | 9 (60) | 6 (40) |
| VPD cases definitions available and displayed | 4 (26,7) | **11 (73)** |

*VPD: Vaccine Preventable Disease; EPI: Expanded Programme of Immunization; AFP: Acute Flaccid Paralysis*

# Discussion

The aim of this study was to assess the implementation of vaccination programs in an urban HD of Yaounde. It informs vaccination program implementation and identifies areas for improvement to enhance vaccination coverage and effectiveness in Efoulan HD.

The geographical coverage of the study areas was satisfactory. Most of the HCWs visited had a designated immunization coordinator and a VPD focal point. However, almost half of the HCWs had less than two years' experience in immunization services. This may be due to the high turnover of HCWs between HFs. Retention of skilled HCWs remains critical to the management of human resources for health in Cameroon [13,14]. HCWs, especially those in private facilities, frequently move between HFS in search of better offers and benefits, hindering the effective use of skills acquired through training and supervision.

Significant gaps have been identified in calculating target population sizes, determining loss factors, and estimating vaccine requirements. This situation results in either shortages or surpluses of vaccine in the HF's cold chain, which subsequently can lead to expiration and vaccines waste. All this leads to difficulties in stock management and implementation of a quality vaccination service. This result confirms the findings in Dschang HD in the West Region of Cameroon, where inadequate immunization service delivery was associated with limited knowledge of vaccine and cold chain management by HCWs [15].

Most participants lacked knowledge of the tracer antigen, general vaccine content (e.g., pentavalent vaccine), and safe vaccination techniques (intramuscular injection angle). This inadequacy may affect the ability of HCWs to provide effective education during immunization and to retain mothers for follow-up immunization. An inappropriate route of administration may also increase the risk of adverse events following immunization. As a result, parents may be more reluctant to continue vaccinating their children, leading to an increase in the Penta 3 dropout rate. In addition to an adequate supply of vaccines, the problem of absent or incomplete vaccination could be overcome by improving social mobilization strategies to sensitize mothers and other stakeholders who are reluctant to vaccinate their children, as reported by studies conducted in Cameroon and elsewhere in the world [8–10].

Most of HCWs were unaware of the five km distance beyond which advanced strategies must be organized to bring vaccination as close to the population as possible. As a result, advanced vaccination sessions were poorly planned or not organized at all. However, it should also be noted that most HFs in urban HDs are private and it is not part of their culture to implement advanced strategies. This justifies the implementation of supervision of private HFs to sensitize managers and ensure that the benefits of this strategy are well understood and integrated into vaccination planning. Medical evidences has shown that advanced strategies bring immunization services close to the population, reduce time waste by parent to get their children vaccinated and increase the adherence to vaccination [19,20].

The observational analysis revealed significant deficiencies in cold chain management, immunization monitoring. Similar trends were reported in Cameroon and Ethiopia [21,22].

The monitoring curves, although present, were not regularly monitored, and most HFs did not store vaccines properly or monitor temperatures on a daily basis. probably due to ignorance or negligence on the part of the people in charge of immunization in these HFs. This situation is attributed to lack of motivation and poor supervision of HFS by operational and intermediate levels. This result corroborates finding in North West Health Region in Cameroon [22,23].

Poor understanding of the definitions of cases of VPD and the low level of involvement of HF managers in surveillance, particularly the unavailability of notification forms, could explain the overall under-reporting of VPD in the district.

Overcoming all these challenges will require strong commitment and significant financial mobilization. The shift in healthcare attention during COVID-19 pandemic can lead to secondary health crises such as outbreaks of infectious diseases such as measles. Restrictions due to COVID-19 impede access to HFs for vaccination [17]. During an outbreak such as COVID-19, public health efforts were more focused on controlling the outbreak, resulting in fewer children receiving vaccines. During periods of quarantine, immunization services for all age groups were interrupted, delayed, reorganized, or suspended altogether in many parts of the world [24]. These challenges may have significantly affected the delivery of routine immunization services in the urban HD of Yaounde.

## Limitations

The study has limitations, particularly the small sample size. A multi-district study would provide a more comprehensive understanding of the issues in the Central Region. Interpretation of the results should take into account the dynamic environment (staff turnover, supervision, and training); therefore, the results may not accurately reflect the current situation in the HD.

## Conclusions

Implementation of routine vaccination remained poor in most of the HFs visited. Significant gaps remained in the EPI-related knowledge of HCWs in the HD. Efforts are still needed to raise the level of observance of EPI technical and organizational guidelines to ensure safe and effective immunization of target populations. These efforts include ongoing technical and practical training on EPI guidelines and monitoring to ensure effective use of the capacity building received by various stakeholders.

## Recommendations

To address various challenges revealed in this study, we suggest to HD and regional level health authorities to enhance training of HCWs, improve the use of vaccination data tools, strengthen cold chain management and surveillance through planning and implementation of supervision of problematic or priority HFs. They should also develop incentive strategies to retain experienced and trained HCWs.

## Declarations

**Authors' Contribution:** Study design & conception: FZLC; Data collection: FZLC; Data analysis, visualization and interpretation: FZLC; Drafting of original manuscript: FZLC; Critical revision of the manuscript: FZLC, AA, BNA. RKNO, AN, EOG, CSN, AM, GSN, MFE, EABBM, LLEN, CM and FKN; Final approval of the manuscript: All authors.

**Ethical Approval Statement**: Ethical clearance for the present study was waived by the Faculty of Medicine of Yaounde ethical review board. Administrative approval was obtained from the Chief Medical Officer of the Efoulan HD. Additionally, participants were required to provide a signed informed consent for their responses to be used for research purposes. The confidentiality,

anonymity and autonomy of research participants were respected throughout the study. All methods were performed in accordance with the relevant guidelines of the Helsinki declaration.

During the field visits, the participants benefited of practical training on the EPI norms and standards. As a result, the various gaps identified in both knowledge and application of the standards were addressed to the best of our ability.

**Consent for publication:** Not applicable.

**Availability of data and materials:** All data generated or analyzed during this study are included in this published article.

**Competing interests:** All authors declare no conflict of interest and have approved the final version of the article.

**Funding source:** This research did not receive any specific grant from funding agencies in the public, commercial or not-for-profit sectors.

**Acknowledgements:** Our gratitude goes to the HCWs of the Efoulan HD who accepted to participated in this study and to the Chief Medical Officer of Efoulan HD, Dr. Ottou Awa Georges Marie who gave authorization for the conduction of the study in his administrative circumscription.

# References

1. Zhou F. Health and Economic Benefits of Routine Childhood Immunizations in the Era of the Vaccines for Children Program — United States, 1994–2023. *MMWR Morb Mortal Wkly Rep*. 2024;73:682–5.

2. Global immunization efforts have saved at least 154 millions lives over the past 50 years. WHO, Geneva. 2024. https://www.who.int/news/item/24-04-2024-global-immunization-efforts-have-saved-at-least-154-million-lives-over-the-past-50-years. Accessed 2024 Nov 4.

3. Shattock AJ, Johnson HC, Sim SY, Carter A, Lambach P, Hutubessy RCW, et al. Contribution of vaccination to improved survival and health: modelling 50 years of the Expanded Programme on Immunization. *The Lancet*. 2024;403(10441):2307–16.

4. WHO/UNICEF estimates of national immunization coverage. WHO, Geneva. 2024. https://www.who.int/teams/immunization-vaccines-and-biologicals/immunization-analysis-and-insights/global-monitoring/immunization-coverage/who-unicef-estimates-of-national-immunization-coverage. Accessed 2024 Nov 4.

5. Amani A, Saidu Y, Asaah CT, Cheuyem FZL, Njoh AA, Kabir HA, et al. Immunization in Cameroon: Uncovering Progress, Confronting Challenges, and Paving the Path towards Achieving Sustainable Development Goal 2030 Targets. *J Vaccines Vaccin*. 2023;14(3):1–10.

6. Reaching Every District (RED). WHO | Regional Office for Africa. 2017. https://www.who.int/teams/immunization-vaccines-and-biologicals/essential-programme-on-immunization/implementation/reaching-every-district-(red). Accessed 2024 Oct 31.

7. Baguune B, Ndago JA, Adokiya MN. Immunization dropout rate and data quality among children 12–23 months of age in Ghana. *Arch Public Health*. 2017;75(1):1–8.

8. Ntenda PAM, Sixpence A, Mwenyenkulu TE, Mmanga K, Chirambo AC, Bauleni A, et al. Determinants of pentavalent and measles vaccination dropouts among children aged 12–23 months in The Gambia. *BMC Public Health*. 2022;22(1):520.

9. Amani A, Mossus T, Cheuyem FZL, Bilounga C, Mikamb P, Basseguin Atchou J, et al. Gender and COVID-19 Vaccine Disparities in Cameroon. *COVID*. 2022;2(12):1715–30.

10. Yoo KJ, Lee Y, Lee S, Friebel R, Shin S ae, Lee T, et al. The road to recovery: impact of COVID-19 on healthcare utilization in South Korea in 2016–2022 using an interrupted time-series analysis. *Lancet Reg Health – West Pac*. 2023 ;41 : 100904.

11. Geomatic Strategy. Analyse géographique de la couverture sanitaire au Cameroun Répartition des formations sanitaires et de la charge du personnel de santé. https://www.geostrategies.net/2019/04/01/strategies-spatiales-n2-mars-avril-2019-analyse-geographique-de-la-couverture-sanitaire-au-cameroun-repartition-des-formations-sanitaires-et-de-la-charge-du-personnel-de-sante/. Accessed 2021 Sep 11.

12. Zia I, Cheema SS, Sheikh NS, Ashraf H. Hand hygiene knowledge, attitudes, and self-reported practices among medical and nursing staff of a tertiary-care military hospital : a cross-sectional study. *Int J Infect Control*. 2022;18:10.3396/ijic.v18.21469.

13. Kufe NC, Metekoua C, Nelly M, Tumasang F, Mbu ER. Retention of health care workers at health facility, trends in the retention of knowledge and correlates at 3rd year following training of health care workers on the prevention of mother-to-child transmission (PMTCT) of HIV–National Assessment. *BMC Health Serv Res*. 2019;19(1):78.

14. Goga AE, Muhe LM. Global challenges with scale-up of the integrated management of childhood illness strategy: results of a multi-country survey. *BMC Public Health*. 2011;11(1):503.

15. Ebile Akoh W, Ateudjieu J, Nouetchognou JS, Yakum MN, Djouma Nembot F, Nafack Sonkeng S, et al. The expanded program on immunization service delivery in the Dschang health district, west region of Cameroon: a cross sectional survey. *BMC Public Health*. 2016;16(1):801.

16. Félicitée ND, Christiane T, Roger D, Sandra T, Stève FWD, Andreas C, et al. Factors Influencing Routine Vaccination of Children of Mothers Live-Stock Retailers in the Markets of Yaoundé. *World J Vaccines*. 2016;6(2):23–33.

17. Ngwa CH, Doungtsop BCK, Bihnwi R, Ngo NV, Yang NM. Burden of vaccine-preventable diseases, trends in vaccine coverage and current challenges in the implementation of the expanded program on immunization: A situation analysis of Cameroon. *Hum Vaccines Immunother*. 2022;18(1):1939620.

18. Wie SH, Jung J, Kim WJ. Effective Vaccination and Education Strategies for Emerging Infectious Diseases Such as COVID-19. *J Korean Med Sci*. 2023;38(44):e371.

19. Guttman M. Adapting Strategies to Realities in Togo for Catch-Up Vaccination. Sabin Vaccine Institute, Washington, D.C. 2023. https://www.sabin.org/resources/adapting-strategies-to-realities-in-togo-for-catch-up-vaccination/. Accessed 2024 Nov 4.

20. Domoina Ratovozanany. Thanks to advanced strategies, parents waste less time immunizing their children. UNICEF, New York. 2021. https://www.unicef.org/madagascar/en/stories/thanks-advanced-strategies-parents-waste-less-time-immunizing-their-children. Accessed 2024 Nov 4.

21. Bogale HA, Amhare AF, Bogale AA. Assessment of factors affecting vaccine cold chain management practice in public health institutions in east Gojam zone of Amhara region. *BMC Public Health*. 2019;19(1):1433.

22. Yakum MN, Ateudjieu J, Walter EA, Watcho P. Vaccine storage and cold chain monitoring in the North West region of Cameroon: a cross sectional study. *BMC Res Notes*. 2015 ;8(1) :145.

23. Ateudjieu J, Kenfack B, Nkontchou BW, Demanou M. Program on immunization and cold chain monitoring: the status in eight health districts in Cameroon. *BMC Res Notes*. 2013;6(1):101.

24. Hamid H, Mallhi TH, Naseer MS, Younas I, Rashid MA, Pervaiz A, et al. The COVID-19 pandemic threatens the Expanded Program on Immunization: recommendations for sustaining vaccination goals. *Drugs Ther Perspect*. 2020;36(11):523.