Autistic Neural 'Shiftability': The Distinct Pharmacological Landscape of the Autistic Brain

Tobias P. Whelan[1], Lucas G. S. França[1,2,3], Mihail Dimitrov[1], Charlotte M. Pretzsch[1], Hester Velthuis[1], Andreia C. Pereira[4,1], Claire L. Ellis[1], Glynis Ivin[5], James L. Findon[6], Robert H. Wichers[7], Francesca M. Ponteduro[1], Johanna Kangas[1], Naoise Mulcrone[1], Nichol M. L. Wong[8,9], Dafnis Batalle[1,2], Nicolaas A. J. Puts[1,10,11], Eileen Daly[1*], Declan G.M. Murphy[1,10,11*], Gráinne M. McAlonan[1,10,11*]

1) Department of Forensic and Neurodevelopmental Science, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, United Kingdom

2) Department of Early Life Imaging, School of Biomedical Engineering & Imaging Sciences, King's College London, London, United Kingdom

3) Department of Computer and Information Sciences, Faculty of Engineering and Environment, Northumbria University, Newcastle upon Tyne, United Kingdom

4) Coimbra Institute for Biomedical Imaging and Translational Research (CIBIT), Institute for Nuclear Sciences Applied to Health (ICNAS), University of Coimbra, Coimbra, Portugal

5) South London and Maudsley NHS Foundation Trust Pharmacy, London, United Kingdom

6) Department of Psychology, School of Mental Health and Psychological Sciences, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, United Kingdom.

7) N=You Neurodevelopmental Precision Center, Department of Child and Adolescent Psychiatry & Psychosocial Care, Division of Woman-Child, Amsterdam UMC, Amsterdam, The Netherlands

8) Department of Psychology, The Education University of Hong Kong, Ting Kok, Hong Kong

9) Centre for Psychosocial Health, The Education University of Hong Kong, Ting Kok, Hong Kong

10) NIHR-Maudsley Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and the Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, United Kingdom

11) MRC Centre for Neurodevelopmental Disorders, King's College London, London, United Kingdom

*Joint senior authors

One Sentence Summary: The autistic brain is differentially responsive to pharmacological modulation of the GABA, serotonin and µ-opioid neurotransmitter systems.

Abstract

Functionally inter-connected large-scale brain networks regulated by multiple neurotransmitter systems underpin complex human behaviour. A wide range of alterations in this neurofunctional landscape have been associated with autism. However, this evidence is mostly derived from cross-sectional analyses and thus assumes that any differences are 'static'. Indeed, large sample sizes are required to identify reproducible baseline differences in functional connectivity between autistic and non-autistic individuals. Instead, we propose that what is different in autism is not necessarily the baseline functional connectivity of resting-state networks, but rather their responsivity or regulation by neurotransmitter systems. We tested this hypothesis using a uniform analytical framework to capture the modulation of functional connectivity by single doses of pharmacological probes targeting three major neurotransmitter systems implicated in autism, namely gamma-amino-butyric-acid (GABA), serotonin (5HT) and μ-opioid systems. Every drug challenge altered resting-state network connectivity differentially in the autistic compared to the non-autistic group. However, regardless of the neurotransmitter system probed, drug challenge elicited increases in between-network connectivity in autistic participants but minimal or decreased between-network connectivity in their non-autistic peers. There was no group difference in the responsivity of within-network connectivity. Thus, there is altered *responsivity* of neurotransmitter systems in the autistic brain. This has important implications for pharmacotherapy in autism because these neurotransmitter systems are the targets of several medications commonly prescribed to manage mental health conditions that frequently co-occur in autism. Investigating if these drug-induced 'shifts' in functional connectivity can help provide better targeted clinical interventions, will be important next translational steps.

Introduction

Autism is the result of biological differences in the brain. It is, however, complex and commonly complicated by the co-occurrence of mental health challenges. To date, no biological measure reliably separates autistic from non-autistic people[1]. This has been explained as a consequence of i) too broad a diagnostic definition of autism[2] and/or ii) cohorts that are inherently heterogeneous; and/or iii) differences masked by mixing ages and genders[3] and/or iv) a lack of harmonized methods[4]. Yet, perhaps more importantly, the brain is a dynamic organ and doesn't simply function 'at rest'; rather it *responds* to the modulatory effect of a *variety* of neurotransmitter systems. This is rarely examined in humans. Instead, most studies examine the brain at rest and not in response to the challenge of key neurotransmitter pathways. In our recent studies, we have begun to examine neural responsivity of the human brain to pharmacological challenge by targeting a specific neurotransmitter system, and found that autistic individuals respond differently than non-autistic individuals on selected neuroimaging parameters, which have included a 'seed-based' approach or during task conditions)[5]. However, we have not examined non-task dependent or 'resting-state' brain responsivity *across more than one neurotransmitter system (i.e. a variety of challenges)* using a uniform analytical framework.

Here we focus on functionally inter-connected large-scale brain networks which are generally accepted to underpin complex human behaviour[6]. Their within- and between-network connectivity can be captured using functional MRI (fMRI) techniques during 'resting-state', a condition uncomplicated by task-based demands. This systems-level communication arises through neuronal activity across several organisational scales. At the circuit-level, binding of receptors alters neuronal firing rate and modifies the 'blood oxygen level dependent' (BOLD) fMRI signal. Various alterations in functional connectivity have been associated with autism. However, identifying reproducible baseline differences in autism, for example incorporating the default mode and somatomotor networks[7], has required extremely large sample sizes (e.g. in excess of n = 1800[7]), often pooling across multiple studies and cohorts for meta-analyses. Even then, these connectivity differences are only moderately associated with clinical traits of autism[8]. Moreover, to achieve such large samples, participants are often combined across age groups and baseline differences are especially limited when analyses are limited to autistic adults[3]. This demand for large sample sizes to identify only modest differences in baseline

functional connectivity differences in autistic relative to non-autistic people, is also impractical for translating findings to clinical settings.

Cross-sectional designs also limit investigation of how neurotransmitter systems modulate functional connectivity in the human brain and prior studies have relied upon indirect correlational approaches to link measures of brain chemistry (for example, using PET or MR spectroscopy) to function. To provide direct experimental evidence for altered responsivity of functional networks in autism, we need to *change* the target system and observe *shifts* in connectivity which are distinct from non-autistic individuals. This concept underpins a series of studies adopting our 'shiftability' paradigm[5], which involves pharmacologically challenging specific neurotransmitter systems and measuring the resultant shift in brain function across multiple organisational levels.

In our prior 'shiftability' studies of autistic and non-autistic adults we applied this paradigm to investigate several neurotransmitter systems with a range of compounds: acting at GABA receptors A (targeted by positive allosteric modulator AZD7325)[9] and B (targeted by agonist arbaclofen)[10], serotonin (5HT; targeted serotonin re-uptake inhibitor citalopram)[11] and μ-opioid receptor (targeted by agonist tianeptine)[12] pathways. We chose these neurotransmitter systems as they have been implicated in autism by genetic, post-mortem and preclinical studies[13,14] (yet direct evidence for a functional difference is limited – see above) and they are also the targets of medications commonly prescribed to autistic people for co-occurring psychiatric and neurological conditions. In this study, we used the fMRI data collected across these individual studies to set-up a uniform analytical framework to test the hypothesis that the pharmacological responsivity of the autistic and non-autistic human brain is different.

First, we assessed the baseline group differences in within- and between-network functional connectivity which, based on extant literature, we predicted to be minimal between autistic and non-autistic participants at this sample size (total non-autistic, n = 67 and autistic, n = 54 across all study samples included)[3]. Next, we investigated within- and between-network responsivity in autistic and non-autistic participants by examining each pharmacological probe under a uniform analytical framework. Finally, we pooled within- and between-network shifts following drug challenge to establish any generalised response profile in autistic and non-autistic people.

Materials and Methods

Experimental Design

All component case-control studies included here were conducted in accordance with the Declaration of Helsinki at the Institute of Psychiatry, Psychology and Neuroscience (IoPPN), King's College London (KCL) in London, UK. Our studies did not address safety or clinical efficacy and the UK Medicines and Health Regulatory Authority (MHRA) confirmed that these studies used a drug as a probe of neurotransmitter systems and none constituted a clinical trial of an Investigational Medicinal Product (IMP) as defined by the EU Directive 2001/20/EC. Nevertheless, our protocols were registered on clinicaltrials.gov for transparency. Ethical approval had been received from the appropriate Institutional Ethics Board or National UK Health Research Authority in the UK.

Data were collated across three distinct studies conducted during different time periods (**Table 1**). All studies had a comparable design in which resting-state data were acquired following placebo and a single acute oral dose of a pharmacological probe (at maximum plasma concentration, t_{max}) on separate visit days. The administration order was randomized across participants in all studies and the researcher and participant were blind to the dose order.

Participants

Participants recruited into the autism group had IQ>70 and a clinically confirmed Autism diagnosis as per the recruitment criteria of the parent study. The non-autistic participants were recruited to match the group age and IQ of the autistic participants. All participants gave written informed consent prior to being included in any experimental procedures. **Table 1** summarizes the parent studies providing data for these analyses. Further demographic information is included in **Supplementary Table 1**.

Table 1. Studies included in the analyses. Values in parentheses indicate the number of female participants, note that only male participants were recruited for the serotonin and opioid study. 'Time of scan' refers to the approximate time of functional MRI scan post-administration of drug. 'Washout time' refers to the minimum time between repeat visits in which drug/placebo was administered and allowed sufficient time for drug to leave the system (half-life, $T_{1/2}$). MRI data were acquired at maximum plasma concentration (T_{max}). Given the different T_{max} of citalopram and tianeptine in the 'serotonin & opioid' study, placebo scans were acquired at the same time post-dose as either the citalopram (180 mins) or tianeptine condition (60 mins). In all studies, not all participants completed every study visit, this 'dropout' explains the discrepancy in the sample sizes between conditions (please see original reports for the details[15–17]).

Study	Condition	Dosage (mg)	n Non-Autistic	n Autistic	Time of Scan (mins)	Washout Time (Days)	Ethical Approval	Clinical Trials Identifier	Primary Molecular Targets	Supplier
GABA	Placebo	0	11 (5)	8 (4)	88	5 (AZD $T_{1/2}$ = 8.5-9 hrs[18] & Arbaclofen $T_{1/2}$ = 4-6 hrs[19])	18/WM/0208	NCT03678129	Selective GABA$_A$ receptor positive allosteric modulator (selective to GABA$_A$ α2/3 subunit-containing receptors)	AstraZeneca PLC, Cambridge, UK (as AZD7325)
	AZD7325	10	13 (5)	10 (4)						
	AZD7325	20	13 (5)	10 (4)						
	Placebo	0	31 (14)	21 (3)			RESCM-18/19-4081	NCT03594552	Selective GABA$_B$ receptor agonist	Clinical Research Associates LLC, Simons Foundation (as STX209)
	Arbaclofen	15	34 (15)	22 (5)						
	Arbaclofen	30	27 (10)	19 (4)						
Serotonin & opioid	Placebo	0	18 (0)	18 (0)	180/ 60	8 (Citalopram $T_{1/2}$ = 35 hrs[20] & Tianeptine $T_{1/2}$ = 2.5 hrs[21])	14/LO/0663	NCT04145076	SERT	Maudsley Hospital Pharmacy, London, UK
	Citalopram	20	16 (0)	17 (0)	180					
	Tianeptine	12.5	19 (0)	19 (0)	60				μ-opioid (MOR) & δ-opioid (DOR) receptor agonist	Servier Laboratories, Suresnes, France

MRI Data Acquisition

MRI acquisition information is provided in **Table 2** and includes the MR scanner, sequences to acquire both structural and functional scans, repetition time and echo time for each study included in analyses. Further information regarding MRI acquisition is in **Supplementary Table 2**.

Table 2. MRI acquisition information included in analyses. TR, repetition time; TE, echo time; ADNI GO, Alzheimer's Disease Neuroimaging Initiative Grand Opportunities[22] & echo planar imaging, EPI. Further information regarding MRI acquisition is in Supplementary Table 2.

Study ID	MR Scanner	Scan	Sequence	TR (ms)	TE (ms)
GABA$_{A/B}$	3T General Electric Discovery MR750, Centre for Neuroimaging Sciences, King's College London, UK	Structural	ADNI GO	7312	3016
		Functional	EPI	2500	12, 28, 44 & 60
Serotonin & opioid		Structural	ADNI GO	7312	3016
		Functional	EPI	2300	12.7, 31 & 48

MRI Data Preprocessing

For all studies the functional fMRI data was pre-processed using AFNI[23] (AFNI v21.1.07 (*https://afni.nimh.nih.gov/pub/dist/doc/htmldoc/#*) and Anaconda v3.8 (*https://anaconda.org/*) and consisted of de-spiking; slice time correction; co-registration of fMRI to T1 structural scan; normalisation to standard (MNI152_T1_2009c) space; tissue segmentation; optimal combination of echoes and motion correction using multi-echo ICA (tedana v0.0.8)[24]; smoothing (6mm³); scaling (to percent signal change with $\mu = 0$); band-pass filter (0.01 - 0.1Hz); regression of white matter and cerebrospinal fluid signal & censoring (if framewise displacement > 3mm, based on previous literature[8] and/or if motion outlier (i.e. >2 σ)), additional information regarding head motion in the sample is in **Supplementary Table 3**. The raw and pre-processed MRI data was inspected and checked for quality issues using simple image visualisation in FSL's FSLeyes in combination with examination of the reports that were automatically generated by AFNI. No scans were excluded from analyses for the AZD7325 (GABA$_A$) study; 6 scans from non-autistic participants (two placebo, 15 mg and 30 mg sessions) and 6 scans from autistic participants (2 placebo, 15 mg and 30 mg sessions) were excluded for the arbaclofen (GABA$_B$) study due to excessive signal loss; 3 scans from non-autistic participants (3 drug sessions) and 3 scans from autistic participants (3 drug sessions) were excluded from the serotonin (citalopram) study due to excessive motion and finally, 4 scans from non-autistic participants (2 placebo and 2 drug sessions) and 3 scans from autistic

participants (2 placebo and 1 drug session) were excluded from the tianeptine (opioid) study due to excessive signal loss (i.e. motion artifacts).

Functional Network Analysis

The pre-processed fMRI data for each drug was parcellated into 100 cortical brain regions of interest (ROIs) according to the Schaefer parcellation scheme[25]. The average time series in each parcel was used to construct functional connectivity matrices of Fisher-transformed Z values (or Fisher-transformed correlation coefficients) between every pair of ROIs for each individual scan using CONN v21a[26]. Each of the 100 ROIs were assigned into one of seven main functional networks according to the Yeo-7 atlas: default mode (DMN); frontoparietal (FPN); ventral attention (VAN) (or salience); dorsal attention (DAN); limbic (LN); somatomotor (SMN) & visual (VN) networks[25]. Based on this parcellation, within-network connectivity was defined as the mean connectivity between each ROI belonging to a Yeo-7 network. Between-network connectivity was defined as the mean connectivity between ROIs that comprises two different networks (**Fig. 1**).

Figure 1. Schematic of the uniform resting-state network functional connectivity analysis pipeline applied to fMRI data from all studies. Data were parcellated into 100 cortical regions (REF); the average BOLD timeseries extracted from every node was then used to construct a functional connectivity matrix (of Fisher transformed Z values, r-to-z) between each of the 100 nodes for every individual scanning session. As each node is assigned a resting-state network (REF), within- and between-network connectivity can be calculated as the mean connectivity of the ROIs belonging to a given network or between ROIs comprising two different networks, respectively.

Statistical Analyses

First, we tested the prediction that there would be no statistically significant differences within and between networks in the placebo condition between the autistic and non-autistic group. Then, for each drug we tested our main hypothesis that there would be a difference in the brain response to drug challenge in autistic and non-autistic individuals i.e. a dose × group interaction. To evaluate the statistical significance of the interaction between dose and group we fit a general linear mixed effects model (GLME) for all networks with two-sided permutation tests with 10,000 repetitions. The GLME used was given by the expression z̄ ~ B0 + Bdose*group + (1|Subject ID) – with Subject ID accounting for the random effect. This approach accommodates missing data as not all participants had a complete dataset including placebo and all drug conditions. Significance was assessed with two-sided permutation tests featuring 10,000 GLME repetitions. We report p values obtained from the permutation tests uncorrected, highlighting those surviving multiple comparison correction using the False Discovery Rate method with $\alpha = 5\%$[27]. Dose × group interaction p values were corrected across all within- and between-networks for each pharmacological probe. While each study was conducted separately, a small number of autistic participants took part in two or more studies (n = 7, 5 male and 2 female) and were included in the analysis. We report the within-network results first, then the between-network results per pharmacological probe.

To investigate the extent to which any interactions were driven predominantly by a shift in the autistic group or non-autistic group, we estimated the Cohen's d effect size[28] for connectivity changes within- and between every functional network against placebo in each group separately; where our placebo baseline included every placebo fMRI scan from all studies (total non-autistic, n = 67 & autistic, n = 54). This is represented by equation XX, where z̄ represents the average Fisher's Z for a given network and $\sigma(Z)$ the standard deviation of such network:

$$D = \frac{\overline{Z_{drug}} - \overline{Z_{placebo}}}{\sigma\left(Z_{placebo}\right)}$$

Finally, we examined whether drug challenge preferentially increases or decreases within- and between-network connectivity in the autistic brain, irrespective of pharmacological probe (and

underlying neurotransmitter system targeted). To do this, we pooled the within- and between-network effect sizes generated in the analysis described above and then performed non-parametric Mann Whitney U tests to compare the aggregate within- and between-network connectivity shift between groups.

All statistical analyses were conducted in R version 4.1.1 (2021-08-10) (R Core Team 2021) with supporting packages ggplot2[29], reshape2, dplyr, tidyr, tibble, shadowtext, lme4, and ptestr.

Results

A comprehensive neuropharmacological profile of changes in the functional connectivity of cortical networks was produced by collecting resting-state fMRI data using four different pharmacological probes of the $GABA_A$, $GABA_B$, serotonin and μ-opioid systems. All fMRI data was parcellated into the same 100 cortical regions[25]. The resulting correlation matrices were Fisher-transformed (r-to-z values) and average connectivity calculated for all within- and between-network interactions using the previously defined atlas (each of the 100 cortical regions from the matrices belongs to a given Yeo resting-state network) of seven cortical functional networks: the default-mode (DMN), frontoparietal (FPN), ventral attention (VAN) (which is also referred to as the 'salience network'), dorsal attention (DAN), limbic (LN), somatomotor (SMN) and visual networks (VN)[30]. An effect size (Cohen's d) was then obtained for within and between each functional network to represent the magnitude of the 'shift' from baseline (i.e. inactive placebo).

Autistic and non-autistic network connectivity is not significantly different at baseline

First, we examined the main effect of group to ascertain if there were baseline differences between autistic and non-autistic participants in our sample. Prior to correction for multiple comparisons (using the FDR method), only somatomotor-visual network between-network connectivity was significantly greater in the autistic group ($p_{unc} = 0.007$) (**Supplementary Fig.**

1). After correction for multiple comparisons, we observed no statistically significant differences between the groups (For the full main effect of group result at baseline (i.e. placebo) see **Supplementary Table 4**).

Neurotransmitter systems differentially 'shift' network connectivity in the autistic brain

Below we report the within-network dose × group interactions (**Fig. 2**) and then the between-network connectivity dose × group interactions for each drug challenge (**Fig. 3**). Key findings of differential shifts in within- and between-network connectivity between the non-autistic and autistic group for each drug challenge are also summarised in **Table 3.**

Figure 2. Within-network functional connectivity 'shifts' by neurotransmitter system targeted for the non-autistic and autistic groups. Values represent the effect sizes (Cohen's d) and significant dose × group interactions are shown with an asterisk (uncorrected p < 0.05*, p obtained with a two-sided permutation test), those that survive FDR correction are shown in bold and with a dagger (p < 0.05†). DAN, dorsal attention; DMN, default mode; FPN, frontoparietal; LN, limbic; VAN, ventral attention & VN, visual network.

Figure 3. Between-network functional connectivity 'shifts' by neurotransmitter system targeted for the non-autistic and autistic groups. Values represent the effect sizes (Cohen's d) and significant dose × group interactions are shown with an asterisk (uncorrected $p < 0.05$*, p obtained with a two-sided permutation test), those that survive FDR correction are shown in bold and with a dagger ($p < 0.05$†). DAN, dorsal attention; DMN, default mode; FPN, frontoparietal; LN, limbic; VAN, ventral attention & VN, visual network.

Targeting the GABA system

Targeting GABA$_A$:

Within-network connectivity:

Within-network connectivity was differentially shifted by GABA$_A$ activation in each group across several networks. In particular, the low dose increased connectivity of the visual network in non-autistic participants but it decreased in autistic participants.

Between-network connectivity:

In general, shifts in between-network connectivity elicited by GABA$_A$ activation were more pronounced in the autism group compared to non-autistic participants.

In non-autistic participants, GABA$_A$ activation by low and high dose AZD7325 elicited broadly similar shifts in between-network connectivity apart from between-network connectivity of the visual network, which was significantly reduced by the high dose only.

In autistic participants, GABA$_A$ activation by low and high dose AZD7325 elicited a broadly similar pattern of shifts in between-network connectivity but again, the exceptions were mostly in between-network connectivity of the visual network, specifically, increases in connectivity with the higher-order default mode and frontoparietal networks at the higher dose.

Together, GABA$_A$ activation primarily (but not exclusively) shifted within and between-network connectivity of the visual network. The most marked shifts in connectivity were present in the autistic group. Distinct decreases in between-network of higher-order networks (e.g. DMN) responses in the autism were evident at the low dose. By contrast, there were increases in connectivity between these networks at the high dose.

Targeting GABA$_B$:

Within-network connectivity:

Within-network responses to GABA$_B$ agonism were qualitatively similar in the autistic and non-autistic groups and at the low and high doses. Lower-order (visual) network connectivity tended to increase, and higher-order network connectivity tended to decrease in response to arbaclofen. However, within-network responses were more pronounced in the non-autistic group, particularly in the frontoparietal network.

Between-network connectivity:

There was a differential response in between network connectivity in autistic and non-autistic groups across the whole brain. In non-autistics, GABA$_B$ activation by low and high doses of arbaclofen predominantly decreased between-network connectivity. In autistics, low and high doses of arbaclofen predominantly increased between-network connectivity.

Thus, GABA$_B$ activation had a more prominent impact within-networks in non-autistic compared to autistic participants. However, there was an opposite effect on between-network connectivity across the brain in autistic (increased connectivity) compared to non-autistic participants (decreased connectivity). These group differences were more marked at the higher dose and particularly pronounced in higher-order and attentional/sensory networks.

Targeting the serotonin system

Within-network connectivity:

Within-network connectivity responses to serotonin activation were broadly similar in each group except for visual and ventral attention network connectivity which was increased in the non-autistic group and decreased in the autistic group.

Between-network connectivity:

Activation of serotonin signalling primarily elicited a decrease in between-network connectivity of higher order networks (such as default mode and frontoparietal) in non-autistic but not and autistic participants; indeed, serotonin activation significantly increased between-network connectivity of the default mode network in autism and did the opposite in non-autistic participants.

Targeting the opioid system

Within-network connectivity:

The direction of response to μ-opioid agonism within-networks was broadly similar in direction in both groups but of a greater magnitude in autistic participants. There was especially more prominent increase in somatomotor and decrease in limbic network connectivity in autistic participants compared to non-autistic participants. In contrast, ventral attention network connectivity was increased by μ-opioid activation in non-autistic participants groups but decreased in the autistic participants.

Between-network connectivity:

Again, there were differential effects in each group on between-network connectivity. In general, μ-opioid activation increased between-network connectivity of somatomotor and limbic networks in the autism group. Whereas it decreased connectivity of the higher order default mode and frontoparietal networks in non-autistic participants.

Thus, activation of the μ-opioid system has similar effects on within-network connectivity and differential effects on between-network connectivity, with differences in autism concentrated upon lower-order (e.g. somatomotor), limbic and attentional within and between-network connectivity.

Table 3. Key dose × group interaction findings for within- and between-network connectivity across all drug challenges.

Dose	Network	Non-Autistic Effect Size	Autistic Effect Size	P_{unc}	P_{FDR}
GABA_A					
Within-networks					
10 mg	VN	0.30	-0.36	<0.001	<0.001
Between-networks					
10 mg	VN-DMN	0.32	-0.20	<0.001	<0.001
20 mg	VN-DMN	-0.18	0.66	<0.001	0.001
20 mg	VN-FPN	-0.26	0.38	<0.001	<0.001
GABA_B					
Within-networks					
15 mg	FPN	-0.55	-0.10	<0.001	0.003
30 mg	FPN	-0.60	0.09	<0.001	<0.001
Between-networks					
30 mg	FPN-SMN	0.15	0.79	0.015	0.025
30 mg	FPN-VAN	-0.11	0.75	<0.001	<0.001
30 mg	FPN-DAN	-0.16	0.73	<0.001	<0.001
30 mg	DMN-SMN	-0.27	0.49	<0.001	<0.001
30 mg	DMN-VAN	-0.17	0.48	<0.001	<0.001
Serotonin					
Within-networks					
20 mg	SMN	0.42	0.81	<0.001	0.002
20 mg	VN	0.29	-0.35	<0.001	<0.001
20 mg	VAN	0.20	-0.19	0.008	0.022
Between-networks					
20 mg	DMN-VN	-0.56	0.43	<0.001	<0.001
μ-opioid					
Within-networks					
12.5 mg	SMN	0.52	1.32	<0.001	<0.001
12.5 mg	VAN	0.45	-0.18	<0.001	<0.001
Between-networks					
12.5 mg	LN-VAN	-0.33	0.42	0.002	0.005
12.5 mg	SMN-DAN	0.05	0.78	<0.001	<0.001

The divergent 'shift' of the autistic brain in response to drug challenge

Next, we investigated whether there was an overarching pattern to the way in which the autistic brain responded to drug challenge overall (irrespective of the neurotransmitter system targeted) by pooling together the effect sizes for each pharmacological probe as reported above. Overall, the autistic brain had a greater magnitude of shift in response to any drug, compared to the non-autistic brain (median: non-autistic, -0.035 & autistic, 0.08; interquartile range: non-autistic, 0.40 & autistic, 0.48; $p < 0.001$). A net positive shift in connectivity was observed in autism (mean effect size 0.11), and a slight decrease in connectivity in non-autistic participants (mean effect size -0.03).

Post-hoc analyses confirmed that this group-level difference in network responsivity was primarily driven by *increases* in *between-network* connectivity in autism compared to a minimal decrease in the non-autistic group (median: non-autistic, -0.09 & autistic, 0.11; interquartile range: non-autistic, 0.41 & autistic, 0.45; $p = <0.0001$); and there was no difference in within-network connectivity shifts between groups (median: non-autistic, 0.07 & autistic, -0.01; interquartile range: non-autistic, 0.39 & autistic, 0.38; $p = 0.20$). Ventral attention, frontoparietal, somatomotor and (to a lesser extent) dorsal attention network connectivity increases are the largest contributors to shifts in between-network connectivity in autism; whereas, default mode, limbic and visual network connectivity remained unchanged (or were slightly increased) (**Fig. 4**).

Figure 4. Aggregate between-network connectivity shifts (values represent Cohen's d effect size) for each resting-state network separately for both groups. A distinct pattern of increased connectivity in autism is observed, and largely reduced connectivity in non-autistic participants across networks in response to drug challenge. DAN, dorsal attention; DMN, default mode; FPN, frontoparietal; LN, limbic; VAN, ventral attention & VN, visual network.

Discussion

We have used a uniform analytical framework to capture the change in functional connectivity within and between seven brain networks following activation of several known targets of the $GABA_A$, $GABA_B$, serotonin and μ-opioid neurotransmitter systems in autistic and non-autistic adults. As predicted, despite minimal baseline group differences, functional connectivity measures were 'shifted' differently by each drug probe in autistic compared to non-autistic individuals.

There were some target-specific patterns; modulation of $GABA_A$ shifted visual network connectivity; $GABA_B$ shifted between-network connectivity across the brain; serotonin shifted sensory and higher order networks (e.g. default mode network); and μ-opioid shifted somatomotor and limbic network connectivity. However, we also found that, regardless of neurotransmitter system targeted, in general drugs elicited significantly greater increases in between-network connectivity in autistic participants compared to their non-autistic peers; especially between-network connectivity of the VAN, FPN and SMN, which are primarily networks which 'allow' bottom-up information to shift attention and control the initiation of behaviour[31,32].

Target-specific patterns of connectivity shift are different in autistic and non-autistic participants

We observed target-specific differences in functional connectivity responses in autism depending on the neurotransmitter system probed. The networks which shifted in response to each drug broadly aligned with the known expression pattern of their receptors but the direction and/or extent of shift in connections within and between these brain regions was different in autism.

GABA response differences

Autistic differences in network responsivity following $GABA_A$ activation were observed in lower-order sensory networks. The effect was most prominent in the visual cortex where $GABA_A$ receptor expression is reported to be particularly high[33]. Altered responsivity in autism may, in part, be explained by reduced levels of $GABA_A$ receptors in the autistic brain[34,35].

However, this evidence comes from post-mortem data, and the reported decrease in GABA receptors has been less consistent in the living human brain using PET imaging. Recent PET data suggests that cortical $GABA_A$ receptor levels may be no different in autistic and non-autistic adults[36,37].

There were more widespread autistic differences in between-network connectivity in response to $GABA_B$ activation, and this may reflect the broader extent of $GABA_B$ expression across the whole brain[33,38]. Again, this may, in part, be explained by reduced levels of $GABA_B$ receptors in the autistic brain[34,35]. For example, $GABA_B$ receptors may be differently expressed in prefrontal and parietal regions that comprise higher-order and attentional networks in which we see the largest differential shifts in response to arbaclofen here[39]. However, there are no reports to-date of altered $GABA_B$ receptor expression in autism from PET data. Thus, minimal baseline differences in GABA receptor availability may not entirely explain the diverging direction of shifts in connectivity of autistic and non-autistic resting-state networks in response to $GABA_{A/B}$ activation.

We suggest that the explanation may be functional differences in the GABA system. This is consistent with other evidence from preclinical animal work and humans. For example, agonism of $GABA_A$ and $GABA_B$ receptors ameliorates autistic features in an autism-relevant mouse model[40]; and we have shown that visual and auditory sensory processing in the same autistic and non-autistic adults who participated in the present study are also differentially modulated by $GABA_B$ receptor agonism[15,41].

Serotonin response differences
There were paradoxical responses to citalopram in the autistic brain. In contrast to non-autistic participants, within-network connectivity of the visual and ventral attention networks was reduced in the autistic brain, but connectivity between the visual and default mode network was increased.

In the visual network, there is evidence for lower SERT levels in the visual cortex in autism[42], several regions of the DMN also express SERT including the medial prefrontal cortex and posterior cingulate cortex (PCC)[43] and lower SERT binding has been reported in these regions in autistic adults[44]. However, it is difficult to align receptor availability differences with the range of within and between-network alterations elicited by citalopram in autism. The increase

in between-network connectivity of the DMN in autism in response to citalopram is noteworthy given that altered DMN connectivity is implicated in depression, which is more common in autistic people[45]. The augmented responses to SERT activation in autistic individuals across the default mode and sensory (e.g. visual) networks observed here aligns with evidence that autistic patients respond at lower doses to SSRIs[45] and SSRIs can be poorly tolerated by autistic people[46]. We do however acknowledge that this is a single dose in which citalopram was used as a pharmacological probe to elevate bulk serotonin levels to understand the role of the serotonin system and not intended to directly extend to the mechanistic underpinnings of anti-depressant efficacy in autistic and non-autistic people – where treatment with citalopram often requires longer term use (6-8 weeks) and may therefore have different network-level effects than those reported here.

μ-opioid response differences

Lastly, in autistic participants, agonism at the μ-opioid receptor by tianeptine altered within and between-network differently in autistics. Within-network connectivity was increased in somatomotor networks and decreased in limbic and ventral attention networks in autistics but not in non-autistics. Between-network connectivity of the ventral attention and limbic network was increased in autism and decreased in non-autistic participants.

μ-opioid receptors are expressed in several components of these networks with particularly high expression in the cingulate cortex of the limbic network[47] and in somatomotor regions[48]. Recent human PET data suggests that baseline differences in cortical μ-opioid receptor expression in autism are restricted to upregulation in the precuneus only[49], so again differences in receptor levels do not explain responsivity differences. Thus, as for other drug probes, what sets the autistic group apart is a distinct functional response of the brain to neurotransmitter perturbation.

These observed differences in the responsivity of resting-state networks across the neurotransmitter systems probed here may need to be understood in the context of neurodevelopment. Indeed, we have evidence for early alterations in functional connectivity in newborn infants who have a higher likelihood of a neurodevelopmental outcome such as autism[50]. This work observed that local functional connectivity in neonates with a higher likelihood of autism was greater in sensory cortices and limbic areas responsible for face processing[51]. We have more recently reported that moment-to-moment dynamic functional

connectivity profile of brain states that engage sensorimotor networks at birth predicts neurodevelopmental and autistic traits at 18 months[52]. Thus, by birth, there are already differences in the functional landscape of brain regions associated with autism. Although, capturing convincing baseline cross-sectional differences in functional connectivity in larger autistic and non-autistic cohorts assessed at later ages has been challenging[3]. We suggest that what persists in autism is not any average baseline difference in functional connectivity, but a true functional difference as defined by altered response to perturbation (in this case by pharmacological probe).

Consistent with this, we see no substantial baseline difference in resting-state network connectivity between groups of adults in our sample. It could be argued that this is unsurprising given our relatively small sample size but one recent mega-analysis used a sample of 1824 individuals, 796 of whom were autistic to report effects which were still relatively small[7]. What is more striking therefore, is that significant differences in functional connectivity responses to drug challenge were observed even with our small sample size.

Between network functional connectivity in autism increases in response to drug challenge

We pooled resting-state connectivity data across all the pharmacological probes to investigate if there was any generalised response profile common to the autistic group. We report that overall, drug challenge *increased between-network connectivity in autism*, but either had little impact or *decreased* between-network connectivity in non-autistic participants. The main drug-elicited increases in functional connectivity were mainly localised to between-network connectivity of the VAN, FPN and SMN, which are implicated in detecting unexpected stimuli to trigger attentional shift, flexible initiation of new tasks and the processes controlling motor output[53], respectively. Thus, to some extent they can be thought of as allowing a flexible response to external stimuli. Consequently, how neurotransmitter systems in autistic individuals regulate the engagement and disengagement of functional networks in response to changing task demands may be altered. By facilitating connectivity between-networks the drugs used here may promote more flexible responses to external stimuli. We emphasise

however that this is highly speculative and whether this profile has clinical meaning remains to be examined.

An additional or alternative explanation for the increased between-network shift induced by drug challenge in autistic participants may be that homeostasis is differently regulated in autism. For example, if networks remain in their shifted state for longer after drug challenge in autistic participants, we may simply be continuing to capture a functional shift in autistics which has moved back towards baseline in non-autistic participants by the time of data acquisition. In support of this, we have previously reported that limbic system activation takes longer to return to baseline in autism following citalopram during a fMRI face processing task[54]. This result was restricted to *between*-networks, so following perturbation local systems may return to baseline quicker compared to those *between*-networks that are more distal and/or more functionally segregated. Our previous work using other modalities to capture brain dynamics supports this position. We have used the aperiodic (1/f) slope of EEG to show that, despite no baseline group differences, autistic participants 'shifted' their 1/f slope at lower doses than non-autistic participants[55]. Placed together with the present analyses, the homeostatic mechanisms which might be at play in non-autistic people to limit shifts in brain function are not so evident in autism. MRI does not allow us to address this question but there is some evidence from cellular models that synaptic scaling to adjust excitatory or inhibitory synaptic strength up or down to stabilise neural firing rates[56] is disrupted in animal models of autism[57].

We acknowledge several limitations to our work. First, in humans we are often limited to non-selective pharmacological probes (i.e. with more than one molecular target). While both GABA probes used here were relatively selective compounds. For example, tianeptine also has agonist activity at delta (δ) opioid receptors[58] and indirect effects on other neurotransmitter systems (e.g. glutamate[59] and serotonin[60]) which may also contribute to its impact on functional connectivity. Second, our approach included different MRI acquisition sequences adopted between studies (SERT/ μ-opioid versus the GABA probes). The impact of this was somewhat mitigated by using a consistent preprocessing pipeline across all studies. Similarly, an atlas-based parcellation approach was used to provide a uniform analysis framework across studies/probes. Third, motion artifacts (i.e. due to participants moving in the scanner) can alter functional connectivity measures[61]. As a result, our preprocessing pipeline included several techniques focused on motion correction specifically (e.g. despiking, multi-echo ICA

correction, censoring). There was also stringent post-processing quality control and datasets with significant movement (or other quality control issues) were omitted from analyses. Fourth, our sample size for each pharmacological probe is relatively small and the sexes are also not balanced (only males were recruited for the studies targeting SERT and the μ-opioid receptor), therefore more work including female needs to be done. This is a recognised problem in past pharmacological research[62] and in our current studies our goal is to recruit an equal number of men and women. Despite a small sample size for each probe our statistical power was increased due to the within-subject repeated-measures design of our studies.

In conclusion, our findings suggest that the modulation of resting-state network connectivity by multiple key neurotransmitter systems is different in autistic and non-autistic participants. These systems are targeted by medications used to treat neurological and psychiatric conditions experienced by neurodivergent people. Our work indicates that the autistic brain is 'pharmacologically distinct'. This is crucial because autistic people have a higher incidence of psychiatric and neurological conditions[63,64] but drug development and trials for mental health problems do not consider neurodivergence. These results suggest this needs to change. Investigating if these drug-induced 'shifts' in functional connectivity can help provide better targeted interventions will be an important next translational step.

References

1. Loth, E. *et al.* The meaning of significant mean group differences for biomarker discovery. *PLoS Comput Biol* **17**, e1009477 (2021).

2. Mottron, L. & Gagnon, D. Prototypical autism: New diagnostic criteria and asymmetrical bifurcation model. *Acta Psychologica* **237**, 103938 (2023).

3. Uddin, L. Q., Supekar, K. & Menon, V. Reconceptualizing functional brain connectivity in autism from a developmental perspective. *Front. Hum. Neurosci.* **7**, (2013).

4. Müller, R.-A. *et al.* Underconnected, but How? A Survey of Functional Connectivity MRI Studies in Autism Spectrum Disorders. *Cerebral Cortex* **21**, 2233–2243 (2011).

5. Whelan, T. P. *et al.* Editorial Perspective: Bridging the translational neuroscience gap in autism – development of the 'shiftability' paradigm. *Child Psychology Psychiatry* **65**, 862–865 (2024).

6. Mišić, B. & Sporns, O. From regions to connections and networks: new bridges between brain and behavior. *Current Opinion in Neurobiology* **40**, 1–7 (2016).

7. Ilioska, I. *et al.* Connectome-wide Mega-analysis Reveals Robust Patterns of Atypical Functional Connectivity in Autism. *Biological Psychiatry* **94**, 29–39 (2023).

8. Holiga, Š. *et al.* Patients with autism spectrum disorders display reproducible functional connectivity alterations. *Sci. Transl. Med.* **11**, eaat9223 (2019).

9. Fan, C. *et al.* Transcriptomics of Gabra4 knockout mice reveals common NMDAR pathways underlying autism, memory, and epilepsy. *Molecular Autism* **11**, 13 (2020).

10. Schuler, V. *et al.* Epilepsy, Hyperalgesia, Impaired Memory, and Loss of Pre- and Postsynaptic GABAB Responses in Mice Lacking GABAB(1). *Neuron* **31**, 47–58 (2001).

11. Muller, C. L., Anacker, A. M. J. & Veenstra-VanderWeele, J. The serotonin system in autism spectrum disorder: From biomarker to animal models. *Neuroscience* **321**, 24–41 (2016).

12. Becker, J. A. *et al.* Autistic-Like Syndrome in Mu Opioid Receptor Null Mice is Relieved by Facilitated mGluR4 Activity. *Neuropsychopharmacol* **39**, 2049–2060 (2014).

13. Pellissier, L. P., Gandía, J., Laboute, T., Becker, J. A. J. & Le Merrer, J. μ opioid receptor, social behaviour and autism spectrum disorder: reward matters. *British J Pharmacology* **175**, 2750–2769 (2018).

14. Marotta, R. *et al.* The Neurochemistry of Autism. *Brain Sciences* **10**, 163 (2020).

15. Huang, Q. *et al.* GABA $_B$ receptor modulation of visual sensory processing in adults with and without autism spectrum disorder. *Sci. Transl. Med.* **14**, eabg7859 (2022).

16. Wichers, R. H. *et al.* Modulation of atypical brain activation during executive functioning in autism: a pharmacological MRI study of tianeptine. *Molecular Autism* **12**, 14 (2021).

17. Wong, N. M. L. *et al.* Serotonin differentially modulates the temporal dynamics of the limbic response to facial emotions in male adults with and without autism spectrum disorder (ASD): a randomised placebo-controlled single-dose crossover trial. *Neuropsychopharmacol.* **45**, 2248–2256 (2020).

18. Chen, X. *et al.* The central nervous system effects of the partial GABA - A $α_{2,3}$ -selective receptor modulator AZD7325 in comparison with lorazepam in healthy males. *Brit J Clinical Pharma* **78**, 1298–1314 (2014).

19. Sanchez-Ponce, R. *et al.* Metabolic and Pharmacokinetic Differentiation of STX209 and Racemic Baclofen in Humans. *Metabolites* **2**, 596–613 (2012).

20. Sangkuhl, K., Klein, T. E. & Altman, R. B. PharmGKB summary: citalopram pharmacokinetics pathway. *Pharmacogenetics and Genomics* **21**, 769–772 (2011).

21. Salvadori, C., Ward, C., Defrance, R. & Hopkins, R. The pharmacokinetics of the antidepressant tianeptine and its main metabolite in healthy humans – influence of alcohol co-administration. *Fundamemntal Clinical Pharma* **4**, 115–125 (1990).

22. Jack, C. R. *et al.* Magnetic resonance imaging in Alzheimer's Disease Neuroimaging Initiative 2. *Alzheimer's & Dementia* **11**, 740–756 (2015).

23. Cox, R. W. AFNI: Software for Analysis and Visualization of Functional Magnetic Resonance Neuroimages. *Computers and Biomedical Research* **29**, 162–173 (1996).

24. Kundu, P. *et al.* Integrated strategy for improving functional connectivity mapping using multiecho fMRI. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 16187–16192 (2013).

25. Schaefer, A. *et al.* Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. *Cerebral Cortex* **28**, 3095–3114 (2018).

26. Whitfield-Gabrieli, S. & Nieto-Castanon, A. *Conn* : A Functional Connectivity Toolbox for Correlated and Anticorrelated Brain Networks. *Brain Connectivity* **2**, 125–141 (2012).

27. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B: Statistical Methodology* **57**, 289–300 (1995).

28. Sawilowsky, S. S. New Effect Size Rules of Thumb. *J. Mod. App. Stat. Meth.* **8**, 597–599 (2009).

29. Wickham, H. *Ggplot2*. (Springer International Publishing, Cham, 2016). doi:10.1007/978-3-319-24277-4.

30. Thomas Yeo, B. T. *et al.* The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *Journal of Neurophysiology* **106**, 1125–1165 (2011).

31. Rosazza, C. & Minati, L. Resting-state brain networks: literature review and clinical applications. *Neurol Sci* **32**, 773–785 (2011).

32. Seeley, W. W. *et al.* Dissociable Intrinsic Connectivity Networks for Salience Processing and Executive Control. *J. Neurosci.* **27**, 2349–2356 (2007).

33. Calver, A. R. *et al.* The expression of GABAB1 and GABAB2 receptor subunits in the cNS differs from that in peripheral tissues. *Neuroscience* **100**, 155–170 (2000).

34. Fatemi, S. H. *et al.* mRNA and Protein Levels for GABAAα4, α5, β1 and GABABR1 Receptors are Altered in Brains from Subjects with Autism. *J Autism Dev Disord* **40**, 743–750 (2010).

35. Oblak, A., Gibbs, T. T. & Blatt, G. J. Decreased GABA $_A$ receptors and benzodiazepine binding sites in the anterior cingulate cortex in autism. *Autism Research* **2**, 205–219 (2009).

36. Fung, L. K. *et al.* Thalamic and prefrontal GABA concentrations but not GABAA receptor densities are altered in high-functioning adults with autism spectrum disorder. *Mol Psychiatry* **26**, 1634–1646 (2021).

37. Horder, J. *et al.* GABA $_A$ receptor availability is not altered in adults with autism spectrum disorder or in mouse models. *Sci. Transl. Med.* **10**, eaam8434 (2018).

38. Billinton, A. *et al.* GABAB receptor heterodimer-component localisation in human brain. *Molecular Brain Research* **77**, 111–124 (2000).

39. Fatemi, S. H., Reutiman, T. J., Folsom, T. D. & Thuras, P. D. GABAA Receptor Downregulation in Brains of Subjects with Autism. *J Autism Dev Disord* **39**, 223–230 (2009).

40. Yang, J.-Q., Yang, C.-H. & Yin, B.-Q. Combined the GABA-A and GABA-B receptor agonists attenuates autistic behaviors in a prenatal valproic acid-induced mouse model of autism. *Behavioural Brain Research* **403**, 113094 (2021).

41. Huang, Q. *et al.* Exploratory evidence for differences in GABAergic regulation of auditory processing in autism spectrum disorder. *Transl Psychiatry* **13**, 320 (2023).

42. Nakamura, K. *et al.* Brain Serotonin and Dopamine Transporter Bindings in Adults With High-Functioning Autism. *Arch Gen Psychiatry* **67**, 59 (2010).

43. Beliveau, V. *et al.* A High-Resolution *In Vivo* Atlas of the Human Brain's Serotonin System. *J. Neurosci.* **37**, 120–128 (2017).

44. Andersson, M. *et al.* Serotonin transporter availability in adults with autism—a positron emission tomography study. *Mol Psychiatry* **26**, 1647–1658 (2021).

45. Howes, O. D. *et al.* Autism spectrum disorder: Consensus guidelines on assessment, treatment and research from the British Association for Psychopharmacology. *J Psychopharmacol* **32**, 3–29 (2018).

46. King, B. H. *et al.* Lack of Efficacy of Citalopram in Children With Autism Spectrum Disorders and High Levels of Repetitive Behavior: Citalopram Ineffective in Children With Autism. *Arch Gen Psychiatry* **66**, 583 (2009).

47. Kantonen, T. *et al.* Interindividual variability and lateralization of μ-opioid receptors in the human brain. *NeuroImage* **217**, 116922 (2020).

48. Sun, L. *et al.* Mu-opioid receptor system modulates responses to vocal bonding and distress signals in humans. *Phil. Trans. R. Soc. B* **377**, 20210181 (2022).

49. Noppari, T. *et al.* Dopamine D2R and opioid MOR availability in autism spectrum disorder. Preprint at https://doi.org/10.1101/2024.04.09.588651 (2024).

50. Ciarrusta, J. *et al.* Emerging functional connectivity differences in newborn infants vulnerable to autism spectrum disorders. *Transl Psychiatry* **10**, 131 (2020).

51. Ciarrusta, J. *et al.* Social Brain Functional Maturation in Newborn Infants With and Without a Family History of Autism Spectrum Disorder. *JAMA Netw Open* **2**, e191868 (2019).

52. França, L. G. S. *et al.* Neonatal brain dynamic functional connectivity in term and preterm infants and its association with early childhood neurodevelopment. *Nat Commun* **15**, 16 (2024).

53. Seitzman, B. A., Snyder, A. Z., Leuthardt, E. C. & Shimony, J. S. The State of Resting State Networks. *Topics in Magnetic Resonance Imaging* **28**, 189–196 (2019).

54. Wong, N. M. L. *et al.* Serotonin differentially modulates the temporal dynamics of the limbic response to facial emotions in male adults with and without autism spectrum disorder (ASD): a randomised placebo-controlled single-dose crossover trial. *Neuropsychopharmacol.* **45**, 2248–2256 (2020).

55. Ellis, C. L. *et al.* The dynamically neurodiverse human brain: Measuring excitatory-inhibitory dynamics and identifying homeostatic differences in autistic and non-autistic people. Preprint at https://doi.org/10.1101/2023.06.19.23291507 (2023).

56. Turrigiano, G. Homeostatic Synaptic Plasticity: Local and Global Mechanisms for Stabilizing Neuronal Function. *Cold Spring Harbor Perspectives in Biology* **4**, a005736–a005736 (2012).

57. Tatavarty, V. *et al.* Autism-Associated Shank3 Is Essential for Homeostatic Compensation in Rodent V1. *Neuron* **106**, 769-777.e4 (2020).

58. Gassaway, M. M., Rives, M.-L., Kruegel, A. C., Javitch, J. A. & Sames, D. The atypical antidepressant and neurorestorative agent tianeptine is a μ-opioid receptor agonist. *Transl Psychiatry* **4**, e411–e411 (2014).

59. Kole, M. H. P., Swan, L. & Fuchs, E. The antidepressant tianeptine persistently modulates glutamate receptor currents of the hippocampal CA3 commissural associational synapse in chronically stressed rats. *Eur J of Neuroscience* **16**, 807–816 (2002).

60. Datla, K. P. & Curzon, G. Behavioural and neurochemical evidence for the decrease of brain extracellular 5-HT by the antidepressant drug tianeptine. *Neuropharmacology* **32**, 839–845 (1993).

61. Mahadevan, A. S., Tooley, U. A., Bertolero, M. A., Mackey, A. P. & Bassett, D. S. Evaluating the sensitivity of functional connectivity measures to motion artifact in resting-state fMRI data. *NeuroImage* **241**, 118408 (2021).

62. Sosinsky, A. Z. *et al.* Enrollment of female participants in United States drug and device phase 1–3 clinical trials between 2016 and 2019. *Contemporary Clinical Trials* **115**, 106718 (2022).

63. Pan, P.-Y., Bölte, S., Kaur, P., Jamil, S. & Jonsson, U. Neurological disorders in autism: A systematic review and meta-analysis. *Autism* **25**, 812–830 (2021).

64. Underwood, J. F. G. *et al.* Neurological and psychiatric disorders among autistic adults: a population healthcare record study. *Psychol. Med.* **53**, 5663–5673 (2023).

Acknowledgements

Competing Interests

T.P.W. holds shares in Compass Pathfinder Ltd. N.A.P. has consulted for Deerfield Discovery and Research. D.G.M.M. has consulted for Jaguar Gene Therapy LLC. G.M.M. has received funding for investigator-initiated studies from GW Pharmaceuticals and Compass Pathfinder Ltd. G.M.M. has consulted for Greenwich Biosciences, Inc. L.G.S.F., M.D., C.M.P., H.V., A.C.P., C.L.E., G.I., J.L.F., R.H.W., F.M.P., J.K., N.M., N.M.L.W., D.B. and E.D. have no conflicts to declare.

Author Contributions

T.P.W. performed data preprocessing and analysis, prepared and drafted the manuscript. L.G.S.F. contributed to data analysis and helped draft the manuscript. M.D. contributed to data collection and preprocessing, and helped draft the manuscript. C.M.P., H.V., A.C.P., C.L.E., J.L.F., R.H.W., F.M.P., J.K., N.M.L.W. contributed to data collection. G.I. provided pharmacy services for the studies. N.M. contributed to preprocessing and data analysis. D.B. and N.A.P. edited the manuscript and provided input on data analysis. E.D., D.G.M.M. and G.M.M. conceived the idea for the studies; contributed to study design and drafted the manuscript. All authors reviewed and approved the final manuscript.

Funding

The authors also receive support from EU-AIMS (European Autism Interventions)/EU AIMS-2-TRIALS, an Innovative Medicines Initiative Joint Undertaking under Grant Agreement No. 777394. In addition, this paper represents independent research part funded by the infrastructure of the National Institute for Health Research (NIHR) Mental Health Biomedical Research Centre (BRC) at South London and Maudsley NHS Foundation Trust and King's College London, and the Medical Research Council Centre for Neurodevelopmental Disorders. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Data and Material Availability

Data from this study is available upon request.

Supplementary Information

Supplementary Table 1. Demographic information of the sample. Values in brackets represent the standard deviation.

GABA: AZD Study				
	Non-Autistic	**Autistic**	**statistic**	**p value**
Age	35 (± 13)	34 (± 10)	t = 0.12	p = 0.91
IQ	120 (± 13)	112 (± 12)	t = 1.53	p = 0.14
GABA: Arbaclofen Study				
	Non-Autistic	**Autistic**	**statistic**	**p value**
Age	28 (± 8)	34 (± 10)	-2.64	p = 0.01
IQ	120 (± 10)	117 (± 10)	1.33	p = 0.19
Serotonin & Opioid Study				
	Non-Autistic	**Autistic**	**statistic**	**p value**
Age	26 (± 7)	29 (± 10)	w = 150.5	p = 0.39
IQ	114 (± 10)	111 (± 16)	t = 0.66	p = 0.52

Supplementary Table 2. Additional MRI acquisition information.

Study ID	Scan	Inversion Time (ms)	Flip Angle (o)	Field of View (mm)	Matrix Size (no. of voxels)	No. of Sagittal Slices	Axial Slices per Volume	Slice Thickness (mm)	Interslice Gap (mm)	Voxel size	No. of volumes
GABA	Structural	400	11	270	256 x 256	196	-	1.2	0	1.05 x 1.05 x 1.2 mm	1
	Functional	0	80	240	64 x 64	-	32	3.0	1.0	3.75 x 3.75 x 3.0 mm	192 (length = 8 mins)
Serotonin & Opioid	Structural	400	11	270	256 x 256	196	-	1.2	0	1.055 x 1.055 x 1.2 mm	1
	Functional	0	90	240	64 x 64	-	33	3.8	0.4	3.75 x 3.75 x 4.5 mm	215 (length = 8.24 mins)

Supplementary Table 3. Degree of head motion in the non-autistic and autistic groups across all studies. Mean FD, mean framewise displacement.

Study	Condition	Dosage (mg)	Non-Autistic Mean FD (mm)	Autistic Mean FD (mm)	Test Statistic	P Value
GABA	Placebo	0	0.07 (± 0.03)	0.07 (± 0.04)	59	1.00
	AZD	10	0.07 (± 0.05)	0.10 (± 0.10)	57	0.642
	AZD	20	0.06 (± 0.03)	0.13 (± 0.14)	62	0.583
	Placebo	0	0.15 (± 0.15)	0.18 (± 0.09)	105	0.117
	Arbaclofen	15	0.15 (± 0.12)	0.22 (± 0.16)	121	0.156
	Arbaclofen	30	0.16 (± 0.15)	0.19 (± 0.15)	212	0.326
Serotonin & opioid	Placebo	0	0.10 (± 0.05)	0.15 (± 0.08)	97	0.026
	Citalopram	20	0.12 (± 0.06)	0.12 (± 0.06)	49	0.528
	Tianeptine	12.5	0.11 (± 0.04)	0.13 (± 0.05)	110	0.106

Supplementary Table 4. The main effect of group showing differences in connectivity of all resting-state networks between autistic and non-autistic participants at baseline (placebo). P values were corrected for multiple comparisons across all networks.

Network		P_{unc}	P_{FDR}
SMN	VN	0.0066	0.1848
SMN	SMN	0.054	0.54145
LN	SMN	0.0717	0.54145
VAN	VN	0.0807	0.54145
VAN	VAN	0.0986	0.54145
DAN	DAN	0.1233	0.54145
SMN	VAN	0.1492	0.54145
DMN	LN	0.1547	0.54145
LN	LN	0.2466	0.6776
LN	VAN	0.2751	0.6776
DAN	LN	0.2929	0.6776
DAN	SMN	0.3099	0.6776
FPN	LN	0.3146	0.6776
DMN	FPN	0.3843	0.726075
DMN	DMN	0.4072	0.726075
FPN	FPN	0.4149	0.726075
DAN	VN	0.4613	0.75978
FPN	VAN	0.5147	0.80064
FPN	VN	0.5708	0.841179
DMN	SMN	0.6525	0.9135
DAN	VAN	0.708	0.923364
FPN	SMN	0.7255	0.923364
DAN	DMN	0.7674	0.934226
VN	VN	0.8265	0.950432
LN	VN	0.8486	0.950432
DAN	FPN	0.9452	0.9738
DMN	VN	0.9518	0.9738
DMN	VAN	0.9738	0.9738

Supplementary Figure 1. Baseline difference in somatomotor-visual network connectivity between the non-autistic and autistic groups.