A multicentre prospective observational study to investigate the
prevalence and short-term impact of frailty, sarcopenia,
anaemia and multiple long term health conditions in chronic
limb threatening ischaemia (CLTI) - The FraiLTI (Frailty in
Chronic Limb Threatening Ischaemia) Study

Ashwin Sivaharan*[1], Blaise Hickson*[2], Tamer El-Sayed[1], Lauren Shelmerdine[1], Sai Wunnava[1], Emily
James[3,4], Craig I Nesbitt[1], Alasdair O'Doherty[5,6], Miles D Witham[7,8], Sandip Nandhra[1,9], and the FraiLTI
collaborative

*Authors share co-first authorship

Author affiliations:

1. Northern Vascular Centre, Freeman Hospital, Freeman Rd, High Heaton, Newcastle upon
Tyne, NE7 7DN
2. The Medical School, Newcastle University, Framlington Place, Newcastle upon Tyne, NE2 4HH
3. NIHR Leicester Biomedical Research Centre, Leicester, UK
4. Diabetes Research Centre, University of Leicester, Leicester, UK
5. Department of Sport and Exercise Sciences, Durham University, Green Ln, Durham, DH1 3LA
6. Department of Sport, Exercise and Rehabilitation, Northumbria University, Newcastle upon
Tyne, NE1 8ST
7. AGE Research Group, Translational and Clinical Research Institute, Faculty of Medical
Sciences, Newcastle University, Newcastle upon Tyne, UK
8. NIHR Newcastle Biomedical Research Centre, Newcastle upon Tyne Hospitals NHS
Foundation Trust, Cumbria Northumberland Tyne and Wear NHS Foundation Trust and
Faculty of Medical Sciences Newcastle University, Newcastle upon Tyne, UK
9. Population Health Sciences Institute, Faculty of Medical Sciences, Newcastle University,
Baddiley-Clark Building, Newcastle upon Tyne, NE2 4AX

Short title: The FraiLTI (Frailty in Chronic Limb Threatening Ischaemia) Study

Funding:

This work was funded by the National Institute for Health and Care Research (NIHR) Newcastle
Biomedical Research Centre (reference: IS_BRC_1215_20001). The views expressed in this
publication are those of the authors and do not necessarily reflect the views of the National Institute
for Health and Care Research or the Department of Health and Social Care.

Newcastle Hospital Charities

Corresponding Author:

Mr Ashwin Sivaharan (a.sivaharan@nhs.net)

Northern Vascular Centre, Freeman Hospital, Freeman Rd, High Heaton, Newcastle upon Tyne, NE7
7DN

Preprint doi: https://doi.org/10.1101/2024.11.11.24317086

Category: Observational study

Keywords:

Chronic limb threatening ischaemia (CLTI), frailty, sarcopenia, multiple long-term health

conditions, anaemia

No conflicts of interest to declare

Word Count

Abstract: 288

Main text: 2766

Article Highlights:

Type of research:
Multi-centre, prospective, observational study.
12
13   Key Findings:
14
15   The prevalence of frailty (51.8%), anaemia (53.8%), and sarcopenia (33.3%) is high in people with
16   CLTI. Most patients with CLTI had multiple long-term conditions (MLTC) (67.9%) or polypharmacy
17   (79.8%).
18
19   Take home Message:
20   This study shows that frailty, sarcopenia and anaemia are highly prevalent in those with CLTI.
21   The management of CLTI patients extends beyond revascularisation alone.
22
23
24   *Table of contents Summary:*

25   This study shows that frailty, sarcopenia and anaemia are highly prevalent in those with CLTI.
26   The management of CLTI patients extends beyond revascularisation alone.

Collaborators

Newcastle upon Tyne Hospitals NHS Foundation Trust

- Mr Bence Baljer (0000-0002-5805-8067)
- Mr Dilraj Bhullar (0000-0001-5190-5545)
- Mr Mohammed Fageer
- Miss Rana Khalil (0000-0002-9383-2986)
- Mr Lawrence Ugwumba (0009-0008-4197-1811)
- Ms Sarah Sillito (0000-0002-5768-5202)
- Mr Simon Jones (0000-0002-0101-3407)
- Dr Edward Buckley

South Tees Hospitals NHS Foundation Trust

- Miss Fatima Mannan (0000-0002-5505-9857)
- Mr Kamel Mekhaeil

University Hospitals Birmingham NHS Foundation Trust

- Mr Hosaam Nasr
- Mr Arooj Syed (0000-0002-1951-2036)
- Miss Emma Valums
- Miss Sarah Harley
- Mr Clive Allen

NHS Lothian

- Mr Andrew Tambyraja (0000-0002-9393-6291)
- Miss Sharon Chan

University Hospitals of Leicester NHS Trust

- Professor Athanasios Saratzis (0000-0002-3399-094X)
- Mr John Houghton (0000-0002-9159-5307)
- Professor Robert Sayers (0000-0002-6889-4659)
- Professor Matthew J Bown (0000-0002-6180-3611)
- Mr Badri Vijaynagar

Hull University Teaching Hospitals NHS Trust

- Professor Ian Chetter (0000-0002-2566-6859)
- Miss Louise Hitchman (0000-0001-9681-6749)
- Mr Abduraheem Mohamed (0000-0002-8892-3293)
- Mr Arthur Lim (0000-0003-0011-1336)

Abstract

Background

CLTI is a life and limb threatening condition that is associated with a gradual overall decline in health which remains under-explored. We explore the prevalence of correlated conditions (frailty, sarcopenia and anaemia) in patients with CLTI and assess their ability to perform activities of daily living (ADLs) alongside multiple long term health conditions (MLTC), polypharmacy and clinical outcomes.

Methods

FraiLTI was a UK multi-centre, prospective, observational study, evaluating the prevalence of frailty (Rockwood Clinical Frailty score of ≥5), sarcopenia based on low handgrip strength (<27kg for men and <16kg for women). and anaemia in patients admitted to hospital with chronic limb threatening ischaemia (CLTI). Other outcomes included prevalence of polypharmacy (≥5 medications) and MLTC (≥2 long-term health conditions). Recruitment took place from October 2021 to October 2022 in six UK centres supported by the Vascular and Endovascular Research Network and funded by the National Institute for Health and Care Research (NIHR) Biomedical Research Centre (BRC) North-East.

Results

84 patients were included in the study (54 [64.3%] men), with a mean (SD) age of 71.5 (11) years. 43/83 (51.8%) patients were living with frailty. 27/81 (33.3%) patients had probable sarcopenia 43/80 (53.8%) patients had anaemia of which 35 (92.1%) had normocytic anaemia. There was no difference in 90-day amputation-free survival for those living with or without frailty (97.6% vs 94.8%, p=0.78), sarcopenia (100% vs 94.4%, p=0.58), or anaemia (95.1% vs 97.3% vs p=0.45). MLTC were present in 57 (67.9%) patients and 67 (79.8%) patients met the definition of polypharmacy.

Conclusion

Frailty, sarcopenia and anaemia are highly prevalent in those with CLTI but none of these factors were associated with amputation free survival in this study. This has highlighted that the management of CLTI patients extends beyond revascularisation alone.

Introduction

Chronic Limb Threatening Ischaemia (CLTI) has been highlighted as a research priority by both patients and clinicians, including in a recent James Lind Alliance (JLA) priority-setting partnership.[1] CLTI is associated with a 3-year mortality rate of up to 40%, an increased risk of limb loss,[2] and has a negative impact on patients' quality of life.[3] Patients with CLTI often have many associated conditions that might worsen outcomes.

Frailty is defined as an age-related multisystem impairment in homeostasis that leaves patients vulnerable to stressors such as illness, trauma, or surgery[4]. Frailty is related to nutritional deficiencies, the presence of multiple chronic conditions, hormonal changes and increased systemic inflammation.[5,6] Frailty has been associated with worse outcomes in a wide range of surgical disciplines[7,8], including vascular surgery.[9,10] Sarcopenia is a distinct, but related concept, that describes age-related loss of muscle strength, muscle mass and muscle function.[11] Sarcopenia forms an important component of the physical frailty syndrome [12] and has been associated with worse survival following endovascular surgery[13] and surgical revascularisation.[14]

Multiple long-term conditions (MLTC; sometimes referred to as multimorbidity) may also be a contributing factor to outcomes in this population, both with respect to cardiovascular and non-cardiovascular conditions and has been associated with poorer outcomes in multiple surgical subspecialities.[15] There are also retrospective data that suggest that anaemia, a potential marker of overall health state, is associated with worse outcomes in lower limb vascular bypass patients.[16]

Currently there exists no prospective data that describes the prevalence of these associated conditions in the population with CLTI.

FraiLTI (Frailty in chronic Limb Threatening Ischaemia) was a UK multi-centre prospective observational study aimed at evaluating the prevalence and impact of frailty (and associated conditions) on outcomes for patients presenting to hospital with CLTI.

Methods

Study design & setting

This was a multicentre, prospective observational study was conducted from October 2021 to October 2022, led by a Newcastle University based Principal Investigator team and supported by the Vascular and Endovascular Research Network (VERN). Recruitment was undertaken in 6 UK Vascular Centres, including, Newcastle upon Tyne Hospitals NHS Foundation Trust, South Tees Hospitals NHS Foundation Trust, University Hospitals Birmingham NHS Foundation Trust, University Hospitals of Leicester NHS Trust, Hull University Teaching Hospitals NHS Trust and NHS Lothian. The study was funded by the National Institute of Health Research Biomedical Research Centre (NIHR-BRC) and the Newcastle Hospitals Charity with full ethical approval (21/PR/0750 REC by the London - Hampstead NHS Research Ethics Committee). The study is registered on the ISRCTN registry (ISRCTN18644880) and the protocol has been published in full.[17] The manuscript has been prepared as per the STROBE (Strengthening the reporting of observational studies in epidemiology) guidelines for the reporting of observational studies.[18]

Participants

All patients admitted to UK vascular centres, with CLTI as per consensus definition (the presence of peripheral artery disease (PAD) in combination with rest pain, gangrene, or a lower limb ulceration >2 weeks duration)[19] were invited to participate in the study across six sites. Patients were excluded if they were younger than 18 years, pregnant, or unable to consent to participation.

Outcomes

The main outcomes assessed included frailty (see definitions), sarcopenia and anaemia. Other outcomes included walk speed, self-reported low physical activity, multiple long-term conditions and polypharmacy. Key clinical outcomes (measured at 90 days) included observed clinical care pathway (revascularisation/amputation etc.), mortality, major adverse limb events (MALEs), major adverse cardiovascular events (MACE), readmissions, reinterventions and other complications (including

50   respiratory, urinary, and wound-related). Interaction between the main outcomes and the clinical

51   outcomes were then analysed (see statistical analysis section).

52   Definitions

Frailty was assessed using the Rockwood CFS. A Rockwood CFS of ≥5 was considered frail (a score of

5 denotes mild frailty).[20] Sarcopenia was assessed using grip strength, measured using a hand

dynamometer (Jamar Smart Digital Hand Dynamometer). Participants were seated with the elbow

flexed at ninety degrees and the wrist positioned with the thumb facing upwards. Six readings were

taken in total (3 from each hand), and the highest reading was used for analysis. Grip strength of

<27kg for men and <16kg for women was indicative of probable sarcopenia.[11]

Anaemia was defined as a haemoglobin of <130g/L in men and <120g/L in women as per the World

Health Organisation (WHO) definition,[21] based on the most recent pre-operative blood tests.

A low walk speed was diagnosed if the speed was <0.8m/s was noted on a 4m walk test. If the

patient was unable to complete the test, this was considered diagnostic of low walk speed.

Low physical activity was measured using four activity questions used in the English Longitudinal

Study of Ageing.[22] The methodology used to diagnose low physical activity based on these questions

has previously been published.[23]

MLTC was defined as ≥2 long-term health conditions (including ischaemic heart disease, previous

stroke/ TIA, heart failure, atrial fibrillation, hypothyroidism, dementia, anxiety and depression,

Parkinson's disease, chronic obstructive pulmonary disease, asthma, osteoarthritis and any other

inflammatory arthropathies).

Polypharmacy was defined as ≥5 medications, with excessive polypharmacy defined as ≥10

medications.[24]

Major adverse limb events (MALE) were defined as above ankle amputation of the index limb

or major vascular reintervention.[25] MACE was defined as a composite of acute myocardial infarction,

stoke or cardiovascular death.[26] Any lower limb reintervention or readmission for any cause within 90

days was recorded.

Data collection

Anonymised data included patient demographics, previous treatments, and hospital admissions from

the past six months. The research teams gathered this information prospectively using both

electronic and paper-based hospital records. The data was then entered into a custom-built, secure,

electronic database, hosted on the Newcastle Joint Research Office's Research Electronic Data

Capture (REDCap) platform.

The following conditions were documented based on the patients' medical histories: ischaemic heart

disease or previous myocardial infarction, atrial fibrillation, hypertension, cerebrovascular diseases

(such as ischaemic stroke, haemorrhagic stroke, or transient ischaemic attack), end-stage renal

failure requiring dialysis, and chronic obstructive pulmonary disease. Pre-operative medications were

noted from pre-assessment clinic records.

Routine clinical laboratory results (including haemoglobin [g/dl], white cell count [10^9/L], albumin

[g/L], creatinine [μmol/L], eGFR [mL/min/1.73 m²], C-reactive protein [mg/L], and HbA1c

[mmol/mol]) were also collected, along with height and weight measurements.

Statistical methods

Statistical analysis was performed using R version 4.5.0 (R Foundation for Statistical Computing,

Vienna, Austria). Descriptive proportional data is presented as number (percentage, 95% confidence

interval (CI), calculated using the exact binomial test). Continuous data is presented as mean (SD),

and hypothesis testing was performed with t-tests or Mann Whitney u-tests as appropriate based on

normality testing with the Shapiro-Wilk test.

Proportional comparisons were performed with Fisher's exact tests. A p-value <0.05 was considered

statistically significant for single comparisons. Log rank tests are reported to compare the overall

amputation-free survival (AFS).

99   Results

100   Eighty-four participants were recruited between October 2021 to October 2022. 54 (64%) were men

101   and the average BMI was 27.4 (6.9). 20 (24%) patients were current smokers, 47 (57%) were ex-

102   smokers, and 16 (19%) were never-smokers. Demographics, co-morbidities and social factors are

103   summarised in Table 1.

Main outcomes (see table 2)

Fifty-two percent (43/83 participants) were living with frailty, with median (IQR) Rockwood CFS of 5(2). Patients who were living with frailty were more likely to be sarcopenic by grip strength (19/43 (46%) vs 8/40 (20%), p=0.018). These results are summarised in Table 3.

A third (33%) of participants (27/81) had sarcopenia by grip strength. Patients who were considered sarcopenic by grip strength were more likely to be anaemic (21/27 (79%) vs 21/54 (41%), p=0.004). These results are summarised in Supplementary Table 1.

Overall, 54% (43/80) were anaemic, with the majority being normocytic (35/38, 92%). Supplementary Table 2 summarises these findings.

For patients who were considered frail, 19/43 (46%) were sarcopenic by grip strength and 26/43 (63%) were anaemic.

Survival analysis
90-day AFS in the whole cohort was 96%. There was no significant difference in 90-day AFS for those

living with frailty (98% vs 95%, p=0.78), sarcopenia (100% vs 94%, p=0.58), or anaemia (95% vs 97%,

p=0.45) or MLTC (96 vs 96%, p=0.67).

Other outcomes
Low walk speed (<0.8m/s) or failure of the 4m walk speed test was observed in 30/82 patients (37%).

The self-reported physical activity index suggested that 56 patients (68%) were classified as having

low activity. MLTC was seen in 57 (68%) patients and 67 (80%) patients met the definition of

polypharmacy.

Patients living with frailty were more likely to fail the walk speed test (21/43 (49%) vs 10/40 (25%),

p=0.040) and self-report low activity (39/43 (91%) vs 17/40 (43%), p<0.001). Anaemic patients had a

greater reduction in activity levels (35/43 (81%) vs 19/37 (51%), p=0.008) and were more likely to fail

the walk speed test (23/43 (54%) vs 8/37 (22%), p=0.005).

Pearson correlation coefficient was computed to determine the relationship between haemoglobin

and grip strength. The results suggest a positive relationship that was statistically significant (r(76) =

0.35, p=0.0017). Figure 2 shows a scatterplot demonstrating this relationship.

Clinical outcomes:

Most patients (64/81; 79.0%) underwent revascularisation, 36 (57.1%) were endovascular and 27

(42.9%) open or hybrid. Seventeen (21%) patients did not undergo revascularisation; 5 (29.4%)

underwent primary amputation, and 12 (70.6%) underwent a conservative or palliative approach.

Figure 2 shows a flowchart summarising this. Those with anaemia were less likely to undergo any

open/ hybrid reinterventions within 90 days (0 (0.0%) vs 4 (10.8%), p=0.046), but more likely to

undergo minor foot debridement (6 (14.6%) vs 0 (0.0%), p=0.027).

During the study period, there were 3 (4%) MACEs, 2 (3%) respiratory complications, 5 (6%) wound

complications and 6 (7%) MALEs. 29 (35%) patients in the study had at least one readmission while in

the study.

Sixteen patients had reinterventions within 90 days of index surgery; 4 (5%) open/ hybrid

interventions, 6 (7%) endovascular interventions, 2 (2%), wound-related procedures and 6 (7%) foot

debridements/ minor amputations.

Relationships between frailty, sarcopenia, anaemia and clinical factors

There were no significant correlations found between the presence of frailty and the complication

rate, readmission rates or the number and type of reinterventions. These results are summarised in

Table 3.

Supplementary tables 3 and 4 summarise the relationships between walk speed test failure and self-

reported low activity, against other characteristics.

Discussion

In the first UK prospective observational study of its kind, we have shown that a high prevalence of frailty, sarcopenia, anaemia, MLTC and polypharmacy exists in the CLTI and PAD population. We have also shown that there were positive associations between frailty and sarcopenia and between anaemia and sarcopenia, among CLTI patients.

The high prevalence of frailty in the population is striking but unsurprising. Indeed, this is in keeping with the literature in the UK; a recent retrospective study has shown that 98/190 (52%) patients were living with frailty by the Rockwood CFS.[27] A meta-analysis of global literature has suggested that 49% of those with PAD are frail, although this study was limited due to significant study heterogeneity.[28] In comparison, the prevalence of frailty in English adults aged >50 years old, was 8.1% [95% CI 7.3-8.8].[29]

Our findings on the prevalence of sarcopenia also matches up reasonably with the literature, with previous retrospective data suggesting prevalence of around 40% (37.8%[30] to 40.8%[31]). It is worth noting that most of these studies only include patients who undergo intervention, in contrast to our study, which includes those managed conservatively. Similarly, this study supports previously published findings of the prevalence of anaemia in the CLTI population of around 50%.[32]

Our study has identified an association between frailty and sarcopenia. The claudication symptoms that often precede CLTI can limit activity levels[33] and by the time patients present with rest pain or tissue loss, this insidious reduction in physical activity may lead to loss of independence, functional muscle loss and sarcopenia.[34,35] This poses the question of whether early revascularisation in those with claudication is warranted; there is reasonable data to support the fact that early surgery in this population leads to a higher risk of MLLA.[36] This, therefore, leaves us with more conservative measures; aggressive lifestyle modification and supervised exercise programmes (usually walking and treadmill based regimes) reduce the progression and severity of claudication symptoms and improve quality of life.[37] Unfortunately, these programmes are rarely available in the National Health

Service.[38] There is also a role for resistance training exercise regimes, as this type of exercise

programme has been shown to be beneficial in those with sarcopenia.[39] This could be incorporated

into both pre-operative and post-operative rehabilitation strategies.

The description of anaemia and its relationship with sarcopenia by grip strength is a new finding in

the UK CLTI population, however, it has been previously described in other non-vascular surgery

populations, particularly in East Asia.[40–42] Both anaemia and sarcopenia have been suggested to

contribute to poor outcomes after vascular surgery.[14,16] There is probably an element of mixed

anaemia in this patient cohort, but there are certainly unanswered questions about the true

mechanism of anaemia in this population.

Frailty and sarcopenia are both well understood to be part of aging and the concept of "inflamm-

aging" has been suggested as a mechanism for this. The imbalance of pro- and anti-inflammatory

processes are thought to drive this and result in inflammation mediated loss of muscle mass, and

inhibition of muscle regeneration.[43] There is potential that the disease of CLTI itself, and the

associated local lower limb inflammation could be further driving this inflammatory state. It could be

hypothesised that this could be the case with earlier stage PAD, where the onset of claudication

triggers a systemic inflammatory response. Cellular senescence and mitochondrial dysfunction have

also been postulated to play a role in the pathophysiology of sarcopenia.[44]

With such high rates of frailty, sarcopenia and anaemia in the vascular surgery population, it is

incumbent that services take steps to manage, and where possible, treat these conditions. The main

challenge to achieve this comes down to the short timeframes required for CLTI intervention (less

than 14 days as an out-patient and less than 5 days as an in-patient).[45] Complex geriatric assessment

(CGA) is an increasingly used service in vascular surgery units and can be implemented via an

inpatient or outpatient basis.[46–48] Indeed, the increasing use of "hot clinics" and outpatient pathways

may alleviate the risk of the deconditioning that often occurs with a prolonged hospital admission.[9]

Limitations

Despite this being a UK-first, there remain some limitations of this study. Primarily, a lack of long-term follow-up limited our ability to draw conclusions regarding frailty, sarcopenia and their effect on long-term outcomes, such as amputation free survival.

In this study, there was no detectable relationship between frailty, sarcopenia or anaemia against survival or other major clinical outcomes such as amputation, readmission or death. This may represent that these factors do not affect mid-term mortality and amputation rates, however, this is more likely to be because our study cohort of 84 patients provides insufficient power to detect statistically significant rates of complications between groups. This issue was compounded by the low rates of complications and death in the study, a fact that probably reflects the short follow-up time and the often-reported problem of study patients not necessarily representing the real-world population.[49]

In addition, some of the planned analyses were limited due to missing data. The Fried Frailty index was a significant outcome which was limited in this way, mostly driven by a lack of data pertaining to unintentional weight loss. Skeletal muscle area at L3 was also affected, and this may represent technical challenges in the measurement of this variable, which could be alleviated in future studies by improved training, or perhaps centralised image analysis. This prevented some of the analyses that were initially outlined in the protocol.[17] and impacted our ability to assess muscle quantity.

Future work to address the issues raised by this study should target the multiple facets of frailty and sarcopenia. Investigation of the mechanisms of sarcopenia and its relationship with anaemia, may lead to novel findings that could provide potential targets to slow down their progression. This approach may also unveil biological processes that are unique to the pathophysiology of CLTI.

Nutritional assessment and post-operative exercise and rehabilitation programmes may help alleviate the negative consequences of frailty and sarcopenia. Correcting anaemia may also be a potential avenue, and our unit is exploring further research on this topic.

Conclusion

This prospective multi-centre study highlights that there is a high prevalence of frailty, sarcopenia and anaemia in a population with CLTI. The positive association between sarcopenia and anaemia is novel in this population, and merits rigorous evaluation in future research. Our results suggest that a focus should be placed on managing these conditions alongside CLTI in a multidisciplinary fashion, and there is scope for further larger-scale prospective studies on these conditions and their impact on outcomes and quality of life after arterial intervention.

Figures and Tables

Figure 1: Scatterplot showing relationship between haemoglobin and grip strength.

Figure 2: Flowchart showing how many patients underwent intervention for CLTI. There were 3

missing data points for revascularisation.

Table 1: Table showing baseline demographics, co-morbidities and social history.

Table 2: Table showing prevalence of main outcomes and the interaction between conditions.

Table 3: Table summarising the relationships between frailty and clinical factors. All numbers are

reported to the nearest whole number (and percentage). Missing data values are included in square

brackets. *Composite including MACE, MALE, respiratory, urinary or wound-related complications.

Supplementary Table 1: Table summarising the relationships between sarcopenia and clinical factors.

All numbers are reported to the nearest whole number (and percentage). Missing data values are

included in square brackets. *Composite including MACE, MALE, respiratory, urinary or wound-

related complications.

Supplementary Table 2: Table summarising the relationships between anaemia and clinical factors.

All numbers are reported to the nearest whole number (and percentage). Missing data values are

included in square brackets. *Composite including MACE, MALE, respiratory, urinary or wound-

related complications.

Supplementary Table 3: Table summarising the relationships between walk speed and clinical factors.

All numbers are reported to the nearest whole number (and percentage). Missing data values are

included in square brackets. *Composite including MACE, MALE, respiratory, urinary or wound-

related complications.

Supplementary Table 4: Table summarising the relationships between self-reported low activity and

clinical factors. All numbers are reported to the nearest whole number (and percentage). Missing

data values are included in square brackets. *Composite including MACE, MALE, respiratory, urinary

or wound-related complications.

Acknowledgements

The authors of this study would like to acknowledge the involvement of the vascular research team (Samantha Mallon, Noala Parr, Esther Olabiyi and Tony Robinson), who provided invaluable assistance in the running of the study, and the clinical surgical teams in all recruitment sites, who supported recruitment to the study. In addition, we would like to recognise the contribution of the Vascular and Endovascular Research Network (VERN) for their support in running the study.

MDW acknowledges support from the NIHR Newcastle Biomedical Research Centre. SN acknowledges support from the NIHR Academy and the Newcastle Health Innovation/Research Partnership.

References

1. Smith GE, Long J, Wallace T, Carradice D, Chetter IC, Collaborative VR, et al. Identifying the research priorities of healthcare professionals in UK vascular surgery: modified Delphi approach. BJS Open. 2020;5(2):zraa025.

2. Duff S, Mafilios MS, Bhounsule P, Hasegawa JT. The burden of critical limb ischemia: a review of recent literature. Vasc Heal Risk Manag. 2019;15:187–208.

3. Menard MT, Farber A, Powell RJ, Rosenfield K, Conte MS, Hamza TH, et al. Quality of Life in Patients With Chronic Limb-Threatening Ischemia Treated With Revascularization. Circulation. 2024;149(16):1241–53.

4. Cesari M, Landi F, Vellas B, Bernabei R, Marzetti E. Sarcopenia and Physical Frailty: Two Sides of the Same Coin. Front Aging Neurosci [Internet]. 2014;6:192. Available from: http://journal.frontiersin.org/article/10.3389/fnagi.2014.00192/abstract

5. Collerton J, Martin-Ruiz C, Davies K, Hilkens CM, Isaacs J, Kolenda C, et al. Frailty and the role of inflammation, immunosenescence and cellular ageing in the very old: Cross-sectional findings from the Newcastle 85+ Study. Mech Ageing Dev. 2012;133(6):456–66.

6. Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in Older Adults: Evidence for a Phenotype. J Gerontol A Biol Sci Med Sci [Internet]. 2001;56(3):M146–57. Available from: https://academic.oup.com/biomedgerontology/article-abstract/56/3/M146/545770

7. Crozier-Shaw G, Joyce WP. Too frail for surgery? A frailty index in major colorectal surgery. Anz J Surg [Internet]. 2018;88:1302–5. Available from: http://doi.wiley.com/10.1111/ans.14792

8. Han B, Li Q, Chen X. Frailty and postoperative complications in older Chinese adults undergoing major thoracic and abdominal surgery. Clin Interv Aging. 2019;14:947–57.

9. Sivaharan A, Williams T, Ng C, Nixon R, Green M, Benton L, et al. Ambulatory vascular clinics provide a safe and effective pathway for management of chronic limb threatening ischaemia. J Vasc Soc Gt Br Irel. 2023;2.

10. Fernando ME, Blanchette V, Mishra R, Zulbaran-Rojas A, Rowe V, Mills JL, et al. Frailty in People with Chronic Limb Threatening Ischemia and Diabetes-Related Foot Ulcers: A Systematic Review. Ann Vasc Surg. 2023;89:322–37.

11. Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T, et al. Sarcopenia: revised European consensus on definition and diagnosis. Age Ageing. 2019;48(1):16–31.

12. Janssen I. The Epidemiology of Sarcopenia. Clin Geriatr Med [Internet]. 2011;27:355–63. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0749069011000176

13. Newton DH, Kim C, Lee N, Wolfe L, Pfeifer J, Amendola M. Sarcopenia predicts poor
long-term survival in patients undergoing endovascular aortic aneurysm repair. J Vasc Surg.
2018;67:453–9.

14. Sivaharan A, Boylan L, Witham MD, Nandhra S. Sarcopenia in Patients Undergoing Lower
Limb Bypass Surgery is Associated with Higher Mortality and Major Amputation Rates. Ann
Vasc Surg. 2021;75:227–36.

15. Ramadan OI, Rosenbaum PR, Reiter JG, Jain S, Hill AS, Hashemi S, et al. Redefining
Multimorbidity in Older Surgical Patients. J Am Coll Surg. 2023;236(5):1011–22.

16. Nandhra S, Boylan L, Prentis J, Nesbitt C. The Influence of Preoperative Anemia on
Clinical Outcomes After Infrainguinal Bypass Surgery. Ann Vasc Surg. 2020;

17. Hickson B, Sivaharan A, El-Sayed T, Baljer B, Sillito S, Shelmerdine L, et al. The FraiLTI
(Frailty in chronic Limb-Threatening Ischaemia) Protocol. J Vasc Soc Gt Br Irel. 2024;3.

18. Elm E von, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, et al. The
Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement:
guidelines for reporting observational studies. Lancet. 2007;370(9596):1453–7.

19. Conte MS, Bradbury AW, Kolh P, White JV, Dick F, Fitridge R, et al. Global vascular
guidelines on the management of chronic limb-threatening ischemia. J Vasc Surg [Internet].
2019;69(6):3S-125S.e40.   Available   from: www.jvascsurg.org.
https://linkinghub.elsevier.com/retrieve/pii/S0741521419303210

20. Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, McDowell I, et al. A global
clinical measure of fitness and frailty in elderly people. Can Méd Assoc J. 2005;173(5):489–
95.

21. McLean E, Cogswell M, Egli I, Wojdyla D, Benoist B de. Worldwide prevalence of
anaemia, WHO Vitamin and Mineral Nutrition Information System, 1993–2005. Public Heal
Nutr. 2008;12(4):444–54.

22. Steptoe A, Breeze E, Banks J, Nazroo J. Cohort Profile: The English Longitudinal Study of
Ageing. Int J Epidemiology. 2013;42(6):1640–8.

23. Witham MD, Wason J, Dodds R, Sayer AA. Developing a composite outcome measure for
frailty prevention trials – rationale, derivation and sample size comparison with other
candidate measures. BMC Geriatr. 2020;20(1):113.

24. Masnoon N, Shakib S, Kalisch-Ellett L, Caughey GE. What is polypharmacy? A systematic
review of definitions. BMC Geriatr. 2017;17(1):230.

25. Conte MS, Geraghty PJ, Bradbury AW, Hevelone ND, Lipsitz SR, Moneta GL, et al.
Suggested objective performance goals and clinical trial design for evaluating catheter-
based treatment of critical limb ischemia. J Vasc Surg. 2009;50(6):1462-1473.e3.

26. Bosco E, Hsueh L, McConeghy KW, Gravenstein S, Saade E. Major adverse cardiovascular
event definitions used in observational analysis of administrative databases: a systematic
review. BMC Méd Res Methodol. 2021;21(1):241.

27. Houghton JS, Nickinson AT, Helm JR, Dimitrova J, Dubkova S, Rayt HS, et al. Associations
of Clinical Frailty with Severity of Limb Threat and Outcomes in Chronic Limb-threatening
Ischaemia. Ann Vasc Surg. 2021;76:406–16.

28. Wang Y, Wu X, Hu X, Yang Y. Prevalence of frailty in patients with lower extremity
peripheral arterial disease: A systematic review and meta-analysis. Ageing Res Rev.
2022;82:101748.

29. Sinclair DR, Maharani A, Chandola T, Bower P, Hanratty B, Nazroo J, et al. Frailty among
Older Adults and Its Distribution in England. J Frailty Aging. 2022;11(2):163–8.

30. Selçuk N, Albeyoğlu Ş, Bastopcu M, Selçuk İ, Barutca H, Şahan H. Sarcopenia is a risk
factor for major adverse cardiac events after surgical revascularization for critical limb
ischemia. Vascular. 2023;31(1):64–71.

31. Cao Z, Zhao B, Jiang T, Zhang T, Yu X, Li Y, et al. Association of Sarcopenia With Mortality
in Patients With Chronic Limb-Threatening Ischemia Undergoing Endovascular
Revascularization. J Surg Res. 2023;289:52–60.

32. Theuma F, Nickinson ATO, Cullen S, Patel B, Dubkova S, Davies RSM, et al. The Impact of
Anemia on One-Year Amputation-Free Survival in Patients Undergoing Revascularization for
Chronic Limb-Threatening Ischemia: A Retrospective Cohort Study. Ann Vasc Surg.
2022;79:201–7.

33. Gerage AM, Correia M de A, Oliveira PML de, Palmeira AC, Domingues WJR, Zeratti AE,
et al. Physical Activity Levels in Peripheral Artery Disease Patients. Arq Bras Cardiol.
2019;113(3):410–6.

34. Sánchez-Sánchez JL, He L, Morales JS, Barreto P de S, Jiménez-Pavón D, Carbonell-Baeza
A, et al. Association of physical behaviours with sarcopenia in older adults: a systematic
review and meta-analysis of observational studies. Lancet Heal Longev. 2024;5(2):e108–19.

35. Hämäläinen O, Tirkkonen A, Savikangas T, Alén M, Sipilä S, Hautala A. Low physical
activity is a risk factor for sarcopenia: a cross-sectional analysis of two exercise trials on
community-dwelling older adults. BMC Geriatr. 2024;24(1):212.

36. Golledge J, Moxon JV, Rowbotham S, Pinchbeck J, Yip L, Velu R, et al. Risk of major
amputation in patients with intermittent claudication undergoing early revascularization. J
Br Surg. 2018;105(6):699–708.

37. Harwood AE, Pymer S, Ingle L, Doherty P, Chetter IC, Parmenter B, et al. Exercise training
for intermittent claudication: a narrative review and summary of guidelines for
practitioners. BMJ Open Sport Exerc Med. 2020;6(1):e000897.

38. Haque A. Few UK vascular centres offer a fully NICE-compliant supervised exercise
programme: a national audit. Ann R Coll Surg Engl. 2022;104(2):130–7.

39. Hurst C, Robinson SM, Witham MD, Dodds RM, Granic A, Buckland C, et al. Resistance
exercise as a treatment for sarcopenia: prescription and delivery. Age Ageing.
2022;51(2):afac003.

40. Dai S, Wang S, He Y, Dai C, Yu J, Ma X. Association between anemia and sarcopenia
among Chinese elderly: A cross-sectional study based on the China health and retirement
longitudinal study. Exp Gerontol. 2023;177:112183.

41. Lee DY, Shin S. Sarcopenia and Anemia in Elderly Koreans: A Nationwide Population-
Based Study. Healthcare. 2023;11(17):2428.

42. Liu Q, You J, Zhong M, Wu Z, Geng Y, Huang C. Hemoglobin level is negatively associated
with sarcopenia and its components in Chinese aged 60 and above. Front Public Heal.
2023;11:1081843.

43. Franceschi C, Campisi J. Chronic Inflammation (Inflammaging) and Its Potential
Contribution to Age-Associated Diseases. J Gerontol Ser A: Biomed Sci Méd Sci.
2014;69(Suppl_1):S4–9.

44. Antuña E, Cachán-Vega C, Bermejo-Millo JC, Potes Y, Caballero B, Vega-Naredo I, et al.
Inflammaging: Implications in Sarcopenia. Int J Mol Sci. 2022;23(23):15039.

45. Boyle JR, Atkins ER, Birmpili P, Pherwani AD, Brooks MJ, Biram RWS, et al. A Best Practice
Clinical Care Pathway for Peripheral Arterial Disease. J Vasc Soc Gt Br Irel [Internet]. 2022;1.
Available from: https://www.vsqip.org.uk/content/uploads/2022/05/PAD-QIF-2022-
Update.pdf

46. Partridge J, Sbai M, Dhesi J. Proactive care of older people undergoing surgery. Aging
Clin Exp Res. 2018;30(3):253–7.

47. Partridge JSL, Healey A, Modarai B, Harari D, Martin FC, Dhesi JK. Preoperative
comprehensive geriatric assessment and optimisation prior to elective arterial vascular
surgery: a health economic analysis. Age Ageing. 2021;50(5):1770–7.

48. Partridge JSL, Harari D, Martin FC, Peacock JL, Bell R, Mohammed A, et al. Randomized
clinical trial of comprehensive geriatric assessment and optimization in vascular surgery. Brit
J Surg. 2017;104(6):679–87.

49. Habibzadeh F. Disparity in the selection of patients in clinical trials. Lancet.
2022;399(10329):1048.

Demographics	
Age (years)	
Mean (SD)	71.5 (11.0)
Range	49.0 – 93.0
Sex	
Female	30 (36%)
Male	54 (64%)
Ethnicity (missing = 14)	
Asian	1 (1%)
Bangladeshi	1 (1%)
Black Caribbean	2 (3%)
Indian	2 (3%)
White British	64 (91%)
Weight (kg)	
Mean (SD)	78.8 (20.4)
Range	47.0 - 152.0
BMI (kg/m2)(missing = 1)	
Mean (SD)	27.4 (6.9)
Range	18.7 - 55.8
Ischaemic heart disease (missing = 1)	37 (45%)
Previous stroke/ TIA (missing = 1)	12 (15%)
Heart failure (missing = 1)	13 (16%)
Type 2 diabetes (missing = 1)	43 (52%)
Atrial Fibrillation (missing = 1)	16 (19%)
Hypothyroidism (missing = 1)	10 (12%)
Dementia (missing = 2)	0 (0%)
Anxiety (missing = 2)	7 (9%)
Depression (missing = 3)	8 (10%)
Parkinsons (missing = 2)	1 (1%)
Chronic obstructive pulmonary disease (missing = 2)	22 (27%)
Asthma (missing = 2)	5 (6%)
Other lung conditions (missing = 1)	4 (5%)
Osteoarthritis (missing = 2)	23 (28%)
Inflammatory arthropathy (missing = 3)	5 (6%)
Smoking status (missing = 1)	
Current smoker	20 (24%)
Ex-smoker (for longer than 2 months)	36 (43%)
Ex-smoker (for shorter than 2 months)	11 (13%)
Never smoker	16 (19%)
Alcohol intake (missing = 4)	
0 drinks daily	57 (71%)
1-2 drinks daily	15 (19%)
3-4 drinks daily	6 (8%)
5+ drinks daily	2 (3%)
Living circumstances	
Bungalow	17 (20%)
Flat	12 (14%)
House	51 (61%)
Other	3 (4%)
Residential Care	1 (1%)

Condition	Prevalence Number affected (percentage)	Subgroup Prevalence Number affected (percentage)
Frailty	43/83 (52%)	Sarcopenia: 19/43 (44%) Anaemia: 26/43 (60%) MLTC: 33/43 (77%)
Sarcopenia	27/81 (33%)	Frailty: 19/27 (70%) Anaemia: 21/27 (78%) MLTC: 21/27 (78%)
Anaemia	43/80 (54%)	Frailty: 26/43 (60%) Sarcopenia: 21/43 (49%) MLTC: 33/43 (77%)

Clinical Frailty scale (missing = 1)	Not Frail (N=40)	Frail (N=43)	p value
Sarcopenic (grip strength)	8 (20%)	19 (46%) [2]	**0.018***
MLTC	24 (60%)	33 (77%)	0.16
Anaemia	17 (44%) [2]	26 (63%) [2]	0.12
Walk speed test failure	10 (25%)	21 (49%)	**0.040***
Low Activity	17 (43%)	39 (91%)	**<0.001***
Polypharmacy	31 (78%)	36 (84%)	0.58
Revascularisation			
Revascularisation performed	33 (85%) [1]	31 (74%) [1]	0.28
Type of revascularisation	[7]	[13]	0.20
Endovascular	16 (49%)	20 (67%)	
Open or Hybrid	17 (52%)	10 (33%)	
Complications			
MACE	0 (0%) [1]	3 (7%) [1]	0.24
MALE	3 (8%) [1]	3 (7%) [1]	1.00
All complications*	4 (10%) [1]	9 (21%) [1]	0.23
Readmissions and reinterventions			
Number of readmission	[4]	[5]	0.14
0	24 (67%)	21 (55%)	
1	9 (25%)	10 (26%)	
2	1 (3%)	5 (13%)	
3	2 (6%)	0 (0%)	
4	0 (0%)	2 (5%) [1]	
Reintervention (Open or hybrid)	3 (8%) [1]	1 (2%) [1]	0.35
Reintervention (Endovascular)	1 (3%) [1]	5 (12%) [1]	0.20
Reintervention (Wound debridement)	1 (3%) [1]	1 (2%) [1]	1.00
Reintervention (Foot debridement)	4 (10%) [1]	2 (5%) [1]	0.42