Highway paving dramatically increased dengue transmission in the Amazon

Alyson L. Singleton*[1], Andres G. Lescano[2,3], Andrew J. MacDonald[4,5], Lisa Mandle[6], Terrell J.

Sipin[4], Kevin S. Martel[7,8], César V. Munayco[2,7], Esteban D. R. Carrera[9], Gustavo A. Choque[9],

Ana S. M. Bautista[9], Stephen P. Luby[10,11], Erin A. Mordecai[11,12]

[1] Emmett Interdisciplinary Program in Environment and Resources, Stanford University,
Stanford, California, United States of America

[2] Clima, Latin American Center of Excellence for Climate Change and Health

[3] Emerge, Emerging Diseases and Climate Change Research Unit, School of Public Health and
Administration, Universidad Peruana Cayetano Heredia, Lima, Peru

[4] Bren School of Environmental Science and Management, University of California, Santa
Barbara, Santa Barbara, California, United States of America

[5] Earth Research Institute, University of California, Santa Barbara, Santa Barbara, CA, United
States of America

[6] Natural Capital Project, Stanford University, Stanford, California, United States of America

[7] Centro Nacional de Epidemiología, Prevención y Control de Enfermedades, Peru Ministry of
Health, Lima, Peru

[8] School of Public Health and Administration, Universidad Peruana Cayetano Heredia, Peru

[9] Dirección Regional de Salud Madre de Dios, Madre de Dios, Peru

[10] Division of Infectious Diseases and Geographic Medicine, Stanford University School of
Medicine, Stanford, California, United States of America

[11] Woods Institute for the Environment, Stanford University, Stanford, California, United States
of America

[12] Department of Biology, Stanford University, Stanford, California, United States of America

Corresponding author: asinglet@stanford.edu (AS)

Preprint submitted to the medRxiv server.

Classification: Biological Sciences (major) and Environmental Sciences (minor).

Keywords: Dengue, highway, infrastructure, infectious disease

Abstract

Human mobility drives the spread of many infectious diseases, yet the health impacts of changes

in mobility due to new infrastructure development are poorly understood and currently not

accounted for in impact assessments. We take a novel quasi-experimental approach to

identifying the link between mobility and infectious disease, leveraging historical road upgrades

as a proxy for regional human mobility changes. We analyzed how highway paving altered

transmission of dengue—a high-burden mosquito-borne disease—via changes in human

movement in the Madre de Dios region of Peru. The paving of the Interoceanic Highway through

a formerly isolated region of the Amazon in 2009 provided a unique opportunity to quantify the

causal impact of road paving on disease transmission. To uncover this relationship, we compared

dengue incidence data from healthcare facilities in Madre de Dios near to versus far from the

newly paved highway before and after paving, while controlling for observable and unobservable

confounding variables (a difference-in-differences causal inference approach). We found that the

paving of the highway caused at least an additional 9,826 (95% CI: 8,562–10,684) dengue cases

since paving, accounting for 45.2% (95% CI: 39.4%–49.2%) of all dengue cases recorded in the

region post highway paving (2009–2022). Our findings demonstrate the impact that

infrastructure can have on dengue transmission, likely via its effects on human mobility. As a

result, we advocate for future road construction plans in tropical regions to account for potential

increases in dengue transmission during impact assessments.

Significance statement

Large-scale infrastructure projects are critical for meeting global development goals, but they

can also have unintended consequences for local human and environmental health. More

evidence attributing human health effects to infrastructure projects is needed to understand how

to mitigate these negative health impacts. Our study contributes to this body of knowledge,

attributing nearly half of recent dengue cases in Madre de Dios, Peru to the paving of the

Interoceanic Highway through the region in 2009. Our results demonstrate the impact that

highway construction can have on human health and call for future infrastructure projects

through tropical forests to consider infectious disease outcomes in their impact assessments.

Introduction

Meeting United Nations Sustainable Development goals requires massive infrastructure build-out, especially in developing countries (1). While there are many benefits promised by these infrastructure projects—such as increased access to labor opportunities, creation of water resources, and potential for economic growth—there are also many costs that have not been accounted for historically, such as harms to local environmental and human health (2, 3). Well-designed roads are critical for the distribution of resources and opportunities, and their growth is widespread; as of 2020, 12,263 km of road construction was planned in the Amazon alone (4). Despite their importance to local development, roads drive changes in human mobility and migration that have been previously associated with increases in infectious disease transmission (5, 6). However, these effects remain a challenge to identify and are often not included in even the most comprehensive road assessment analyses (4), likely due to the lack of causal analyses that disentangle disease from complex social-environmental processes. Current methods linking mobility and infectious disease also typically rely on cell phone or survey-based mobility data, which can be expensive, inaccessible, and/or biased due to lack of representativeness or recall bias (7, 8). To bridge these gaps, we take a novel quasi-experimental approach where we demonstrate a causal link between highway construction and dengue virus transmission, leveraging the rapid paving of a highway through the Peruvian Amazon as a proxy for a discrete and significant increase in regional human movement and connectivity.

The Madre de Dios region of Peru is an ideal case study to investigate the relationships between human mobility and infectious disease given its recent infrastructure development, history of vector-borne disease, and high environmental suitability for vectors (9, 10). Madre de Dios is an 85,000 km^2 department in the southwest Amazon in the southeast of Peru. In this

region, most human population centers are distributed along rivers, secondary unpaved roads,

and the only major paved road, the Interoceanic Highway, a controversial construction project

completed in 2009 that intensified regional in-migration (11) and legal and illegal development

(12) (**Fig 1**). Touted as a project that would have socio-economic benefits for rural Peruvian

communities, there have thus far been limited reports of economic or trade-based benefits in the

area (13, 14). However, the transformation of an unpaved road into a paved highway spurred

poverty-driven environmental degradation, such as accelerated deforestation and illegal gold

mining (12). There has been limited quantification of the health impacts of paving this section of

highway; studies to date have been qualitative or correlational analyses (14, 15). Causal analyses

of how highway construction may have influenced human health could help to build the evidence

base on impacts of infrastructure projects in the Amazon.

We hypothesized that the highway paving most immediately affected dengue

transmission through increases in mobility and connectivity both into and within the region, due

to dengue's specific epidemiology and ecology. Dengue virus is transmitted to humans by biting

mosquitoes (16). It is the most rapidly spreading mosquito-borne disease worldwide (17) and has

caused multiple record-breaking outbreaks over the last two years across the globe (18).

Qualitative analyses have reported that Madre de Dios community members believe that the

highway construction facilitated increased dengue transmission (14). Dengue was present

throughout Madre de Dios well before the paving in 2009 (10) (**Fig 1B, SFig S1A**), and its

primary vector—*Aedes aegypti*—inhabits urban and peri-urban areas that line highway corridors

(19). Dengue is a highly localized disease, appearing in clusters at the household and

neighborhood level (10). Previous studies have established that diffusion of dengue through

regional scales relies on human movement and migration (20), given that its vector typically

does not fly more than 1km (21). We hypothesized that when the construction of the Interoceanic

Highway further connected Madre de Dios to the bordering dengue-endemic regions of Acre,

Brazil and Pando, Bolivia, increased human movement along the highway amplified virus

importation and circulation (**SFig S2**).

In contrast to dengue, leishmaniasis—a parasitic disease caused by infection with

Leishmania parasites transmitted by sandflies—was also well established in the region prior to

road paving (**Fig 1C, SFig S1B**) but is not expected to be tightly linked to human mobility and,

by extension, to road paving (22, 23). Although sandflies have similar flight dispersal limitations

(24), they occur in rural, forested, and frontier ecosystems farther from highway corridors (23)

and become infected by feeding on infectious animal reservoir hosts rather than humans (humans

are dead-end hosts (25)). Therefore, while regional leishmaniasis transmission might have

increased somewhat due to highway-driven land conversion, its eco-epidemiology minimizes the

potential for it to be impacted by human movement along the highway. We use leishmaniasis as

a 'negative control' when testing impacts of the paving, in that we do not expect it to respond

immediately to highway construction after controlling for land-use changes, but it might capture

any impacts of highway paving on diagnostic capacity or healthcare seeking behavior.

To establish whether the paving of the Interoceanic Highway in Madre de Dios caused an

increase in dengue incidence rates, we conducted a difference-in-differences panel regression

(26), comparing dengue incidence rates in healthcare facility clusters ($n = 71$) within 5km of the

newly paved highway ('exposed') and those outside 10km ('unexposed'), before and after the

highway was paved in 2009 (**Fig 1**). This is a causal inference framework that approximates a

before-after treatment-control experimental design and incorporates (1) fixed effects, which

control for unobserved confounders (e.g., baseline dengue burden, large-scale political changes),

and (2) observed control variables, which control for known dengue drivers (i.e., climate and

land-use changes). We also tested whether or not the paving similarly affected leishmaniasis

incidence rates, with the hypothesis that there was no impact on leishmaniasis because it is not

directly affected by human mobility.

Results

We found that the paving of the highway increased dengue incidence rates by 184%

(95% CI: 65%–389%) in clusters within 5km of the highway compared to those outside 10km,

after incorporating fixed effects and accounting for potentially confounding spatial, temporal,

and environmental variation (**Table 1**). Yearly increases ranged from 20% to 458% across the

14-year study period (**Fig 2A**, **STable S1**). Overall, we attributed 9,826 (95% CI: 8,562–10,684)

dengue cases to the paving of the highway, accounting for 45.2% (95% CI: 39.4%–49.2%) of all

dengue cases recorded in the region post highway construction (**Fig 2A**). There was no

systematic difference in trends in dengue incidence between healthcare facility clusters near to

the highway and those far from the highway before paving (coefficients remain at zero prior to

2008, **Fig 2A**), supporting the parallel trends assumption and implying that far clusters represent

reasonable controls (i.e., counterfactuals) for clusters near the road. These findings were robust

to various model specifications (**STable S2, SFig S3**) and produced p-values < 0.01 when

compared to permuted samples in permutation inference analyses (**SFig S4**). Increases in urban

area were associated with increased dengue incidence, aligning with biological knowledge of

Aedes vector habitat (**Table 1**). Seasonal (biannual) models aggregated across the study period

revealed larger differences in dengue incidence between healthcare facility clusters near and far

from the highway during rainy seasons (173%, [95% CI: 73–334%]), when the majority of cases

occur, as compared to dry seasons (68%, [95% CI: 12–152%]) (**Fig 2B**).

Table 1: Dengue and leishmaniasis incidence long-differences estimation results.

	Dengue Yearly	Dengue Biannual Dry	Dengue Biannual Rainy	Leish Yearly	Leish Biannual Dry	Leish Biannual Rainy
5km x Post-2008 Yearly	1.0* (0.28)			0.40 (0.24)		
5km x Post-2008 Biannual		0.52* (0.21)	1.0* (0.23)		0.47* (0.20)	0.16 (0.15)
log(Urban)	0.09 (0.10)	0.10 (0.07)	0.06 (0.09)	-0.005 (0.07)	0.03 (0.04)	-0.05 (0.04)
log(Ag)	-0.23 (0.14)	-0.17 (0.12)	-0.24 (0.15)	0.23 (0.15)	0.14 (0.14)	0.21 (0.11)
Sum Precip	0.09 (0.24)	0.73 (0.58)	0.02 (0.24)	-0.34* (0.13)	-0.60* (0.30)	-0.22* (0.10)
Mean Temp	-0.05 (0.47)	0.25 (0.25)	-0.05 (0.53)	0.86* (0.43)	0.45* (0.20)	0.28 (0.34)
FEs						
Cluster	X	X	X	X	X	X
Year	X			X		
Biannual		X	X		X	X
Cor^2	0.54541	0.3425	0.54752	0.53686	0.59255	0.55846
N	974	1,040	1,040	975	1,040	1,040
Clusters	65	65	65	65	65	65

151
Estimation results from Equation 2 for yearly and biannual models. Yearly models include cluster and
yearly fixed effects, and biannual models include cluster and biannual fixed effects. Standard errors are
clustered at the unit level for all models and are displayed inside parentheses next to each estimated
coefficient. P-values indicating a statistical significance level below 0.05 are designated with an asterisk
(*). N increases for biannual models because the reference time step is a six month period, resulting in the
inclusion of the corresponding six months as an additional data point. **STable S3** displays results for
models with quadratic climate covariates, finding no major differences.

As a 'negative control' that we do not expect to be immediately affected by increased

human mobility, we used the same approach to examine impacts on leishmaniasis incidence

before and after road paving. We found that leishmaniasis incidence did not differ systematically

between healthcare facility clusters near and far from the highway either before or after highway

paving (**Fig 3A**). Although we saw small differences between healthcare facilities near and far

from the highway almost a decade after paving and during the dry season, the differences were

minimal and were not significant when aggregated across the study period (**Table 1, Fig 3B**).

Increases in agricultural area were associated with increased leishmaniasis incidence, aligning

with biological knowledge of sandfly vector habitat (**Table 1**).

To better understand (1) which communities are at heightened risk of increased dengue

transmission due to highway paving and (2) the extent of potential spatial spillover effects of

exposure, we compared groups of healthcare facility clusters at increasing distances from the

highway—1km, 5km, 10km, 15km, and 20km—to a control group of healthcare facility clusters

>20km from the highway (**Fig 4B**). Healthcare facility clusters within 1km, 5km, and 10km all

had significantly higher dengue incidence than healthcare facility clusters >20km from the

highway post paving (**Fig 4A**). Differences attenuated to zero and were not statistically

significant for healthcare facility clusters farther than 10km from the highway.

Discussion

We found large and lasting impacts of the paving of the Interoceanic Highway on dengue

incidence in a previously isolated region of the Peruvian Amazon: incidence rates increased by

184% (95% CI: 65%–389%) in healthcare facility clusters near to versus far from the highway

(**Table 1**, **Fig 2**). While we do not directly test the mechanism, we hypothesize that the

significant increase in dengue burden was driven by increased human mobility into and within

the region, leading to increased virus importation and circulation (**SFig S2**). We extend previous

research (5, 6) that associates highway construction with increased dengue transmission by

offering causal estimates of the impact of road paving on dengue while controlling for potentially

confounding variables. These estimates are robust to a suite of alternative model specifications

(**SFig S3**). We also conducted a 'negative control' analysis as a way to check for systematic

changes that could confound our estimates, such as healthcare-seeking or case-reporting

behaviors, and find that for leishmaniasis, a disease not expected to change with human mobility,

there is minimal effect of road paving on incidence (**Fig 3**).

Our primary hypothesized mechanism for this effect is that highway paving facilitated

regional human movement changes that caused additional dengue outbreaks to be seeded along

the highway corridor. The highway increased connectivity to Brazil and Bolivia, both of which

are dengue hyper-endemic, with regular co-circulation of multiple serotypes and genotypes (5,

27) (**SFig S2**). Another relevant dynamic is the in-migration that occurred after the highway was

built, with people moving to Madre de Dios from the nearby, newly connected Andes mountains

and other non-endemic regions (11) (**SFig S5**). This increased the number of susceptible people

living in the area. Although we directly account for yearly population changes when calculating

incidence (**SFig S5**) and determine that dengue cases increased more than what would have been

expected just due to population growth, we do not explicitly include susceptibility dynamics in

our model. We hypothesize susceptibility dynamics are also what might be responsible for the

small increases in leishmaniasis incidence later in the study period (**Fig 3**).

Effects of highway paving on dengue were significantly larger during rainy seasons as

compared to dry, even after explicitly controlling for precipitation and including biannual fixed

effects (**Fig 2B**). This could be due to the larger dengue burden during the rainy season enabling

larger differences to occur between healthcare facilities near and far from the highway.

Alternatively, this could be due to increased differences in mobility during the rainy season;

rivers and unpaved roads often become impassable during the rainy season, while the highway is

less affected by increased precipitation (28). Therefore, mobility continues to a higher degree

along the highway compared to farther away, potentially causing larger dengue differences

between healthcare facility clusters near and far from the highway in the rainy season. Further

supporting the role of movement and accessibility of the road, we found that communities as far

as 10km from the highway experienced increased dengue transmission after highway paving,

while communities farther than 10km from the highway did not (**Fig 4**). This helps to define the

spatial scale of impacts of highway development projects on dengue, with applications for

designing control interventions.

Although this work isolates a plausible causal impact of highway paving on dengue

transmission, our quasi-experimental approach does not allow us to definitively identify the

specific mechanism that caused this effect. We hypothesize the main mechanism to be the

increase in human movement along the corridor, supported by the fact that (1) we only see an

immediate effect on dengue and not on leishmaniasis, (2) the effect is large even when explicitly

controlling for resulting in-migration and land-use changes, and (3) the estimated effect

attenuates at distances (>10km) unlikely to be affected by mobility around the highway. Direct

mobility data, such as surveys or mobile phone data, could further bolster our understanding of

the mechanistic effects of highway paving on mobility, though they would likely be subject to

recall bias and/or lack of representativeness. While changes in mobility and connectivity are

parsimonious and plausible mechanisms behind the observed increases in dengue following

highway paving, there are a few alternative phenomena that could have also contributed to the

identified effect. We are not able to include susceptibility dynamics in our model, which are

known to be important for dengue due to the cross-reactivity of various dengue serotypes.

However, it is important to note that large interannual changes in dengue outbreaks, for example

following the introduction of a new serotype, are accounted for in the year fixed effects except to

the extent that they differentially affect near versus far healthcare clusters. We also were not able

to control for hyper-localized habitat changes that might have implications for cluster-level

vector abundance and transmission capabilities, such as water storage and garbage practices that

can influence breeding habitat.

Other limitations of this work include that different communities were exposed to their

stretch of highway being paved at different times. Therefore, some units may have experienced

exposure before 2009, resulting in an underestimation of the true effect size. Additionally, even

after removing some units to create a 'buffer' against spatial spillover, some spillover likely

remains throughout the unexposed units, also resulting in potential underestimation. Finally,

although we attempt to control for development differences between exposed and unexposed

groups through measuring yearly changes in urban area, it is possible that healthcare facilities

near the highway developed increased capacity for diagnosing dengue compared to those far

from the highway post paving. However, we show our results are robust to the inclusion of

"probable" cases (**STable S2, SFig S3**), alleviating concerns of potential distortions due to the

availability of test supplies.

We demonstrate the importance of including human health outcomes, such as infectious

disease, in impact assessments when planning infrastructure development projects (3, 29). Our

analysis establishes that highway construction through dengue-endemic areas makes

communities uniquely vulnerable to dengue transmission. This work is especially timely because

two additional highway construction projects through Madre de Dios were approved in 2023,

over the protest of local communities and Indigenous groups (28, 30). Outside of the Amazon,

other dengue-endemic tropical regions are also planning highway construction projects,

including in Central America, Central Africa, and Southeast Asia (31). Given the measurable and

significant impact that road construction had on local dengue transmission in the Madre de Dios

region, our findings contribute to understanding the social, environmental, and economic

tradeoffs of this proposed infrastructure development. Should these infrastructure projects be

approved, our work highlights the importance of coordinating with local, regional, and national

healthcare officials at the outset of infrastructure project planning to forecast health impacts and

target public health resources and interventions. Potential human health interventions could

include vector control strategies (32), investment in water and sanitation infrastructure, and

health system strengthening along highway corridors, such as increased staffing, priority access

to new dengue vaccines once approved (33, 34), and deployment of dengue rapid tests for

improved diagnostics and notification (35, 36).

We contribute to a growing body of literature that demonstrates the impact of

infrastructure on human health, but dengue is only one of many infectious diseases that may be

impacted by changes in human movement and development. Future work can help to identify the

responses of other diseases to road building and other infrastructure development in the Amazon

and across the globe. Generating this evidence will be key for guiding informed decision-making

on meeting our development goals, while proactively mitigating any potential harms to local

human and environmental health.

Materials and methods

Ethics statement

The Stanford IRB office determined that this analysis did not qualify as Human Subjects

Research as defined in 45 CFR 46.102(e)(5), given the data did not contain any identifiable

information.

Data collection and preparation

De-identified, anonymized dengue and leishmaniasis case data reported from 2000–2022

were provided by the Regional Health Directorate of Madre de Dios (DIRESA, (37)). Both

dengue and leishmaniasis are reportable infectious diseases and are included in mandatory

surveillance for each healthcare facility ($n = 105$) in Madre de Dios. Case data included

diagnosis code, week and year of diagnosis, method of diagnosis, locality name, and healthcare

facility code.

Of 25,211 total dengue cases, 94.1% were laboratory confirmed and 5.9% were probable

(see **Supplementary Information** for diagnosis methods). We included only confirmed cases in

the main model and conducted a sensitivity analysis that included both confirmed and probable

cases (**STable S2**, **SFig S3**). Of 15,507 total leishmaniasis cases, 99.4% of cases were laboratory

confirmed and 0.6% were probable. We similarly included only confirmed leishmaniasis cases.

86.6% of leishmaniasis cases were cutaneous leishmaniasis and 13.4% were mucocutaneous.

Healthcare facilities were clustered to create non-overlapping spatial boundaries to extract

climate and land use covariate data, resulting in $n = 71$ spatial units (**SFig S7**). We tested

alternative clustering specifications that increased the number of spatial units ($n = 78$ and $n = 84$)

and found no substantial differences in main model results (**SFig S8**, see **Supplementary**

Information for clustering methods).

We calculated cluster-specific, yearly population measures by integrating locality

population data from DIRESA and WorldPop (37, 38) (**SFig S5A**, see **Supplementary**

Information). Remotely-sensed environmental covariate data was included explicitly in models

as control variables. First, we calculated the total yearly urban and agricultural area using

Mapbiomas—a publicly available dataset of classified annual land-use maps of the Amazon

basin with a spatial resolution of $30m^2$ (**SFigs S5B and S5C** (39)). Second, we calculated

monthly mean temperature and accumulated precipitation using the ERA5 Land Monthly

Aggregated dataset, which provides climate data at a spatial resolution of $11km^2$ (40). Monthly

values were averaged or summed annually and biannually depending on each model's temporal

resolution.

Interoceanic Highway

We leveraged the paving of the highway as a natural experiment, or shock, hypothesized

to result in a sudden increase in mobility-driven dengue transmission in areas close to the

highway. This paving is plausibly exogenous to our system (i.e., the decision to pave and

location of the road paving did not depend on dengue incidence at baseline), given that the road

was paved as a part of the larger Interoceanic Highway construction project, spanning from the

Atlantic to the Pacific coast (41). The road paving site was selected to pass through the tri-border

area at the northeast end of the region where Peru borders Brazil and Bolivia and then minimize

costs to reach Puerto Maldonado, the region's capital, and eventually Cuzco (41) (**SFig S2**).

The construction of the Interoceanic Highway through Peru took place from 2006 to 2010

(42). Further investigation confirmed the section of highway specific to Madre de Dios (Tramo

3, Inambari–Iñapari) was completed in 2009 (43). While paving through Madre de Dios occurred

from 2007 to 2009 (43–45), our goal was to capture when regional human connectivity and

movement shifted regionally, rather than the moment households individually experienced

paving adjacent to their community. Therefore, we designate 2009 as the year when exposure

began for all spatial units and 2008 as our 'baseline' year (i.e., the last year before exposure

began). The paving of the highway shortened the time it takes to cross the region from a multi-

day trip to five hours (46).

This study design designates healthcare facility clusters within 5km of the highway as

'exposed' and those further than 10km from the road as 'unexposed,' using distance as a proxy

for connectivity to the highway. Understanding there is likely spatial spillover of exposure, we

removed healthcare facility clusters between 5-10km from the highway (**Fig 1A**). Given distance

is only a proxy for accessibility to the highway, we tested the sensitivity of the model to both the

boundary and buffer specifications (**SFig S3**). The key identifying assumption in the difference-

in-differences design is that both groups of healthcare facility clusters would have seen dengue

incidence develop along parallel trends in the absence of the road paving. Visually, the parallel

trends assumption appears defensible, with dengue and leishmaniasis incidence following

parallel trajectories in clusters designated exposed versus unexposed prior to paving (i.e., pre-

exposure; **Figs 1B and 1C**). Additionally, all healthcare facility clusters far from the highway are

connected to the region's larger transportation network either by unpaved roads or rivers, making

their connection to the region's transportation network comparable to that of the healthcare

facility clusters near to the highway prior to exposure (**Fig 1A**).

Statistical models

To identify the effect of paving the Interoceanic Highway on dengue transmission, we

used a difference-in-differences panel regression (26), comparing healthcare facility clusters

within 5km of the newly paved highway ('exposed') and those outside 10km ('unexposed'),

before and after the highway was paved in 2009. The model takes the form (Equation 1):

$$\log(dengue_{i,t}) = \sum_{\tau \in \{2000,\dots,2020\},\tau!=2008} \beta_\tau \, (DIST)_i * 1\{t = \tau\} + \gamma_i + \lambda_t + X_{i,t}\theta + \epsilon_{i,t}$$

where *i* is unit (healthcare facility cluster) and *t* is year. Our outcome of interest, $\log(dengue_{i,t})$, is

the logged dengue incidence in healthcare facility cluster *i* for year *t*. Our exposure variable,

DIST, is a dummy (binary) variable that equals one for exposed clusters (<5km) and zero for

unexposed clusters (>10km). The DIST dummy variable interacts with year dummy variables

(one for each year of interest and zero otherwise). We omit 2008, the year before the highway

was paved, as the baseline year. The coefficients, β's, on the interaction term then estimate the

response in our outcome of interest following the road paving. Each coefficient estimates the log

of the incidence rate ratio between near and far healthcare facilities for each year in our set of

τ's. We include unit (i.e., healthcare facility cluster) fixed effects (FEs), γ_i, to help account for

any time-invariant healthcare facility characteristics, such as differences in their invasion history

of *Aedes* mosquitoes, proximity to protected areas, and baseline dengue burden at the community

level. FEs are a set of dummy variables that each represent a group (e.g., healthcare facility

clusters), demeaning the data within each group. We also include year FEs, λ_t, to account for

large-scale changes that might impact dengue comparably across all healthcare facility clusters

over time, such as El Niño events, economic recessions, and global pandemics. Finally, we

include four observable controls, noted by the matrix $X_{i,t}$, to explicitly control for any variation

due to changes in urban area, agricultural area, temperature, or precipitation. The Θ vector

recovers our covariate coefficients. Any remaining unobserved variation is captured by the error

term, $\epsilon_{i,t}$.

If our hypothesis that the highway paving increased dengue transmission in healthcare

facility clusters near to the highway is correct, we expect the β coefficients will diverge from

zero and become positive following 2008. Coefficients remaining at zero prior to 2008 would

support the parallel trends assumption and the validity of using the unexposed group as controls.

As a 'negative control' test, we conducted the same analysis with leishmaniasis incidence as our

outcome, with the expectation of no significant differences in outcome between exposed and

unexposed groups post-road paving.

Models were run as Poisson regressions using the *fixest* package in R with standard errors

clustered at the unit-level to account for autocorrelation between pre- and post-road paving in the

same healthcare facility cluster. We tested the sensitivity of model estimates to population

weighting, excluding Puerto Maldonado (the region's capital), changing the spatial boundary

delineating exposed and unexposed groups, changing the buffer zone size between exposed and

unexposed groups, including both confirmed and probable cases, and only including spatial units

reporting disease case data both pre- and post-road paving (**STable S2, SFig S3**). We also

compared linear climate variables to quadratic forms (**STable S3**). Finally, we conducted

permutation inference analysis, where we randomized our exposure variable using three

methods—randomizing across clusters and years, randomizing across clusters while maintaining

temporal structure, and randomizing across years while maintaining spatial structure—to

establish that there are no spurious cross-sectional or temporal phenomena contributing to the

observed effect (**SFig S4**).

Because rainfall is highly seasonal in this region and dengue transmission peaks in the

rainy season (**SFig S9**), we also ran biannual models to account for heterogeneous effects across

seasons. These models were equivalently parameterized except t was biannual time periods

(October–March, April–September for each year) that were constructed to align with the rainy

season (months with above average monthly accumulated precipitation from 2000–2022, **SFig**

S9). Biannual models included biannual fixed effects in place of yearly fixed effects.

Finally, we estimated aggregated versions of Equation 1 to summarize average treatment

effects and the total number of attributable cases (e.g., a long-differences model). We define a

post-paving dummy variable equal to zero in 2008 and one from 2009 to 2022. The aggregated

model takes the form (Equation 2):

$$log(dengue_{i,t}) = \beta \ (DIST)_i * 1\{t \in [2009,2022]\}_t + \gamma_i + \lambda_t + X_{i,t}\theta + \epsilon_{i,t}$$

This is also the model specification used for aggregating biannual model average

treatment effects (with biannual fixed effects substituted for yearly fixed effects, **Fig 2B** and **Fig**

3B). We also used the aggregated model to compare groups of healthcare facility clusters at

increasing distances from the highway—1km, 5km, 10km, 15km, and 20km—to a control group

of healthcare facility clusters >20km from the highway (**Fig 4A**). We calculated bootstrapped

standard errors for this analysis given the small sample size of some of the distance subgroups

(i.e., 1km, 5km, etc.). We bootstrapped by sampling healthcare facility clusters with replacement,

retaining the full time series for each cluster in the bootstrapped sample and maintaining the ratio

of exposed to unexposed units for each subgroup. In all other analyses we used analytic standard

errors.

Attributable cases

We calculated percent changes in dengue incidence due to highway paving as (Equation 3):

$$percent \ change \ dengue_t = (\exp{(\beta)_t} - 1) * 100$$

where *t* is year and β_t is the log of the incidence rate ratio between near and far healthcare

facilities for each year (i.e., coefficients from Equation 1 that account for fixed effects and

control variables). We similarly calculated the number of cases attributable to the highway

paving by exponentiating the β coefficient recovered from our aggregated model (i.e.,

calculating incidence rate ratios that account for fixed effects and control variables) and using the

following approximation (Equation 4):

$$attributable\ cases\ =\ pcet\ \left(\frac{exp(\beta)\ -\ 1}{exp(\beta)}\right)$$

410   where $pcet$ is the proportion of cases exposed to treatment (47).

Acknowledgements

We thank all the health personnel of the Madre de Dios region who contributed to generate this valuable data, as well as the Peruvian National Epidemiology Network (RENACE). We acknowledge and thank the members of the Mordecai and Luby Labs of Stanford University for their support and contribution to this work. We would also like to acknowledge computational resources from Google Cloud for Earth Engine. ALS was supported by the Jim and Gaye Pigott Fellowship through the Stanford Interdisciplinary Graduate Fellowship program at Stanford University. AGL is sponsored by Emerge, the Emerging Diseases Epidemiology Research Training grant D43 TW007393 awarded by the Fogarty International Center of the US National Institutes of Health. AJM acknowledges the United States Department of Agriculture National Institute of Food and Agriculture (2023-68016-40683) and the National Science Foundation and Fogarty International Center (DEB-2011147; DEB-2339209). LM was supported by the National Science Foundation (DEB-2011147 with Fogarty International Center). EAM was supported by the National Science Foundation (DEB-2011147 with Fogarty International Center), the National Institutes of Health (R35GM133439, R01AI168097, R01AI102918), the Stanford Woods Institute for the Environment, Center for Innovation in Global Health, and King Center on Global Development.

References

1. S. Thacker, *et al.*, Infrastructure for sustainable development. *Nat. Sustain.* **2**, 324–331

(2019).

2. R. K. Morgan, Environmental impact assessment: the state of the art. *Impact Assess. Proj.*

Apprais. **30**, 5–14 (2012).

3. J. A. Patz, *et al.*, Unhealthy Landscapes: Policy Recommendations on Land Use Change

and Infectious Disease Emergence. *Environ. Health Perspect.* **112**, 1092–1098 (2004).

4. T. Vilela, *et al.*, A better Amazon road network for people and the environment. *Proc. Natl.*

Acad. Sci. **117**, 7095–7102 (2020).

5. R. M. Lana, M. F. da C. Gomes, T. F. M. de Lima, N. A. Honório, C. T. Codeço, The

introduction of dengue follows transportation infrastructure changes in the state of Acre,

Brazil: A network-based analysis. *PLoS Negl. Trop. Dis.* **11**, e0006070 (2017).

6. Q. Li, W. Cao, H. Ren, Z. Ji, H. Jiang, Spatiotemporal responses of dengue fever

transmission to the road network in an urban area. *Acta Trop.* **183**, 8–13 (2018).

7. C. Buckee, A. Noor, L. Sattenspiel, Thinking clearly about social aspects of infectious

disease transmission. *Nature* **595**, 205–213 (2021).

8. A. Wesolowski, *et al.*, Multinational patterns of seasonal asymmetry in human movement

influence infectious disease dynamics. *Nat. Commun.* **8**, 2069 (2017).

9. J. J. Scullion, K. A. Vogt, A. Sienkiewicz, S. J. Gmur, C. Trujillo, Assessing the influence

of land-cover change and conflicting land-use authorizations on ecosystem conversion on

the forest frontier of Madre de Dios, Peru. *Biol. Conserv.* **171**, 247–258 (2014).

10. G. Salmón-Mulanovich, *et al.*, Individual and Spatial Risk of Dengue Virus Infection in

Puerto Maldonado, Peru. *Am. J. Trop. Med. Hyg.* **99**, 1440–1450 (2018).

11. K. E. Jensen, *et al.*, Small scale migration along the interoceanic highway in Madre de

Dios, Peru: an exploration of community perceptions and dynamics due to migration. *BMC*

Int. Health Hum. Rights **18**, 12 (2018).

12. A. M. Sánchez-Cuervo, *et al.*, Twenty years of land cover change in the southeastern

Peruvian Amazon: implications for biodiversity conservation. *Reg. Environ. Change* **20**, 8

(2020).

13. A. S. Oliveira, *et al.*, Bringing economic development for whom? An exploratory study of

the impact of the Interoceanic Highway on the livelihood of smallholders in the Amazon.

Landsc. Urban Plan. **188**, 171–179 (2019).

14. A. R. Riley-Powell, *et al.*, The Impact of Road Construction on Subjective Well-Being in

Communities in Madre de Dios, Peru. *Int. J. Environ. Res. Public. Health* **15**, 1271 (2018).

15. J. F. Sanchez, *et al.*, Unstable Malaria Transmission in the Southern Peruvian Amazon and

Its Association with Gold Mining, Madre de Dios, 2001–2012. *Am. J. Trop. Med. Hyg.* **96**,

304–311 (2017).

16. T. P. Monath, Dengue: the risk to developed and developing countries. *Proc. Natl. Acad.*

Sci. **91**, 2395–2400 (1994).

17. WHO, Dengue: the fastest growing mosquito-borne disease in the world. (2010). Available

at: https://www.who.int/news/item/29-10-2010-dengue-the-fastest-growing-mosquito-

borne-disease-in-the-world [Accessed 26 May 2023].

18. Dengue - Global situation. (2024). Available at: https://www.who.int/emergencies/disease-

outbreak-news/item/2024-DON518 [Accessed 18 July 2024].

19. S. A. Guagliardo, *et al.*, Patterns of Geographic Expansion of Aedes aegypti in the Peruvian

Amazon. *PLoS Negl. Trop. Dis.* **8**, e3033 (2014).

20. S. T. Stoddard, *et al.*, The Role of Human Movement in the Transmission of Vector-Borne

Pathogens. *PLoS Negl. Trop. Dis.* **3**, e481 (2009).

21. L. C. Harrington, *et al.*, Dispersal of the dengue vector Aedes aegypti within and between

rural communities. *Am. J. Trop. Med. Hyg.* **72**, 209–220 (2005).

22. J. P. Guthmann, *et al.*, Patients' associations and the control of leishmaniasis in Peru. *Bull.*

World Health Organ. **75**, 39–44 (1997).

23. V. Zorrilla, *et al.*, Distribution and identification of sand flies naturally infected with

Leishmania from the Southeastern Peruvian Amazon. *PLoS Negl. Trop. Dis.* **11**, e0006029

(2017).

24. A. C. Morrison, C. Ferro, A. Morales, R. B. Tesh, M. L. Wilson, Dispersal of the sand fly

Lutzomyia longipalpis (Diptera: Psychodidae) at an endemic focus of visceral leishmaniasis

in Colombia. *J. Med. Entomol.* **30**, 427–435 (1993).

25. E. K. Saliba, O. Y. Oumeish, Reservoir hosts of cutaneous leishmaniasis. *Clin. Dermatol.*

17, 275–277 (1999).

26. J. B. Dimick, A. M. Ryan, Methods for Evaluating Changes in Health Care Policy: The

Difference-in-Differences Approach. *JAMA* **312**, 2401–2402 (2014).

27. Y. Orba, *et al.*, Diverse mosquito-specific flaviviruses in the Bolivian Amazon basin. *J.*

Gen. Virol. **102**, 001518 (2021).

28. P. Moore, A decade on, Peru's corruption-laden Interoceanic Highway troubles

communities. *Dialogo Chino* (2022). Available at:

https://dialogochino.net/en/infrastructure/52497-peru-corruption-interoceanic-highway-

trouble-communities/ [Accessed 26 May 2023].

29. L. Flynn, R. Bery, A. E. Kaitano, Emerging Infectious Diseases and Impact Assessments.

(2013).

30. Peru ends era of 'roadless wilderness' in its Amazon rainforests. *The Independent* (2018).

Available at: https://www.independent.co.uk/climate-change/news/peru-amazon-

rainforests-road-building-roadless-wilderness-madre-de-dios-a8196971.html [Accessed 22

July 2024].

31. F. Kleinschroth, J. R. Healey, Impacts of logging roads on tropical forests. *Biotropica* **49**,

620–635 (2017).

32. S. Durand, *et al.*, Cost of controlling the dengue vector Aedes aegypti in the Peruvian

amazon. *Rev. Peru. Med. Exp. Salud Pública* **41**, 46–53 (2024).

33. Dengue is raging in Brazil. A promising local vaccine is at least a year away. Available at:

https://www.science.org/content/article/dengue-raging-brazil-promising-local-vaccine-

least-year-away [Accessed 22 July 2024].

34. M. Lenharo, Brazil's record dengue surge: why a vaccine campaign is unlikely to stop it.

Nature **627**, 250–251 (2024).

35. A. C. Morrison, *et al.*, Potential for community based surveillance of febrile diseases:

Feasibility of self-administered rapid diagnostic tests in Iquitos, Peru and Phnom Penh,

Cambodia. *PLoS Negl. Trop. Dis.* **15**, e0009307 (2021).

36. B. Valdivia-Conroy, *et al.*, Diagnostic performance of the rapid test for the detection of

NS1 antigen and IgM and IgG anti-antibodies against dengue virus. *Rev. Peru. Med. Exp.*

Salud Pública **39**, 434–441 (2023).

37. Dirección Regional de Salud Madre de Dios - Diresa Madre de Dios. (2021). Available at:

https://www.gob.pe/regionmadrededios-diresa [Accessed 18 July 2024].

38. WorldPop, WorldPop □:: Population Counts. (2023). Available at:

https://hub.worldpop.org/project/categories?id=3 [Accessed 26 May 2023].

39. Mapbiomas Brasil. Available at: https://mapbiomas.org/ [Accessed 26 May 2023].

40. J. Muñoz Sabater, ERA5-Land monthly averaged data from 1950 to present. *Copernic.*

Clim. Change Serv. C3S Clim. Data Store CDS (2019). Available at:

https://cds.climate.copernicus.eu/cdsapp#!/dataset/10.24381/cds.68d2bb30?tab=overview

[Accessed 18 July 2024].

41. Peru's Interoceanic: the Most Corrupt Highway in the World. *Up. Amaz. Conserv. EN*

(2018). Available at:

https://www.upperamazon.org/news/fz8vmhz9ytujotmqksrxoex5rljfcv-mnsk7 [Accessed 21

May 2023].

42. UNASUR - COSIPLAN. (2017). Available at:

https://www.iirsa.org/proyectos/detalle_proyecto.aspx?h=319 [Accessed 18 July 2024].

43. S. Bravo Orellana, Carretera Interoceánica Sur del Perú. Retos e innovación. (2013).

44. Oficina de Estadística - OGPP, Ministerio de Transportes y Comunicaciones, MAPA VIAL

DEL PERU. (2007).

45. Oficina de Estadística, Oficina General de Planeamiento y Presupuesto - MTC, MAPA

VIAL - MADRE DE DIOS. (2009).

46. M. Johanson, A highway in the Amazon ushers in a new era of deforestation and social

upheaval. *World Wildl. Fund* (2024). Available at:

https://www.worldwildlife.org/stories/how-the-interoceanic-highway-ushered-in-a-new-era-

of-deforestation-and-social-upheaval-in-the-amazon [Accessed 29 August 2024].

47. B. Rockhill, B. Newman, C. Weinberg, Use and misuse of population attributable fractions.

Am. J. Public Health **88**, 15–19 (1998).

Figure legends

Figure 1: A map of Madre de Dios (A) displaying the Interoceanic highway (black line),

unpaved roads (grey lines), rivers (blue lines), and the locations of all healthcare facility clusters

colored by their exposure group (points). Clusters within 5km of the road are considered

exposed, those >10km are considered unexposed, and those between 5-10km from the road were

excluded from analyses as a buffer against spillover effects of exposure to the highway. Average

dengue (B) and leishmaniasis (C) yearly incidence rates (cases per person per year) over the time

period of interest colored by exposure group (buffer clusters are not included). Vertical dotted

line at 2008 represents the last year before the highway was paved, which is the baseline year in

our analyses.

Figure 2: Dengue incidence increased substantially in healthcare facility clusters near to the

highway compared to those far from the highway due to highway paving. Panel A displays

estimation results from Equation 1. Panel B displays results for the biannual models aggregated

across the full time period (Equation 2). Average treatment effects are represented by points, and

95% confidence intervals are represented by error bars (standard errors clustered at the unit

level). The vertical dotted line at 2008 represents the last year before the highway was paved

(baseline year). A table displaying yearly results (**STable S1**) and a figure displaying

disaggregated biannual estimates (**SFig S6**) can be found in the **Supplementary Information**.

Figure 3: Leishmaniasis incidence remains stable post highway paving in healthcare facility

clusters near to the highway compared to those far from the highway. Panel A displays

estimation results from Equation 1. Panel B displays results for the biannual models aggregated

across the full time period (Equation 2). Average treatment effects are represented by points, and

95% confidence intervals are represented by error bars (standard errors clustered at the unit

level). The vertical dotted line at 2008 represents the last year before the highway was paved

(baseline year). A table displaying yearly results (**STable S1**) and a figure displaying

disaggregated biannual estimates (**SFig S6**) can be found in the **Supplementary Information**.

Figure 4: Effect of the highway attenuates when communities are at distances of greater than

10km from the highway. Panel A displays estimation results from Equation 2, where a model

was run for each distance subgroup (colors). Average treatment effects are represented by points,

and 95% bootstrapped confidence intervals are represented by error bars (bootstrapped standard

errors clustered at the unit level). Panel B displays a map of the Madre de Dios region with

healthcare facility clusters colored by their distance subgroup (i.e., 1km, 5km, etc.).

Highway paving dramatically increased dengue transmission in the Amazon

Supplementary Information

Supplementary Text

Disease diagnosis

As defined by the Peruvian Ministry of Health, laboratory confirmed dengue cases are cases that produce a positive result from one or more of the following tests: "1) DENV isolation by cell culture, 2) qRT-PCR, 3) ELISA antigen NS1, 4) detection of IgM antibodies for dengue in a single sample by ELISA in dengue endemic areas, and/or 5) evidence of IgM seroconversion in paired samples, where the second sample should be taken after 14 days of the onset of symptoms in areas where there is no transmission of dengue (and these cases should have an epidemiological investigation)" (1). Probable dengue cases are clinically diagnosed and must either meet case definition criteria set by the World Health Organization (2) or have a direct epidemiological link to a confirmed case (1).

Laboratory confirmed leishmaniasis cases are defined as cases producing a positive result from "1) a parasitological examination, 2) an immunological or histopathological clinical , and/or 3) a culture, whether or not in the form of a pruritic nodule." Probable cases of cutaneous leishmaniasis were clinically diagnosed and defined as, "any person with progression to ulcerative or ulcerous- crusted lesions, shallow, rounded, non-painful, with well-defined borders and inflammatory signs, with evolution time of no less than four weeks and lack of response to conventional treatment. With a history of origin or residence in … areas endemic to leishmaniasis." Probable cases of mucocutaneous leishmaniasis were clinically diagnosed and defined as, "any person with a clinical picture characterized by raised granulomatous or ulcerative lesions of the nasal mucosa, mouth, white palate, pharynx, larynx or trachea, with a history of active or healed skin lesions, origin or residence in endemic areas (3, 4)."

Clustering

We identified latitude and longitude for all 105 healthcare facilities using a combination of government sources and the *ubigeo* package in R (5, 6) (**SFig S7A**). We then grouped healthcare facilities into 71 clusters to standardize the size of their catchment areas and created non-overlapping boundaries to extract climate and land use data (**SFig S7B**). Clustering was done using an iterative hierarchical clustering approach that joined healthcare facilities in groups within a pre-specified distance (*hclust* and *cutree* functions from the *stats* R package). We used a pre-specified distance of 7.5km and visually checked that this provided sufficient separation between clusters (**SFig S7B**). Finally, we created non-overlapping shapefile boundaries with the goal of capturing all populated and non-forested areas near each of the healthcare facility clusters. We built these boundaries by creating circular buffer zones with a radius of 7.5km and then assigning each pixel within the buffer zones to the closest healthcare facility cluster, eliminating any overlap between clusters (done using the *distanceFromPoints* function from the *raster* package in R). A radius of 7.5km was chosen to ensure all population centers and non-forested land would be captured (determined via visualization of satellite images and WorldPop

raster data in Google Earth Engine (7)).

Population calculation

Locality population data was also provided by DIRESA for the years 2009–2017 and 2020–2022. This population data was linked to case data through the locality names in the disease case dataset and then clustered as described above. In years without DIRESA-provided population data, we imputed missing values with WorldPop data (7) by multiplying each cluster-year WorldPop value by the corresponding cluster-specific average ratio between DIRESA and WorldPop values. WorldPop is a remotely-sensed population dataset that estimates the number of people living in each 100m pixel from 2000–2019 (7). It is important to measure annual changes in each cluster's population due to the rapid in-migration and population growth in the area during the study period (8) (**SFig S5**). We use these population measures to derive incidence rates from disease case counts as a measure of changes in per-capita dengue risk, above and beyond changes in raw case counts that would be expected due to population growth alone.

Supplementary Table S1: Dengue and leishmaniasis yearly estimation results (Equation 1). Standard errors are clustered at the unit level for all models and are displayed inside parentheses next to each estimated coefficient. P-values indicating a statistical significance level below 0.05 are designated with an asterisk (*).

	Dengue Yearly	Leish Yearly
5km x 2000	-0.47* (0.16)	-0.60* (0.27)
5km x 2001	-0.25 (0.22)	0.09 (0.29)
5km x 2002	-0.15 (0.21)	-0.84* (0.28)
5km x 2003	-0.14 (0.21)	-0.12 (0.29)
5km x 2004	-0.04 (0.16)	-0.50 (0.26)
5km x 2005	0.02 (0.17)	-0.02 (0.24)
5km x 2006	0.03 (0.18)	-0.44 (0.23)
5km x 2007	0.08 (0.12)	-0.64* (0.23)
5km x 2008	----	----
5km x 2009	0.56* (0.23)	0.14 (0.21)
5km x 2010	1.2* (0.37)	-0.20 (0.20)
5km x 2011	0.91* (0.31)	-0.09 (0.24)
5km x 2012	1.2* (0.34)	0.42 (0.30)
5km x 2013	0.92* (0.35)	0.60* (0.29)
5km x 2014	0.57 (0.30)	0.31 (0.28)
5km x 2015	0.45 (0.23)	0.40 (0.36)
5km x 2016	0.38 (0.22)	0.44 (0.23)
5km x 2017	0.18 (0.25)	0.34 (0.38)
5km x 2018	1.1* (0.39)	0.71 (0.38)
5km x 2019	1.7* (0.51)	0.78 (0.45)
5km x 2020	1.5* (0.30)	0.44 (0.31)
5km x 2021	0.98* (0.34)	0.38 (0.29)
5km x 2022	0.96* (0.42)	0.60 (0.33)
log(Urban)	0.17 (0.10)	-0.03 (0.08)
log(Ag)	-0.22 (0.17)	0.38* (0.14)
Mean Precip	0.05 (0.19)	-0.06 (0.13)
Mean Temp	0.22 (0.39)	0.40 (0.30)
FEs		
Cluster	X	X
Year	X	X
Cor^2	0.561	0.606
N	1,494	1,495
Clusters	65	65

STable S2: Dengue long-differences estimation results (Equation 2) are displayed for the following model specifications: main model, model excluding Puerto Maldonado (the region's largest city), population-weighted model also excluding Puerto Maldonado, model designating <1km from the highway as exposed and >1km from the highway as unexposed, model designating <10km as exposed and >10km as unexposed, model with no spatial buffer (<5km exposed, >5km unexposed), model with a larger spatial buffer (<5km exposed, >20km unexposed), model including both confirmed and probable cases, and model including only spatial units reporting at least one disease case both pre and post road paving. All models include unit and year fixed effects. Standard errors are clustered at the unit level for all models and are displayed inside parentheses next to each estimated coefficient. P-values indicating a statistical significance level below 0.05 are designated with an asterisk (*).

	Main model	No PM	Pop weight no PM	1km boundary	10km boundary	No buffer (<5km v >5km)	Large buffer (<5km v >20km)	Conf & prob cases	Units reporting pre & post
5km x Post-2008	1.0* (0.28)	0.98* (0.28)	1.3* (0.60)			0.91* (0.28)	1.1* (0.27)	1.1* (0.28)	1.2* (0.35)
1km x Post-2008				0.62* (0.31)					
10km x Post-2008					1.1* (0.26)				
log(Urban)	0.09 (0.10)	0.09 (0.10)	0.29 (0.33)	0.13 (0.12)	0.11 (0.10)	0.14 (0.12)	0.28 (0.18)	0.09 (0.11)	0.07 (0.15)
log(Ag)	-0.23 (0.14)	-0.27 (0.14)	0.32 (0.24)	-0.20 (0.14)	-0.20 (0.13)	-0.20 (0.13)	-0.35* (0.15)	-0.25 (0.15)	-0.27 (0.15)
Sum Precip	0.09 (0.24)	0.04 (0.24)	-0.53 (0.33)	0.06 (0.22)	0.07 (0.22)	0.08 (0.22)	0.15 (0.26)	0.07 (0.24)	-0.03 (0.25)
Mean Temp	-0.05 (0.47)	-0.10 (0.48)	1.2* (0.61)	0.11 (0.46)	0.14 (0.46)	0.16 (0.46)	0.08 (0.48)	-0.06 (0.53)	0.12 (0.48)
Cor^2	0.54541	0.55571	0.5468	0.55196	0.55612	0.55421	0.56119	0.58093	0.61472
N	974	959	959	1,064	1,064	1,064	854	974	735
Clusters	65	64	64	71	71	71	57	65	49

STable S3: Dengue and leishmaniasis incidence long-differences estimation results (Equation 2) with precipitation and temperature covariates as quadratic functions (rather than linear). Corresponds to **Table 1** in main text. Yearly models include cluster and yearly fixed effects, and biannual models include cluster and biannual fixed effects. Standard errors are clustered at the unit level for all models and are displayed inside parentheses next to each estimated coefficient. P-values indicating a statistical significance level below 0.05 are designated with an asterisk (*).

	Dengue Yearly	**Dengue Biannual Dry**	**Dengue Biannual Rainy**	**Leish Yearly**	**Leish Biannual Dry**	**Leish Biannual Rainy**
5km x Post-2008 Yearly	1.0* (0.28)			0.42 (0.24)		
5km x Post-2008 Biannual		0.52* (0.21)	1.0* (0.24)		0.47* (0.20)	0.17 (0.15)
log(Urban)	0.09 (0.10)	0.10 (0.07)	0.06 (0.09)	-0.004 (0.07)	0.03 (0.04)	-0.04 (0.04)
log(Ag)	-0.23 (0.14)	-0.17 (0.12)	-0.24 (0.15)	0.23 (0.15)	0.14 (0.14)	0.21 (0.11)
Sum Precip^2	0.01 (0.03)	0.19 (0.27)	-0.002 (0.04)	-0.05* (0.01)	-0.36* (0.12)	-0.03* (0.01)
Mean Temp^2	-9.6e-5 (0.0008)	0.0003 (0.0004)	-0.0001 (0.0008)	0.002* (0.0007)	0.0009* (0.0004)	0.0006 (0.0006)
FEs						
Cluster	X	X	X	X	X	X
Year	X			X		
Biannual		X	X		X	X
Cor^2	0.54538	0.33792	0.54779	0.53878	0.5937	0.55868
N	974	1,040	1,040	975	1,040	1,040
Clusters	65	65	65	65	65	65

Supplementary Figure S1: Bar plots displaying disease case data coverage: number of healthcare facility clusters (i.e., spatial units) reporting at least one dengue (Panel A) or leishmaniasis (Panel B) case by year and exposure group (color). All clusters are included in this plot to display dengue dynamics throughout the region (i.e., clusters >5km and <10km from the highway are not excluded). Prior to paving, dengue was reported in Puerto Maldonado, Iberia, Mavila, Mazuko, Iñapari, Planchon, Loero, Laberinto, and Boca Colorado.

Supplementary Figure S2: Time series plots of yearly dengue incidence rates for regions bordering and otherwise relevant to Madre de Dios (9–11). Interoceanic Highway colored in red. Clockwise from top right: Acre, Brazil, dengue-endemic pre-2009 and bordering to the Northeast; Pando, Bolivia, dengue-endemic pre-2009 and bordering to the East (no data pre 2014); Madre de Dios, Peru, study setting; Cusco, Peru, only a small portion of the department is suitable for dengue, bordering to the West; Loreto, Peru, dengue-endemic and the only other substantially populated Amazonian department in Peru, not bordering Madre de Dios.

Supplementary Figure S3: Model results are robust to various model specifications. Estimation results from Equation 1 for the following model specifications (colors): main model, model excluding Puerto Maldonado (the region's largest city), population-weighted model also excluding Puerto Maldonado, model designating <1km from the highway as exposed and >1km from the highway as unexposed, model designating <10km as exposed and >10km as unexposed, model with no spatial buffer (<5km exposed, >5km unexposed), model with a larger spatial buffer (<5km exposed, >20km unexposed), model including both confirmed and probable cases, and model including only spatial units reporting at least one disease case both pre and post road paving. Average treatment effect (points) with 95% confidence interval (error bars). Standard errors clustered at the unit level. Vertical dotted line at 2008 representing the last year before the highway was paved (baseline year).

Supplementary Figure S4: Permutation inference analysis. Histograms of bootstrapped effect estimates from three methods of randomly shuffling the treatment variable (grey bars) compared to the estimated effect from the non-bootstrapped sample (vertical red line). The three methods include: A) Random, the entire treatment column is randomly shuffled across clusters and years, B) Full, randomly assign clusters to treatment, maintaining temporal structure, and C) Within, randomly assign years as treated within clusters, maintaining spatial structure.

Supplementary Figure S5: Time series plots displaying yearly measures of population, urban area, and agricultural area by exposure group (color). Yellow line displays the 'exposed' group (pink) excluding Puerto Maldonado (PM), the largest city in the region. Sources and calculations of these plots are discussed in the **Methods** section and in the **Supplementary Information**.

<u>Supplementary Figure S6:</u> Plots display disaggregated biannual model results for dengue (A) and leishmaniasis (B) from Equation 1, colored by season (rainy v dry). Average treatment effects are represented by points, and 95% confidence intervals are represented by error bars (standard errors clustered at the unit level). The vertical dotted line at 2008 represents the last year before the highway was paved (baseline year).

Supplementary Figure S7: Panel A shows all healthcare facilities (*n* = 105, blue points) mapped onto the study region. Panel B shows healthcare facility clusters (*n* = 71, pink points) and constructed spatial boundaries (grey polygons) mapped onto the study region.

— Unpaved Roads — Highway — Rivers

<u>Supplementary Figure S8:</u> Panel A shows healthcare facility clusters clustered with a radius of 7.5km ($n = 71$, pink points, main model). Alternative clustering specifications with radii 6km and 5km are shown in Panels B ($n = 78$, yellow points) and C ($n = 84$, blue points), respectively. Panel D shows estimation results from Equation 1 for the differing clustering specifications (colors same). Average treatment effect (points) with 95% confidence interval (error bars). Standard errors clustered at the unit level. Vertical dotted line at 2008 representing the last year before the highway was paved (baseline year).

Supplementary Figure S9: Seasonality of monthly accumulated precipitation (A) and dengue cases (B) in Madre de Dios, Peru during the study period (2000–2022). The red horizontal line in panel A shows the monthly accumulated precipitation across the entirety of the study period. The red vertical lines show the months that intersect with the red horizontal line delineating rainy (October – March) versus dry season (April – September).

References

1. Ministerio de Salud del Perú, Norma técnica de salud para la vigilancia epidemiológica Y diagnóstico de laboratorio de dengue, chikungunya, zika y otras arbovirosis en el Perú. (2017). Available at: https://www.dge.gob.pe/portal/docs/tools/arbovirosis18.pdf.

2. World Health Organization (WHO), Special Programme for Research and Training in Tropical Diseases (TDR), Dengue guidelines, for diagnosis, treatment, prevention and control. (2009).

3. Normas y procedimientos para el control de las leishmaniasis en el Perú. Available at: https://www.gob.pe/institucion/minsa/informes-publicaciones/353504-normas-y-procedimientos-para-el-control-de-las-leishmaniasis-en-el-peru [Accessed 22 August 2024].

4. Ministerio de Salud del Perú, Leishmaniasis - Módulos Téchnicos. (2000). Available at: https://bvs.minsa.gob.pe/local/OGEI/795_MS-OGE106.pdf.

5. J. Castagnetto, ubigeo. (2021). Available at: https://github.com/jmcastagnetto/ubigeo/?tab=readme-ov-file [Accessed 18 July 2024].

6. Plataforma Nacional de Datos Abiertos. *Plataforma Nac. Datos Abiertos* (2024). Available at: https://www.datosabiertos.gob.pe/dataset/establecimientos-de-salud [Accessed 18 July 2024].

7. Open Spatial Demographic Data and Research. *WorldPop* (2024). Available at: https://www.worldpop.org/ [Accessed 19 June 2024].

8. K. E. Jensen, *et al.*, Small scale migration along the interoceanic highway in Madre de Dios, Peru: an exploration of community perceptions and dynamics due to migration. *BMC Int. Health Hum. Rights* **18**, 12 (2018).

9. C. Cabezas, *et al.*, Dengue en el Perú: a un cuarto de siglo de su reemergencia. *Rev. Peru. Med. Exp. Salud Pública* 146–156 (2015). https://doi.org/10.17843/rpmesp.2015.321.1587.

10. L. A. Gutiérrez, https://www.facebook.com/pahowho, PAHO/WHO Data - Bolivia - Casos de dengue | OPS/OMS. *Pan Am. Health Organ. World Health Organ.* (2015). Available at: https://www3.paho.org/data/index.php/es/temas/indicadores-dengue/dengue-subnacional/536-bol-dengue-casos-es.html [Accessed 16 August 2024].

11. (DATASUS, Ministério da Saúde do Brasil, 2022), Doenças e Agravos de Notificação – 2007 em Diante (SINAN). Available at: https://datasus.saude.gov.br/acesso-a-informacao/doencas-e-agravos-de-notificacao-de-2007-em-diante-sinan/ [Accessed 30 October 2024].

A

B Dengue incidence

C Leish incidence

● Exposed (<5km)   ● Unexposed (>10km)   ○ Buffer (removed)   — Unpaved Roads   **▬** Highway   — Rivers

A

Percent change dengue incidence

B

Legend

● 1km ● 10km ● 20km
● 5km ● 15km ● Control

100 km

N