Reassessing the link between adiposity and head and neck cancer: a Mendelian randomization study

Fernanda Morales-Berstein[1,2], Jasmine Khouja[1,2], Mark Gormley[1-3], Elmira Ebrahimi[4], Shama Virani[4], James McKay[4], Paul Brennan[4], Tom G Richardson[1,2], Caroline L Relton[1,2,5], George Davey Smith[1,2], M Carolina Borges[1,2], Tom Dudding[1-3], Rebecca C Richmond[1,2]

[1]MRC Integrative Epidemiology Unit, University of Bristol, Bristol, United Kingdom.
[2]Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, United Kingdom.
[3]University of Bristol Dental School, 1 Trinity Walk, Avon Street, Bristol, United Kingdom.
[4]Genomic Epidemiology Branch, International Agency for Research on Cancer, World Health Organization, Lyon, France.
[5]London School of Hygiene & Tropical Medicine, Keppel Street, London WC1E 7HT, United Kingdom.

For correspondence: dy20206@bristol.ac.uk

Keywords: Head and neck cancer, adiposity, body mass index, waist-to-hip ratio, waist circumference, Mendelian randomization.

1 Abstract

Background: Adiposity has been associated with an increased risk of head and neck cancer

(HNC). Although body mass index (BMI) has been inversely associated with HNC risk among

smokers, this is likely due to confounding. Previous Mendelian randomization (MR) studies

could not fully discount causality between adiposity and HNC due to limited statistical power.

Hence, we aimed to revisit this using the largest genome-wide association study (GWAS) of

HNC available, which has more granular data on HNC subsites.

Methods: We assessed the genetically predicted effects of BMI (N=806,834), waist-to-hip

ratio (WHR; N=697,734) and waist circumference (N=232,101) on the risk of HNC (N=12,264

cases and 19,259 controls) and its subsites (oral, laryngeal, hypopharyngeal and

oropharyngeal cancers) using a two-sample MR framework. We used the inverse variance

weighted (IVW) MR approach and multiple sensitivity analyses including the weighted

median, weighted mode, MR-Egger, MR-PRESSO, and CAUSE approaches. We also used

multivariable MR (MVMR) to explore the direct effects of the adiposity measures on HNC,

while accounting for smoking behaviour, a well-known HNC risk factor.

Results: In univariable MR, higher genetically predicted BMI increased the risk of overall

HNC (IVW OR=1.17 per 1 standard deviation [1-SD] higher BMI, 95% CI 1.02–1.34, p=0.03),

with no heterogeneity across subsites (Q p=0.78). However, the effect was not consistent in

sensitivity analyses. The IVW effect was attenuated when smoking was included in the

MVMR model (OR accounting for comprehensive smoking index=0.96 per 1-SD higher BMI,

95% CI 0.80–1.15, p=0.64) and CAUSE indicated the IVW results could be biased by

correlated pleiotropy. Furthermore, we did not find a link between genetically predicted

WHR (IVW OR=1.05 per 1-SD higher WHR, 95% CI 0.89–1.24, p=0.53) or waist circumference

and HNC risk (IVW OR=0.83 per 1-SD higher waist circumference, 95% CI 0.56–1.23, p=0.35).

Conclusions: Our findings suggest that adiposity does not play a role in HNC risk.

Introduction

Head and neck cancer (HNC) is among the ten most common cancers in Europe, with an age standardised incidence rate of 10.3 per 100,000 person-years[1]. Around 90% of HNCs are classed as squamous cell carcinomas of the oral cavity, pharynx or larynx[2]. Tobacco smoking and alcohol consumption are well-established HNC risk factors[3-9]. High-risk human papillomavirus (HPV) infection has also been causally linked to the risk of HNC, especially oropharyngeal cancer[10-12]. In contrast, the role of adiposity in the development of HNC is less well understood.

The World Cancer Research Fund/American Institute for Cancer Research (WCRF/AICR) Continuous Update Project (CUP) Expert Report published in 2018 determined higher body fatness (i.e., body mass index [BMI], waist-to-hip ratio [WHR] and waist circumference) likely increases the risk of HNC[13]. The CUP panel reached this conclusion even though a higher BMI has been associated with a decreased risk of HNC[13], since they noted the inverse association appears to be limited to current smokers. They concluded the association between BMI and HNC risk may be biased among individuals who smoke (because smoking is a HNC risk factor associated with lower weight). It is thought that nicotine consumption could lead to appetite suppression and increased energy expenditure, which could in turn, lead to weight loss (and spurious inverse associations between BMI and HNC risk[14]) among smokers[15]. Among never smokers, BMI has been positively associated with HNC risk, in line with the evidence observed for measures of central adiposity (i.e., WHR and waist circumference)[14, 16].

However, the relationship between adiposity and smoking is complex, with evidence from

Mendelian randomization (MR) studies suggesting higher adiposity increases the risk of

smoking[17, 18] while simultaneously suggesting smoking may lead to lower adiposity[17,

19-22]. Additionally, excess adiposity, socioeconomic deprivation and (both active and

passive) smoking are often strongly correlated[23-26]. Thus, the positive associations

between adiposity (i.e., BMI among non-smokers, WHR and waist circumference) and HNC

risk may not be as unbiased as they appear.

It is important to acknowledge that previous MR studies on adiposity (i.e., BMI, WHR, waist

circumference) and HNC risk were relatively small (maximum N=6,034 cases) and could not

fully discount causality due to limited statistical power[27-29]. Therefore, the aim of this MR

study was to revisit the link between adiposity and HNC risk using data from a HNC genome-

wide association study (GWAS) that includes over two times the number of cases than the

Genetic Associations and Mechanisms in Oncology (GAME-ON) GWAS[30] used by Gormley

et al.[28] (the largest to date; N=12,619, including 6,034 cases and 6,585 controls) and has

more granular data on HNC subsites (i.e., oral cavity, hypopharynx, oropharynx and larynx).

We also aimed to use multivariable MR (MVMR) to explore the direct effects of the adiposity

measures on HNC, while accounting for smoking behaviour.

Methods

Study design

We used a two-sample MR framework to assess the genetically predicted effects of BMI, WHR and waist circumference on the risk of HNC and its subsites (oral, laryngeal, hypopharyngeal and oropharyngeal cancers) among individuals of European ancestry. Genetic variants associated with these adiposity traits were used as instrumental variables to estimate causal effects under the three core MR assumptions[31]: 1) the genetic variants are strongly associated with the adiposity trait of interest (relevance assumption); 2) the distribution of the genetic variants in the population is not influenced by factors that also influence HNC risk, such as population stratification, assortative mating and dynastic effects (independence assumption); and 3) the genetic variants can only influence HNC risk via their effect on the adiposity trait of interest (exclusion restriction assumption). This work was conducted and reported according to the STROBE-MR guidelines[32] (**Additional File 1: Appendix**).

Head and neck cancer GWAS

GWAS summary statistics for HNC were obtained from a European HEADSpAcE consortium GWAS that excluded UK Biobank participants (N=31,523, including 12,264 cases and 19,259 controls) to avoid overlapping samples across the exposure and outcome datasets. It includes the European GAME-ON data used by Gormley et al.[30] and has more granular data on HNC subsites (i.e., oral cavity [N=21,269, including 3,091 cases and 18,178 controls], hypopharynx [N=18,652, including 474 cases and 18,178 controls], HPV positive oropharynx [N=20,146, including 1,980 cases and 18,166 controls], HPV negative oropharynx [N=19,114,

including 948 cases and 18,166 controls] and larynx [N=20,668, including 2,490 cases and

18,178 controls])[33].

HNC was defined based on the 10[th] revision of the International Classification of Diseases

(ICD-10)[34]. It included cancers of the oral cavity (C00.3, C00.4, C00.5, C00.6, C00.8, C00.9,

C02.0–C02.9 [except C02.4], C03.0–C03.9, C04.0–C04.9, C05.0–C06.9 [except C05.1, C05.2]),

the oropharynx (C01-C01.9, C02.4, C05.1, C05.2, and C09.0–C10.9), the hypopharynx (C12.0-

C13.0), the larynx (C32), and overlapping or not otherwise specified sites (C14, C05.8, C02.8,

C76.0)[33].

Further detail on the HEADSpAcE GWAS has been published elsewhere[33]. In brief,

genotype data were obtained using nine different genotyping arrays. They were

subsequently converted to genome build 38 for consistency across datasets. Quality control

(QC) procedures were conducted by genotyping array rather than by study. Samples were

excluded for the following reasons: sex mismatch (heterozygosity <0.8 for males and >0.2 for

females), autosomal heterozygosity (>3 standard deviation [SD] units from the mean),

missingness (>0.03), and cryptic relatedness (identity-by-decent >0.185). Single nucleotide

polymorphisms (SNPs) were removed due to genotype missingness (>0.01), deviations from

Hardy-Weinberg equilibrium (p <1e-05) and low minor allele count (<20). Imputation was

performed using the TOPMed Imputation Server. Only SNPs with an imputation score r^2 >0.3

and a minor allele frequency (MAF) >0.005 were included in the GWAS. The analyses were

conducted in PLINK using logistic regressions adjusted for sex, the top principal components

and imputation batch (six in total, which account for both genotyping array and study

differences).

Genetic instruments for adiposity

GWAS summary statistics for waist circumference (N=232,101) in SD units were obtained

from the Genetic Investigation of Anthropometric Traits (GIANT) meta-analysis by Shungin et

al.[35] available via the IEU OpenGWAS platform (id: ieu-a-61). GWAS summary statistics for

BMI (N=806,834) and WHR (N=697,734) in SD units were obtained from the latest GIANT

consortium's GWAS meta-analysis by Pulit et al.[36] available at

https://zenodo.org/records/1251813. The meta-analysis is the biggest to date, as it

combines the meta-analysis by Shungin et al.[35] with UK Biobank data. The UK Biobank

GWAS[36] was conducted using imputed data and the BOLT-LMM software[37]. The linear

mixed models (LMMs) were solely adjusted for genotyping array. GIANT and UK Biobank

data were meta-analysed[36] using an inverse-weighted fixed-effect meta-analysis in

METAL[38].

We extracted GWAS-significant SNPs for waist circumference using the standard threshold

(p<5e-08). For BMI and WHR, we extracted them according to the stringent threshold

recommended by Pulit et al.[36] to account for denser imputation data (p<5e-09). We then

performed LD-clumping to select independent lead SNPs for each exposure (r^2=0.001,

10,000 kb). In total, 458 and 283 and 41 SNPs remained for BMI, WHR and waist

circumference, respectively.

Data harmonisation

We extracted HNC GWAS summary statistics that corresponded to the list of SNPs selected

as instruments for the exposures. Proxy SNPs (r^2>0.8) were used when the instrumental

SNPs were not available in the outcome datasets. Proxies were identified using the

"extract_outcome_data" function of the "TwoSampleMR" R package and the 1000 Genomes

Project European reference panel. We harmonised the exposure and outcome datasets

using the "harmonise_data" function of the "TwoSampleMR" R package[39]. Positive strands

were inferred using allele frequencies and ambiguous palindromic SNPs with MAFs ≥0.3

were removed. The harmonised data used in the analyses are available in **Supplementary**

Tables 1 to 6 (Additional File 2).

We calculated mean F-statistics and total R^2 values to assess the strength of our genetic

instruments after data harmonisation[40, 41].

Statistical analysis

Main analyses

156 The multiplicative random effects inverse-variance weighted (IVW) MR approach[42] (the

157 default IVW method of the "TwoSampleMR" package[39]) was used to investigate the

158 genetically predicted effects of BMI, WHR and waist circumference on HNC risk. We did not

159 correct our results for multiple testing, as all our exposures are strongly correlated[35, 36].

Sensitivity analyses

Because the IVW method assumes all genetic variants are valid instruments[42], which is

unlikely the case, three pleiotropy-robust two-sample MR methods (i.e., MR-Egger[43],

weighted median[44] and weighted mode[45]) were used in sensitivity analyses. We also

performed tests for SNP heterogeneity (i.e., Q statistic test)[46] and directional horizontal

pleiotropy (i.e., MR-Egger intercept test)[43]. When directional horizontal pleiotropy was

identified, we used the intercept value to evaluate the extent of the bias. The MR-

PRESSO[47] method was used to identify outliers (outlier test p<0.05) and calculate outlier-

corrected causal estimates when there was evidence of SNP heterogeneity. The MR-PRESSO

distortion test was used to evaluate differences between the outlier-corrected and IVW

estimates.

In addition, we ran MVMR[48] analyses to evaluate the direct effects of adiposity measures

with evidence of a total effect in our main analyses. The aim of the MVMR analyses was to

separate the effect of adiposity from smoking behaviour (a well-known HNC risk factor

which has a complex relationship with adiposity) in the development of HNC. We obtained

genetic instruments for smoking behaviour from two different sources: a smoking initiation

GWAS (N=805,431 excluding 23andme) derived by the GWAS and Sequencing Consortium of

Alcohol and Nicotine use (GSCAN)[49] and a comprehensive smoking index (CSI; a measure

of lifetime smoking that captures smoking heaviness, duration and cessation) GWAS

(N=462,690) conducted by Wootton et al.[50]. Each was separately investigated in a MVMR

framework. For each smoking trait, we selected SNPs that passed the GWAS-significance and

independence thresholds (p<5e-08, r^2=0.001, 10,000 kb) and combined them with the list of

SNPs identified as instruments for the relevant exposure. We then performed LD-clumping

across the combined list of SNPs, to then use these independent SNPs in MVMR analyses.

The exposure and outcome datasets were harmonised to the same effect allele using the

"harmonise_data" function of the "TwoSampleMR" R package[39]. We formatted the data

using the "format_mvmr" function of the "MVMR" R package, calculated the conditional F-

statistics for the MVMR instruments using the "strength_mvmr" function and ran the MVMR

analyses using the "ivw_mvmr" function.

Since we used large GWAS datasets for the selection of our genetic instruments, our

analyses are at an increased likelihood of being biased due to correlated horizontal

pleiotropy (sometimes referred to as heritable confounding), which occurs when the genetic

instruments are associated with the exposure through their effect on confounders of an

exposure-outcome association[51, 52]. To mitigate this bias, we used Steiger filtering[53] to

remove SNPs that are more strongly associated with smoking behaviour (a confounder of

the exposure-outcome association) than the exposure of interest, as proposed by Sanderson

et al.[54]. We also used Causal Analysis using Summary Effect Estimates (CAUSE)[52],

another pleiotropy-robust MR method, to further investigate whether our results could be

biased by correlated horizontal pleiotropy. CAUSE uses Bayesian expected log pointwise

posterior predictive densities (ELPDs) to compare null, sharing and causal models. A higher

ELPD represents a better model fit, so a positive delta ELPD (where delta ELPD = ELPD model

1 - ELPD model 2) suggests model 1 fits the data better than model 2, while a negative delta

ELPD suggests the opposite. If we find evidence to reject the null hypothesis that the sharing

model (i.e. causal effect fixed at zero) fits the data at least as well as the causal model (i.e.

causal effect can differ from zero), our findings would be consistent with a causal effect.

Steiger filtering and CAUSE analyses were only conducted for adiposity measures with

evidence of a total effect in our main IVW analyses.

Moreover, we used the MR-Clust algorithm[55] to find distinct SNP clusters underlying the

relationship between adiposity measures with evidence of a total effect in our main analyses

and HNC. The identification of substantial clusters could provide insight into potential causal

mechanisms. It could also flag pleiotropic variables that are associated with SNPs in each

cluster. We filtered SNPs with conditional probabilities <0.8. At least four SNPs needed to

remain per cluster for a substantial cluster to be reported.

Secondary analyses

In secondary analyses, we investigated the role of BMI, WHR and waist circumference on the

risk of HNC by subsite (i.e., oral cavity, hypopharynx, HPV positive oropharynx, HPV negative

oropharynx and larynx). We used a Cochran's Q test to examine heterogeneity across HNC

subsites.

We also explored the role of other adiposity-related anthropometric measures on the risk of

HNC and its subsites. These anthropometric measures included: 1) four body shape principal

components[56], 2) childhood and adulthood body size[57], 3) metabolically favourable and

unfavourable adiposity[58], 4) body fat percentage, and 5) brain and adipose tissue-specific

BMI[59]. The data sources for these traits are summarised in **Table 1**.

Table 1. Data sources and instruments for other adiposity-related anthropometric measures.

Study	Year	Data source	Trait	Unit	Download link or OpenGWAS ID
Ried et al.[56]	2016	GIANT	Body shape PC1 (overall adiposity)	SD	https://www.joelhirschhornlab.org/giant-consortium-results
			Body shape PC2 (tall and slim vs short and plump)	SD	
			Body shape PC3 (tall with small hip vs short with big hip)	SD	
			Body shape PC4 (high BMI and weight with small hip and waist vs low BMI and weight with big hip and waist)	SD	
Richardson et al.[57]	2020	UKB	Childhood body size	Change in body size category	"ieu-b-5107"
			Adulthood body size	Change in body size category	"ieu-b-5118"
Martin et al.[58]	2021	UKB	Metabolically favourable adiposity	SD	https://doi.org/10.2337/figshare.14555463

			Metabolically unfavourable adiposity	SD	.v1
MRC-IEU (Elsworth)	2018	UKB	Body fat percentage	SD	"ukb-b-8909"
Leyden et al.[59]	2022	GIANT + UKB	Brain tissue-specific BMI	SD	https://www.ncbi.nlm .nih.gov/pmc/articles /PMC8874216/bin/m mc2.xlsx
			Adipose tissue-specific BMI	SD	

231 Abbreviations: BMI, body mass index; GIANT, Genetic Investigation of Anthropometric Traits; N, number; PC, principal
232 component; SD, standard deviation; SNP, single-nucleotide polymorphism; UKB, UK Biobank.
233
234

Statistical software

236 We completed all MR analyses using R software version 4.4.0 and the "TwoSampleMR"

237 v0.6.3, "MRPRESSO" v1.0, "MVMR" v0.4, "cause" v1.2.0 and "mrclust" v0.1.0 R packages.

238 The "ggplot2" v3.5.1 and "ggforestplot" v0.1.0 R packages were used to create forest plots.

239 The code used to run the MR analyses is available at

240 http://github.com/fernandam93/adiposity_HNC_MR.

Results

Genetic instruments for BMI, WHR and waist circumference

After data harmonisation and the removal of ambiguous palindromic SNPs, 442 genetic

variants remained as instruments for BMI, while 267 remained for WHR and 37 for waist

circumference (**Additional File 2: Supplementary Table 1**). The mean F-statistic for BMI was

77 (range 33–844) and the total variance explained was 4.8%. For WHR, the mean F-statistic

was 73 (range 33–820) and the total variance explained was 3.1%. For waist circumference,

the mean F-statistic was 50 (range 29–144) and the total variance explained was 0.8%.

The F-statistics and R^2 values for the other adiposity-related anthropometric measures have

been summarised in **Table 2**.

Table 2. F-statistics and variance explained for other adiposity-related anthropometric
measures.

Trait	N SNPs before/after harmonisation	Total R^2	Mean F-statistics (range)
Body shape PC1 (overall adiposity)	29/28	16%	54 (28–302)
Body shape PC2 (tall and slim vs short and plump)	84/81	3.4%	54 (30–211)
Body shape PC3 (tall with small hip vs short with big hip)	28/27	0.9%	41 (30–82)
Body shape PC4 (high BMI and weight with small hip and waist vs low BMI and weight with big hip and waist)	10/ 10	24.7%	42 (30–98)
Childhood body size	206/198	3.4%	78 (28–1102)
Adulthood body size	339/324	4.2%	59 (30–1109)
Metabolically favourable adiposity	34/31	0.4%	64 (25–400)
Metabolically unfavourable adiposity	29/27	0.8%	131 (25–400)
Body fat percentage	377/360	4.7%	59 (30–682)
Brain tissue-specific BMI	140/133	1.2%	61 (29–270)
Adipose tissue-specific BMI	86 /81	0.7%	63 (30–270)

256 Abbreviations: BMI, body mass index; N, number; PC, principal component; SNP, single-nucleotide polymorphism.
257
258

Genetically predicted effects of BMI, WHR and waist circumference on HNC

risk

In univariable MR, higher genetically predicted BMI increased the risk of overall HNC (IVW

OR=1.17 per 1 standard deviation [1-SD] higher BMI, 95% CI 1.02–1.34, p=0.03), with no

heterogeneity across subsites (Q p=0.78) (**Figure 1, Supplementary Figure 1 and Additional**

File 2: Supplementary Table 7). However, the positive relationship between genetically

predicted BMI and HNC risk was not consistent across the MR-Egger, weighted median and

weighted mode analyses, with point estimates in opposing directions and confidence

intervals including the null. The Q statistic and MR-Egger intercept tests suggested that there

was heterogeneity across individual SNP estimates (Q=609, p<0.001) and a minor degree of

unbalanced horizontal pleiotropy (intercept=0.007, p=0.03) that could have biased the main

IVW results (**Additional File 2: Supplementary Tables 8 and 9**). Although the MR-PRESSO

analysis identified two outliers (i.e., rs11611246 and rs9603697), the distortion test

suggested the outlier-corrected estimate (outlier-corrected IVW OR=1.14 per 1-SD higher

BMI, 95% CI 1.00–1.30, p=0.05) was not statistically different to the main IVW estimate

(p=0.94) (**Additional File 2: Supplementary Table 10**).

Furthermore, we did not find a link between genetically predicted WHR and HNC risk (IVW

OR=1.05 per 1-SD higher WHR, 95% CI 0.89–1.24, p=0.53) and there was no heterogeneity

across subsites (Q p=0.15) (**Figure 2, Supplementary Figure 2 and Additional File 2:**

Supplementary Table 7). MR-Egger, weighted median and weighted mode results were

consistent with a null effect. The Q statistic and MR-Egger intercept tests suggested that

there was some evidence of SNP heterogeneity (Q=332, p=0.004) and unbalanced horizontal

pleiotropy (intercept=0.008, p=0.03) (**Additional File 2: Supplementary Tables 8 and 9**). The

MR-PRESSO analysis did not identify any significant outliers (**Additional File 2:**

Supplementary Table 10).

Similarly, we did not find a genetically predicted effect of waist circumference on HNC risk

(IVW OR=0.83 per 1-SD higher waist circumference, 95% CI 0.56–1.23, p=0.35) or evidence

of heterogeneity across subsites (Q p=0.60) (**Figure 3, Supplementary Figure 3 and**

Additional File 2: Supplementary Table 7). The MR-Egger, weighted median and weighted

mode consistently suggested the absence of an effect of waist circumference on HNC risk.

The Q statistic and MR-Egger intercept tests suggested SNP heterogeneity (Q=89, p<0.001)

but no unbalanced horizontal pleiotropy (intercept=-0.019, p=0.30) (**Additional File 2:**

Supplementary Tables 8 and 9). The MR-PRESSO analysis identified three outliers (i.e.,

rs3127553, rs6440003 and rs806794) but the distortion test suggested the outlier-corrected

estimate (outlier-corrected IVW OR=0.77 per 1-SD higher waist circumference, 95% CI 0.59–

1.01, p=0.07) was not statistically different to the main IVW estimate (p=0.73) (**Additional**

File 2: Supplementary Table 10).

MVMR estimates for BMI on HNC risk after accounting for smoking behaviour

The effect of BMI on HNC risk was attenuated when smoking behaviour was included in the

MVMR model (OR accounting for CSI=0.93 per 1-SD higher BMI, 95% CI 0.78–1.12, p=0.47;

OR accounting for smoking initiation=1.09 per 1-SD higher BMI, 95% CI 0.88–1.34, p=0.43)

(**Figures 4 and 5 and Additional File 2: Supplementary Table 11**). Genetically predicted

smoking behaviour increased the risk of HNC even after accounting for BMI (CSI OR

accounting for BMI=4.25 per 1-SD higher CSI, 95% CI 3.18–5.67, p<0.001; SI OR accounting

for BMI=2.10 per 1-SD higher SI, 95% CI 1.61–2.73, p<0.001). The conditional F-statistics for

the BMI estimates were 30.5 and 30.3 in the CSI and smoking initiation analyses,

respectively (**Additional File 2: Supplementary Table 11**). They were slightly lower for the

smoking behaviour estimates conditioning on BMI (13.4 and 19.5 in the CSI and smoking

initiation analyses, respectively).

MR estimate for BMI on HNC risk after Steiger filtering SNPs more strongly

associated with smoking behaviour than BMI

After removing six SNPs (i.e., rs10002111, rs2503185, rs264941, rs10858334, rs225882,

rs2273175) that were more strongly associated with smoking behaviour (i.e., CSI or smoking

initiation) than BMI, the genetically predicted effect of BMI on HNC risk slightly attenuated

towards the null (Steiger filtered IVW OR=1.14 per 1-SD higher BMI, 95% CI 1.00–1.31,

p=0.05) (**Additional File 2: Supplementary Table 12**).

CAUSE estimate for BMI on HNC risk

We did not find evidence against bias due to correlated pleiotropy, since the causal model did not fit the data much better than the sharing model (CAUSE OR 1.12 per 1-SD higher BMI, 95% credible interval 0.93–1.34, delta ELPD for sharing vs causal=-0.07, p=0.47) (**Supplementary Figure 4**). Interestingly, neither the sharing nor the causal model fitted the data much better than the null model (delta ELPD for null vs sharing=-0.39, p=0.36; and delta ELPD for null vs causal=-0.46, p=0.41).

MR-Clust estimates for the relationship between BMI and HNC risk

After filtering SNPs with conditional probabilities <0.8 and clusters with fewer than four SNPs (e.g., cluster 1, as only three of 17 SNPs remained after probability filtering), only a null cluster including 372 SNPs (424 before filtering) remained in the MR-Clust output for BMI and HNC risk (**Supplementary Figure 5 and Additional File 2: Supplementary Table 13**). Hence, the MR-Clust analysis did not reveal any mechanistic pathways underlying the effect observed.

Genetically predicted effects of other adiposity-related anthropometric measures on HNC risk

We did not find consistent evidence of genetically predicted effects of other anthropometric measures on HNC risk (**Supplementary Figures 6–10 and Additional File 2: Supplementary Table 14**). The IVW estimate for PC2 capturing a combination of taller height and slimmer waist suggested this body shape decreased HNC risk (OR=0.86, 95% CI 0.75–0.99, p=0.04) (**Supplementary Figure 6b**). Similarly, the IVW estimate for PC3 capturing a combination of taller height and narrower hips suggested this body shape also reduced HNC risk (OR=0.73,

344   95% CI 0.55–0.97, p=0.03) (**Supplementary Figure 6c**). However, these inverse relationships

345   were not consistent with results obtained using pleiotropy-robust methods (i.e., MR-Egger,

346   weighted median and weighted mode).

348   Discussion

In this MR study, we reaffirmed that there is no clear evidence of a genetically predicted

effect of adiposity (i.e., BMI, WHR and waist circumference) or related anthropometric

measures on the risk of HNC or its subsites. Although we found higher genetically predicted

BMI increased the risk of overall HNC in the main univariable MR analysis, this was not

consistent across the sensitivity analyses. Notably, the MVMR results suggested the main

analysis may have been biased by smoking (and/or related traits), as the effect disappeared

after accounting for smoking behaviour. The results obtained after Steiger filtering SNPs

more strongly associated with smoking behaviour than BMI suggested correlated pleiotropy

may have been partly biasing the BMI-HNC estimate. CAUSE, which is more robust to

correlated pleiotropy than the IVW method, further supported this hypothesis.

Our results are consistent with previous MR studies that suggest adiposity does not

influence HNC risk[27-29]. Gormley et al.[28] did not find a genetically predicted effect of

adiposity on combined oral and oropharyngeal cancer when investigating either BMI

(OR=0.89 per 1-SD, 95% CI 0.72–1.09, p=0.26), WHR (OR=0.98 per 1-SD, 95% CI 0.74–1.29,

p=0.88) or waist circumference (OR=0.73 per 1-SD, 95% CI 0.52–1.02, p=0.07) as risk factors.

Similarly, a large two-sample MR study by Vithayathil et al.[29] including 367,561 UK

Biobank participants (of which 1,983 were HNC cases) found no link between BMI and HNC

risk (OR=0.98 per 1-SD higher BMI, 95% CI 0.93–1.02, p=0.35). Larsson et al.[27] meta-

analysed Vithayathil et al.'s[29] findings with results obtained using FinnGen data to increase

the sample size even further (N=586,353, including 2,109 cases), but still did not find a

genetically predicted effect of BMI on HNC risk (OR=0.96 per 1-SD higher BMI, 95% CI 0.77–

1.19, p=0.69).

In our study, we found some evidence that the genetically predicted effect of BMI on HNC

risk was influenced by smoking. This could be due to the bidirectional relationship between

smoking and adiposity reported in previous MR studies[17-22] or due to their shared genetic

architecture[60, 61]. A strength of our study was that it was the first to exploit MVMR to

disentangle the effects of BMI and smoking behaviour on the risk of HNC and its subsites. An

advantage of our approach compared to conducting univariable MR analyses stratified by

smoking status is that the former does not induce collider bias and provides estimates of

direct effects irrespective of horizontal pleiotropy or mediation. Yet, we acknowledge the

smoking behaviour traits used in our MVMR analyses likely capture more than just smoking,

since some of the SNPs used to instrument these traits have been associated with risk-taking

phenotypes and socioeconomic factors[62-65]. This places limits in the inferences that can

be made about smoking in the context of mediation.

An important strength of our study was that the HEADSpAcE consortium GWAS used had a

large sample size which conferred more statistical power to detect effects of adiposity on

HNC risk compared to previous MR analyses[27-29], increasing one's confidence in the null

nature of the estimates. Furthermore, the availability of data on more HNC subsites,

including oropharyngeal cancers by HPV status, allowed us to investigate the relationship

between adiposity and HNC risk in more detail than previous MR studies which limited their

subsite analyses to oral cavity and overall oropharyngeal cancers[28, 66]. This is relevant

because distinct HNC subsites are known to have different aetiologies[67], although we did

not find evidence of heterogeneity across subsites in our study.

We acknowledge that a major limitation of MR studies, including ours, is that several

untestable assumptions are required to make accurate causal inferences. It is unlikely that

our findings were influenced by weak instrument bias (i.e., violating the relevance

assumption) because we used strong genetic instruments to proxy our adiposity traits.

However, the independence assumption of no genetic confounding and the exclusion

restriction assumption of no horizontal pleiotropy could have been violated.

While our study contributes valuable evidence on the role of adiposity in the development

of HNC, we recognise there is a need for additional research on the subject. Our study was

limited to individuals of European ancestry, so our findings should be replicated in other

ancestry groups before being generalised to non-European populations. Moreover, further

research is needed to understand the biology underlying the complex relationship between

smoking and adiposity, especially since it may be difficult to intervene on one without

influencing the other[17].

Conclusions

In conclusion, this study indicates that adiposity does not play a role in HNC risk. Although

we did not find strong evidence of a causal effect of adiposity on HNC, obesity is an

established risk factor for multiple cancers and other chronic diseases[27, 68, 69]. Hence

there is still value in aiming to reduce the levels of excess adiposity in the population.

References

418   1.   **Global Cancer Observatory: Cancer Today (version 1.1)**
419   [https://gco.iarc.who.int/today]
420   2.   Curado MP, Hashibe M: **Recent changes in the epidemiology of head and neck**
421   **cancer**. *Curr Opin Oncol* 2009, **21**(3):194-200.
422   3.   Hashibe M, Brennan P, Benhamou S, Castellsague X, Chen C, Curado MP, Dal Maso L,
423   Daudt AW, Fabianova E, Fernandez L *et al*: **Alcohol drinking in never users of tobacco,**
424   **cigarette smoking in never drinkers, and the risk of head and neck cancer: pooled**
425   **analysis in the International Head and Neck Cancer Epidemiology Consortium**. *J*
426   *Natl Cancer Inst* 2007, **99**(10):777-789.
427   4.   Lee YC, Boffetta P, Sturgis EM, Wei Q, Zhang ZF, Muscat J, Lazarus P, Matos E, Hayes
428   RB, Winn DM *et al*: **Involuntary smoking and head and neck cancer risk: pooled**
429   **analysis in the International Head and Neck Cancer Epidemiology Consortium**.
430   *Cancer Epidemiol Biomarkers Prev* 2008, **17**(8):1974-1981.
431   5.   Purdue MP, Hashibe M, Berthiller J, La Vecchia C, Dal Maso L, Herrero R, Franceschi S,
432   Castellsague X, Wei Q, Sturgis EM *et al*: **Type of alcoholic beverage and risk of head**
433   **and neck cancer--a pooled analysis within the INHANCE Consortium**. *Am J*
434   *Epidemiol* 2009, **169**(2):132-142.
435   6.   Hashibe M, Brennan P, Chuang SC, Boccia S, Castellsague X, Chen C, Curado MP, Dal
436   Maso L, Daudt AW, Fabianova E *et al*: **Interaction between tobacco and alcohol use**
437   **and the risk of head and neck cancer: pooled analysis in the International Head and**
438   **Neck Cancer Epidemiology Consortium**. *Cancer Epidemiol Biomarkers Prev* 2009,
439   **18**(2):541-550.
440   7.   Marron M, Boffetta P, Zhang ZF, Zaridze D, Wunsch-Filho V, Winn DM, Wei Q,
441   Talamini R, Szeszenia-Dabrowska N, Sturgis EM *et al*: **Cessation of alcohol drinking,**
442   **tobacco smoking and the reversal of head and neck cancer risk**. *Int J Epidemiol* 2010,
443   **39**(1):182-196.
444   8.   Humans IWGotEoCRt: **Tobacco smoke and involuntary smoking**. *IARC Monogr Eval*
445   *Carcinog Risks Hum* 2004, **83**:1-1438.
446   9.   Gormley M, Dudding T, Sanderson E, Martin RM, Thomas S, Tyrrell J, Ness AR,
447   Brennan P, Munafo M, Pring M *et al*: **A multivariable Mendelian randomization**
448   **analysis investigating smoking and alcohol consumption in oral and oropharyngeal**
449   **cancer**. *Nat Commun* 2020, **11**(1):6071.
450   10.   Sabatini ME, Chiocca S: **Human papillomavirus as a driver of head and neck cancers**.
451   *Br J Cancer* 2020, **122**(3):306-314.
452   11.   Gillison ML, Koch WM, Capone RB, Spafford M, Westra WH, Wu L, Zahurak ML,
453   Daniel RW, Viglione M, Symer DE *et al*: **Evidence for a causal association between**
454   **human papillomavirus and a subset of head and neck cancers**. *J Natl Cancer Inst*
455   2000, **92**(9):709-720.

456   12.   de Martel C, Georges D, Bray F, Ferlay J, Clifford GM: **Global burden of cancer**
457   **attributable to infections in 2018: a worldwide incidence analysis.** *Lancet Glob*
458   *Health* 2020, **8**(2):e180-e190.
459   13.   Research WCRFAIfC: **Diet, nutrition, physical activity and cancers of the mouth,**
460   **pharynx and larynx.** In: *Continuous Update Project Expert Report*. 2018.
461   14.   Gaudet MM, Kitahara CM, Newton CC, Bernstein L, Reynolds P, Weiderpass E,
462   Kreimer AR, Yang G, Adami HO, Alavanja MC *et al*: **Anthropometry and head and**
463   **neck cancer:a pooled analysis of cohort data.** *Int J Epidemiol* 2015, **44**(2):673-681.
464   15.   Jo YH, Talmage DA, Role LW: **Nicotinic receptor-mediated effects on appetite and**
465   **food intake.** *J Neurobiol* 2002, **53**(4):618-632.
466   16.   Watts EL, Moore SC, Gunter MJ, Chatterjee N: **Adiposity and cancer: meta-analysis,**
467   **mechanisms, and future perspectives.** *medRxiv*
468   2024:https://doi.org/10.1101/2024.1102.1116.24302944.
469   17.   Taylor AE, Richmond RC, Palviainen T, Loukola A, Wootton RE, Kaprio J, Relton CL,
470   Davey Smith G, Munafo MR: **The effect of body mass index on smoking behaviour**
471   **and nicotine metabolism: a Mendelian randomization study.** *Hum Mol Genet* 2019,
472   **28**(8):1322-1330.
473   18.   Carreras-Torres R, Johansson M, Haycock PC, Relton CL, Davey Smith G, Brennan P,
474   Martin RM: **Role of obesity in smoking behaviour: Mendelian randomisation study**
475   **in UK Biobank.** *BMJ* 2018, **361**:k1767.
476   19.   Asvold BO, Bjorngaard JH, Carslake D, Gabrielsen ME, Skorpen F, Smith GD,
477   Romundstad PR: **Causal associations of tobacco smoking with cardiovascular risk**
478   **factors: a Mendelian randomization analysis of the HUNT Study in Norway.** *Int J*
479   *Epidemiol* 2014, **43**(5):1458-1470.
480   20.   Freathy RM, Kazeem GR, Morris RW, Johnson PC, Paternoster L, Ebrahim S, Hattersley
481   AT, Hill A, Hingorani AD, Holst C *et al*: **Genetic variation at CHRNA5-CHRNA3-CHRNB4**
482   **interacts with smoking status to influence body mass index.** *Int J Epidemiol* 2011,
483   **40**(6):1617-1628.
484   21.   Taylor AE, Morris RW, Fluharty ME, Bjorngaard JH, Asvold BO, Gabrielsen ME,
485   Campbell A, Marioni R, Kumari M, Hallfors J *et al*: **Stratification by smoking status**
486   **reveals an association of CHRNA5-A3-B4 genotype with body mass index in never**
487   **smokers.** *PLoS Genet* 2014, **10**(12):e1004799.
488   22.   Morris RW, Taylor AE, Fluharty ME, Bjorngaard JH, Asvold BO, Elvestad Gabrielsen M,
489   Campbell A, Marioni R, Kumari M, Korhonen T *et al*: **Heavier smoking may lead to a**
490   **relative increase in waist circumference: evidence for a causal relationship from a**
491   **Mendelian randomisation meta-analysis. The CARTA consortium.** *BMJ Open* 2015,
492   **5**(8):e008808.
493   23.   Yang S, Lynch J, Schulenberg J, Diez Roux AV, Raghunathan T: **Emergence of**
494   **socioeconomic inequalities in smoking and overweight and obesity in early**
495   **adulthood: the national longitudinal study of adolescent health.** *Am J Public Health*
496   2008, **98**(3):468-477.
497   24.   Marmot M, Bell R: **Social determinants and non-communicable diseases: time for**
498   **integrated action.** *BMJ* 2019, **364**:l251.
499   25.   Marmot M: **The health gap: Doctors and the social determinants of health.** *Scand J*
500   *Public Health* 2017, **45**(7):686-693.
501   26.   Filippidis FT, Agaku IT, Girvalaki C, Jimenez-Ruiz C, Ward B, Gratziou C, Vardavas CI,
502   Tobacco Control Committee of the European Respiratory S: **Relationship of**

secondhand smoke exposure with sociodemographic factors and smoke-free
legislation in the European Union. *Eur J Public Health* 2016, **26**(2):344-349.
27.   Larsson SC, Burgess S: **Causal role of high body mass index in multiple chronic**
diseases: a systematic review and meta-analysis of Mendelian randomization
studies. *BMC Med* 2021, **19**(1):320.
28.   Gormley M, Dudding T, Thomas SJ, Tyrrell J, Ness AR, Pring M, Legge D, Davey Smith
G, Richmond RC, Vincent EE *et al*: **Evaluating the effect of metabolic traits on oral**
and oropharyngeal cancer risk using Mendelian randomization. *Elife* 2023, **12**.
29.   Vithayathil M, Carter P, Kar S, Mason AM, Burgess S, Larsson SC: **Body size and**
composition and risk of site-specific cancers in the UK Biobank and large
international consortia: A mendelian randomisation study. *PLoS Med* 2021,
18(7):e1003706.
30.   Lesseur C, Diergaarde B, Olshan AF, Wunsch-Filho V, Ness AR, Liu G, Lacko M, Eluf-
Neto J, Franceschi S, Lagiou P *et al*: **Genome-wide association analyses identify new**
susceptibility loci for oral cavity and pharyngeal cancer. *Nat Genet* 2016,
48(12):1544-1550.
31.   Davies NM, Holmes MV, Davey Smith G: **Reading Mendelian randomisation studies:**
a guide, glossary, and checklist for clinicians. *BMJ* 2018, **362**:k601.
32.   Skrivankova VW, Richmond RC, Woolf BAR, Yarmolinsky J, Davies NM, Swanson SA,
VanderWeele TJ, Higgins JPT, Timpson NJ, Dimou N *et al*: **Strengthening the**
Reporting of Observational Studies in Epidemiology Using Mendelian
Randomization: The STROBE-MR Statement. *JAMA* 2021, **326**(16):1614-1621.
33.   Ebrahimi E, Sangphukieo A, Park H, Gaborieau V, Ferreiro-Iglesias A, Diergaarde B,
Ahrens W, Alemany L, Arantes L, Betka J *et al*: **Cross-ancestral GWAS identifies 29**
novel variants across Head and Neck Cancer subsites. *medRxiv*
2024:https://doi.org/10.1101/2024.1111.1118.24317473.
34.   World Health O, World Health O: **ICD-10 : international statistical classification of**
diseases and related health problems, Tenth revision, Fifth edition. edn. Geneva:
World Health Organization; 2016.
35.   Shungin D, Winkler TW, Croteau-Chonka DC, Ferreira T, Locke AE, Magi R, Strawbridge
RJ, Pers TH, Fischer K, Justice AE *et al*: **New genetic loci link adipose and insulin**
biology to body fat distribution. *Nature* 2015, **518**(7538):187-196.
36.   Pulit SL, Stoneman C, Morris AP, Wood AR, Glastonbury CA, Tyrrell J, Yengo L, Ferreira
T, Marouli E, Ji Y *et al*: **Meta-analysis of genome-wide association studies for body**
fat distribution in 694 649 individuals of European ancestry. *Hum Mol Genet* 2019,
28(1):166-174.
37.   Loh PR, Tucker G, Bulik-Sullivan BK, Vilhjalmsson BJ, Finucane HK, Salem RM,
Chasman DI, Ridker PM, Neale BM, Berger B *et al*: **Efficient Bayesian mixed-model**
analysis increases association power in large cohorts. *Nat Genet* 2015, **47**(3):284-
290.
38.   Willer CJ, Li Y, Abecasis GR: **METAL: fast and efficient meta-analysis of genomewide**
association scans. *Bioinformatics* 2010, **26**(17):2190-2191.
39.   Hemani G, Zheng J, Elsworth B, Wade KH, Haberland V, Baird D, Laurin C, Burgess S,
Bowden J, Langdon R *et al*: **The MR-Base platform supports systematic causal**
inference across the human phenome. *Elife* 2018, **7**.

40. Palmer TM, Lawlor DA, Harbord RM, Sheehan NA, Tobias JH, Timpson NJ, Davey
Smith G, Sterne JA: **Using multiple genetic variants as instrumental variables for**
modifiable risk factors. *Stat Methods Med Res* 2012, **21**(3):223-242.
41. Burgess S, Thompson SG: **Bias in causal estimates from Mendelian randomization**
studies with weak instruments. *Stat Med* 2011, **30**(11):1312-1323.
42. Burgess S, Butterworth A, Thompson SG: **Mendelian randomization analysis with**
multiple genetic variants using summarized data. *Genet Epidemiol* 2013, **37**(7):658-
665.
43. Bowden J, Davey Smith G, Burgess S: **Mendelian randomization with invalid**
instruments: effect estimation and bias detection through Egger regression. *Int J*
Epidemiol 2015, **44**(2):512-525.
44. Bowden J, Davey Smith G, Haycock PC, Burgess S: **Consistent Estimation in**
Mendelian Randomization with Some Invalid Instruments Using a Weighted
Median Estimator. *Genet Epidemiol* 2016, **40**(4):304-314.
45. Hartwig FP, Davey Smith G, Bowden J: **Robust inference in summary data Mendelian**
randomization via the zero modal pleiotropy assumption. *Int J Epidemiol* 2017,
46(6):1985-1998.
46. Bowden J, Del Greco MF, Minelli C, Zhao Q, Lawlor DA, Sheehan NA, Thompson J,
Davey Smith G: **Improving the accuracy of two-sample summary-data Mendelian**
randomization: moving beyond the NOME assumption. *Int J Epidemiol* 2019,
48(3):728-742.
47. Verbanck M, Chen CY, Neale B, Do R: **Detection of widespread horizontal pleiotropy**
in causal relationships inferred from Mendelian randomization between complex
traits and diseases. *Nat Genet* 2018, **50**(5):693-698.
48. Sanderson E, Davey Smith G, Windmeijer F, Bowden J: **An examination of**
multivariable Mendelian randomization in the single-sample and two-sample
summary data settings. *Int J Epidemiol* 2019, **48**(3):713-727.
49. Saunders GRB, Wang X, Chen F, Jang S-K, Liu M, Wang C, Gao S, Jiang Y,
Khunsriraksakul C, Otto JM *et al*: **Genetic diversity fuels gene discovery for tobacco**
and alcohol use. *Nature* 2022, **612**(7941):720-724.
50. Wootton RE, Richmond RC, Stuijfzand BG, Lawn RB, Sallis HM, Taylor GMJ, Hemani G,
Jones HJ, Zammit S, Davey Smith G *et al*: **Evidence for causal effects of lifetime**
smoking on risk for depression and schizophrenia: a Mendelian randomisation
study. *Psychol Med* 2020, **50**(14):2435-2443.
51. Darrous L, Mounier N, Kutalik Z: **Simultaneous estimation of bi-directional causal**
effects and heritable confounding from GWAS summary statistics. *Nat Commun*
2021, **12**(1):7274.
52. Morrison J, Knoblauch N, Marcus JH, Stephens M, He X: **Mendelian randomization**
accounting for correlated and uncorrelated pleiotropic effects using genome-wide
summary statistics. *Nat Genet* 2020, **52**(7):740-747.
53. Hemani G, Tilling K, Davey Smith G: **Orienting the causal relationship between**
imprecisely measured traits using GWAS summary data. *PLoS Genet* 2017,
13(11):e1007081.
54. Sanderson E, Rosoff D, Palmer T, Tilling K, Smith GD, Hemani G: **Bias from heritable**
confounding in Mendelian randomization studies. *medRxiv*
2024:2024.2009.2005.24312293.

55.   Foley CN, Mason AM, Kirk PDW, Burgess S: **MR-Clust: clustering of genetic variants**
595   **in Mendelian randomization with similar causal estimates**. *Bioinformatics* 2021,
596   **37**(4):531-541.
597   56.   Ried JS, Jeff MJ, Chu AY, Bragg-Gresham JL, van Dongen J, Huffman JE, Ahluwalia TS,
598   Cadby G, Eklund N, Eriksson J *et al*: **A principal component meta-analysis on**
599   **multiple anthropometric traits identifies novel loci for body shape**. *Nat Commun*
600   2016, **7**:13357.
601   57.   Richardson TG, Sanderson E, Elsworth B, Tilling K, Davey Smith G: **Use of genetic**
602   **variation to separate the effects of early and later life adiposity on disease risk:**
603   **mendelian randomisation study**. *BMJ* 2020, **369**:m1203.
604   58.   Martin S, Cule M, Basty N, Tyrrell J, Beaumont RN, Wood AR, Frayling TM, Sorokin E,
605   Whitcher B, Liu Y *et al*: **Genetic Evidence for Different Adiposity Phenotypes and**
606   **Their Opposing Influences on Ectopic Fat and Risk of Cardiometabolic Disease**.
607   *Diabetes* 2021, **70**(8):1843-1856.
608   59.   Leyden GM, Shapland CY, Davey Smith G, Sanderson E, Greenwood MP, Murphy D,
609   Richardson TG: **Harnessing tissue-specific genetic variation to dissect putative**
610   **causal pathways between body mass index and cardiometabolic phenotypes**. *Am J*
611   *Hum Genet* 2022, **109**(2):240-252.
612   60.   Thorgeirsson TE, Gudbjartsson DF, Sulem P, Besenbacher S, Styrkarsdottir U,
613   Thorleifsson G, Walters GB, Consortium TAG, Oxford GSKC, consortium E *et al*: **A**
614   **common biological basis of obesity and nicotine addiction**. *Transl Psychiatry* 2013,
615   **3**(10):e308.
616   61.   Wills AG, Hopfer C: **Phenotypic and genetic relationship between BMI and cigarette**
617   **smoking in a sample of UK adults**. *Addict Behav* 2019, **89**:98-103.
618   62.   Gage SH, Sallis HM, Lassi G, Wootton RE, Mokrysz C, Davey Smith G, Munafo MR:
619   **Does smoking cause lower educational attainment and general cognitive ability?**
620   **Triangulation of causal evidence using multiple study designs**. *Psychol Med* 2022,
621   **52**(8):1578-1586.
622   63.   Schellhas L, Haan E, Easey KE, Wootton RE, Sallis HM, Sharp GC, Munafo MR, Zuccolo
623   L: **Maternal and child genetic liability for smoking and caffeine consumption and**
624   **child mental health: an intergenerational genetic risk score analysis in the ALSPAC**
625   **cohort**. *Addiction* 2021, **116**(11):3153-3166.
626   64.   Khouja JN, Wootton RE, Taylor AE, Davey Smith G, Munafo MR: **Association of**
627   **genetic liability to smoking initiation with e-cigarette use in young adults: A cohort**
628   **study**. *PLoS Med* 2021, **18**(3):e1003555.
629   65.   Reed ZE, Wootton RE, Khouja JN, Richardson TG, Sanderson E, Davey Smith G,
630   Munafo MR: **Exploring pleiotropy in Mendelian randomisation analyses: What are**
631   **genetic variants associated with 'cigarette smoking initiation' really capturing?**
632   *Genet Epidemiol* 2024.
633   66.   Gui L, He X, Tang L, Yao J, Pi J: **Obesity and head and neck cancer risk: a mendelian**
634   **randomization study**. *BMC Med Genomics* 2023, **16**(1):200.
635   67.   Thomas SJ, Penfold CM, Waylen A, Ness AR: **The changing aetiology of head and**
636   **neck squamous cell cancer: A tale of three cancers?** *Clin Otolaryngol* 2018,
637   **43**(4):999-1003.
638   68.   Lauby-Secretan B, Scoccianti C, Loomis D, Grosse Y, Bianchini F, Straif K, International
639   Agency for Research on Cancer Handbook Working G: **Body Fatness and Cancer--**
640   **Viewpoint of the IARC Working Group**. *N Engl J Med* 2016, **375**(8):794-798.

641 69.   Mariosa D, Carreras-Torres R, Martin RM, Johansson M, Brennan P: **Commentary:**
642   **What can Mendelian randomization tell us about causes of cancer?** *Int J Epidemiol*
643   2019, **48**(3):816-821.

Availability of data and materials

All the GWAS datasets used in our study are publicly available. The GWAS summary statistics

for waist circumference by Shungin et al. are available via the IEU OpenGWAS platform (id:

ieu-a-61). The GWAS summary statistics for BMI and WHR by Pulit et al.[36] can be

downloaded from https://zenodo.org/records/1251813. The data sources for the other

adiposity-related measures have been specified in Table 1. The smoking behaviour traits

GWAS data were downloaded from

https://data.bris.ac.uk/data/dataset/10i96zb8gm0j81yz0q6ztei23d (for CSI) and

https://doi.org/10.13020/przg-dp88 (for smoking initiation). The outcome datasets used in

our analyses have been uploaded to the IEU OpenGWAS project platform for reproducibility.

However, because the data was originally in build GRCh38, some multiallelic SNPs that could

not be aligned with GRCh37 Human Genome reference sequence were dropped when lifting

the data to build HG19/GRCh37 (which was required at the time of upload: April 2024). The

following IEU OpenGWAS id's were assigned to the European HEADSpAcE HNC GWAS

datasets including/excluding UK Biobank: ieu-b-5129/ieu-b-5123 for overall HNC, ieu-b-

5132/ieu-b-5126 for oral cavity cancer, ieu-b-5130/ieu-b-5124 for hypopharynx cancer, ieu-

b-5134/ieu-b-5128 for HPV positive oropharynx cancer, ieu-b-5133/ieu-b-5127 for HPV

negative oropharynx cancer, and ieu-b-5131/ieu-b-5125 for larynx cancer. The R code used

to run the MR analyses is available at http://github.com/fernandam93/adiposity_HNC_MR.

Acknowledgements

We thank Richard Wilkinson for proofreading several versions of the manuscript. We would also like to thank Weili (Jason) Qiu, the IEU Data Manager, for his help debugging code and uploading the HNC GWAS summary statistics to the IEU OpenGWAS platform.

Funding

FMB was supported by a Wellcome Trust PhD studentship in Molecular, Genetic and Lifecourse Epidemiology (224982/Z/22/Z). RCR was supported by a Cancer Research UK grant (C18281/A29019). MCB is supported by a University of Bristol Vice Chancellor's Fellowship, the British Heart Foundation (AA/18/1/34219) and the UK Medical Research Council (MC_UU_00032/05). GDS works within the MRC Integrative Epidemiology Unit at the University of Bristol, which is supported by the Medical Research Council (MC_UU_00011/1). CLR was supported by the Medical Research Council (MC_UU_00011/5) and by a Cancer Research UK (C18281/A29019) programme grant (the Integrative Cancer Epidemiology Programme). SV was funded by an EU Horizon 2020 grant (agreement number 825771). JK works in a unit that receives support from the University of Bristol and the UK Medical Research Council (grant number: MC_UU_00032/7).

Competing interests

TGR works as a full-time employee for GlaxoSmithKline outside of this research. All other authors declare no conflicts of interest.

IARC disclaimer

Figures

Figure 1. Forest plot for the genetically predicted effects of body mass index on the risk of head and neck cancer and its subsites.

Figure 2. Forest plot for the genetically predicted effects of waist-to-hip ratio on the risk of head and neck cancer and its subsites.

Figure 3. Forest plot for the genetically predicted effects of waist circumference on the risk of head and neck cancer and its subsites.

Figure 4. Forest plot for the genetically predicted effects of BMI on the risk of HNC and its subsites, before (univariable-black) and after (multivariable-blue) accounting for comprehensive smoking index (CSI).

Figure 5. Forest plot for the genetically predicted effects of BMI on the risk of HNC and its subsites, before (univariable-black) and after (multivariable-blue) accounting for smoking initiation (SI).