Molnupiravir clinical trial simulation suggests that polymerase chain reaction underestimates antiviral potency against SARS-CoV-2

Authors

Shadisadat Esmaeili,[1]* Katherine Owens,[1] Jessica Wagoner,[2] Stephen J. Polyak,[2] Shengyuan Zhang[3], Joseph F. Standing[3,4], David M. Lowe[5,6], Joshua T. Schiffer[1,7]

Affiliations

[1] Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Center; Seattle, WA, USA.

[2] Department of Laboratory Medicine & Pathology, University of Washington; Seattle, WA, USA.

[3] Infection, Immunity and Inflammation, Great Ormond Street Institute of Child Health, University College London, London, UK

[4] Great Ormond Street Hospital for Children NHS Trust, London, UK

[5] Department of Clinical Immunology, Royal Free London NHS Foundation Trust, London, UK.
[6] Institute of Immunity and Transplantation, University College London, London, UK

[7] Department of Medicine, University of Washington; Seattle, WA, USA.

*Corresponding Author: sesmaeil@fredhutch.org

Abstract

Molnupiravir is an antiviral medicine that induces lethal copying errors into SARS-CoV-2 RNA. Molnupiravir reduced hospitalization in one pivotal trial by 50% and had variable effects on reducing viral RNA levels in three separate trials. We used mathematical models to simulate these trials and closely recapitulated their virologic outcomes. Model simulations suggest lower antiviral potency against pre-omicron SARS-CoV-2 variants than against omicron. We estimate that in vitro assays underestimate in vivo potency 7-8 fold against omicron variants. Our model suggests that because polymerase chain reaction detects molnupiravir mutated variants, the true reduction in genetically intact viral RNA is underestimated by ~0.5 log in the two trials, which included participants with omicron. Our results reinforce past work suggesting that in vitro assays are unreliable for estimating in vivo antiviral drug potency and suggest that virologic endpoints for respiratory virus clinical trials should be catered to the drug mechanism of action.

Introduction

Molnupiravir is an antiviral drug that induces errors in the SARS-CoV-2 genome, which typically renders the virus unable to replicate further.[1] In the randomized double-blinded MOVe-OUT trial, which enrolled unvaccinated individuals when delta, mu, and gamma variants of concern (VOC) were circulating, molnupiravir reduced hospitalization by 50% and viral load after treatment by 0.3 log relative to placebo.[2] In the platform adaptive PANORAMIC trial, which enrolled vaccinated individuals when omicron VOCs were circulating, molnupiravir did not lower hospitalization rates, which were only 1% in both arms, but lowered viral load after treatment by 0.94 log relative to usual care.[3] In the platform adaptive PLATCOV trial, which enrolled low-risk individuals when omicron VOCs were circulating, molnupiravir lowered viral load after treatment by 1.09 log relative to ususal care.[4] Taken together, these trials demonstrate that molnupiravir has both clinical and virologic efficacy which varied across trials and viral variants.

Overall, use of molnupiravir has been lower than that of nirmatrelvir / ritonavir based on lower reduction in hospitalization in MOVe-OUT relative to the EPIC-HR trial for nirmatrelvir / ritonavir.[5] A concern has also been raised that molnupiravir's mechanism of action could generate novel mutants that persist after cessation of treatment,[6] and then spread in the population.[7] Nevertheless, PANORAMIC and PLATCOV results suggest high potency, and molnupiravir is still considered in individuals in whom nirmatrelvir / ritonavir is contraindicated, and in combination with other drugs in immunocompromised hosts.[8, 9] There is currently no explanation for the disparate antiviral effects in MOVe-OUT versus PANORAMIC and PLATCOV. Moreover, the fact that polymerase chain reaction (PCR) detects drug-altered viral RNA molecules[6] has not always been considered in the analysis of trial outcomes.

We previously used clinical trial simulation to reproduce results from nirmatrelvir / ritonavir trials for SARS-CoV-2.[10, 11] We first validated a viral immune dynamic model (VID) against a very large prospective cohort of infections that included multiple VOCs.[11] We used diverse virologic output from this model to create simulated cohorts for the control arms of trials. We then layered pharmacokinetic (PK) and pharmacodynamic (PD) models for nirmatrelvir / ritonavir on the VID models to simulate treatment arms.[10] This approach recapitulated mean viral load reduction in the EPIC-HR and PLATCOV trials, as well as individual viral load trajectories in PLATCOV. The validated model was then used to explain the high frequency of virologic and concurrent symptomatic rebound with use in the community,[12] despite very low levels of virologic and symptomatic rebound in the EPIC-HR trial.[13, 14] Model output suggests that extending therapy from 5 to 10 days would nearly eliminate rebound,[10] a result confirmed with modeling of separate data by another group.[15]

Here we expand this approach to develop a new joint VID-PK-PD model to account for the unique mechanism of action of molnupiravir. We simulate by fitting the model to results from the MOVe-OUT, PLATCOV and PANORAMIC trials. Our results suggest that quantitative viral PCR likely underestimates the reduction in non-mutated viral RNA and therefore the true potency of molnupiravir during omicron infections.

88
89
90 **Results**
91
Viral immune dynamic, pharmacokinetic and pharmacodynamic clinical trial simulation
models.
We previously described our viral immune dynamic (VID) model that was fit to diverse viral
loads from 1510 SARS-CoV-2 infected individuals in the National Basketball Association
cohort.[11, 16] The model assumes a finite number of susceptible cells. An eclipse phase delays viral
production by infected cells. In keeping with an early innate immune response, susceptible cells
become refractory to infection in the presence of infected cells but also revert to a susceptible
state at a constant rate. Infected cells are cleared by cytolysis and delayed acquired immunity,
which is activated in a time-dependent fashion (**Fig 1a**). We used a mixed-effect population
approach implemented in Monolix to estimate model parameters.[17]
102

103
104 **Fig 1. Schematic of the (a) viral dynamic model and (b) molnupiravir PK-PD model.**
105
To reproduce levels of molnupiravir, we used a three-compartment pharmacokinetic (PK) model
(**Fig. 1b**). Using Monolix and the mixed-effect population approach, we estimated parameter
values by fitting the model to the average plasma concentration of healthy subjects.[18] The model
closely recapitulated observed drug levels following multiple doses of 50, 100, 200, 300, 400,
600, and 800 mg given twice daily for five days (**Fig S1, Table S1**). The estimated value for the
transition rate from plasma to peripheral compartment (was dose-dependent in the form of
112   . All other PK parameters were dose independent. For the pharmacodynamic
(PD) model, we assumed drug efficacy follows a Hill equation with respect to concentration. We
parameterized the model using in vitro efficacy data collected at different concentrations (details
in **Materials and Methods**, **Fig S2, Table S2**).[19]

We combined the VID and PKPD models by using treatment efficacy to convert non-mutated
virus to mutated virus, both of which are assumed to be detected with polymerase chain reaction
(PCR) assays, given the low probability of drug-induced mutations in the PCR primer region.
This assumption is based on the observed drug-induced mutation of approximately 1 in 2000
sites.[20] Given the average length of most PCR primers of ~25 base pairs, the chance of the
primer remaining unmutated after treatment is $(1999/2000)^{25}$, or 98.76%. A limitation of viral
load data from the included clinical trials is that it lacks early pre-symptomatic endpoints to
estimate the viral expansion slope. To further train the model, we included 1023 Omicron-
infected participants from the NBA cohort to inform rates of viral upslope in the trials.[11] We first
fit the combined model to individual viral load data from 149 low-risk, symptomatic vaccinated
participants infected with omicron VOCs in the PLATCOV trial (65 treated and 84 controls) and
from 80 high-risk, symptomatic vaccinated participants infected with omicron VOCs in the
PANORAMIC trial (38 treated and 42 controls) (**Fig 2a, 2b, Fig S3, S4, S5, S6**). We next fit the
combined model to trial endpoint data (mean viral load drop from baseline) reported in three
randomized, controlled trials: PLATCOV (**Fig 3a, 4a-d**),[4] PANORAMIC (**Fig 3b, 4e-h**),[3] and
the MOVe-OUT trial with 1093 high-risk unvaccinated symptomatic individuals infected with
pre-omicron VOCs (549 treated + 544 placebo, **Fig 3c, 4i-k**).[2] All model fitting was using
Monolix with non-linear mixed effects approaches described in the **Materials and Methods**.

Fig 2. Mathematical model fits of SARS-CoV-2 viral load over time to a subset of study participants
in PLATCOV and PANORAMIC receiving no treatment (control) or molnupiravir. (a) Model fits
to 9 control and 9 treatment participants in PLATCOV**. (b)** Model fits to 9 control and 9 treatment
participants in PANORAMIC. **(c)** Individual estimates for potency adjustment factor (ratio of *in vivo : in*
vitro EC50) in the two trials.

Model fitting to individual viral load trajectories in PLATCOV and PANORAMIC.

For each participant, we defined the in vivo EC_{50} as the plasma drug concentration required to
inhibit viral replication by 50% and the potency adjustment factor (paf) as the ratio between the
in vivo and in vitro EC_{50}.[21, 22] To estimate the paf, we fit the combined VID-PKPD model to
individual viral load data of both arms of PLATCOV and PANORAMIC trials and Omicron
infected individuals in the NBA cohort using the population mixed effect approach in Monolix.
We achieved good model fit to individual viral load trajectories in the control and treatment arms
of PLATCOV **(Fig 2a, Fig S3, S4)** and PANORAMIC **(Fig 2b, S5, S6)**. The model projected
higher levels of total detected SARS-CoV-2 RNA in most participants relative to non-mutated
viral RNA **(Fig 2a,b)**. We estimated a range of individual paf values with similar mean and
median values estimated for both trials **(Fig 2c)**. These values suggest that in vivo potency of
molunpiravair is on average 7-8 fold higher than in in vitro estimates. Each participant had an
estimated paf less than one indicating that enhanced potency is necessary to include to optimize
fit to the data.
154

Figure 3. Mean viral load reduction in (a) PLATCOV, (b) PANORAMIC, and (c) MOVe-OUT
which are targets for model fitting.

Model fit to trial virologic endpoint data from PLATCOV, PANORAMIC, and MOVe-
OUT.
As a second approach, we assessed whether a virtual cohort strategy where control participants
are modeled from pre-existing cohorts can predict virologic trial endpoints. This approach is
necessary for situations where individual viral load data are not available as with MOVe-OUT
and is important as it demonstrates that the model can reproduce the primary virologic endpoint
of the study. We simulated virtual cohorts using the combined VID-PKPD model and fit results
to viral load decay from baseline in the three trials. For each trial arm, we randomly selected 400
individuals from the NBA cohort with the closest matching viral variant, symptom, and vaccine
status and used their estimated individual viral dynamic parameters in simulations. To address

variability in timing of baseline viral load measurement relative to infection, we randomly
assigned all individuals an incubation period selected from a variant-specific gamma distribution
found in the literature.[23, 24] Treatment start day was randomly selected from a distribution based
on observed enrollment windows in the three trials. Due to the lack of individual PK data, the
same estimated population PK parameters were used for all simulated treated individuals
(**Table S1**). PD parameters were randomly selected from a log-normal distribution with
estimated mean and standard error (**Table S2**).

Our model closely reproduced kinetics of viral decay in PLATCOV in control (**Fig 3a, 4a**) and
treatment arms (**Fig 3a, 4b**) and estimated a paf=0.13 (**Fig 4c**) similar to our mean estimate using
individual fits (**Fig 2c**). The model also predicted individual-level variability in virologic
responses observed in PLATCOV, including instances of increased viral load following therapy
(**Fig 4d**). We compared simulated and actual distributions of viral load change among trial
participants in the control and treatment arms. On most post-treatment days, simulated and actual
distributions were not statistically dissimilar. Wider distributions of observed versus simulated
viral load change were noted on post-randomization days 1 and 2 for control and treatment (**Fig**
4d), likely due to noise in viral load data from oral swabs: differences of 1–2 logs were often
noted between replicates collected from PLATCOV participants at equivalent timepoints,
particularly on days 1 and 2.[10]
187

188
Figure 4. Model fit to virologic trial outcomes for (a-d) PLATCOV, (e-h) PANORAMIC, (i-k)
MOVe-OUT. Results include (**a, e, i**) control groups, (**b, f, j**) treatment groups, (**d,h**) comparing
individual variability of data vs simulation in control and treatment arms, and (**c, g, k**) estimate for
potency adjustment factor. To only capture the effect of treatment and address potential identifiability
issues, data from the first seven days after baseline were used to estimate the paf. Therefore, the crossed-
out data points were not included in the calculation of the R^2.
195
Similarly, our model closely reproduced kinetics of viral decay in PANORAMIC in control (**Fig**
3b, 4e) and treatment arms (**Fig 3b, 4f**) and estimated a paf=0.19 (**Fig 4g**) similar to our mean
estimate using individual fits (**Fig 2c**). The model also predicted individual-level variability in

virologic responses observed in PANORAMIC, including instances of increased viral load
following therapy **(Fig 4h)**. We compared simulated and actual distributions of viral load change
among trial participants in the control and treatment arms. On all post-treatment days other than
day 2 control, simulated and actual distributions were not statistically dissimilar **(Fig 4h).** This
likely indicates less noise in viral load data from nasopharyngeal swabs collected in
PANORAMIC relative to oral swabs in PLATCOV.
205
206 Finally, the model reproduced kinetics of viral decay in MOVe-OUT in control **(Fig 4i)** and
207 treatment arms **(Fig 4j)** but estimated a higher paf=2.64 **(Fig 4k).** The higher paf maps to the far
208 less substantial viral load reduction in MOVe-OUT relative to the other two trials which in turn
209 may be explained by less potency against pre-omicron variants which has been observed
210 experimentally.[25]
211
As a further validation step, we performed counterfactual simulations which assess viral kinetics
of control study participants assuming treatment and treatment participants assuming
placebo/usual care. Counterfactual control simulations slightly overestimated late viral loads for
PLATCOV **(Fig S7a)** and PANORAMIC **(Fig S8a)**. This may be because therapy suppresses
acquired immune responses, which is not captured in our model.[15] Counterfactual treatment
simulations fit the data well for PLATCOV **(Fig S7b)** and PANORAMIC **(Fig S8b)**. Simulations
occasionally predicted viral rebound following treatment **(Fig S7c,d and S8c,d).**
219
220

221
Figure 5. PCR underestimates the true reduction in non-mutated SARS-CoV-2 RNA in PLATCOV
and PANORAMIC but not MOVe-OUT. Simulated mean viral loads including non-mutated viral RNA
in (a) PLATCOV, (b) PANORAMIC and (c) MOVe-OUT. (d) Individual reduction at day 5 in the
simulated control group (blue), simulated total viral RNA (grey) and simulated non-mutated viral RNA
(pink) in the three trials showing no statistical difference between total and non-mutated viral RNA

despite a lower median. (e) Individual viral area under the curve from the start of the treatment through
day 5 in the simulated control group (blue), simulated total viral RNA (grey), and simulated non-mutated
viral RNA (pink) in the three trials showing a statistical difference between total and non-mutated viral
RNA. Boxplots include the interquartile range (IQR) with whiskers equaling 1.5 the IQR. (f) Table of
mean viral load reductions in the trials and simulations including the predicted mean difference in total
versus mutated viral RNA.
233
Estimates of reduction in fully mutated viruses versus non-mutated SARS-CoV-2 RNA.
We used our optimized model with solved paf to project the trajectory of non-mutated viral RNA
during treatment relative to values measured with PCR, which detects viral RNA with drug-
induced mutations.[6] In PLATCOV (**Fig 5a,d,f**) and PANORAMIC (**Fig 5b,d,f**), owing to higher
drug potency, total SARS-CoV-2 viral RNA on treatment exceeded non-mutated viral RNA by
~0.48 and 0.59 log respectively, suggesting that measured endpoints underestimate the drug's
true antiviral effect. However, these differences did not achieve statistical significance, perhaps
because estimated total and non-mutated viral RNA levels converge at drug trough. In MOVe-
OUT (**Fig 5c,d,f**), there was no significant difference between total SARS-CoV-2 viral RNA on
treatment and non-mutated viral RNA.
244

245
Figure 6. Relationship of drug pharmacokinetics and pharmacodynamics to in vivo potency and
viral load reduction. (a) Molnupiravir plasma concentration during five days of treatment with 800 mg
molnupirvir given twice daily. The dashed lines mark the EC_{50} with different paf values which differ by
trial. For Paf = 0.13, drug levels are almost entirely above EC_{50}. (b) Dynamic shifts in molnupiravir
efficacy for different paf values which differ by trial. Efficacy only drops minimally at trough levels when
paf is low (i.e. 0.14 and 0.13 in PLATCOV and PANORAMIC) but drops significantly at trough levels in
MOVe-OUT. (c) Drug potency of SARS-CoV-2 antivirals according to trial. The in-vivo efficacy of
molnupiravir in PLATCOV and PANORAMIC trials is close to the in vivo efficacy of
nirmatrelvir/ritonavir in the PLATCOV trial and higher than EPIC-HR. MOVe-OUT potency is
significantly lower due to a higher paf and higher in vivo EC50 value. (d) Simulated mean drops in total
viral RNA from baseline relative to counterfactual placebo/usual care arms on day 5 in the three
molnupiravir trials and two nirmatrelvir/ritonavir trials. (e) Simulated mean drops in non-mutated viral
RNA from baseline relative to counterfactual placebo on day 5 in the three molnupiravir trials and two
nirmatrelvir/ritonavir trials. In the molnupiravir trials, total viral RNA drops less than non-mutated viral

RNA due to PCR detection of drug-mutated viral RNA. Total possible reduction in non-mutated SARS-
CoV-2 RNA is less for MOVe-OUT than PLATCOV and PANORAMIC due to higher initial viral loads
and lower values of detection in the trials.
263
In all 3 trials, the models suggest that non-mutated viral loads during treatment may be lowest at
drug peak, reflecting the short plasma half-life of molnupiravir. Therefore, the cumulative
effective of drug is best estimated with viral area under the curve which accounts for highly
variable drug activity over time due to short drug half-life. By this estimate, the reduction in non-
mutated viral RNA far exceeds that of total measured viral RNA **(Fig 5e)**. In the case of MOVe-
OUT, this may explain why significant clinical benefit is associated with only a marginal
reduction in observed decline in total viral RNA. This also suggests that there might be utility to
measure viral load after drug peak and trough for agents with short half-life as viral loads may
differ by a full order of magnitude according to drug level.
273
Differences in trial participants and model parameters.
We also compared features of each trial as they related to model predictions by assessing the
viral dynamic range in each trial **(Fig S9a)**. Control participants in PLATCOV had lower mean
viral loads throughout the course of infection relative to PANORAMIC and MOVe-OUT **(Fig
S9b)**. Given PLATCOV and PANORAMIC enrolled participants with the omicron variant, we
surmise that these differences relate to demographic differences in study participant
demographics, slightly shorter estimated time to treatment in PANORAMIC versus PLATCOV
estimated by our model (t_0 in **Fig S10**), and/or characteristics of the PCR assay used in the
studies. The trials also employed different limits of detection which impacted observed
reductions in viral load **(Fig S9b).** Model parameters were largely equivalent between studies
and between treatment and control arms, reflecting the flexibility of the model **(Fig S10)**. The
parameter governing transition of susceptible cells to a refractory state was higher in PLATCOV
relative to PANORAMIC which likely was necessary to achieve lower viral loads overall and
may reflect the younger study participants **(Fig S10)**.
288
**Antiviral potency, viral load assessment and trial design all impact observed antiviral
reduction.**
We next combined PK and PD models to assess the average efficacy of the drug during days 0-5
in all three trials **(Fig 6a,b)** and noted an efficacy of 53% in MOVe-OUT **(Fig 6c)**. The efficacy
of molnupiravir in PLATCOV (94%) and PANORAMIC (95%) was similar to that of
nirmatrelvir in PLATCOV (94%) and EPIC-HR (82%) owing to a much lower paf for
molnupiravir relative to nirmatrelvir / ritonavir **(Fig 6c)**.
296
These potencies mapped to different reductions in viral load relative to placebo/usual care. Total
viral RNA reduction in PANORMIC and PLATCOV exceeded that in MOVe-OUT owing to
lower paf **(Fig 6d),** but also due to a larger viral dynamic range (defined as the distance from
baseline viral load to the lower threshold PCR (LOD) **(Fig S9b)**), which allows for a greater
observed reduction in viral load. PANORAMIC used LOD = 100 (imputed as 50), and
PLATCOV had LOD≅18 (copies/ml), versus LOD = 500 copies/ml for MOVe-OUT.
PANORAMIC also had much higher average starting viral loads (7.4 log10 (copies/ml)) versus
PLATCOV (5.8 log10 copies/ml) and MOVe-OUT (6.8 log10 copies/ml) **(Fig S9b)**.
Molnupiravir approached maximal possible total viral load viral RNA reduction in

PANORAMIC and PLATCOV, whereas protease inhibitors could still achieve greater viral load
reduction at lower paf **(Fig 6d)** as recently observed with ensitrelvir.[26]
308
The greater possible reduction in total viral load for protease inhibitors relative to molnupiravir
owes to different mechanisms of action**.** Model projected reductions in non-mutated viral RNA
reduction in PANORMIC and PLATCOV approximated viral load reductions observed in EPIC-
HR and PLATCOV on nirmatrelvir / ritonavir **(Fig 6e)**, suggesting that PCR detection of mutated
viruses underestimates true molnupiravir potency and that a more potent mutagenic agent could
accrue further virologic benefit.
315
316 **Optimization of molnupiravir therapy to avoid viral rebound.**
317 Instances of viral rebound were observed in PLATCOV and PANORAMIC and have been
318 observed following molnupiravir treatment.[4, 27] We analyzed higher doses and prolonged therapy
319 and noted that as with nirmatrelvir, prolonging therapy is a better method to prevent rebound
320 than increasing dose **(Fig S11).**
321
322
323

Discussion

We recapitulated the virologic results of three clinical trials of molnupiravir with our combined clinical trial simulation VID/PK/PD models. Model output highlights key differences in viral load reduction between the trials and identifies mechanistic differences to explain this. The MOVe-OUT trial was associated with significantly less reduction in viral load between treatment and control arms than the other two trials. Accordingly, the drug efficacy (0.53) was lower in this trial, and our estimate for paf was 2.64, signifying marginally lower potency in vivo than in vitro. This result is compatible with 0.4-fold lower median EC50 values for molnupiravir in vitro against omicron relative to prior variants. [25] The higher paf in MOVe-OUT permitted drug troughs below the EC50, limiting potency throughout the dosing interval. In PLATCOV and PANORAMIC, our model suggests drug levels remain above the in vivo EC50 throughout the dosing interval though potency does fluctuate according to drug level.

The estimated drug efficacy in PLATCOV and PANORAMIC was considerably higher (0.94 and 0.95, respectively), and the paf was estimated to be 0.14 and 0.13, indicating greater potency in vivo than in vitro. We have applied our clinical trial simulation technique to multiple drugs for SARS-CoV-2,[10] HSV-2,[21, 22] and HIV,[28] and this is the first time we have identified this trend. Allowing molnupiravir to be more potent in vivo in our modeling of PLATCOV and PANORAMIC was necessary to capture the much greater reduction in total viral RNA relative to off-treatment in these studies (1.09 log and 0.94 log respectively versus 0.3 log in MOVe-OUT).

A key outcome of our analysis is that SARS-CoV-2 PCR likely underestimates molnupiravir potency because it still detects drug-mutated viral RNA.[6] This appears to be most significant when antiviral potency is higher, as in PANORAMIC and PLATCOV, leading to a 0.48-0.59 log underestimation of reduction in non-mutated virus. Our results suggest that use of standard PCR for assessing SARS-CoV-2 levels may lead to underestimation of drug potency. Multi probe assays as have been used for the HIV reservoir may improve specificity for viruses that remain intact and replication competent.[29] Viral culture is potentially a useful metric but lacks sufficient sensitivity and precision and is too labor intensive to serve as a viable trial endpoint.[12]

Our results suggest that viral loads may vary according to drug level given molnupiravir's short plasma half-life. A sub-study within future trials comparing viral loads between drug trough and peak would be useful for the field. This could validate our model's prediction that even small reductions in viral RNA may be associated with substantial reductions in total viral area under the curve (a surrogate for infection surface area) particularly with a short half-life drug. Even minor reductions in viral load may be associated with substantial clinical benefit in this case. These fluctuations may be less evident if the intra-cellular half-life of the drug is longer or if PK measures in our model underestimate true drug levels in PANORMIC due to older age or impaired renal clearance. Model accuracy would be improved if viral load and PK data were available from the same patients in similar trials.

Another key practical outcome is that as for nirmatrelvir, extension of molnupiravir therapy to 10 days is likely to prevent rebound though our simulations do not suggest any benefit from increasing dose.[10] This suggests that prolonging therapy or using longer half-life agents is ideal for treating SARS-CoV-2.[30]

Each trial represented a unique set of issues for model fitting. In MOVe-OUT, because the treatment arm mean viral kinetics curve differed only slightly from the control arm, the model without drug provided reasonable fit to the treatment arm. Nevertheless, the paf was identifiable for this trial indicating that the model was able to detect and specify the very limited potency of the drug. The fact that the drug's potency and clinical efficacy appears to have increased with introduction of the omicron variant demonstrates a massive challenge for the therapeutics field: as with vaccines, trials performed when prior variants were circulating may prove less relevant as new variants continually emerge. A priority should be retesting existing agents against newly emerging variants in small nimble trials such as PLATCOV, with viral load endpoints.

For PLATCOV, the model for the treatment arm matched the trial data very precisely and identified the paf. The drug achieved nearly maximal observed viral load reduction in this trial. We identified a similar trend for PANORAMIC. It is notable that the model had the flexibility to account for different viral loads between these trials by predicting more rapid innate immune responses in PLATCOV.

We arrived at similar estimate for paf in PLATCOV and PANORAMIC, while using two separate methods: fitting to individual viral loads and fitting to mean viral load reduction trial endpoints. This suggests that our approach using in silico cohorts and fitting to population level outcomes produces reliable results.[10] In many cases, it remains challenging for academics to obtain individualized data from industry sponsored trials. Therefore, it is important that the endpoint fitting approach should be considered when this is the only data available.

Finally, our results highlight challenges in trial design associated with the selection of PCR assays and limits of detection. Different limits of detection were selected for each trial which in turn impacts the degree of viral load decrease that can be observed. Initial viral loads were notably higher in PANORAMIC and MOVe-OUT than PLATCOV. The equivalent viral loads between PANORAMIC and PLATCOV may reflect a more sensitive PCR in the PANORAMIC study as past immunity has consistently predicted lower viral loads and more rapid viral elimination for omicron variants.[11, 31] On the other hand, PANORMIC viral loads could have been higher than in PLATCOV despite both enrolling omicron infections due to an older and less healthy population, Ideally, equivalent internationally standardized PCR quantitation would be used across all trials.

Our study has a couple of limitations. The estimated paf is based on the in vitro assay data against delta variant in Calu-3 cells. In vitro EC50 is sensitive to assay conditions, including cell type, the variant of concern, the multiplicity of infection (MOI), and specific lab. In general, the inability of in vitro assays to match in vivo conditions makes it an unreliable proxy for the in vivo potency of the drug and explains the necessity of incorporating the paf parameter when simulating an antiviral clinical trial.

In our model, we assumed all mutated viruses are noninfectious. While most drug-induced random mutations reduce the fitness/viability of the virus, the molnupiravir mutation signature has been detected in circulating variants, especially in regions where the drug has been used most commonly.[7] Furthermore, in the PANORAMIC study, no statistical difference was observed

416 between the rate of culturable samples in the treatment and control arms which implies some
417 mutated viruses might be transmissible.[32]
418
PK parameters were estimated using the data from the plasma concentration of healthy
individuals with the age range of 19-60 (mean 39.6) years old. The clearance rate of renally
cleared drugs often increase with age.[33] This implies that the paf may be larger in an older
population, such as in PANORAMIC participants. Further, we used the plasma concentration in
the PD model to calculate the drug efficacy. However, using the drug's intracellular
concentration with a longer half-life, represented by the central compartment of the PK model,
would also likely lead to a larger estimated paf.
426
427 In summary, we further demonstrate the utility of clinical trial simulation using models that
428 capture drug PK and PD, as well as infection dynamics. In the case of molnupiravir, our results
429 suggest that final viral endpoints should be adjusted based on the drug's mechanism of action.
430
431

Materials and Methods

Study Design.

We developed a viral dynamics model recapitulating viral load data collected from symptomatic individuals in the NBA (National Basketball Association) cohort.[11] We used a two-compartment model to reproduce the PK data of molnupiravir.[18] For clinical trial simulation, we constructed a virtual cohort by randomly selecting 400 individuals from the NBA cohort, trying to match the trial populations regarding vaccine status and history of infection. Due to lack of individual PK data, we used estimated population PK parameters for all individuals in the virtual cohort. The PD parameters for each individual were randomly selected from a log-normal distribution with their estimated means and standard errors. We fit the combined viral dynamic and PKPD model to the average change in viral load from the baseline of the control and treatment arms of three previously published molnupiravir clinical trials.[2-4, 34] Comparing our model to the control arms validated our viral dynamic model and demonstrated how well our virtual cohorts represent the trial control arms. We used the average data from the treatment arms to estimate the potency adjustment factor (paf) by maximizing the R^2 of the fit.[10] We also fit to individual viral load trajectories in PLATCOV and PANORAMIC using the mixed-effect population approach implemented in Monolix[35-37] and obtained both individual paf values and a population distribution.

Viral load data

The NBA cohort dataset published by Hay et al consists of 2875 documented SARS-CoV-2 infections in 2678 people detected through frequent PCR testing regardless of symptoms.[16] We used the viral load data from 1510 infections in 1440 individuals that had at least 4 positive quantitative samples to estimate viral dynamic parameters. We used parameter sets estimated for the symptomatic subpopulation of these individuals to construct virtual cohorts.[11]

Clinical trial data

We used viral load data from three molnupiravir clinical trials. MOVe-OUT by Jayk Bernal et al. included 544 and 549 symptomatic high-risk individuals in the control and treatment arms, respectively.[2] We obtained the average change in viral load data of the control and treatment arms as shared in the supplementary material of their published manuscript in Table S6. Nasal viral load was measured using PCR assay on days 0, 3, 5, 10, and 14 after the treatment start day and adjusted by the baseline viral load. The lower limit of detection (LOD) of 500 copies/ml were used in this trial. The treatment was started within five days of symptom onset.

PLATCOV by Schilling et al. is an open-label, randomized, controlled adaptive trial with 85 and 58 symptomatic, young, healthy individuals in the control and molnupiravir treatment arms, respectively.[4] The oropharyngeal samples from each participant were collected daily on days 0 through 7 and on day 14 after the treatment start day, and viral load was measured using PCR assay. We used the individual viral load data published by the authors. From PLATCOV, we averaged over the two oral samples collected from each individual and calculated viral load drop from baseline or used the individual-level viral load data. We used the maximum LOD reported in the published data (~1.26 log).

PANORAMIC is a platform adaptive randomized, controlled trial with 42 and 38 symptomatic, vaccinated individuals with at least one risk factor in the control and treatment arms,

respectively.[3] Nasal viral load was measured using PCR and samples were collected on days 0
through 6 and on day 13. We used individual viral load data shared by the authors and adjusted
by baseline viral load to obtain mean viral load drop from the baseline. Mean days since
symptom onset at baseline (sd) were 2.4 (0.78) for the treatment arm and 2.5(1.12) for the
control arm. The lower limit of detection of 100 copies/ml (imputed as 50 copies/ml) was used.

In all three trials, the study participants were treated with 800mg molnupiravir twice per day, for
five days.

PK data

Mean plasma concentration data of molnupiravir were obtained by digitizing Figure 3 of the
phase I trial by Painter et al. using WebPlotDigitizer.[18] The data used in this paper belong to the
multiple-ascending part of the phase I trial where six participants were given 50, 100, 200, 300,
400, 600, and 800 mg of molnupiravir twice daily for 5.5 days, and their plasma concentrations
were measured after the first and last doses.

PD data

The data on drug efficacy was obtained from experiments performed at the University of
Washington. The efficacy of molnupiravir was measured against the delta variant of SARS-
CoV2 in Calu-3 cells (human lung epithelial). Briefly, Calu 3 cells were treated with varying
concentrations of molnupiravir prior to infection with SARS-CoV-2 (delta isolate) at a
multiplicity of infection of 0.01. Antiviral efficacy and cell viability (of non-infected cells treated
with drugs) were assessed as described.[9, 10] There were five replicates per condition, pooled from
2 independent technical experimental repeats (one experiment with triplicate conditions, one
experiment in duplicate conditions).

Viral dynamics model

We used our model of SARS-CoV-2 dynamics to model the viral load of symptomatic
individuals with SARS-CoV-2 infection.[11] Our model assumes that susceptible cells (S) are
infected at rate βVS by SARS-CoV-2 virions. The infected cells go through a non-productive
eclipse phase (I_E) before producing viruses and transition to becoming productively infected
(I_P) at rate κI_E. When encountering productively infected cells, the susceptible cells become
refractory to infection (R) at the rate $\phi I_P S$. Refractory cells revert to a susceptible state at rate
ρR. The productively infected cells produce virus at the rate πI_P and are cleared at rate δI
representing cytolysis and the innate immune response that lacks memory and is proportional to
the amount of ongoing infection. If the infection persists longer than time τ, then cytotoxic
acquired immunity is activated, which is represented in our model by the rate $m I_P$. Finally, free
virions are cleared at the rate γ. Of note, this model, previously proposed by Ke et al. was
selected against other models based on superior fit to data and parsimony.[38] The model written as
a set of differential equations has the form,

$$\frac{dS}{dt} = -\beta SV - \phi I_P S + \rho R \tag{1a}$$

$$\frac{dR}{dt} = \phi I_P S - \rho R \tag{1b}$$

$$\frac{dI_E}{dt} = \beta SV - \kappa I_E \tag{1c}$$

$\quad \frac{dI_P}{dt} = \kappa I_E - \delta I_P - m(t)I_P$ (1d)

$\quad \frac{dV}{dt} = \pi I_P - \gamma V$ (1e)

$\quad$ where $\begin{cases} m(t) = 0 & t < \tau \\ m(t) = m & t \geq \tau \end{cases}$ (1f)

To estimate parameter values, we fit the model to viral load data from the NBA cohort using a
mixed-effect population approach implemented in Monolix. Details on the model selection and
fitting process can be found in Owens et al.[11]

We start the simulations with 10^7 susceptible cells. The initial value of the refractory cells is
assumed to be zero since the interferon signaling is not active prior to infection. We further
assume there are no infected cells (eclipse or productive) at the beginning of the infection. We
fix the level of inoculum (V_0) at 97 copies/ml for each individual.

To resolve identifiability issues, we fixed two parameter values, setting the inverse of the eclipse
phase duration to $\kappa = 4$, and the rate of clearance of virions to $\gamma = 15$.[11]

PK model

We used a two-compartmental PK model which includes the amount of drug in the GI tract
(A_{GI}), the plasma compartment (A_p), and the respiratory tract (A_L). The drug is administered
orally, passes through the GI tract and gets absorbed into the blood at the rate κ_a. The drug then
transfers from the blood into the peripheral compartment (or the lung) at the rate κ_{PL}. The
metabolized drug transfers back into the plasma at the rate κ_{LP} from where it clears from the
body at the rate κ_{CL}. The model in the form of ordinary differential equations is written as,

$\quad \frac{dA_{GI}}{dt} = -\kappa_a A_{GI}$ (2a)

$\quad \frac{dA_P}{dt} = \kappa_a A_{GI} + \kappa_{LP}A_L - (\kappa_{CL} + \kappa_{PL})A_P$ (2b)

$\quad \frac{dA_L}{dt} = \kappa_{PL}A_P - \kappa_{LP}A_L$ (2c)

We used Monolix and a mixed-effect population approach to estimate the parameters and their
standard deviations. With the initial condition of ($A_{GI} = \text{Dose}, A_p = 0, A_L = 0$); we fit $C_P = \frac{A_P}{Vol}$
to the plasma concentration data where Vol is the estimated plasma volume. Details on
parameter values and the error model provided in **Table S1**.

PD model

For the pharmacodynamics model we used Hill equation, $\epsilon(t) = \frac{E_{max}C(t)^n}{C(t)^n + EC_{50}^n}$, where $C(t)$ is the
drug concentration in plasma, E_{max} is the maximum efficacy, n is the Hill coefficient, and EC_{50}
is the drug concentration in plasma required for 50% efficacy. We used least-squared fitting to

obtain the three parameters (E_{max}, n, and EC_{50}) and their standard deviations. The average drug
efficacy is measured using,

$$E_{ave} = \frac{1}{t_{start} - t_{end}} \int_{t_{start}}^{t_{end}} \epsilon(t) dt \qquad (3)$$

Where t_{start} and t_{end} are the treatment start day and end day, respectively.

Combined PKPD and VL models

The plasma concentration of molnupiravir obtained from the PK model is used in the PD model
to obtain time-dependent efficacy. Since molnupiravir imposes lethal mutations during the virus
replication process, in our model, a portion of produced viruses, measured by $\epsilon(t)$, are mutated
(V_m) and therefore assumed to be non-infectious, with the addition that most detected viral RNA
pre-treatment is also non-infectious. The production rate of non-mutated viruses is decreased by
a factor of $(1 - \epsilon(t))$. Equation 1e is written as,

$$\frac{dV}{dt} = (1 - \epsilon(t)) \pi I_P - \gamma V \qquad (4a)$$

$$\frac{dV_m}{dt} = (\epsilon(t)) \pi I_P - \gamma V_m \qquad (4b)$$

Total viral load ($V + V_m$) is used to fit the trial data.

Fitting the combined model to individual viral load data in the PLATCOV trial

We used the population mixed-effect approach and Monolix to estimate each individual's viral
dynamics parameters and the potency adjustment factor (paf). Due to the lack of data from the
initial phase of infection in the PLATCOV and PANORAMIC trials, we include the data from
Omicron-infected individuals in the NBA cohort to inform the model about the initial phase of
infection. We fixed the PK parameters to the estimated population values (**Table S1**), and the PD
parameters other than EC50 to the in vitro estimated population values (**Table S2**). We used the
study category (NBA vs PLATCOV and PANORAMIC) as a covariate for t_0 (timing of
infection) and τ (timing of the adaptive immune response) since the first recorded positive test is
likely much later for the clinical trials. In the NBA study, samples were collected almost daily
regardless of symptoms often leading to pre-symptomatic detection, while in the PLATCOV
study, the baseline measurement occurred after symptom onset, trial enrollment and consent.

Construction of a virtual cohort

To generate a cohort for our simulated clinical trials, we randomly selected 400 individuals (for
each arm of the simulated trials) from the unvaccinated symptomatic subpopulation of the NBA
cohort for MOVe-OUT and the vaccinated, Omicron infected subpopulation for PLATCOV and
PANORAMIC and used their individual viral load parameters estimated by fitting our viral
dynamics model to the data. A sample size of n=400 (out of 822 vaccinated individuals with
Omicron infection) was used to mimic a large-scale clinical trial and maintain relatively low
overlap between virtual cohorts used in each arm of the simulations and between different
simulations. Since the time of symptom onset is not available for all individuals in the NBA data,
we randomly drew an incubation period for each individual from gamma distributions with
variant-specific parameters estimated by Gamiche et al.[39] The start of treatment relative to
symptom onset was randomly selected according to a uniform distribution for MOVe-OUT and

PLATCOV, and a logit normal distribution for PANORAMIC with limits of [0,5] days and mean
and standard deviation reported in the PANORAMIC trial for control and treatment arms. The
same population PK parameters were assigned to each individual. The relevant dose in each
scenario was added to the A_{GI} compartment (the absorption equation) of the PK model (**eq 2a**) at
each dosing timepoint (t=0, 0.5, 1, 1.5, …., 4.5 days). For each dose, the appropriate PK
parameter were used (**Table S1**). PD parameters were also randomly drawn from a log-normal
distribution with the estimated mean and standard deviation. The standard deviation of the PD
parameters represents the accuracy of the assays and not individual variability.

Potency adjustment factor (paf)

The paf is defined as,

597
$$\text{paf} = \frac{EC_{50,in\ vivo}}{EC_{50,in\ vitro}} \tag{5}$$

We estimated the paf by maximizing R^2 when fitting the change in viral load of the treatment
arm of our simulation to the data from day 0 to day 7 of the treatment arm of the clinical trial.

Measuring rebound probability

A viral load rebound in the treatment arm was defined when the viral load at any time after
treatment ended exceeded the viral load at the end of the treatment by 1 log. In the control group,
viral rebound was defined in patients who had at least two peaks with minimum height of 1000
copies/ml in their viral load trajectories and the second peak was 1 log higher than its preceding
local minimum.

Software:

All the model fittings in this study were performed using Monolix version 2023R1.

The data analysis and simulations were performed using Python 3.9.12.

Data availability:

The data analyzed in this work was previously published by Hay et al. and Schilling et al. and is
available on GitHub at

https://github.com/gradlab/SC2-kinetics-immune-history and
https://github.com/jwatowatson/PLATCOV-Molnupiravir/tree/V1.0

Pharmacodynamics data is available on GitHub at
https://github.com/sEsmaeili/MolnupiravirModeling
619
620 <u>**Code availability:**</u>

621 All codes and materials used in the analysis are available on GitHub

622 https://github.com/sEsmaeili/MolnupiravirModeling

623 **Funding:**

This work is supported by NIAID R01AI77512-01 (JTS, SP).

References

1. Abdelnabi R, Foo CS, De Jonghe S, Maes P, Weynand B, Neyts J. Molnupiravir Inhibits Replication of the Emerging SARS-CoV-2 Variants of Concern in a Hamster Infection Model. J Infect Dis. 2021;224(5):749-53. doi: 10.1093/infdis/jiab361. PubMed PMID: 34244768; PMCID: PMC8408768.

2. Jayk Bernal A, Gomes da Silva MM, Musungaie DB, Kovalchuk E, Gonzalez A, Delos Reyes V, Martín-Quirós A, Caraco Y, Williams-Diaz A, Brown ML, Du J, Pedley A, Assaid C, Strizki J, Grobler JA, Shamsuddin HH, Tipping R, Wan H, Paschke A, ..., Group M-OS. Molnupiravir for Oral Treatment of Covid-19 in Nonhospitalized Patients. N Engl J Med. 2022;386(6):509-20. Epub 20211216. doi: 10.1056/NEJMoa2116044. PubMed PMID: 34914868; PMCID: PMC8693688.

3. Butler CC, Hobbs FDR, Gbinigie OA, Rahman NM, Hayward G, Richards DB, Dorward J, Lowe DM, Standing JF, Breuer J, Khoo S, Petrou S, Hood K, Nguyen-Van-Tam JS, Patel MG, Saville BR, Marion J, Ogburn E, Allen J, ..., Group PTC. Molnupiravir plus usual care versus usual care alone as early treatment for adults with COVID-19 at increased risk of adverse outcomes (PANORAMIC): an open-label, platform-adaptive randomised controlled trial. Lancet. 2023;401(10373):281-93. Epub 20221222. doi: 10.1016/S0140-6736(22)02597-1. PubMed PMID: 36566761; PMCID: PMC9779781.

4. Schilling WHK, Jittamala P, Watson JA, Boyd S, Luvira V, Siripoon T, Ngamprasertchai T, Batty EM, Cruz C, Callery JJ, Singh S, Saroj M, Kruabkontho V, Ngernseng T, Tanglakmankhong N, Tubprasert J, Abdad MY, Madmanee W, Kouhathong J, ..., Group PC. Antiviral efficacy of molnupiravir versus ritonavir-boosted nirmatrelvir in patients with early symptomatic COVID-19 (PLATCOV): an open-label, phase 2, randomised, controlled, adaptive trial. Lancet Infect Dis. 2024;24(1):36-45. Epub 20230928. doi: 10.1016/S1473-3099(23)00493-0. PubMed PMID: 37778363; PMCID: PMC7615401.

5. Hammond J, Leister-Tebbe H, Gardner A, Abreu P, Bao W, Wisemandle W, Baniecki M, Hendrick VM, Damle B, Simón-Campos A, Pypstra R, Rusnak JM, Investigators E-H. Oral Nirmatrelvir for High-Risk, Nonhospitalized Adults with Covid-19. N Engl J Med. 2022;386(15):1397-408. Epub 20220216. doi: 10.1056/NEJMoa2118542. PubMed PMID: 35172054; PMCID: PMC8908851.

6. Fountain-Jones NM, Vanhaeften R, Williamson J, Maskell J, Chua IJ, Charleston M, Cooley L. Effect of molnupiravir on SARS-CoV-2 evolution in immunocompromised patients: a retrospective observational study. Lancet Microbe. 2024;5(5):e452-e8. Epub 20240322. doi: 10.1016/S2666-5247(23)00393-2. PubMed PMID: 38527471.

7. Sanderson T, Hisner R, Donovan-Banfield I, Hartman H, Løchen A, Peacock TP, Ruis C. A molnupiravir-associated mutational signature in global SARS-CoV-2 genomes. Nature. 2023;623(7987):594-600. Epub 20230925. doi: 10.1038/s41586-023-06649-6. PubMed PMID: 37748513; PMCID: PMC10651478.

8. Rosenke K, Lewis MC, Feldmann F, Bohrnsen E, Schwarz B, Okumura A, Bohler WF, Callison J, Shaia C, Bosio CM, Lovaglio J, Saturday G, Jarvis MA, Feldmann H. Combined molnupiravir-nirmatrelvir treatment improves the inhibitory effect on SARS-CoV-2 in macaques. JCI Insight. 2023;8(4). Epub 20230222. doi: 10.1172/jci.insight.166485. PubMed PMID: 36574296; PMCID: PMC9977490.

9.   Wagoner J, Herring S, Hsiang TY, Ianevski A, Biering SB, Xu S, Hoffmann M,
Pöhlmann S, Gale M, Aittokallio T, Schiffer JT, White JM, Polyak SJ. Combinations of Host-
and Virus-Targeting Antiviral Drugs Confer Synergistic Suppression of SARS-CoV-2. Microbiol
Spectr. 2022;10(5):e0333122. Epub 20221003. doi: 10.1128/spectrum.03331-22. PubMed PMID:
36190406; PMCID: PMC9718484.
10.   Esmaeili S, Owens K, Wagoner J, Polyak SJ, White JM, Schiffer JT. A unifying model to
explain frequent SARS-CoV-2 rebound after nirmatrelvir treatment and limited prophylactic
efficacy. Nat Commun. 2024;15(1):5478. Epub 20240628. doi: 10.1038/s41467-024-49458-9.
PubMed PMID: 38942778; PMCID: PMC11213957.
11.   Owens K, Esmaeili S, Schiffer JT. Heterogeneous SARS-CoV-2 kinetics due to variable
timing and intensity of immune responses. JCI Insight. 2024;9(9). Epub 20240404. doi:
10.1172/jci.insight.176286. PubMed PMID: 38573774; PMCID: PMC11141931.
12.   Edelstein GE, Boucau J, Uddin R, Marino C, Liew MY, Barry M, Choudhary MC,
Gilbert RF, Reynolds Z, Li Y, Tien D, Sagar S, Vyas TD, Kawano Y, Sparks JA, Hammond SP,
Wallace Z, Vyas JM, Barczak AK, ..., Siedner MJ. SARS-CoV-2 Virologic Rebound With
Nirmatrelvir-Ritonavir Therapy : An Observational Study. Ann Intern Med. 2023;176(12):1577-
85. Epub 20231114. doi: 10.7326/M23-1756. PubMed PMID: 37956428; PMCID:
PMC10644265.
13.   Harrington PR, Cong J, Troy SB, Rawson JMO, O'Rear JJ, Valappil TI, McGarry
Connelly S, Farley J, Birnkrant D. Evaluation of SARS-CoV-2 RNA Rebound After
Nirmatrelvir/Ritonavir Treatment in Randomized, Double-Blind, Placebo-Controlled Trials -
United States and International Sites, 2021-2022. MMWR Morb Mortal Wkly Rep.
2023;72(51):1365-70. Epub 20231222. doi: 10.15585/mmwr.mm7251a2. PubMed PMID:
38127674; PMCID: PMC10754264.
14.   Anderson AS, Caubel P, Rusnak JM, Investigators E-HT. Nirmatrelvir-Ritonavir and
Viral Load Rebound in Covid-19. N Engl J Med. 2022;387(11):1047-9. Epub 20220907. doi:
10.1056/NEJMc2205944. PubMed PMID: 36069818; PMCID: PMC9513855.
15.   Perelson AS, Ribeiro RM, Phan T. An explanation for SARS-CoV-2 rebound after
Paxlovid treatment. medRxiv. 2023:2023.05.30.23290747. doi: 10.1101/2023.05.30.23290747.
16.   Hay JA, Kissler SM, Fauver JR, Mack C, Tai CG, Samant RM, Connolly S, Anderson DJ,
Khullar G, MacKay M, Patel M, Kelly S, Manhertz A, Eiter I, Salgado D, Baker T, Howard B,
Dudley JT, Mason CE, ..., Grad YH. Quantifying the impact of immune history and variant on
SARS-CoV-2 viral kinetics and infection rebound: A retrospective cohort study. Elife. 2022;11.
Epub 20221116. doi: 10.7554/eLife.81849. PubMed PMID: 36383192; PMCID: PMC9711520.
17.   Clairon Q, Pasin C, Balelli I, Thiébaut R, Prague M. Parameter estimation in nonlinear
mixed effect models based on ordinary differential equations: an optimal control approach.
Computational Statistics. 2023. doi: 10.1007/s00180-023-01420-x.
18.   Painter WP, Holman W, Bush JA, Almazedi F, Malik H, Eraut NCJE, Morin MJ,
Szewczyk LJ, Painter GR. Human Safety, Tolerability, and Pharmacokinetics of Molnupiravir, a
Novel Broad-Spectrum Oral Antiviral Agent with Activity Against SARS-CoV-2. Antimicrob
Agents Chemother. 2021;65(5). Epub 20210301. doi: 10.1128/AAC.02428-20. PubMed PMID:
33649113; PMCID: PMC8092915.
19.   Shen L, Peterson S, Sedaghat AR, McMahon MA, Callender M, Zhang H, Zhou Y, Pitt E,
Anderson KS, Acosta EP, Siliciano RF. Dose-response curve slope sets class-specific limits on
inhibitory potential of anti-HIV drugs. Nature Medicine. 2008;14(7):762-6. doi:
10.1038/nm1777.

20.   Zhou S, Long N, Rosenke K, Jarvis MA, Feldmann H, Swanstrom R. Combined
Treatment of Severe Acute Respiratory Syndrome Coronavirus 2 Reduces Molnupiravir-Induced
Mutagenicity and Prevents Selection for Nirmatrelvir/Ritonavir Resistance Mutations. J Infect
Dis. 2024. Epub 20240708. doi: 10.1093/infdis/jiae213. PubMed PMID: 38973065.
21.   Schiffer JT, Swan DA, Corey L, Wald A. Rapid viral expansion and short drug half-life
explain the incomplete effectiveness of current herpes simplex virus 2-directed antiviral agents.
Antimicrob Agents Chemother. 2013;57(12):5820-9. Epub 2013/09/11. doi:
10.1128/AAC.01114-13. PubMed PMID: 24018260; PMCID: PMC3837890.
22.   Schiffer JT, Swan DA, Magaret A, Corey L, Wald A, Ossig J, Ruebsamen-Schaeff H,
Stoelben S, Timmler B, Zimmermann H, Melhem MR, Van Wart SA, Rubino CM, Birkmann A.
Mathematical modeling of herpes simplex virus-2 suppression with pritelivir predicts trial
outcomes. Sci Transl Med. 2016;8(324):324ra15. Epub 2016/02/05. doi:
10.1126/scitranslmed.aad6654. PubMed PMID: 26843190; PMCID: PMC4880060.
23.   Backer JA, Eggink D, Andeweg SP, Veldhuijzen IK, van Maarseveen N, Vermaas K,
Vlaemynck B, Schepers R, van den Hof S, Reusken CB, Wallinga J. Shorter serial intervals in
SARS-CoV-2 cases with Omicron BA.1 variant compared with Delta variant, the Netherlands,
13 to 26 December 2021. Euro Surveill. 2022;27(6). doi: 10.2807/1560-
7917.ES.2022.27.6.2200042. PubMed PMID: 35144721; PMCID: PMC8832521.
24.   Wu Y, Kang L, Guo Z, Liu J, Liu M, Liang W. Incubation Period of COVID-19 Caused
by Unique SARS-CoV-2 Strains: A Systematic Review and Meta-analysis. JAMA Netw Open.
2022;5(8):e2228008. Epub 20220801. doi: 10.1001/jamanetworkopen.2022.28008. PubMed
PMID: 35994285; PMCID: PMC9396366.
25.   Li P, Wang Y, Lavrijsen M, Lamers MM, de Vries AC, Rottier RJ, Bruno MJ,
Peppelenbosch MP, Haagmans BL, Pan Q. SARS-CoV-2 Omicron variant is highly sensitive to
molnupiravir, nirmatrelvir, and the combination. Cell Res. 2022;32(3):322-4. Epub 20220120.
doi: 10.1038/s41422-022-00618-w. PubMed PMID: 35058606; PMCID: PMC8771185.
26.   Yotsuyanagi H, Ohmagari N, Doi Y, Yamato M, Bac NH, Cha BK, Imamura T, Sonoyama
T, Ichihashi G, Sanaki T, Tsuge Y, Uehara T, Mukae H. Efficacy and Safety of 5-Day Oral
Ensitrelvir for Patients With Mild to Moderate COVID-19: The SCORPIO-SR Randomized
Clinical Trial. JAMA Netw Open. 2024;7(2):e2354991. Epub 20240205. doi:
10.1001/jamanetworkopen.2023.54991. PubMed PMID: 38335000; PMCID: PMC10858401.
27.   Cruciani M, Pati I, Masiello F, Piccinini V, Pupella S, De Angelis V. SARS-CoV-2
infection rebound among patients receiving antiviral agents, convalescent plasma, or no
treatment: a systematic review with meta-analysis. Blood Transfus. 2024. Epub 20240527. doi:
10.2450/BloodTransfus.764. PubMed PMID: 38814880.
28.   Reeves DB, Mayer BT, deCamp AC, Huang Y, Zhang B, Carpp LN, Magaret CA, Juraska
M, Gilbert PB, Montefiori DC, Bar KJ, Cardozo-Ojeda EF, Schiffer JT, Rossenkhan R, Edlefsen
P, Morris L, Mkhize NN, Williamson C, Mullins JI, ..., Edupuganti S. High monoclonal
neutralization titers reduced breakthrough HIV-1 viral loads in the Antibody Mediated
Prevention trials. Nat Commun. 2023;14(1):8299. Epub 20231214. doi: 10.1038/s41467-023-
43384-y. PubMed PMID: 38097552; PMCID: PMC10721814.
29.   Reeves DB, Gaebler C, Oliveira TY, Peluso MJ, Schiffer JT, Cohn LB, Deeks SG,
Nussenzweig MC. Impact of misclassified defective proviruses on HIV reservoir measurements.
Nat Commun. 2023;14(1):4186. Epub 20230713. doi: 10.1038/s41467-023-39837-z. PubMed
PMID: 37443365; PMCID: PMC10345136.

30.   Weinreich DM, Sivapalasingam S, Norton T, Ali S, Gao H, Bhore R, Musser BJ, Soo Y,
Rofail D, Im J, Perry C, Pan C, Hosain R, Mahmood A, Davis JD, Turner KC, Hooper AT,
Hamilton JD, Baum A, ..., Investigators T. REGN-COV2, a Neutralizing Antibody Cocktail, in
Outpatients with Covid-19. N Engl J Med. 2021;384(3):238-51. Epub 20201217. doi:
10.1056/NEJMoa2035002. PubMed PMID: 33332778; PMCID: PMC7781102.
31.   Watson JA, Kissler SM, Day NPJ, Grad YH, White NJ. Characterizing SARS-CoV-2
Viral Clearance Kinetics to Improve the Design of Antiviral Pharmacometric Studies. Antimicrob
Agents Chemother. 2022;66(7):e0019222. Epub 20220623. doi: 10.1128/aac.00192-22. PubMed
PMID: 35736134; PMCID: PMC9295592.
32.   Standing JF, Buggiotti L, Guerra-Assuncao JA, Woodall M, Ellis S, Agyeman AA, Miller
C, Okechukwu M, Kirkpatrick E, Jacobs AI, Williams CA, Roy S, Martin-Bernal LM, Williams
R, Smith CM, Sanderson T, Ashford FB, Emmanuel B, Afzal ZM, ..., Group PV. Randomized
controlled trial of molnupiravir SARS-CoV-2 viral and antibody response in at-risk adult
outpatients. Nat Commun. 2024;15(1):1652. Epub 20240223. doi: 10.1038/s41467-024-45641-0.
PubMed PMID: 38396069; PMCID: PMC10891158.
33.   Lonsdale DO, Kipper K, Baker EH, Barker CIS, Oldfield I, Philips BJ, Johnston A,
Rhodes A, Sharland M, Standing JF. β-Lactam antimicrobial pharmacokinetics and target
attainment in critically ill patients aged 1 day to 90 years: the ABDose study. J Antimicrob
Chemother. 2020;75(12):3625-34. doi: 10.1093/jac/dkaa363. PubMed PMID: 32989452.
34.   Singh S, Boyd S, Schilling WHK, Watson JA, Mukaka M, White NJ. The relationship
between viral clearance rates and disease progression in early symptomatic COVID-19: a
systematic review and meta-regression analysis. J Antimicrob Chemother. 2024. Epub 20240222.
doi: 10.1093/jac/dkae045. PubMed PMID: 38385479.
35.   Prague M, Lavielle M. SAMBA: A novel method for fast automatic model building in
nonlinear mixed-effects models. CPT Pharmacometrics Syst Pharmacol. 2022;11(2):161-72.
Epub 20220201. doi: 10.1002/psp4.12742. PubMed PMID: 35104058; PMCID: PMC8846636.
36.   Bing A, Hu Y, Prague M, Hill AL, Li JZ, Bosch RJ, De Gruttola V, Wang R. Comparison
of empirical and dynamic models for HIV viral load rebound after treatment interruption. Stat
Commun Infect Dis. 2020;12(Suppl 1). Epub 20200821. doi: 10.1515/scid-2019-0021. PubMed
PMID: 34158910; PMCID: PMC8216669.
37.   Chan PL, Jacqmin P, Lavielle M, McFadyen L, Weatherley B. The use of the SAEM
algorithm in MONOLIX software for estimation of population pharmacokinetic-
pharmacodynamic-viral dynamics parameters of maraviroc in asymptomatic HIV subjects. J
Pharmacokinet Pharmacodyn. 2011;38(1):41-61. Epub 2010/11/19. doi: 10.1007/s10928-010-
9175-z. PubMed PMID: 21088872; PMCID: PMC3020311.
38.   Ke R, Zitzmann C, Ho DD, Ribeiro RM, Perelson AS. In vivo kinetics of SARS-CoV-2
infection and its relationship with a person's infectiousness. Proc Natl Acad Sci U S A.
2021;118(49). doi: 10.1073/pnas.2111477118. PubMed PMID: 34857628.
39.   Galmiche S, Cortier T, Charmet T, Schaeffer L, Chény O, von Platen C, Lévy A, Martin
S, Omar F, David C, Mailles A, Carrat F, Cauchemez S, Fontanet A. SARS-CoV-2 incubation
period across variants of concern, individual factors, and circumstances of infection in France: a
case series analysis from the ComCor study. Lancet Microbe. 2023;4(6):e409-e17. Epub
20230418. doi: 10.1016/S2666-5247(23)00005-8. PubMed PMID: 37084751; PMCID:
PMC10112864.
804
805