Development of a dynamic counterfactual risk stratification strategy for newly diagnosed acute myeloid leukemia patients treated with venetoclax and azacitidine

Nazmul Islam[1*], PhD; Justin L. Dale[1], BBA; Jamie S. Reuben[1], MPH; Karan Sapiah[1], MS; James W. Coates[1], MS; Frank R. Markson[1], BA; Jingjing Zhang[2], MD, PhD; Lezhou Wu[1], PhD; Maura Gasparetto[3], MD; Brett M. Stevens[1,3], PhD; Sarah E. Staggs[1,3], PhD; William M. Showers[1], MS; Monica R. Ransom[1,3], PhD; Jairav Desai[1], MS; Ujjwal V. Kulkarni[1], MS; Krysta L. Engel[1], PhD; Craig T. Jordan[3], PhD; Michael Boyiadzis[4], MD; Clayton A. Smith[1, 3,4], MD

[1]RefinedScience, Aurora, Colorado

[2]Department of Pathology, University of Colorado Anschutz Campus, Aurora, Colorado

[3]Department of Medicine, University of Colorado Anschutz Campus, Aurora, Colorado

[4]OncoVerity, Aurora, Colorado

*Communicating Author:

Nazmul Islam, PhD

RefinedScience, 2115 N Scranton Street, Suite 2-70

Aurora Colorado, 80045, USA

+1-919-604-9705

nazmul.islam@refinedscience.com

Key words: Acute Myeloid Leukemia; Venetoclax; Azacitidine; Risk Stratification; Machine Learning

Abstract count: 240

Word count: 3,299 (Tables: 1, Figures: 6)

Article type: Original Report (Research and Applications)

CONTEXT SUMMARY

Key Objective

What is the empirical risk stratification strategy for de-novo Acute Myeloid Leukemia (AML) patients treated with venetoclax plus azacitidine (ven/aza)?

Knowledge Generated

The proposed tunable machine learning (ML)-based risk-stratification approach addresses real-world data (RWD) challenges and enables clinical implementation by adapting to site-specific preferences for context-specific parameters and analytical samples of the corresponding site. The proposed strategy produces a series of risk models and exhibits superior, but clinically meaningful, numerical performance compared to the state-of-the-art static one-size-fits-all approach proposed by the European Leukemia Net (ELN).

ABSTRACT

Objective

The objective of this study was to develop a flexible risk model (RM) stratification strategy for Acute Myeloid Leukemia (AML) that is specific for the new standard of care venetoclax plus azacitidine (ven/aza), captures disease heterogeneity, and addresses a range of real-world data issues.

Materials and Methods

A series of tunable RMs based on a dynamic counterfactual machine learning (ML) strategy that utilized next-generation sequencing, cytogenetics, flow cytometry, and other features of diagnostic AML samples were developed and tested on a single institutional cohort of 316 newly diagnosed patients treated initially with ven/aza.

Results

Favorable, Intermediate, and Adverse risk groups were identified in a series of novel RMs derived using ML models for overall survival (OS) and event-free survival (EFS). Most, but not all, models demonstrated equitable patient distribution into different risk categories (~20%-40% in each group) with significant separation between categories (log-rank based P-values <0.001), and with predictability computed by survival AUC values in the 0.60-0.70 range.

Discussion and Conclusion

The general strategy employed here is specific for AML patients treated with ven/aza, is based on a wide range of diagnostic AML pathology features, considers feature interactions, and addresses data missingness, sparsity, and the confounding effects of allogeneic hematopoietic cell transplant. It is also readily tunable through simple coding and context-specific parameter updates and is adaptable to reflect different use-case needs. This strategy represents a new approach to developing more effective RMs for AML and possibly other diseases.

INTRODUCTION

Acute myeloid leukemia (AML) is an aggressive hematopoietic malignancy diagnosed in ~20,000

individuals annually in the United States (US)[1]. For young and fit patients, initial treatment typically

involves intensive chemotherapy (IC), followed by consolidation with allogeneic hematopoietic cell

transplant (allo-HCT) or additional high-dose chemotherapy[2,3]. For older and less-fit patients,

venetoclax (ven) combined with a hypomethylating agent (HMA) such as azacitidine (aza) has become

a standard of care[4-6]. For IC therapies, various prognostic models have been developed based on

patient- and AML-specific features to stratify patients into subgroups with varying outcomes[7-15]. The

European Leukemia Network (ELN) developed a widely used risk categorization approach for AML in

2017 (ELN17), with an update in 2022 (ELN22), which was subsequently adopted and modified by the

National Comprehensive Cancer Network (NCCN) and others[16,17]. ELN categories are divided into

Favorable, Intermediate, and Adverse groups associated with overall survival (OS) based on AML

cytogenetic (CYT), fluorescence *in situ* hybridization (FISH), and next-generation sequencing (NGS)

features[16,17]. However, it was recently shown that the ELN17 and ELN22 risk categorization

approaches performed sub-optimally and did not effectively stratify ven/HMA-treated patients, likely

because the ELN criteria are primarily based on treatment outcomes following IC[18-22]. Here, we

describe the development of a flexible AML risk stratification strategy for newly diagnosed AML patients

treated with ven/aza based on a wide range of AML diagnostic pathology features, while addressing the

real-world data (RWD) challenges that can alter the effectiveness of risk models (RMs).

METHODS

Patient populations and outcomes definitions

The dataset included 316 adult patients from the University of Colorado (CU) with newly diagnosed AML treated with ven/aza as standard care or in a clinical trial between January 2015 and March 2024. Patients with acute promyelocytic leukemia (APL), bi-phenotypic leukemia, blast crisis chronic myeloid leukemia, or central nervous system involvement with AML at diagnosis were excluded. The details of the patient cohort and subset management are shown in Figure 1. The patient demographics are summarized in Supplemental Table 1A. Key AML diagnostic pathology features for developing the RMs are summarized in Supplemental Tables 1B-F, including NGS, CYT, FISH, PCR results, flow cytometric (FC) data, and "AML mutations," which included composite mutations from various tests (patients with multiple positive tests were counted once). For example, "p53m" composite mutation is a combination of p53 abnormalities by NGS and losses or gains of 17p by FISH. Complex cytogenetics were defined as ≥ 3 karyotype abnormalities. Data, including NGS and FC results, were simplified into binary values as described previously[23]. FISH data were simplified by combining all chromosomal specific probes into "Chr Abn". FISH data for DEK and MECOM were not routinely available before mid-2019 (0 DEK results, 3/7 positive MECOM results) and thus were not used before that date. Additional feature nomenclature is provided in the respective figure legends and Supplemental Table 2. Responses (overall survival (OS), event-free survival (EFS), best response (BR)) are defined in Figure 1[17, 23]. This retrospective study was approved by the institutional review board (IRB) and used a limited dataset with a waiver of consent.

Exploratory analyses

Descriptive statistics were provided for baseline covariates and outcomes[24]. Kaplan-Meier (KM) analyses were performed assuming right censoring for OS and EFS based on the Full Analytical Set (FAS), dataset censoring allo-HCT recipients, and set excluding allo-HCT recipients. The equality of survival curves was evaluated using various testing approaches[25-31] and noise variables were filtered to

select a parsimonious feature list; further details are provided in the Supplemental Methods. The resulting list of confounding variables (Supplemental Tables 1B-F) was used for both multivariable OS and EFS models.

Risk model development

Table 1 summarizes the overall RM specifications, and Figure 2 shows the RM development strategy. The key components include the following.

Survival regressions

Machine learning (ML) models (multivariable penalized (ridge) Cox proportional hazard (mCOXr)[32,33]) were fitted adjusting for age and gender for OS and EFS. Separate models were fitted adjusting for confounding variables selected by null hypothesis significance testing (NHST) (Step 1, Figure 2) to identify influential variables within each group (Step 3, Figure 2). Missing values were treated as separate categories (Step 2, Figure 2). The tuning parameters were selected via cross-validation (CV). The corresponding 95% confidence intervals (CIs) of adjusted hazard ratios (aHRs) were constructed using the fractional random weight bootstrap (FRWB)[34] approach of 2,000 runs (Supplemental Table 3-4).

Counterfactual effects estimation

Biomarker-specific mean predicted marginal risks (MRs) over time were calculated using "*recycled*" predictions with counterfactual arguments for the corresponding mCOXr models (Step 4, Figure 2)[35,36]. A fixed patient cohort was used for each feature to estimate risk differences (RDs) adjusting for risk factor mix and severity while reducing potential bias from skewed distributions of patient characteristic. At each time point, the MRs of a biomarker being positive or negative were calculated by averaging the predicted marginal risk probabilities for all patients as if they were (hypothetically) (+ve) or (-ve) for genetic features, or if they expressed a marker by FC. Bootstrap-based MR profiles were constructed by averaging the FRWB estimates, and the corresponding percentile-based 95% CIs were reported. The Kolmogorov-Smirnov test assessed the differences in counterfactual risk distributions[37].

Multivariable penalized (ridge) logistic regressions[33] were used to model BR confirming the consistency of the mCOXr estimates.

Variations in risk models

Varying endpoints: RMs for OS and EFS were developed separately in contrast to the *one-size-fits-all* approach.

Treatment variation: Allo-HCT is a potentially curative therapy that can profoundly affect ven/aza treated patient outcomes[38]. To address this confounding factor, RMs were developed by censoring patients with allo-HCT. In a sample with a high proportion of allo-HCT recipients (92/316), as in CU, survival probabilities may be overestimated if not adjusted for allo-HCT, leading to improper feature selection and suboptimal characterization of the relationship between response and features.

Dynamic behavior: Counterfactual RDs were computed dynamically over 36 months (Steps 4-5, Figure 2, and Supplemental Figures 1A-1D). Landmarks of 14.7 and 9.8 months were chosen to find optimal feature-specific RD for OS and EFS based on the median time-to-event in the VIALE-A study, which established ven/aza as a new standard of care[6].

Prevalence: Variables with low and high prevalence for (+ve) cases may cause unstable parameter estimates and imbalances in subject-level stratification. Rule-I (Step 5, Figure 2) is flexible and does not exclude features with respect to prevalence. Three additional rules were considered with varying degrees of stringency – Rule-II excluded any variables containing less than 2% and more than 40% (+ve) cases in risk stratification even if they exhibited more than 5% RD. Rule-III used FRWB based MR profiles to calculate RD, and Rule-IV combined Rule-II and Rule-III; see Supplemental Table 5A.

Covariate-level classification: Two levels of classification, covariate and subject, were performed. The magnitude and direction of marginal RDs between (+ve) and (-ve) at a time point of interest were used to classify each feature into either "Adverse" or "Favorable". Positive RD values of 0-5%, 5-10%, and >10% classified a feature as *slightly*, *moderately*, or *strongly* "Adverse," respectively. Similarly, negative RD values determined *slightly*, *moderately*, or *strongly* "Favorable" features (Step 6, Figure 2). A cut-off

value of 5% was assumed to be the clinically relevant minimum RD. Supplemental Tables 5B-5E depict covariate-level classifications under different combinations of parameters.

Subject-level classification: Patient-specific risk stratification ("Adverse", "Intermediate", and "Favorable") was determined by combining covariate-level classifications using a 'Boolean' operator (AND/OR). This assigns a patient to the "Favorable" risk group without any "Adverse" features and vice versa. A patient was assigned to the "Intermediate" risk group without any "Favorable" and "Adverse" features or with at least one "Favorable" and one "Adverse" feature (Step 7, Figure 2).

Model specification: RMs using genetic and genetic-plus-phenotypic features were developed to reflect different scenarios and their implications on model performance (Table 1).

Missingness: Primary analyses treated missing values as a separate category for simplicity, avoiding imputation or exclusion. Reported results were based on this approach unless otherwise specified. Additional analyses were performed to check the robustness of different missing data handling techniques at the covariate-level classification step including: (A) Biases stemming from complete cases (CC) analyses were adjusted by incorporating inverse probability weights (IPW) into mCOXr, where subject-specific weights signaling the propensity of having any missing data were estimated from multivariable logistic regression models[39]. (B) Multiple imputations (five) were generated for each missing data element by exploiting multivariate data sequentially (multivariate imputation by chained equations (MICE))[40,41]. The corresponding aHRs were averaged over five samples and 2,000 FRWB runs, and percentile-based 95% CIs were reported. Subsequently, the marginal RDs were calculated as before and averaged over five samples. (C) Covariate-level classifications were performed based on different missing methods, with final stratification obtained by synthesizing with respect to *majority-votes*.

Evaluation of numerical performance

Multiple evaluation criteria were used to provide a holistic view of numerical performance in terms of *equitability* (extent of risk group size variation), *separability* (extent of outcome variation by risk strata),

conformity (extent of risk group overlap between methods), and *predictability* (degree of prognostic relationship with outcomes). Internal validation of *predictability* of the RMs and ELN22 for OS and EFS was conducted using 15-fold CV; the corresponding steps are illustrated in Figure 3. Further details of the computation and evaluation criteria are provided in the Supplemental Methods.

All statistical tests, except for the comparison of MR curves and survival AUC values, were two-sided with 5% significance level. As the analyses were exploratory, no multiplicity adjustments were made. All analyses were performed using R, version 4.2.3.

RESULTS

OS risk models

Using the strategy described above, we first explored whether a novel genetic feature-based RM, termed RM_G, could be developed. Covariate-level classification Rule-II demonstrated superior numerical performance with respect to *equitability* (Figure 4A), *separability* (Figure 4B-D), *conformity* (Supplemental Figure 5), and *predictability* (Supplemental Table 6B). Risk features identified using this variant of RM_G, termed $RM_{G,II}$, and the distribution of patients in different risk categories are summarized in Figure 4A. $RM_{G,II}$ demonstrated equitable distributions of patients and effective separation of Adverse, Intermediate, and Favorable subgroups based on OS and BR in both the allo-HCT excluded and FAS cohorts (Figures 4B and 4C). Similar behavior was noted when applied to the patient cohort where allo-HCT recipients were censored (Figure 4D, left panel). In the subset of only the patients who underwent allo-HCT, the RM did not distinguish Favorable and Intermediate risk patients, however it did distinguish the Adverse risk population from the other two categories (Figure 4D, right panel). A similar comparison was performed for the RM_G variants using Rule-I ($RM_{G,I}$), Rule-III ($RM_{G,III}$), and Rule-IV ($RM_{G,IV}$) which demonstrated similar performance characteristics, except for reduced equitability in the assignment of patients to different risk categories (Supplemental Figure 2-4 as well as Supplemental Tables 6A, 6C, and 6D). The other OS models summarized in Table 1 provided similar results (Supplemental Tables 5B, 5D-5E). The application of the ELN22 model to the FAS demonstrated a relatively skewed distribution of patients assigned to the Adverse risk category and less effective separation of OS curves compared to the $RM_{G,II}$ model for both the allo-HCT excluded cohort and the FAS (Supplemental Figure 5A-B). Pairwise comparison of the allo-HCT excluded cohort between $RM_{G,II}$ and ELN22 demonstrated substantial OS differences in the Adverse risk groups, moderate differences in the Intermediate groups, and relatively similar behavior in the Favorable risk groups (Supplemental Figure 5C). Inter-group conformity analyses using Fleiss kappa demonstrated the highest level of concordance in patient assignment for the Favorable risk category (Supplemental Figure 5D). Predictive evaluations (time-varying AUCs and iBrier) favored the $RM_{G,II}$ model

(Supplemental Table 6B bottom). Next, we explored whether the same RM strategies could be applied to a feature set comprising genetic-plus-phenotypic features defined by FC termed as RM_{GP}. The impacts of applying Rule-I, II, III, and IV to the RM_{GP} development strategy were explored as described above (Supplemental Figures 6-9 and Supplemental Tables 6A-6D top). In general, the overall performance of the RM_{GP} models paralleled that of the RM_G models, but with less equitable assignment of patients to the Favorable risk category depending on models, rules, and assumptions. Together, these findings demonstrate that the proposed ML-based strategy may be effective in generating AML ven/aza-specific RMs for OS and highlight the impact of different assumptions and rules on overall RM performance.

EFS risk models

Next, we explored whether applying the general RM development strategy to EFS was feasible, as this is another key metric in AML analysis[42]. RMs with features based on the genetics and genetics-plus-phenotypic features set were developed and tested as above. The covariate-level classification Rule-II for the EFS-specific model using only genetic features ($RM_{G,II-EFS}$) demonstrated the best performance (Figure 5A-5C and Supplemental Table 7B). The comparison between the $RM_{G,II-EFS}$ and ELN22 was similar to that for OS (Supplemental Figure 13). Rule-I, Rule-III, and Rule-IV were also explored in the context of genetic features-based EFS RM. $RM_{G,I-EFS}$ resulted in skewed distributions among risk strata with few Favorable and Adverse risk patients leading to suboptimal numerical performance (Supplemental Figure 10 and Supplemental Table 7A). In contrast, the $RM_{G,EFS}$ models utilizing Rule-III and Rule-IV behaved more similarly to that of Rule-II (Supplemental Figures 11-12 and Supplemental Table 7C-7D bottom). When utilizing the genetics-plus-phenotypic features, the RM_{GP} strategy applied to EFS roughly paralleled the behavior of the RM_G strategy (Supplemental Figures 14-17 and Supplemental Tables 7A-7D top). Overall, the findings indicate that EFS can be effectively stratified using the appropriate EFS-specific RM_G and RM_{GP} models but again highlight the impact of different assumptions and rules on model performance.

A series of heatmaps summarizing patient-specific risk classifications of 316 patients for OS and EFS are depicted in Figure 6 and Supplemental Figure 18, where moderate differences in classifications were observed across different modeling assumptions further emphasizing the impact of context-specific parameters on endpoints.

DISCUSSION

The study objective was to develop a risk stratification strategy reflecting RWD challenges that could be readily tunable for specific use-cases, objectives, and resources without loss of generality. This approach is inherently more flexible and adaptive than the ELN17 and ELN22 risk categories, where the risk stratification remains relatively fixed until the accumulated published data gets updated periodically after several years. Specifically, these traditional approaches can lag rapidly evolving changes in practices, diagnoses, and treatments, such as the advent of ven/HMAs as a new standard of care treatment for patients unfit for IC. Recently, lower intensity specific RMs have been developed to address some of these concerns[18-21,43]. However, several of these RMs were derived from patients treated with a range of agents and schedules, and were based on a limited number of NGS features that may not comprehensively reflect AML heterogeneity and biology. Additionally, these newer RMs may not have captured critical RWD challenges including missingness, sparsity, scarcity, endpoint types, collinearity, and the impact of allo-HCT on OS[19-21,43,44]. Lastly, as with the ELN17 and ELN22, many of these models are not readily adaptable as technologies and findings evolve. In contrast, the proposed RMs (1) were based on probabilistic RDs, (2) dealt with feature interactions (using a logic-based approach at the level of two-factor interactions), (3) addressed the dynamic effects of features, (4) were developed separately for different endpoints, (5) were treatment specific, (6) accounted for the confounding effects of allo-HCT, and 7) included a range of features (genetic and genetic-plus-phenotypic) adjusting for collinearity. The assessments using multiple evaluation criteria demonstrated clinical and numerical effectiveness, while showing that different modeling assumptions, adjustments, and rules can influence covariate- and subject-level classifications. In the initial analyses, the $RM_{G,II}$ appeared to have better performance than the ELN22 (Supplemental Figure 5 and 13 for *equitability*, *conformity*, and *separability*, and Supplemental Table 6B for *predictability*). If further validated, this could have important consequences for assessing the impact of novel AML therapies and diagnostics in clinical trials and real-world settings.

The proposed RMs are tunable through coding, parameters, and feature updates by readily accommodating changes in diagnostics, treatments, assumptions, definitions, and rules to reflect evolving changes in information, technology, and use-cases. This capability is particularly important in AML, as new insights and advancements in clinical care, treatment, disease biology, and testing modalities have led to complex datasets, which introduce rapid changes in RM performance and utility.

A series of RMs were proposed emphasizing various aspects of real-world and clinical trial data, objectives, assumptions, and requirements. The "best" performing model is typically subjective, depending on the context and dataset where it is applied. The proposed approach differs philosophically from the conventional one-size-fits-all approach, which assumes the model works well for all scenarios regardless of underlying differences in datasets. In contrast, the proposed strategy requires investigators to consider nuances in the respective analytical dataset and choose model parameters accordingly. For example, for a site like CU, where the proportion of allo-HCT is high, adjustments for allo-HCT needs to be made. However, this may be redundant for sites where the proportion of allo-HCT is low or negligible. One needs to choose how rigorously to account for missingness in the dataset which primarily depends on the magnitude and pattern of missingness. Furthermore, a user may prefer phenotypic-and-genetic models over genetic models alone. The proposed approach enables potential clinical implementation of a risk stratification methodology adapted to site-specific preferences and underlying patient population. Such a single, simple, and practical version of the RMs described here are underway, and the current status of this work is described in a preprint[45] which also discusses further development, external validation, and additional comparison with the ELN22.

There are several limitations to this study. This is an exploratory retrospective study and thus the findings need to be validated in confirmatory prospective studies. Although genetic, FC, FISH, and composite AML mutation features were simplified into binary variables for numerical convenience and

interpretability in counterfactual estimates, these steps may result in critical information loss and miss important biological aspects of AML. Similarly, the composite mutation category for RUNX and other features (Supplemental Table 1E) treated all abnormalities as potentially biologically relevant, which may be an oversimplification. Reintroducing the complexity of these features' constituents may improve RM stratification ability providing additional clinical insights; however, future studies with larger datasets are needed for validation. While we developed RMs combining genetic and phenotypic features, FC remains unstandardized between centers; therefore, the proposed models may be best suited for single-center or small consortium settings with consistent FC datasets. There is also a risk of underreporting samples with dimly expressed FC features, which may occur with markers such as CD45. Some RMs showed suboptimal stratification for allo-HCT recipients, which may improve by developing RMs with relevant features specific to this cohort. The interplay between genetic and AML features was accounted for on a logical rather than parametric basis. An alternative flexible modeling approach (e.g., tree-based ensembles) trained on a larger sample could alleviate this concern. Using the same analytical dataset for variable selection, risk classifications, and internal validation may induce "*data-leakage*" leading to over-fitting. Additionally, all models, particularly those with higher stringency, risk missing important features due to underestimated effects, small sample size, and prespecified cut-offs for prevalence rates and RDs. In contrast, less stringent RMs may incorporate noisy features with spurious associations. Therefore, to understand the generalizability of the RMs, validating findings with external, independent, and diverse data from multiple institutions, and through comparison to published findings and newly emerging RMs[20,43,44] for ven/aza and other low intensity treatments is critical. A simplified version of the RMs described here was developed and validated with a separate external dataset; details are provided in Islam et al.[45].

CONCLUSION

In summary, we developed and internally validated a risk stratification strategy and provided a series of RMs for AML patients treated with ven/aza, addressing the limitations in previous AML risk

classification models. The simplified version of these novel models could be useful for upfront patient allocation to risk groups in clinical trials and post hoc analysis of novel treatments for different AML risk groups. The proposed strategies are disease-agnostic and may be adaptable to other diseases.

ACKNOWLEDGMENTS

The authors gratefully thank Austin E. Gillen from the Department of Medicine, University of Colorado Anschutz Campus, Aurora, Colorado; in addition, Bin Yao and Adam George from the OncoVerity, Aurora, Colorado; and Grant Weller from the RefinedScience, Aurora, Colorado for their feedback and reviews during this project.

CONFLICT OF INTEREST

Both CAS and MB are employees of and hold equity in OncoVerity. In addition, CAS is a consultant to RefinedScience. All other authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

CAS and NI designed the study and drafted the manuscript. JLD, JSR, KS, JWC, FRM, and UVK processed and pulled the structured analytical datasets. CAS, NI, JSR, JLD, JZ, KS, JWC, and LW assessed the validity and quality of data. NI performed numerical analyses. CAS, NI, CTJ, JSR, JLD, JZ, MB, SES, MG, BMS, WMS, MRR, and KLE interpreted the results of the analyses. All authors reviewed, provided constructive comments, and agreed to its publication.

FUNDING

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

SUPPLEMENTARY MATERIAL

Supplementary material is available along with the submission detailing additional numerical results pertinent to the study.

DATA AVAILABILITY

The raw, individual patient data are protected and not available due to data privacy laws. The processed data are available at reasonable request to the corresponding author.

CODE SHARING

Controlled access to the code may be provided to qualified researchers upon request under a confidentiality and data use agreement, subject to institutional approval. For request, please email at nazmul.islam@refinedscience.com.

ABBREVIATIONS

Acute myeloid leukemia	AML
Intensive chemotherapy	IC
Allogeneic hematopoietic cell transplant	allo-HCT
European Leukemia Network	ELN
Complete response	CR
Best response	BR
CR with incomplete hematologic recovery	CRi
CR with partial hematologic recovery	CRh
Morphologic leukemia free state	MLFS
Partial remission	PR
Cytogenetic	CYT
Flow cytometric	FC
Fluorescence in situ hybridization	FISH
Next generation sequencing	NGS
Polymerase chain reaction	PCR
Venetoclax plus azacytidine	ven/aza
Full analytical set	FAS
Event-free survival	EFS
Overall survival	OS
Institutional review board	IRB
University of Colorado	UC
Kaplan-Meier	KM
Confidence interval	CI
Null hypothesis significance testing	NHST
Log-rank	LR
Tarone-Ware	TW
Fleming-Harrington	FH
Weighted multiple direction	mdir

Max-Combo	MC
K-sample omnibus non-proportional hazard	KONP
Restricted mean survival times	RMST
Machine learning	ML
Multivariable Cox proportional hazard models with ridge (L2-norm) penalty	mCOXr
Proportional hazard	PH
Fractional random weight bootstrap	FRWB
Adjusted hazard ratio	aHR
Cross-validation	CV
Kolmogorov-Smirnov	KS
Area under the curve	AUC
Receiver operative characteristics	ROC
Cumulative case dynamic control ROC	cAUC
Incident case dynamic control ROC	iAUC
Integrated Brier scores	IBS
Complete cases	CC
Inverse probability weights	IPW
Multivariate imputation by chained equations	MICE
Marginal risk	MR
Risk model	RM
Risk difference	RD
RM based on genetic-plus-phenotypic features with respect to OS	RM_{GP}
RM based on genetic features with respect to OS	RM_G
RM based on genetic-plus-phenotypic features with respect to EFS	RM_{GP-EFS}
RM based on genetic features with respect to EFS	RM_{G-EFS}
Real world data	RWD
Rule-I	I
Rule-II	II
Rule-III	III
Rule-IV	IV

REFERENCES

Shallis, R. M., Wang, R., Davidoff, A., Ma, X. & Zeidan, A. M. Epidemiology of acute myeloid leukemia: Recent progress and enduring challenges. *Blood Rev* **36**, 70-87 (2019). https://doi.org/10.1016/j.blre.2019.04.005

Bittencourt, M. C. B. & Ciurea, S. O. Recent Advances in Allogeneic Hematopoietic Stem Cell Transplantation for Acute Myeloid Leukemia. *Biol Blood Marrow Transplant* **26**, e215-e221 (2020). https://doi.org/10.1016/j.bbmt.2020.06.007

Blum, W. G. & Mims, A. S. Treating acute myeloid leukemia in the modern era: A primer. *Cancer* **126**, 4668-4677 (2020). https://doi.org/10.1002/cncr.32904

Chua, C. C. *et al.* Chemotherapy and Venetoclax in Elderly Acute Myeloid Leukemia Trial (CAVEAT): A Phase Ib Dose-Escalation Study of Venetoclax Combined With Modified Intensive Chemotherapy. *J Clin Oncol*, JCO2000572 (2020). https://doi.org/10.1200/JCO.20.00572

DA, P., M, A., P, S. & MY, K. Venetoclax for AML: changing the treatment paradigm. Blood Adv. 2019;3(24):4326-4335. *Blood Adv* **4**, 1020 (2020). https://doi.org/10.1182/bloodadvances.2020001769

DiNardo, C. D. *et al.* Azacitidine and Venetoclax in Previously Untreated Acute Myeloid Leukemia. *N Engl J Med* **383**, 617-629 (2020). https://doi.org/10.1056/NEJMoa2012971

Kantarjian, H. *et al.* Results of intensive chemotherapy in 998 patients age 65 years or older with acute myeloid leukemia or high-risk myelodysplastic syndrome: predictive prognostic models for outcome. *Cancer* **106**, 1090-1098 (2006). https://doi.org/10.1002/cncr.21723

Malfuson, J. V. *et al.* Risk factors and decision criteria for intensive chemotherapy in older patients with acute myeloid leukemia. *Haematologica* **93**, 1806-1813 (2008). https://doi.org/10.3324/haematol.13309

Kantarjian, H. *et al.* Intensive chemotherapy does not benefit most older patients (age 70 years or older) with acute myeloid leukemia. *Blood* **116**, 4422-4429 (2010). https://doi.org/10.1182/blood-2010-03-276485

Pastore, F. *et al.* Combined molecular and clinical prognostic index for relapse and survival in cytogenetically normal acute myeloid leukemia. *J Clin Oncol* **32**, 1586-1594 (2014). https://doi.org/10.1200/JCO.2013.52.3480

DiNardo, C. D. & Cortes, J. E. Mutations in AML: prognostic and therapeutic implications. *Hematology Am Soc Hematol Educ Program* **2016**, 348-355 (2016). https://doi.org/10.1182/asheducation-2016.1.348

Daneshbod, Y., Kohan, L., Taghadosi, V., Weinberg, O. K. & Arber, D. A. Prognostic Significance of Complex Karyotypes in Acute Myeloid Leukemia. *Curr Treat Options Oncol* **20**, 15 (2019). https://doi.org/10.1007/s11864-019-0612-y

Estey, E. H. Acute myeloid leukemia: 2021 update on risk-stratification and management. *Am J Hematol* **95**, 1368-1398 (2020). https://doi.org/10.1002/ajh.25975

Sasaki, K. *et al.* Prediction of survival with intensive chemotherapy in acute myeloid leukemia. *Am J Hematol* **97**, 865-876 (2022). https://doi.org/10.1002/ajh.26557

Eckardt, J. N. *et al.* Prediction of complete remission and survival in acute myeloid leukemia using supervised machine learning. *Haematologica* **108**, 690-704 (2023). https://doi.org/10.3324/haematol.2021.280027

Dohner, H. *et al.* Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel. *Blood* **129**, 424-447 (2017). https://doi.org/10.1182/blood-2016-08-733196

Dohner, H. *et al.* Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood* **140**, 1345-1377 (2022). https://doi.org/10.1182/blood.2022016867

Hoff, F. W. *et al.* Beat-AML 2024 ELN-Refined Risk Stratification for Older Adults with Newly Diagnosed AML Given Lower-Intensity Therapy. *Blood Adv* (2024). https://doi.org/10.1182/bloodadvances.2024013685

Dohner, H. *et al.* Genetic Risk Stratification and Outcomes Among Treatment-Naive Patients With AML Treated With Venetoclax and Azacitidine. *Blood* (2024). https://doi.org/10.1182/blood.2024024944

Dohner, H. *et al.* Genetic risk classification for adults with AML receiving less-intensive therapies: the 2024 ELN recommendations. *Blood* (2024). https://doi.org/10.1182/blood.2024025409

Bataller, A. *et al.* Prognostic risk signature in patients with acute myeloid leukemia treated with hypomethylating agents and venetoclax. *Blood Adv* **8**, 927-935 (2024). https://doi.org/10.1182/bloodadvances.2023011757

Dohner, H. *et al.* ELN Risk Stratification Is Not Predictive of Outcomes for Treatment-Naïve Patients with Acute Myeloid Leukemia Treated with Venetoclax and Azacitidine. *Blood* **140 (Supplement 1)**, 1441-1444 (2022).

Islam, N. *et al.* Machine Learning-Based Exploratory Clinical Decision Support for Newly Diagnosed Patients With Acute Myeloid Leukemia Treated With 7 + 3 Type Chemotherapy or Venetoclax/Azacitidine. *JCO Clin Cancer Inform* **6**, e2200030 (2022). https://doi.org/10.1200/CCI.22.00030

Austin, P. C. An Introduction to Propensity Score Methods for Reducing the Effects of Confounding in Observational Studies. *Multivariate Behav Res* **46**, 399-424 (2011). https://doi.org/10.1080/00273171.2011.568786

Ditzhaus, M. & Friedrich, S. More powerful logrank permutation tests for two-sample survival data. arXiv:1807.05504 (2018). <https://ui.adsabs.harvard.edu/abs/2018arXiv180705504D>.

Fleming, T. R., Harrington, D. P. & O'Sullivan, M. Supremum Versions of the Log-Rank and Generalized Wilcoxon Statistics. *Journal of the American Statistical Association* **82**, 312-320 (1987). https://doi.org/10.1080/01621459.1987.10478435

Gorfine, M., Schlesinger, M. & Hsu, L. K-sample omnibus non-proportional hazards tests based on right-censored data. *Stat Methods Med Res* **29**, 2830-2850 (2020). https://doi.org/10.1177/0962280220907355

TARONE, R. E. & WARE, J. On distribution-free tests for equality of survival distributions. *Biometrika* **64**, 156-160 (1977). https://doi.org/10.1093/biomet/64.1.156

Tian, L., Zhao, L. & Wei, L. J. Predicting the restricted mean event time with the subject's baseline covariates in survival analysis. *Biostatistics* **15**, 222-233 (2014). https://doi.org/10.1093/biostatistics/kxt050

Uno, H. *et al.* Moving beyond the hazard ratio in quantifying the between-group difference in survival analysis. *J Clin Oncol* **32**, 2380-2385 (2014). https://doi.org/10.1200/JCO.2014.55.2208

Horiguchi, M. & Uno, H. On permutation tests for comparing restricted mean survival time with small sample from randomized trials. *Stat Med* **39**, 2655-2670 (2020). https://doi.org/10.1002/sim.8565

Simon, N., Friedman, J., Hastie, T. & Tibshirani, R. Regularization Paths for Cox's Proportional Hazards Model via Coordinate Descent. *J Stat Softw* **39**, 1-13 (2011). https://doi.org/10.18637/jss.v039.i05

Gareth, J., Witten, D., Hastie, T. & Tibshirani, R. *An introduction to statistical learning*. Vol. 112 (Springer, 2013).

Xu, L., Gotwalt, C., Hong, Y., King, C. B. & Meeker, W. Q. Applications of the Fractional-Random-Weight Bootstrap. *The American Statistician* **74**, 345-358 (2020). https://doi.org/10.1080/00031305.2020.1731599

Stoltz, D. & Williams, R.   (London, 2020).

Zhang, Z. *et al.* Estimate risk difference and number needed to treat in survival analysis. *Ann Transl Med* **6**, 120 (2018). https://doi.org/10.21037/atm.2018.01.36

Marsaglia, G., Tsang, W. W. & Wang, J. Evaluating Kolmogorov's Distribution. *Journal of Statistical Software* **8**, 1 - 4 (2003). https://doi.org/10.18637/jss.v008.i18

Pollyea, D. A. *et al.* Venetoclax and azacitidine followed by allogeneic transplant results in excellent outcomes and may improve outcomes versus maintenance therapy among newly diagnosed AML patients older than 60. *Bone Marrow Transplant* **57**, 160-166 (2022). https://doi.org/10.1038/s41409-021-01476-7

Seaman, S. R. & White, I. R. Review of inverse probability weighting for dealing with missing data. *Stat Methods Med Res* **22**, 278-295 (2013). https://doi.org/10.1177/0962280210395740

van Buuren, S. & Groothuis-Oudshoorn, K. mice: Multivariate Imputation by Chained Equations in R. *Journal of Statistical Software* **45**, 1 - 67 (2011). https://doi.org/10.18637/jss.v045.i03

Rubin, D. *Multiple imputation for nonresponse in surveys*. (John Wiley & Sons, Inc., 1987).

Mustafa Ali, M. K. *et al.* Outcomes of Newly Diagnosed Acute Myeloid Leukemia Patients Treated With Hypomethylating Agents With or Without Venetoclax: A Propensity Score-Adjusted Cohort Study. *Front Oncol* **12**, 858202 (2022). https://doi.org/10.3389/fonc.2022.858202

Gangat, N. *et al.* Venetoclax and hypomethylating agent combination therapy in newly diagnosed acute myeloid leukemia: Genotype signatures for response and survival among 301 consecutive patients. *Am J Hematol* **99**, 193-202 (2024). https://doi.org/10.1002/ajh.27138

Othman, J. *et al.* Real-world outcomes of newly diagnosed AML treated with venetoclax and azacitidine or low-dose cytarabine in the UK NHS. *Blood Neoplasia* **1**, 100017 (2024). https://doi.org/https://doi.org/10.1016/j.bneo.2024.100017

Islam, N. *et al.* A simplified risk model for pretreatment stratification of newly diagnosed acute myeloid leukemia patients treated with venetoclax and azacitidine. *medRxiv*, 2024.2012.2002.24318344 (2024). https://doi.org/10.1101/2024.12.02.24318344

TABLE LEGEND

Table 1. Model specifications for risk stratification models with respect to censoring assumptions, methodologies to deal with missing data, end points, variable selection rules, and feature sets. Primary models were denoted by (M1) – (M16) and reported results were based on (M2) and (M8) (highlighted in red).

Table 1. Specifications of risk stratification models.

		Missing data adjustment	Censoring	Survival
	(M1)	Category	FAS	OS
	(M2)	Category	allo-HCT as censored	OS
	(M3)	CC with IPW	FAS	OS
	(M4)	CC with IPW	allo-HCT as censored	OS
	(M5)	Imputation by MICE	FAS	OS
	(M6)	Imputation by MICE	allo-HCT as censored	OS
	(M7)	Category	FAS	EFS
	(M8)	Category	allo-HCT as censored	EFS
	(M9)	CC with IPW	FAS	EFS
	(M10)	CC with IPW	allo-HCT as censored	EFS
	(M11)	Imputation by MICE	FAS	EFS
	(M12)	Imputation by MICE	allo-HCT as censored	EFS
	(M13)	Ensemble of (M1, M3, M5)	FAS	OS
	(M14)	Ensemble of (M2, M4, M6)	allo-HCT as censored	OS
	(M15)	Ensemble of (M7, M9, M11)	FAS	EFS
	(M16)	Ensemble of (M8, M10, M12)	allo-HCT as censored	EFS

Covariate-level classification rules

[**Rule-I**] Use point-estimates to compute mean marginal risks for each variable at t

[**Rule-II**] Use point-estimates to compute mean marginal risks for each variable at t. We excluded variables that are sparsely (< 2% (+ve) cases) or overly (> 40% (+ve) cases) represented.

[**Rule-III**] Use bootstrap-based point-estimates averaged over 2000 runs to compute mean marginal risks for each variable at t.

[**Rule-IV**] Combining Rule-II and Rule-III

Type of markers

[**Type-1**] Genetic

[**Type-2**] Genetic + Phenotypic

Total risk stratification models (covariate and subject level)

16 X 4 X 2 = 128

Remarks:
CC – Complete case analysis
IPW – Inverse probability of weights
MICE – Multiple imputation by chained equations
FAS – Full analytical dataset
allo-HCT – allogeneic hematopoietic cell transplant

FIGURE LEGENDS

Figure 1. Summary of the de-novo AML ven/aza treated patient cohort. The flow diagram summarizes the patient populations in the CU dataset and how they were utilized in generating the risk stratification models for OS/BR and EFS.

Figure 2. Graphical representation of the machine-learning methodology used to develop risk models with respect to overall survival and event free survival. Details of each step 1-7 are described in the Methods section.

Figure 3. Internal validation of predictive performance for proposed risk stratification variables. Architecture of internal validation steps assessing predictive performance for proposed risk stratification variables.

Figure 4. Overall survival (OS) following treatment based on the genetic features-based risk model. The genetic features-based covariate-level classification Rule-II risk model ($RM_{G,II}$) is depicted. A) Definitions of the feature identification strategy and genetic features in $RM_{G,II}$; B) OS (left panel) and best response (BR) (right panel) based on $RM_{G,II}$ applied to the cohort excluding allo-HCT recipients; C) OS (left panel) and BR (right panel) based on $RM_{G,II}$ applied to the full analytical set (FAS); and D) $RM_{G,II}$ based on OS treating allo-HCT recipients as censored (left panel) and within the cohort comprised solely of allo-HCT recipients. Note BCOR, DNMT3A, DDX41, SF3B1, KRAS, KMT2A, PTPN11, SMC1A, JAK2 are mutations defined by NGS. IDH1, IDH2, NPM1 and p53m are defined by composite molecular tests as defined in Methods and Supplemental Table 1E for "composite mutation features". Chr 7 abn:loss, Chr 8 abn:loss, DEK:loss, MECOM:gain, RARA:loss are FISH features. The remainder including Del7q, T(9;11), Minus5, Minus17, Complex cytogenetics are karyotype features.

Figure 5. Event free survival (EFS) based on the genetic model using covariate-level classification. The covariate-level classification Rule-II Risk Model based on genetic features and specific to EFS ($RM_{G,II-EFS}$) is depicted**.** A) Feature identification strategy and definitions of the features of $RM_{GP,II-EFS}$; B) EFS based on $RM_{GP,II-EFS}$ applied to the full analytical set (FAS) (left panel) and set excluding (right panel) allo-HCT recipients; and C) EFS based on $RM_{GP,II-EFS}$ treating allo-HCT recipients as censored (left panel) and only within the cohort comprised of allo-HCT recipients. Note BCOR, KRAS, KMT2A, PTPN11, JAK2, were mutations defined by NGS. IDH1, IDH2, NPM1 and p53m are defined by composite molecular tests as defined in Methods and Supplemental Table 1E for "composite mutation features", Chr 7 abn:loss, Chr 8 abn:loss, RARA:loss, MECOM:rearrangement, KMT2A:rearrangement are FISH features. The remainder including Del 5q Del7q, T(9;11), Minus5, Minus17, Complex cytogenetics, Indeterminate cytogenetics and Normal cytogenetics are karyotype features.

Figure 6. Heatmap of risk stratification. Subject-specific risk stratification by various risk models (16 models) described in this report are summarized in the two panels and are denoted as Adverse risk (black), Intermediate risk (red), and Favorable risk (green).

Figure 1. Summary of the de-novo AML ven/aza treated patient cohort.

(A) (Uni)-Multivariate hazard and counterfactual risk-stratification model for overall survival (OS)
(B) (Uni)-Multivariate hazard and counterfactual risk-stratification model for event free survival (EFS)
(C) Validation analyses by transporting risk models on best responses.

i. **Overall survival** (OS): Time from ven/aza treatment initiation to all-cause mortality
ii. **Event free survival** (EFS): Time from ven/aza treatment initiation to the first documented "relapse from CR/CRi/MLFS", "no response", "partial remission", and all-cause mortality whichever occurs first
iii. **Best response** (BR): Defined as the best response recorded within the cycles containing the first treatment course of ven/aza.

Figure 2. Strategy and methodologies used to develop risk models with respect to overall and event-free survival.

i : Subscript defines a subject; $i = 1, \ldots, N$
j : Superscript defines type of OMICs data (i.e., FC, CYT, FISH, NGS, AML mutation)
b : Subscript denotes a bootstrap sample; $b = 1, \ldots, B$
$(c, t, \mathcal{X})$: Observed data where $y = (c, t)$ refers to survival data and $X \subseteq \mathcal{X}$: Set of features obtained from feature space $\mathcal{X}$
t_r : Grid of time-points of interest; $r = 1, \ldots, T$
M^j : j-th Cox-PH model with ridge (i.e., $L2$ norm) penalty
β^j : Parameter estimates
Λ_i : Set of tuning parameters corresponding to M^j-th model
$\hat{O}(M)^j_{it(+ve)}$: Marginal estimation of relative risks for i-th subject at t-th time based on M^j-th model for (+ve)
$\hat{\bar{O}}(M)^j_{t(+ve)}$: Mean marginal estimation of relative risks at t-th time based on M^j-th model for (+ve)
$\hat{\bar{\bar{O}}}(M)^j_{t(+ve)}$: Mean marginal estimation of relative risks at t-th time averaged over B bootstrap runs based on M^j-th model for (+ve)

Steps summary:

[S1] Select features (i.e., FC, CYT, FISH, NGS, AML mutation) based on univariate KMs with respect to OS, EFS, MRD, and RFS under three censoring mechanisms (for allo-HCT) and get $X = \{X_{FC}, X_{CYT}, X_{FISH}, X_{RD}, X_{MUT}\}$ from $\mathcal{X}$.

[S2] Adjust for missing data via three methods. IPWs were estimated based on GLM for complete case analyses. Missing elements were imputed five times via MICE exploiting X under "*impute then-select*" approach for each biomarker.

[S3] Fit mCOXr for each type of biomarkers separately. Corresponding tuning parameters are selected via cross-validation.

[S4] Mean marginal probabilities of risk over time were obtained utilizing "*counterfactual*" arguments for each subject hypothetically had each biomarker been (+ve) and (-ve).

[S5] Obtain bootstrap-based 95% CI for estimates, marginal risks, and risk differences.

[S6] Calculate risk differences $((1/N) \sum_i \{\hat{O}(M^1)^{FC}_{it_{14.7}(+ve)} - \hat{O}(M^1)^{FC}_{it_{14.7}(-ve)}\}$ were calculated at t_r. Next obtain covariate-level stratification using 5% risk difference rules.

[S7] Apply logic to determine subject specific risk stratification.

Figure 3. Internal validation of predictive performance of risk models.

i : Subscript defines a subject; $i = 1, \ldots, N$

j : Superscript defines type of OMICs data (i.e., FC, CYT, FISH, NGS, AML mutation)

b : Subscript denotes a bootstrap sample; $b = 1, \ldots, B$

$(c, t, \mathcal{X})$: Observed data where $y = (c, t)$ refers to survival data and $X \subseteq \mathcal{X}$: Set of features obtained from feature space $\mathcal{X}$

v : Subscript denotes a cross-validation sample; $v = 1, \ldots, V$

t_r : Grid of time-points of interest; $r = 1, \ldots, T$

$(y^{(v)}, X^{(v)})$: Observed data leaving v-th CV fold out

M^j: j-th Cox-PH model with ridge (i.e., $L2$ norm) penalty

β^j: Parameter estimates

Λ_j : Set of tuning parameters corresponding to M^j-th model

$\hat{O}(M)^j_{it_r(+ve)}$: Marginal estimation of relative risks for i-th subject at t-th time based on M^j-th model with respect to (+ve)

$\hat{\bar{O}}(M)^j_{t(+ve)}$: Mean marginal estimation of relative risks at t-th time based on M^j-th model with respect to (+ve)

$\hat{\bar{\bar{O}}}(M)^j_{t(+ve)}$: Mean marginal estimation of relative risks at t-th time averaged over B bootstrap based on M^j-th model with respect to (+ve)

SR_i : Subject-level risk stratification for i-th subject

Steps summary:

[S1-S7] Steps 1-7 are as same as before. Note 15-fold CV is performed by splitting datasets into mutually exclusive training and validation sets [S2.0].

[S8] Step 8 evaluates predictive performance over fifteen CV runs by different covariate-level classification approaches.

Figure 4. Overall survival (OS) based on the genetic features specific risk model using Rule-II (RM$_{G,II}$).

    A. Feature identification strategy and RM$_{G,II}$ features.

Covariate-level risk classification

RD$_{(+ve)}$ − RD$_{(-ve)}$	%
Strongly adverse	> 10
Moderately adverse	(5, 10]
Slightly adverse	(0, 5]
Slightly favorable	(-5, 0]
Moderately favorable	(-10, -5]
Strongly favorable	≤ -10

Subject-level risk stratification

Groups	Definitions
Adverse	Having at-least one 'Strongly adverse' and/or 'Moderately adverse' risk factor in the absence of any 'Strongly favorable' and/or 'Moderately favorable' factor
Favorable	Having at-least one 'Strongly favorable' and/or 'Moderately favorable' factor in the absence of any 'Strongly adverse' and 'Moderately adverse' risk factor.
Intermediate	Otherwise

Category	Genetic features	Frequency†	Frequency¶
Favorable	Any of ***BCOR, DNMT3A, DDX41, SF3B1, IDH1, IDH2, NPM1*** (in the absence of any Adverse features)	**71/224 (32%)**	**101/316 (32%)**
Intermediate	No Favorable or Adverse features, or ≥1 Favorable features with ≥1 Adverse features	**70/224 (31%)**	**106/316 (34%)**
Adverse	Any of ***KRAS, KMT2A, PTPN11, SMC1A, JAK2, p53m, Del7q, t(9;11), Minus5, Minus17, Complex cytogenetics, Chr 7 abn:loss, Chr 8 abn:loss, DEK:loss, MECOM:gain, RARA:loss*** (in the absence of any Favorable features)	**83/224 (37%)**	**109/316 (34%)**

† Counts (proportions) of patients after excluding allo-HCT
¶ Counts (proportions) of patients without excluding allo-HCT

Figure 4, cont'd

B. RM$_{G,II}$ model for OS (left) and best response (right), excluding allo-HCT recipients.

C. RM$_{G,II}$ model for OS (left) and best response (right), with respect to the FAS.

Figure 4, cont'd

D. RM$_{G,II}$ on OS censoring allo-HCT recipients (left) and only within allo-HCT cohort. (right).

Figure 5. Event free survival (EFS) based on the genetic features specific risk model using Rule-II (RM$_{G,II-EFS}$).

A. Feature identification strategy and RM$_{G,II-EFS}$ features.

Covariate-level risk classification

RD$_{(+ve)}$ − RD$_{(-ve)}$	%
Strongly adverse	> 10
Moderately adverse	(5, 10]
Slightly adverse	(0, 5]
Slightly favorable	(-5, 0]
Moderately favorable	(-10, -5]
Strongly favorable	≤ -10

Subject-level risk stratification

Groups	Definitions
Adverse	Having at-least one 'Strongly adverse' and/or 'Moderately adverse' risk factor in the absence of any 'Strongly favorable' and/or 'Moderately favorable' factor
Favorable	Having at-least one 'Strongly favorable' and/or 'Moderately favorable' factor in the absence of any 'Strongly adverse' and 'Moderately adverse' risk factor.
Intermediate	Otherwise

Category	Genetic features	Frequency[†]	Frequency[¶]
Favorable	Any of **BCOR, Normal cytogenetics, Monosomal karyotype, Del5q, IDH1, IDH2, NPM1** (in the absence of any Adverse features)	69/211 (33%)	100/303 (33%)
Intermediate	No Favorable or Adverse features, or ≥1 Favorable features with ≥1 Adverse features	85/211 (40%)	126/303 (42%)
Adverse	Any of **KRAS, KMT2A, PTPN11, JAK2, p53m, Del7q, t(9;11), Minus5, Minus17, Complex cytogenetics, Indeterminant cytogenetics, Chr 7 abn:loss, Chr 8 abn:loss, RARA:loss MECOM:rearrangement, KMT2A:rearrangement** (in the absence of any Favorable features)	57/211 (27%)	77/303 (25%)

[†] Counts (proportions) of patients after excluding allo-HCT
[¶] Counts (proportions) of patients without excluding allo-HCT

Figure 5, cont'd

B. RM$_{G,II\text{-}EFS}$ on EFS, with respect to the FAS (left) and dataset excluding allo-HCT patients (right).

C. RM$_{G,II\text{-}EFS}$ on EFS censoring allo-HCT (left) and only within allo-HCT recipients (right).

Figure 6. Subject-specific risk stratification of AML patients by risk models.