Reduced brain structural similarity is associated with maturation, neurobiological features, and clinical status in schizophrenia

Natalia García-San-Martín[1,2], Richard AI Bethlehem[2], Patricia Segura[1], Agoston Mihalik[3], Jakob Seidlitz[4,5,6], Isaac Sebenius[3,7], Claudio Alemán-Morillo[1], Lena Dorfschmidt[4,5,6], Golia Shafiei[5,6], Sarah E. Morgan[3,7,8], Miguel Ruiz-Veguilla[9,10,11], Rosa Ayesa-Arriola[9,12], Javier Vázquez-Bourgon[9,12], Bratislav Misic[13], John Suckling[3,14], Benedicto Crespo-Facorro[9,10,11,12], Rafael Romero-García[1,3,9,11*]

Author affiliations:

[1]Department of Medical Physiology and Biophysics, Faculty of Medicine, University of Seville; Seville, Spain.

[2]Department of Psychology, University of Cambridge; Cambridge, UK.

[3]Department of Psychiatry, University of Cambridge; Cambridge, UK.

[4]Department of Child and Adolescent Psychiatry and Behavioral Science, The Children's Hospital of Philadelphia; Philadelphia, PA, USA.

[5]Lifespan Brain Institute, The Children's Hospital of Philadelphia and Penn Medicine; Philadelphia, PA, USA.

[6]Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania; Philadelphia, Pennsylvania, USA.

[7]Department of Computer Science and Technology, University of Cambridge; Cambridge, UK.

[8]School of Biomedical Engineering and Imaging Sciences, King's College London; London, UK.

[9]Biomedical Research Center in Mental Health Network (CIBERSAM), Health Institute Carlos III; Madrid, Spain.

[10]Mental Health Service, Virgen del Rocío University Hospital; Seville, Spain.

[11]Instituto de Biomedicina de Sevilla (IBiS) HUVR/CSIC, University of Seville; Seville, Spain.

[12]Department of Psychiatry, Marqués de Valdecilla University Hospital, IDIVAL, School of Medicine, University of Cantabria; Santander, Spain.

[13]Montreal Neurological Institute, McGill University; Montreal, QC, Canada.

[14]Peterborough NHS Foundation Trust; Peterborough, UK.

Correspondence to: Rafael Romero García

Full address: Avda. Doctor Fedriani S/N, 419009, Seville (Spain)

E-mail: rr480@cam.ac.uk

Abstract

Schizophrenia spectrum disorders (SSD) are characterized by atypical brain maturation, including alterations in structural similarity between regions. Using structural MRI data from 195 healthy controls (HC) and 352 individuals with SSD, we construct individual Morphometric INverse Divergence (MIND) networks. Compared to HC, individuals with SSD mainly exhibit reduced structural similarity in the temporal, cingulate, and insular lobes, being more pronounced in individuals exhibiting a 'poor' clinical status

(more impaired cognitive functioning and more severe symptomatology). These alterations are associated with cortical hierarchy and maturational events, locating MIND reductions in higher-order association areas that mature later. Finally, we map 46 neurobiological features onto MIND networks, revealing a high presence of neurotransmitters and astrocytes, along with decreased metabolism and microstructure, in regions with reduced similarity in SSD. These findings provide evidence on the complex interplay between structural similarity, maturational events, and the underlying neurobiology in determining clinical status of individuals with SSD.

Running title: MIND and maturation in schizophrenia

Keywords: structural connectivity, volumetric centiles, disease epicenters, cognition, positive and negative symptoms, regional vulnerability

1 Introduction

Multiple MRI studies have detected brain structural alterations across several cortical

areas in individuals with schizophrenia spectrum disorders (SSD), including reductions

in volume, surface, thickness, and gyrification[1–5]. The widespread, multicentric pattern

of these structural alterations likely reflects dysconnectivity within brain specific

networks[6]. The structural connectome, traditionally derived from diffusion-weighted

imaging (DWI), has provided valuable insights into connectivity alterations associated

with SSD[7,8]. For example, a more efficiently wired connectome has demonstrated a

better response to antipsychotic treatment[8]. However, this tractography technique

generates a substantial amount of false-positive fiber bundles[9]. Alternatively, structural

networks can be constructed using within-subject cortical similarity approaches, which

have shown a correspondence with tract-tracing similar to that observed in previous

reports on DWI-based networks in macaques[10,11]. Additionally, similarity networks

have demonstrated potential for clinical translation due to their reduced acquisition

times, as networks derived from a single, routinely acquired T_1-weighted MRI scan

have been shown to correspond to those generated from a broader set of multimodal

MRI features[12]. These networks have also demonstrated to be sensitive to the duration

of psychosis[13], the expression of schizophrenia-related genes[6], and the prediction of

treatment response[14], providing novel insights into the role of brain structure in

mediating genetic and neuronal factors that underlie clinical outcomes and the

progression of the disease. Moreover, a recent method based on Morphometric Inverse

Divergence (MIND) has emerged, providing an estimate of within-subject similarity

between cortical areas based on multiple MRI features[15]. These novel similarity

networks have demonstrated increased inter-individual consistency, stronger inter-

hemispheric and within-class connections, and increased coupling with gene co-

expression compared to DWI tractography[15]. Furthermore, MIND networks have also

shown greater reliability, better consistency with cortical cytoarchitectonics and

symmetry, and stronger correlations with tract-tracing measures of axonal connectivity

compared to previous approaches[15]. Exploring MIND networks among individuals with

SSD may shed light on the complex structural alterations that manifest as diverse

cognitive and symptomatologic status.

Extensive research has linked volumetric reductions and structural network alterations

to severe symptomatology in first-episode psychosis (FEP) and chronic

schizophrenia[5,16,17]. Investigations on drug-naïve patients are scarcer, but they have

reported decreases in cortical thickness and structural connectivity associated with

cognitive and functioning deficits[18–20]. In this context, baseline reductions of structural

connectivity have successfully predicted worse functioning outcomes in antipsychotic-

naïve FEP individuals, which could not be attributed to antipsychotic medication[19].

Additionally, reduced overall DWI connectivity and longer path length have also

predicted lower general cognitive ability in individuals with schizophrenia[21]. Positive

symptoms have also been associated with a widespread pattern of decreased gray matter

volume in bilateral frontotemporal and occipital regions in FEP individuals[5]. On the

other hand, structural alterations associated with negative symptomatology have shown

inconsistencies[22]. The controversial impact of antipsychotic drugs on symptomatology

and brain architecture[23–25] highlights the relevance of research on drug-naïve or

minimally medicated patients, which could help elucidate the specific contribution of

disease-related alterations to brain structure.

A growing body of literature supports the hypothesis that psychiatric and neurological

conditions initially manifest as abnormal brain structures in specific regions, referred to

as epicenters[26,27]. These abnormalities subsequently spread to other areas through the

underlying network architecture, involving genetic, trans-neuronal, structural, and

functional processes[16,28–30]. These mechanisms may disrupt the transport of trophic

factors and facilitate the spread of misfolded proteins or inflammatory markers linked to

cell death and atrophy, contributing to the structural and functional progression of

multiple neurodegenerative diseases[31–33]. Moreover, epicenters have been found to

correlate with more severe symptom profiles, suggesting that network organization may

guide cortical alterations throughout the disease process of schizophrenia and may be

influenced by cognitive impairment and symptom severity, among other factors[16].

However, the pathological spread of these abnormalities from disease epicenters to

connected brain regions, resulting in network-wide morphological changes, remains

unclear. Investigating the relationship between network architecture and atypical

development in SSD may be key to understanding abnormal brain maturation and

functioning[34].

A key aspect of human brain maturation is the consolidation of cortical hierarchy, where

higher-order association areas exhibit faster rates of shrinkage and myelination

compared to low-order sensorimotor areas during the course of adolescence, facilitating

efficient connectivity throughout the structural network[35]. Population models have

become a fundamental tool for accurately delineating these brain maturational events,

including determining the ages at which regions reach their peak volume and maximum

growth velocity[36]. These approaches based on normative data also enable benchmarking

regional brain volumes to identify patterns of atypical neuroanatomical maturation after

accounting for age-, sex-, and site-effects[36]. This allows for constructing population-

relative 'centiles', a measure of brain structure atypicality across the lifespan. A recent

study based on this methodology demonstrated common abnormal trajectories of

cortical volume across different psychosis-related conditions, supporting the notion that

regions with similar structural profiles are co-vulnerable to psychosis (i.e., exhibit

similar psychosis-related centile alterations)[37]. Although advanced structural brain

ageing[38] and alterations in higher-order areas[39,40] have been implicated in schizophrenia,

no study has yet explored the complex interaction between maturational events, brain

hierarchy, and structural connectivity derived from similarity networks in SSD.

Understanding how structural alterations in psychiatric conditions relate to the

underlying neurobiological substrates has been particularly challenging due to limited

human data[41,42]. However, recent studies have explored the spatial distribution of

several neurobiological properties, including molecular and cellular markers[43,44]. One

study revealed that, while psychotic-related conditions exhibit distinct patterns of gray

matter reductions, there is a major overlap in the neurobiological features characterizing

atypically maturing regions, including a higher density of serotonin receptors, among

others[37]. No study to date has explored the biological profiles associated with structural

alterations in network connectivity and their potential implication for SSD clinical

status[6].

Here, we exploit a method based on MIND networks to assess how structural markers

are associated with abnormal development in SSD and with their clinical status. We

constructed MIND networks by estimating within-subject similarities between cortical

areas in both SSD and healthy control (HC) subjects. Firstly, we compared the

differences in MIND similarity (also referred here as connectivity) between SSD and

HC, and then after stratifying individuals with SSD into 'poor' or 'good' clinical status.

Secondly, we examined whether any decrease in MIND connectivity of patients with

SSD was linked to centile reductions or other maturational events. For this purpose, we

first computed the regional connectivity to disease epicenters of these individuals to

assess how clinical status was influenced by the regions preferentially linked to

atypically maturing areas. Next, we explored potential associations between MIND

networks and the hierarchical organization of the cortex. Finally, we conducted a

multivariate analysis to examine the associations between regional neurobiological

features and MIND alterations in relation to clinical status.

Results

Decreased MIND connectivity in schizophrenia is associated with clinical status

MIND networks were computed as the within-subject similarity between cortical areas

based on the Kullback-Leibler divergence of multiple MRI features (see *Methods* for

details). To determine how MIND varies between SSD and HC, we computed the effect

size (Cohen's d) between networks (for edges and nodes separately). Compared to HC,

individuals with SSD exhibited significant reductions in MIND connectivity in edges

targeting the temporal, cingulate, and insular cortices ($P_{\text{perm}} < 0.05$), while edges

interconnecting the occipital cortices showed higher connectivity (Fig. 1A, left and

right; see Source Data for the full matrix). Frontal and parietal connecting regions

exhibited both increases and decreases in MIND connectivity. At the regional level,

patients with SSD exhibited a significant reduction in MIND degree in the insular,

cingulate, and temporal cortices, along with an increase in the occipital lobe of the right

hemisphere (Fig. 1A middle and right). The sensitivity analyses revealed that MIND

degree remained highly consistent across both scanning sites and sexes (see

Supplementary Fig. 1), and was robust to inter-individual variations, image quality

(Euler index), and parcellation schemes (Supplementary Fig. 2A-C). Effect sizes were

also significantly correlated with those reported by Morgan *et al.*[6] in Morphometric

Similarity (MS) networks derived from individuals with chronic schizophrenia ($r_{partial}$ =

0.35, P_{spin} = 0.004; Supplementary Fig. 2D).

Effect sizes of MIND edges and degrees were analyzed based on the general

psychopathology and individual scores of global cognitive functioning and clinical

symptoms (BPRS, SAPS, and SANS scales) of patients with SSD, which were

classified with a 'poor' or 'good' status (see *Methods* and Supplementary Fig. 3 for

details on the score distributions across each status; see Supplementary Fig. 4 for

regional correlations between MIND degree and clinical status). Instead of reporting

individual scale scores for cognition, we used an average score for simplicity. This

decision was supported by the Principal Component Analysis (PCA) performed on these

scales, where the first principal component explained 58.65% of the variance and

showed highly similar weights across scales (ranging from 0.27 to 0.44). Compared to

HC, alterations in MIND were observed only in patients with more impaired general

psychopathology ('poor' general psychopathology; Fig. 1B). Consistent with these

findings, reductions in MIND degrees (cingulate, fusiform, and temporal regions) were

observed only in patients with more impaired cognitive functioning ('poor' cognition).

Accordingly, patients with more severe symptomatology ('poor' clinical status)

exhibited a significant reduction in MIND degree in the left rostral anterior cingulate

region across all three scales (Cohen's d < -0.32), but not in individuals with less severe

clinical symptomatology ('good' clinical status), except for SANS. Additionally,

individuals with higher BPRS total scores showed several significant reductions in

MIND edges connecting frontal, temporal, and cingulate regions (Cohen's d < -0.28),

along with few increases in edges connecting frontal and occipital regions (Cohen's d >

0.28). However, the lowest score individuals did not show any significant difference

compared to HC. While individuals with higher SAPS scores presented the temporal

and cingulate edges significantly reduced (Cohen's d < -0.27), as well as the frontal and

occipital edges increased (Cohen's d > 0.27), individuals with lower SAPS scores

barely presented differences compared to HC. Conversely, individuals with lower

SANS scores presented a greater number of significantly reduced regions (temporal,

insular, and cingulate cortices) compared to individuals with higher scores. Similar

results were obtained when Desikan-Killiany-318 atlas parcellation was employed

(Supplementary Fig. 5). The number of significant effect sizes shared across scales was

higher for individuals with 'poor' status (Fig. 1C; see Supplementary Fig. 6 for the

distinct effect sizes of MIND edges and degrees of each scale for a 'poor' and 'good'

status).

Additionally, the association between MIND degree and clinical status was also

explored without stratifying individuals into 'poor' or 'good' status, supporting the

finding that regions with the greatest reductions in MIND degree in SSD were those

showing the strongest correlations with cognition (Supplementary Fig. 7).

163
Figure 1 MIND connectivity in relation to clinical status. (**A**) Effect sizes (Cohen's

d) computed between SSD and HC MIND edges (left) and degrees (middle), where blue

indicates a significant reduction of edge and degree in SSD, yellow and red indicate a

significant increase, and black indicates no significant differences. Spatial 3D

representation of significantly different edges and degrees in SSD compared to HC

(right). Supplemental data are provided as a Source Data file. (**B**) Effect sizes of MIND

edges and degrees stratified by general psychopathology, cognition, and

symptomatology. The Cohen's d of MIND degree for those individuals with SSD whose

general psychopathology is more impaired, whose cognition is lower, and whose

symptomatology (BPRS, SAPS, and SANS) is equal or higher than the median of the

sample (top row). The Cohen's d for those whose general psychopathology is less

impaired, whose symptomatology is lower, and whose cognition is equal or higher than

the median of the sample is shown in the bottom row. The highlighted regions show

those regional degrees that exhibit significant differences from HC after tail area-based

false discovery rate (FDR) correction (permutation test, $P < 0.05$). (**C**) Number of

significant effect sizes shared across scales for a 'poor' and 'good' status. BPRS, Brief

Psychiatric Rating Scale; HC, healthy control; MIND, Morphometric INverse

Divergence; SANS, Scale for the Assessment of Negative Symptoms; SAPS, Scale for

the Assessment of Positive Symptoms; SSD, schizophrenia spectrum disorder.

183   **Connectivity to disease epicenters is associated with clinical**

184   **status**

In addition to the MIND network analyses, we identified the regions with the most

pronounced morphological alterations at SSD onset (centiles). Regional centiles were

computed for each participant by benchmarking regional MRI volumes against

previously established normative trajectories[36]. These centiles were highly consistent

across diagnostic groups (Supplementary Fig. 8). The SSD group exhibited a

generalized significant centile reduction in almost all regions (Fig. 2A left), which were

negatively correlated with HC degrees (r = -0.45; P_{spin} = 0.001). However, this

association was no longer significant after accounting for regional volumes (Fig. 2A

middle; $r_{partial}$ = -0.20; P_{spin} = 0.122). To assess the relationship between MIND edges

and the inter-regional susceptibility to psychosis, we computed the structural co-

vulnerability matrix by correlating the regional effect sizes of the centiles for different

psychosis-related groups (see *Methods* for details). Results revealed a significant

association between MIND connectivity and inter-regional co-vulnerability, indicating

that pairs of regions that were more structurally similar exhibited higher co-

vulnerability to psychosis (Fig. 2A right; r = 0.33; $P_{Betzel} < 10^{-4}$; P_{Moran} = 0.001; see

Subjects and Methods for Betzel rewiring null model and Moran spectral

randomization). Building on this edge-level association between MIND and centile-

based alterations, we further analyzed the connectivity to disease epicenters.

Disease epicenters were defined as the set of regions with the most reduced centiles at

SSD onset (see Supplementary Fig. 9 for correspondence with gray matter thickness

abnormalities in chronic schizophrenia derived from ENIGMA[45]). According to Fig. 2A

middle, these epicenters were significantly anti-correlated with the functional degree

derived from an ENIGMA study in schizophrenia[16] (r = -0.54, P_{spin} = $2 \cdot 10^{-4}$, and r =

0.03, P_{spin} = 0.82, for functional and DWI connectivity, respectively; see Supplementary

Fig. 10). To assess the structural similarity to these atypically maturating regions in

SSD, the connectivity to epicenters was calculated for each region as the Pearson

correlation between its edge (SSD or HC) and the corresponding SSD centiles (see

Subjects and Methods for further details). Unlike HC epicenters, which did not reveal

any significant association, several regions showed a significant negative correlation

between edge strength and SSD centiles, indicating that strong edges are preferentially

connected to SSD epicenters (Fig. 2B; r_{global} = -0.27, computed as the average

correlation across regions; see Source Data for correlation values along with their

regional labels). After stratifying individuals by general psychopathology, individuals

with 'poor' status exhibited a slightly stronger tendency to be connected to disease

epicenters (r_{global} = -0.29) compared to a 'good' condition (r_{global} = -0.23; Fig. 2C).

Accordingly, a moderately higher association between MIND and (r_{global} = -0.33)

disease epicenters was found in patients with 'poor' cognition compared to individuals

with better cognitive functioning (r_{global} = -0.16; see Supplementary Fig. 11 for regional

correlations between disease epicenters and clinical status). Similarly, patients with

more severe BPRS, SAPS, and SANS symptomatology exhibited the same tendency to

be connected to epicenters (r_{global} = -0.28, r_{global} = -0.27, and r_{global} = -0.28, respectively)

compared to those with milder symptoms (r_{global} = -0.21, r_{global} = -0.22, and r_{global} = -

0.24).

228
Figure 2 Connectivity to disease epicenters. (**A**) Regional MRI brain volumes from

subjects with SSD converted into centiles, where blue and red indicate a significant

centile reduction and increase, respectively, in SSD compared to the expected regional

volume for age- and sex-matched normative individuals (left). Correlation between HC

degrees and SSD (middle; two-sided test). Association between HC edges and the

structural co-vulnerability to psychosis matrix (right; two-sided test). (**B**) Correlation

between the corresponding edges to each region and the mean HC (top) and SSD

centiles (bottom), where blue and red indicate a negative and positive correlation,

respectively. (**C**) Connectivity to SSD epicenters stratified by patients with 'poor' and

'good' general psychopathology, cognition, and symptomatology (BPRS, SAPS, and

SANS). The highlighted areas indicate regions where connectivity is significantly

correlated with epicenters after tail area-based FDR correction (P_{Betzel} < 0.05; see

Supplementary Fig. 12 for Moran significance). Supplemental data are provided as a

Source Data file. BPRS, Brief Psychiatric Rating Scale; HC, healthy control; MIND,

Morphometric INverse Divergence; SANS, Scale for the Assessment of Negative

Symptoms; SAPS, Scale for the Assessment of Positive Symptoms; SSD, schizophrenia

spectrum disorder.

Decreased MIND connectivity in schizophrenia is associated with cortical hierarchy and maturational events

We found no association between SSD centiles and the effect sizes of either MIND

degree (Fig. 3A top; $r_{partial}$ = -0.13; P_{spin} = 0.295) or edges (Supplementary Fig. 13; r = -

0.11; P_{Moran} = 0.43), revealing that atypical volumetric maturation and alterations in

MIND network do not co-localize in the same cortical regions. Specifically, after

correlating MIND degree with a sensorimotor-association regional ranking[46], results

showed that regions with reduced MIND degree in SSD were preferentially located in

higher-order association areas (Fig. 3A middle; $r_{partial}$ = -0.25; P_{spin} = 0.049), while SSD

centiles were reduced in lower-order sensorimotor regions (Fig. 3A bottom; $r_{partial}$ =

0.33; P_{spin} = 0.009). To determine whether MIND networks are related to maturational

events, we exploited brain maps indicating the age at which maximum peaks of volume

and velocity occur for each region[36]. Regions that reach maximum volume of

maturation at an early age (e.g., frontal pole, superior parietal, and cuneus) exhibited

stronger structural connectivity (hubs) in HC (Fig. 3B top; $r_{partial}$ = -0.41; P_{spin} = 0.022).

On the other hand, regions that mature later showed the strongest reduction in MIND

degree in SSD compared to HC (Fig. 3B middle; $r_{partial}$ = -0.53; P_{spin} = 0.002), but

higher centiles in SSD (Fig. 3B bottom; $r_{partial}$ = 0.63; P_{spin} = 0.001). Interestingly,

although the peak age of regional velocity did not exhibit a significant correlation with

HC degrees (Fig. 3C top; $r_{partial}$ = 0.15; P_{spin} = 0.418), a significant association was

found with the reduction in MIND degree of patients with SSD compared to HC (Fig.

3C middle; r_{partial} = -0.56; P_{spin} = 0.001), but not with SSD centiles (Fig. 3C bottom;

r_{partial} = 0.10; P_{spin} = 0.602). All associations were calculated after accounting for mean

regional volumes. Additionally, these findings remained highly consistent after

stratifying individuals into 'poor' or 'good' status (Supplementary Figs. 14-17).

Altogether, these findings reveal that reductions in regional structural similarity in

individuals with SSD were co-localized with maturational events and higher-order

association areas.

274
Figure 3 MIND connectivity in relation to sensorimotor-association hierarchy and

maturational events. (**A**) Correlations between effect sizes of degrees and SSD centiles

(top); effect sizes of degrees and cortical hierarchy (middle); SSD centiles and cortical

hierarchy (bottom). (**B**) Correlations between the cortical map of age at regional volume

peak (derived from normative trajectories) and HC degree (top); effect size of degree

(middle); and SSD centiles (bottom). (**C**) Correlations between the normative cortical

map of age at regional velocity peak and HC degree (top); effect size of degree

(middle); and SSD centiles (bottom). All regional correlations (two-sided test)

accounted for mean regional volumes. HC, healthy control; MIND, Morphometric

INverse Divergence; SSD, schizophrenia spectrum disorder.

Neurobiological features are co-localized with alterations in MIND connectivity and related to clinical status

Neurobiological features are co-localized with alterations in

MIND connectivity and related to clinical status

To investigate the neurobiological features potentially linked to MIND networks, we

exploited previously acquired datasets obtained from different modalities consisting of

molecular and micro-architectural data clustered under different neurobiological types

(neurotransmitter, cell type, layer thickness, microstructure, expansion, and metabolism;

see *Subjects and Methods* for additional information)[47]. HC edges significantly

correlated with the neurobiological similarity matrix, indicating that the more connected

a pair of regions is, the more neurobiologically similar they are (Fig. 4A top; $r = 0.47$;

$P_{\text{Betzel}} < 10^{-4}$; $P_{\text{Moran}} < 10^{-4}$). On the contrary, the effect size between SSD and HC edges

did not correlate with neurobiology (Fig. 4A bottom; $r = 0.11$; $P_{\text{Moran}} = 0.33$). However,

when this analysis was conducted separately for each neurobiological type, significant

associations were identified with both HC and effect sizes (Supplementary Fig. 18). At

the regional level, we performed a combined PCA and Canonical Correlation Analysis

(CCA) approach to investigate the regional associations of MIND degree with the

neurobiological features (see *Subjects and Methods* and Supplementary Fig. 19 for

details). The PCA-CCA models were significant (FDR-corrected) for both HC (Fig. 4B

top-left; $r = 0.67$; $P_{\text{spin}} = 0.001$) and effect sizes of degree (Fig. 4B bottom-left; $r = 0.71$;

$P_{spin} = 0.001$). The predicted HC degrees and effect sizes resembled their corresponding

empirical values (Fig. 4B middle). Both models exhibited significant loadings (Fig. 4B

right; $P_{spin} < 0.05$). Cell types (Neuro-Ex/In), cortical expansion (Scaling PNC and NIH,

developmental and evolutionary exp.), neurotransmitters (5-HT$_{2A}$, 5-HT$_4$), and

metabolism (CMRGlu, Glycolytic index, CMRO$_2$) demonstrated a significant positive

contribution to HC degrees, indicating that the presence of these neurobiological

features is highly co-localized with MIND hubs. The negative contribution of

neurotransmitters (5-HTT, DAT, VAChT, 5-HT$_{1A}$, NMDA, H$_3$) and cell types (OPC,

Micro, Astro) indicated a high presence of these features in less connected regions. The

features that most strongly accounted for explaining the regional reductions in MIND

degree between SSD and HC included neurotransmitters (D$_2$, 5-HT$_{1A}$, CB$_1$, MOR, 5-

HT$_4$, DAT, 5-HTT, D$_1$, mGluR$_5$, NMDA, 5-HT$_6$), cell types (OPC, Astro), and layer

thickness (Layer I-III), indicating a high presence in regions with reduced SSD

connectivity. Similar loadings were found when the cohort was randomly split into two

halves (see Supplementary Fig. 20). On the other hand, a high presence of

microstructural features (Gene PC1, Neurotransmitter PC1) and metabolism (CMRO$_2$,

CMRGlu, CBF) contributed to explaining the increased structural connectivity of

individuals with SSD compared to HC. The neurobiological loadings that contributed to

explaining the SSD centiles were shown in a previous study[37].

Finally, we performed two additional PCA-CCA analyses on the regional maps that

represented the positive association between cognition and both MIND degree and

centiles in SSD (Fig. 4C left, top and bottom, respectively). These maps indicated that

lower cognition scores were associated with lower regional degrees and centiles in SSD.

No significant associations were found between symptomatology and degree or centiles

(see Supplementary Figs. 4 and 11). The PCA-CCA models were significant for both

association maps of MIND (Fig. 4C top-middle; r = 0.52; P_{spin} = 0.001) and centiles

(Fig. 4C bottom-middle; r = 0.51; P_{spin} = 0.002), where the predicted maps resembled

their corresponding empirical values. Both models exhibited significant loadings (Fig.

4C right). Cortical expansion (Evolutionary and Developmental exp., Scaling PNC) and

neurotransmitters (MOR, CB_1, $mGluR_5$, 5-HT_4, 5-HT_{2A}, 5-HT_{1B}) demonstrated a

significant positive contribution to the MIND-cognition association, indicating that the

presence of these neurobiological features is highly co-localized with regions where

lower cognition scores are associated with lower MIND degree. The features that

positively accounted for explaining the regional associations between cognition and

SSD centiles were cortical expansion (Evolutionary exp., Scaling NIH, Scaling PNC),

neurotransmitters (5-HT_4, 5-HT_{2A}, CB_1, M_1, D_2), and layer thickness (Layer I-III, V,

and VI), indicating a high presence of these features in regions with lower centiles and

cognition scores. As a sensitivity test, we performed Partial Least Square (PLS)

analyses to assess whether the loading estimations were influenced by the PCA-CCA

method used here. A high correlation (r > 0.95) between loadings derived from both

methods was found for significant models (Supplementary Fig. 21). Additionally,

loadings remained consistent when we re-ran the PCA-CCA analyses with an explained

variance of 60% to prevent overfitting (Supplementary Fig. 22).

Figure 4 Neurobiological characterization of MIND networks. (**A**) Association

(two-sided test) between the neurobiological similarity matrix and HC edges (top) and

the effect sizes of MIND edges between SSD and HC (bottom), where yellow indicates

higher dots density than blue. (**B**) Correlation between empirical and predicted regional

HC degrees (top-left) and effect sizes of degrees (bottom-left); empirical and predicted

regional maps of HC degrees (top-middle) and effect sizes of degrees (bottom-middle);

and significant ($P_{spin} < 0.05$) PCA-CCA neurobiological loadings of HC degrees (top-

right) and effect sizes of degrees, where data ($n = 1,000$ bootstrapped samples) are

presented as mean values +/- standard deviation (bottom-right). (**C**) Regional

associations between cognition and both MIND degree and centiles in SSD, where red

indicates a positive correlation (left, top and bottom, respectively). Correlation between

empirical and predicted regional MIND-cognition (top-middle) and centile-cognition

(bottom-middle) associations in SSD, along with their empirical and predicted regional

maps, where blue and red indicate a negative and positive correlation, respectively.

Significant ($P_{spin} < 0.05$) PCA-CCA neurobiological loadings of MIND-cognition (top-

right) and centile-cognition (bottom-right) associations in SSD. 5-HT$_{1A}$, 5-HT$_{1B}$, 5-

HT$_{2A}$, 5-HT$_4$, 5-HT$_6$, serotonin receptors; 5-HTT, serotonin transporter; Astro,

astrocytes; CB$_1$, cannabinoid receptor; CBF, cerebral blood flow; CBV, cerebral blood

volume; CMRGlu, cerebral metabolic rate of glucose; CMRO$_2$, cerebral metabolic rate

of oxygen; D$_1$, D$_2$, dopamine receptors; DAT, dopamine transporter; Developmental

exp., developmental expansion; Evolutionary exp., evolutionary expansion; Gene PC1,

gene expression PC1; H$_3$, histamine receptor; HC, healthy control; M$_1$, acetylcholine

receptor; Micro, microglia; MIND, Morphometric INverse Divergence; MOR, opioid

receptor; Neuro-Ex, excitatory neurons; Neuro-In, inhibitory neurons; NMDA,

glutamate receptor; OPC, oligodendrocyte precursor cells; Scaling NIH, allometric

scaling from National Institutes of Health; Scaling PNC, allometric scaling from

Philadelphia Neurodevelopmental Cohort; SSD, schizophrenia spectrum disorder;

Thickness, cortical thickness; VAChT, acetylcholine transporter.

Discussion

In the present work, we investigated brain structure and organization by characterizing

Morphometric Inverse Divergence (MIND) networks in individuals with schizophrenia

spectrum disorders (SSD). We first assessed the differences between MIND networks of

individuals with different SSD clinical status and healthy controls (HC). We

additionally determined the regional connectivity to disease epicenters for each

individual to assess how clinical status was influenced by the regions preferentially

linked to atypical maturing areas. Finally, we explored how MIND organization in SSD

relates to maturational events and neurobiological features.

Extensive evidence has shown that schizophrenia impacts both brain structure and

connectivity[48]. In this context, recent morphometric studies have investigated individual

structural similarity patterns to assess how the disease is associated with the expression

of schizophrenia-related genes, leading to a widespread dysconnectivity pattern[6,49,50].

Here, we exploited the MIND method, identifying reductions in the cingulate, temporal,

and insular lobes in individuals with SSD, indicating greater dissimilarity between these

regions compared to HC. These findings align with previous research, providing new

evidence on the dysconnectivity hypothesis of psychosis[6,49,51,52]. This dysconnectivity

may result from the disintegration of nodes or hubs linked to structural changes in

association fibers and aberrant control of synaptic plasticity, ultimately leading to

abnormal functional connectivity in schizophrenia[51]. However, MIND increases were

also found in regions connecting the frontal, parietal, and occipital lobes, indicating

greater similarity between these regions, potentially reflecting increases in myelination

that facilitates the trans-synaptic propagation of pathology[53]. To our knowledge, only

one study has examined MIND alterations in individuals with early-onset schizophrenia,

reporting significant changes in degree, though these were limited to a reduced number

of regions[50]. However, studies in chronic schizophrenia have mainly reported increasing

structural covariance strength as the disease progresses[54], which might be attributed to

disease progression, treatment effects, or medication use.

Abnormal connectivity has been proposed as a fundamental factor underlying cognitive

deficits and clinical symptoms in patients with psychotic disorders[55]. In this study,

individuals with more impaired general psychopathology showed reductions in edges

connecting the temporal, insular, and cingulate lobes, while those with less impaired

general psychopathology showed no significant differences. Accordingly, individuals

with deeper cognitive impairment showed reductions in the temporal and cingulate

lobes, while those with more preserved cognition did not. This could reflect a cognitive

deficit associated to structural disconnections in psychosis-related individuals[56]. On the

other hand, individuals with higher symptomatology reflected common nodal degree

reductions in the rostral anterior cingulate region, which aligns with the notion that

symptoms are associated with a reduction in thalamocortical connectivity to the middle

frontal gyrus, anterior cingulate, and insula regions[55].

Extensive research supports the notion that psychiatric disorders originate from

alterations in specific brain regions, known as epicenters, which then propagate to other

areas through existing connectivity patterns[27,57], involving genetic, trans-neuronal,

structural and functional processes[16,28]. This model aligns with findings in neurological

conditions, where epicenter frameworks based on structural and functional connectivity

and network diffusion models have successfully predicted the progression of

neurodegenerative changes[58–60]. One proposed mechanism for network-specific

degeneration is that axonal connectivity (connectome) guides the spread of misfolded

proteins into new regions[58,61], leading to cell death and atrophy[28]. Alternatively,

dysfunction in one region may trigger abnormal activity in connected areas, which, over

time, could lead to structural changes as a result of disrupted neurotransmission or

diminished trophic support[7,59,60]. A further possibility is that genetic variants of genes

involved in schizophrenia, such as G72 and DISC1, may modulate regional changes

during SSD progression[62,63]. Here, although MIND degree was not associated with SSD

epicenters (centiles), we reported that MIND edges were stronger (i.e., higher structural

regional resemblance) between pair of regions exhibiting similar co-vulnerability to

psychosis[37]. This widespread distribution of regions showing significant connectivity to

disease epicenters likely reflects the globally distributed nature of schizophrenia, rather

than being indicative of focal regional abnormalities[64]. Given this broad network-level

disruption, even moderate alignment across widely distributed regions can yield a large

number of statistically significant associations. This finding cannot be attributed to a

specific permutation method, as both Moran spectral randomization, which accounts for

spatial autocorrelation[65], and a rewired null model that preserves the empirical degree

sequence and edge length distribution[66] produced similar results.

Maturation of higher-order cognitive and affective functions is disrupted in

schizophrenia[67]. Here, while centile decreases affected lower-order primary regions

associated with sensorimotor functions (as observed in antipsychotic-naïve first-episode

psychosis individuals)[68], connectivity reductions impacted higher-order association

areas linked to associative functions[46]. Accordingly, the hierarchical organization of the

cortex has also been demonstrated to be altered in different fMRI studies of

schizophrenia, affecting the secondary visual-to-sensorimotor axis rather than the

primary sensorimotor-to-association axis[40,69]. Taken together, these findings suggest

that SSD-related impairments target specific hierarchical processes, where focal

disruptions in somatosensory-motor areas may extend into higher-order regions[69].

Understanding the interaction between brain maturation and abnormal structural

connectivity is essential for gaining deeper insights into the neurodevelopmental origins

of psychiatric conditions[70,71]. Here, we found that regions maturing earlier tend to

exhibit higher MIND degree, aligning with studies indicating that rapidly maturing hubs

are located in high-level functional networks, potentially facilitating their functional and

anatomical maturation[72]. Additionally, the significant negative association between SSD

degree effect sizes and both volume and velocity age peaks indicated that late-maturing

regions are particularly affected by the disease. This aligns with fMRI connectivity and

DWI studies of schizophrenia[73,74], which suggest that dysmaturation of neural

connectivity is a key component of its pathophysiology[75]. We also found that local

atypical maturation, as indicated by centile analyses, is more prominent in early-maturing regions, consistent with the previously observed lower hierarchical role of these regions. Altogether, these findings suggest a complex interaction between local and network-level SSD alterations, cortical processing hierarchy, and maturational events.

Recent research has co-localized maturational events and structural disease-related alterations with the underlying neurobiology[26]. Particularly, abnormal patterns of MIND connectivity in major depressive disorder have been associated to the cannabinoid receptor and the brain-wide expression of a genetic combination[76]. In this study, we identified a common pattern of neurobiological features across connected regions, suggesting that the macroscopic structural properties used to estimate MIND edges are heavily influenced by underlying microstructural organization[77], neurotransmitter systems[43], and metabolism[78]. This finding aligns with the higher gene co-expression found in highly MIND-connected regions[15]. However, the differences between SSD and HC in MIND edges were not linked to overall shared neurobiology, but rather with specific neurobiological types that were either co-localized or anti-co-localized with differences in SSD edges. At the regional level, the positive contribution of excitatory and inhibitory neurons to MIND edges based on CCA loading estimates provided further evidence of the fundamental role that neuron density plays in shaping the architecture and connectivity of the primate cerebral cortex[79]. The high presence of OPC, microglia, and astrocytes in weakly-connected regions aligns with the role these cell types play in actively guiding brain development[80–83]. Furthermore, these glial cells also contributed to explaining the reduction in MIND degree of patients with SSD. This disruption in astrocyte differentiation may lead to abnormal glial coverage and hypomyelination, resulting in insufficient synaptic support, compromised white matter

integrity, and transmission desynchronization, ultimately causing network

dysconnectivity[84]. Neurotransmitters and layer thickness also contributed to these

alterations in MIND degree due to SSD. The hypothesis of anatomical disconnection in

schizophrenia has also been linked to synaptic damage, as well as dysfunction of

dopaminergic, cholinergic, or glutamatergic systems[85]. A principal component of

cortical gene expression architecture has been found to be specifically associated with

marker genes of layers II and III, and supragranular thinning in schizophrenia[86].

A low presence of myelin and cerebral blood volume was related to regions where

lower MIND degrees were associated with lower cognition scores, aligning with the

disrupted relationship found between myelination and neurocognition in patients with

schizophrenia[87]. Chronic low cerebral blood volume in schizophrenia may lead to neural

hypoactivation, ultimately disrupting cortical cytoarchitectural organization and

potentially contributing to brain atrophy and associated cognitive impairments[88].

However, a high presence of neurotransmitters and cortical expansion features were

associated with regions where both lower MIND degrees and centiles were associated

with lower cognition scores, which aligns with the previous studies relating the

imbalance in glutamate, acetylcholine, GABA, dopamine, and histamine with cognitive

deficits in patients with schizophrenia[89]. Thus, MIND allows for a more precise and

personalized biologically grounded representation of an individual's brain architectome,

potentially serving as a biomarker for diagnostic and stratification purposes, the

identification of biologically defined subtypes, and the development of more targeted

and effective therapeutic strategies.

The present findings must be interpreted with several considerations. First, MRI data

were obtained from different 1.5 and 3T MRI scanners. Nevertheless, site

harmonization was performed to MIND networks, after which MIND degree was highly

correlated between scanners (r = 0.96; Supplementary Fig. 1). Second, the current

implementation of MIND networks is limited to vertex-level data derived from MRI-

based cortical surface reconstructions[15]. As a result, subcortical structures were

excluded from this study, despite their well-established clinical relevance. Third, MIND

networks derived from cross-sectional data were associated with dynamic processes of

maturation. Analyzing longitudinal trajectories of MIND changes would offer a more

precise understanding of how maturational processes interact with disease progression

and treatment effects. Fourth, given the limited sample size scenario (68 regions), the

CCA multivariate approach has a risk of over-fitting[90]. To address this issue, we first

applied PCA prior to CCA. Fifth, the maturational events and neurobiological data used

here were derived from normative non-psychotic individuals, which limits the capacity

to capture patient-specific variability. A recent PET study, however, has demonstrated

that reduced synaptic density in schizophrenia is also spatially aligned with altered

normative distributions of neurotransmitter levels, suggesting that the synaptic deficits

are shaped by the molecular similarity between regions[91]. Nevertheless, relying on

external cellular and molecular datasets limits our ability to capture the biological

attributes underlying between-participant variability in brain structure. In this context, a

recent study that combined antemortem neuroimaging, molecular, and genetic data with

postmortem dendritic spine density and morphometric attributes demonstrated that

modules of synaptic protein and expression can explain individual differences in both

functional connectivity and structural covariation[92]. These findings highlight the

feasibility of cross-scale data integration to enhance our understanding of brain

connectivity. Future work with neurobiological data of individuals with SSD would

provide valuable insights into the molecular alterations causally associated with

abnormal brain alterations.

In summary, using MIND networks, we identified reductions in regional structural

similarity within the temporal, cingulate, and insular lobes of individuals with SSD,

alongside an increase in the occipital lobe. These decreases in MIND connectivity were

particularly pronounced in individuals with more severe clinical status. Moreover, these

MIND reductions were co-localized with maturational events and were more prominent

in regions characterized by a high normative presence of neurotransmitters, specific

cortical layer thickness, and astrocytes, as well as decreased microstructure and

metabolism. Altogether, these findings contribute to a better understanding of the

disease process, highlighting the role of brain structure, maturational events, and

neurobiology in shaping clinical status during the early stages of the disease.

Methods

Subjects

Magnetic Resonance Imaging (MRI) data from individuals who were determined to

have experienced a first-episode non-affective psychosis, confirming the presence of a

schizophrenia spectrum disorder (SSD; $n = 352$; $n_{female} = 138$; $age = 31.43 \pm 8.78$),

including schizophrenia ($n = 169$), schizophreniform ($n = 102$), brief psychotic ($n = 49$),

unspecified psychotic ($n = 26$), and schizoaffective ($n = 6$) disorders, along with a group

of healthy controls (HC; $n = 195$; $n_{female} = 75$; $age = 30.64 \pm 7.66$), were obtained from

Programa de Atención a las Fases Iniciales de Psicosis (PAFIP)[93]. Individuals with

SSD had no prior antipsychotic medication or were minimally medicated. All diagnoses

were made by an experienced psychiatrist using the Structured Clinical Interview for

DSM-IV (SCID-I), confirming the presence of schizophrenia or other psychotic

disorder after 6 months of the first visit. Participants who met criteria and provided

written informed consent were entered into PAFIP. Additional inclusion and exclusion

criteria can be found in *Supplementary Subjects and Methods*.

This program has been approved by the Ethics Committee for Clinical Research (CEIC

- Cantabria) in accordance with international standards (NCT0235832 and

NCT02534363 clinical trial numbers).

MRI acquisition, parcellation, and volume extraction

149 SSD and 80 HC MRI scans were obtained using a 1.5 T General Electric SIGNA

System (GE Medical Systems, Milwaukee, WI), and 203 SSD and 115 HC scans using

a 3 T Philips Medical Systems MRI scanner (Achieva, Best, The Netherlands) at the

Hospital Universitario Marqués de Valdecilla (HUMV). The parameters for 1.5 T were:

TE = 5 ms, TR =24 ms, NEX = 2, rotation angle = 45°, FOV = 26×19.5 cm, slice

thickness = 1.5 mm, and a matrix of 256×192; whereas for 3 T were: TE = 3.7 ms, TR =

8.2 ms, flip angle = 8°, acquisition matrix = 256×256, voxel size = 0.94×0.94×1 mm

and 160 contiguous slices. T_1-weighted images were first visually inspected for artefacts

and gross anatomical abnormalities, and were processed using FreeSurfer 6.0.0

(http://surfer.nmr.mgh.harvard.edu) applying the recon-all pipeline to enhance gray-

white matter boundary delineation.

Cortical parcellation was performed using Desikan-Killiany atlas, and volumetric

measurements were derived for each of the 68 region-of-interest. Quality control

procedures were implemented to ensure the accuracy and reliability of the cerebral

volumes derived. Thirteen HC subjects exhibiting visual artifacts or processing errors

were excluded from further analysis. No further participants were excluded, as their

global degree fell within the range of the average global degree ± 5 standard deviations

(0.166 ± 0.034).

MIND edges and degrees estimation

For each subject, a Morphometric INverse Divergence (MIND) network was constructed by computing the Kullback-Leibler (or Jeffrey's) divergence of multiple MRI features (cortical thickness, mean curvature, sulcal depth, surface area, and gray matter volume) between each pair of regions[15]. Thus, each cortical region was characterized by a multidimensional distribution of these features measured at sub-voxel vertex resolution on a cortical surface mesh. To calculate the multivariate divergence between pairs of regions directly from the observed vertex-level data, this method leverages a *k*-nearest neighbor approach, where a decrease in the divergence of these multivariate distributions (or an increase in their similarity) indicated an increase in MIND. This resulted in a 68×68 edge matrix (bounded between 0 and 1), representing the structural similarity between regions (also referred here as structural connectivity). To correct site-related batch effects while preserving age, sex, and psychosis-related effects including sex-psychosis interaction, we used ComBatLS[94], a recent extension of ComBat method that preserves biological variance, while reducing sex biases. To regress the effect of age, sex, and the estimated total intracranial volume (eTIV), we analyzed the residuals (rather than the raw MIND networks) derived from the linear model shown in Equation (1).

$$\text{MIND edge} \sim 1 + \text{sex} + \text{eTIV} + \text{age} + \text{sex} * \text{age} \qquad (1)$$

All non-binary variables were z-scored prior to regression. Individual MIND nodal degrees were computed by averaging their corresponding MIND residuals (referred here as edges), resulting in an average within-subject similarity degree for each region. Thus, while MIND edges represent how structurally similar each pair of regions is, MIND

degrees reflect the average structural similarity of a given region with the rest of the

brain.

MIND edges and degrees of patients with SSD were compared to HC by calculating the

effect size (Cohen's d) between each SSD and HC edge and region, respectively. To

statistically assess these differences, a permutation test was applied by randomly

shuffling group assignment to create a null distribution (10,000 permutations).

Additionally, tail area-based false discovery rate (FDR) correction[95] across brain

regions and edges was applied to account for multiple comparisons (with an alpha set at

< 0.05).

Assessment of cognition and symptomatology

Effect sizes of MIND edges and degrees were stratified based on the general

psychopathology, the average global cognitive function (i.e., cognition), and the clinical

symptomatology of drug-naïve first-episode psychosis (FEP) individuals. Overall

cognitive score was assessed for the following domains: (1) verbal memory: Rey

Auditory Verbal Learning Test (RAVLT) long-term recall score; (2) visual memory:

Rey Complex Figure test (RCFT) long-term recall score; (3) motor dexterity: grooved

pegboard (GP), time to complete with dominant hand; (4) executive functions: Trail

Making Test part B (TMTb); (5) working memory: WAIS III-Backward Digits (BD)

total score; (6) speed of processing: WAIS III-Digit Symbol (DS) standard total score;

(7) attention: Continuous Performance Test Degraded-Stimulus (CPT-DS), the total

number of correct responses[96]. Symptomatology was assessed with the Brief Psychiatric

Rating Scale (BPRS) (expanded version of 24 items), and the clinical symptoms of

psychosis were assessed with the Scale for the Assessment of Positive Symptoms

(SAPS) and the Scale for the Assessment of Negative Symptoms (SANS). General

psychopathology was assessed by performing a *k*-means clustering based on the

different scales considered (global cognitive function, BPRS, SAPS, and SANS),

stratifying individuals as having a 'good' or 'poor' general psychopathological status.

Individuals with SSD with a 'poor' status in cognition or symptomatology were defined

as those scoring below the median score for cognition and at or above the median for

symptomatology (cognition, $n = 151$; BPRS, $n = 165$; SAPS, $n = 168$; SANS, $n = 168$;

see Supplementary Fig. 3 for mean and median values across scales). Conversely,

individuals with a 'good' status were those who had scores at or above the median score

for cognition and below the median for symptomatology.

Computing connectivity to disease epicenters

Regional centiles were computed for each participant by benchmarking regional MRI

volumes against previously established normative trajectories (from approximately

120,000 participants), which adjusts for age-, sex-, and site-effects[36]. ComBatLS was

also applied to centiles to correct for site-effects. SSD centile distributions were first

compared to HC using the Wilcoxon rank-sum test (FDR-corrected) and subsequently

correlated with the average regional structural similarity (nodal degree) of HC

individuals to assess whether a stronger structural connectivity was associated with

atypically maturing regions in SSD. The structural co-vulnerability to psychosis

matrix (34×34 averaged across hemispheres), derived in our recent study[37], was

constructed by correlating the regional effect sizes of the centiles across different

psychosis-related groups, including first-degree relatives of patients with schizophrenia

or schizoaffective disorder, individuals who had psychotic experiences, individuals

from this cohort (SSD), and patients with chronic schizophrenia or schizoaffective

disorder (see Supplementary Table 1 for demographic details of these cohorts).

Regions with the most pronounced morphological alterations at disease onset (i.e., most

reduced centiles in SSD) were defined as disease epicenters[16,28]. For each individual

with SSD, the connectivity to disease epicenters for each region was computed as the

Pearson correlation r between the SSD edge of that region and the average regional

centiles in SSD. Consequently, regions that were preferentially associated with reduced

centiles were considered connected to disease epicenters. For each region, the P-value

(tail area-based FDR-corrected) was computed as the proportion of permuted r-values,

averaged across individuals, that were more extreme than the observed average values.

The null distributions of r-values were derived from a rewiring null model (Betzel) that

preserves the exact degree sequence and edge weight distribution, while also

approximates its edge length distribution (based on geodesic distance between regions)

and the relationship between edge weights and lengths[66]. Moran spectral randomization,

a method for generating spatially constrained null models that accounts for spatial

dependencies[65], was additionally employed.

As a sensitivity analysis, connections to epicenters were computed for HC subjects

correlating the HC edges of each region with the average regional HC centiles under the

assumption that no preferential connectivity with 'healthy' epicenters would be

revealed.

MIND association with cortical maturation

Centiles and effect sizes were associated with a sensorimotor-association regional

ranking reported by Sydnor *et al.*[46], which ranges from lower-order sensorimotor to

higher-order associative functions. Two additional maturational features derived from

the same normative trajectories as centiles were considered: volume peak, representing

the age at which regional volume reaches its highest point; and velocity peak, indicating

the age at which regional volume growth is the fastest. These features were compared to

MIND networks (both HC and effect sizes) and SSD centiles. All regional-based

associations, computed after accounting for regional volumes, were statistically

assessed by applying a spin test ($n = 10,000$) where spatial autocorrelation was

preserved (see *Supplementary Subjects and Methods* for details). The Betzel rewiring

null model preserving the degree sequence and approximately the edge length

distribution, as well as Moran spectral randomization, were employed on the MIND

edge-based associations to account for basic network properties and spatial

dependencies, respectively.

685   Identifying neurobiological features associated with MIND

686   networks stratified by clinical status

The spatial maps of 46 molecular and micro-architectural features, including

neurotransmitters, cell types, layer thickness, microstructure, cortical expansion, and

metabolism, were collected across multiple studies (see Supplementary Table 2 for a

complete list of neurobiological features, and refs. [42,97,98] for population-specific

characteristics)[47]. These maps were obtained in surface or volumetric spaces

using *neuromaps* toolbox, available at https://github.com/netneurolab/neuromaps, and

were parcellated according to Desikan-Killiany atlas. In cases where multiple maps

were available for a single feature, a weighted average was taken. The neurobiological

similarity matrix was computed by correlating the 46 spatial maps between regions. We

assessed whether the neurobiological similarity was co-localized with MIND edges

using both the degree- and edge length-preserving null model, as well as Moran spectral

randomization.

Canonical Correlation Analysis (CCA) is a powerful multivariate method for capturing

associations between two modalities of data (e.g., brain and behavior;

see *Supplementary Subjects and Methods* for details)[99]. A combined Principal

Component Analysis (PCA) and CCA approach was employed to investigate the

regional associations of MIND degree with the neurobiological features. The

contribution of each feature to this neurobiology-MIND association was calculated by

identifying the model loadings[99]. Statistical significance of the models and loadings was

assessed using the spin test.

To identify the neurobiological features associated with phenotypes that were linked to

clinical status, we performed PCA-CCA analyses on the regional maps that represented

the associations between clinical status and both MIND degrees and centiles. Since

symptomatology did not show any significant association with MIND or centiles (see

Supplementary Figs. 4 and 11), we did not assess the neurobiological association with

these maps.

Data availability

According to the informed consent, MRI data from PAFIP dataset is available from the

corresponding author on request. All maturational features, including centiles and the

volume and velocity peaks, were obtained from https://github.com/brainchart/Lifespan.

The neurobiological maps were obtained using *neuromaps* toolbox,[97] available at

https://github.com/netneurolab/neuromaps. The Desikan–Killiany parcellation atlas

used for the neurobiological mapping was obtained from netneurotools

(https://github.com/netneurolab/netneurotools). Source data are provided with this

paper.

Code availability

All code used to perform the analyses can be found at https://github.com/NeuroimagingBrainNetworks/NeurobiologyMINDPsychosis and https://doi.org/10.5281/zenodo.16090993. The code used for computing MIND networks can be found at https://github.com/isebenius/MIND. The code used for the PCA-CCA analyses is available at https://github.com/NeuroimagingBrainNetworks/cca_pls_toolkit.

References

1. Wright, I. C. *et al.* Meta-Analysis of Regional Brain Volumes in Schizophrenia. *AJP* **157**, 16–25 (2000).

2. de Moura, A. M. *et al.* Investigating brain structural patterns in first episode psychosis and schizophrenia using MRI and a machine learning approach. *Psychiatry Research: Neuroimaging* **275**, 14–20 (2018).

3. Radua, J. *et al.* Multimodal meta-analysis of structural and functional brain changes in first episode psychosis and the effects of antipsychotic medication. *Neuroscience & Biobehavioral Reviews* **36**, 2325–2333 (2012).

4. Crespo-Facorro, B. *et al.* Global and regional cortical thinning in first-episode psychosis patients: relationships with clinical and cognitive features. *Psychological Medicine* **41**, 1449–1460 (2011).

5. Vieira, S. *et al.* Neuroanatomical abnormalities in first-episode psychosis across independent samples: a multi-centre mega-analysis. *Psychological Medicine* **51**, 340–350 (2021).

6. Morgan, S. E. *et al.* Cortical patterning of abnormal morphometric similarity in

psychosis is associated with brain expression of schizophrenia-related genes. *Proc*

Natl Acad Sci U S A **116**, 9604–9609 (2019).

7. Fornito, A., Zalesky, A. & Breakspear, M. The connectomics of brain disorders. *Nat*

Rev Neurosci **16**, 159–172 (2015).

8. Crossley, N. A. *et al.* Connectomic correlates of response to treatment in first-

episode psychosis. *Brain* **140**, 487–496 (2017).

9. Maier-Hein, K. H. *et al.* The challenge of mapping the human connectome based on

diffusion tractography. *Nat Commun* **8**, 1349 (2017).

10. Donahue, C. J. *et al.* Using Diffusion Tractography to Predict Cortical Connection

Strength and Distance: A Quantitative Comparison with Tracers in the Monkey. *J.*

Neurosci. **36**, 6758–6770 (2016).

11. van den Heuvel, M. P. *et al.* Comparison of diffusion tractography and tract-tracing

measures of connectivity strength in rhesus macaque connectome. *Human Brain*

Mapping **36**, 3064–3075 (2015).

12. King, D. J. & Wood, A. G. Clinically feasible brain morphometric similarity

network construction approaches with restricted magnetic resonance imaging

acquisitions. *Netw Neurosci* **4**, 274–291 (2020).

13. Joo, S. W. *et al.* Topological abnormalities of the morphometric similarity network

of the cerebral cortex in schizophrenia. *Schizophr* **10**, 1–9 (2024).

14. Homan, P. *et al.* Structural similarity networks predict clinical outcome in early-

phase psychosis. *Neuropsychopharmacol.* **44**, 915–922 (2019).

15. Sebenius, I. *et al.* Robust estimation of cortical similarity networks from brain MRI.

Nat Neurosci **26**, 1461–1471 (2023).

16. Georgiadis, F. *et al.* Connectome architecture shapes large-scale cortical alterations

in schizophrenia: a worldwide ENIGMA study. *Mol Psychiatry* 1–13 (2024)

doi:10.1038/s41380-024-02442-7.

17. Chand, G. B. *et al.* Two distinct neuroanatomical subtypes of schizophrenia

revealed using machine learning. *Brain* **143**, 1027–1038 (2020).

18. Jessen, K. *et al.* Patterns of Cortical Structures and Cognition in Antipsychotic-

Naïve Patients With First-Episode Schizophrenia: A Partial Least Squares

Correlation Analysis. *Biological Psychiatry: Cognitive Neuroscience and*

Neuroimaging **4**, 444–453 (2019).

19. Chopra, S. *et al.* Brainwide Anatomical Connectivity and Prediction of Longitudinal

Outcomes in Antipsychotic-Naïve First-Episode Psychosis. *Biological Psychiatry*

(2024) doi:10.1016/j.biopsych.2024.07.016.

20. Dwyer, D. B. *et al.* Psychosis brain subtypes validated in first-episode cohorts and

related to illness remission: results from the PHENOM consortium. *Mol Psychiatry*

28, 2008–2017 (2023).

21. Yeo, R. A. *et al.* Graph Metrics of Structural Brain Networks in Individuals with

Schizophrenia and Healthy Controls: Group Differences, Relationships with

Intelligence, and Genetics. *Journal of the International Neuropsychological Society*

22, 240–249 (2016).

22. Zhu, T. *et al.* Meta-analysis of structural and functional brain abnormalities in

schizophrenia with persistent negative symptoms using activation likelihood

estimation. *Front. Psychiatry* **13**, (2022).

23. Lesh, T. A. *et al.* A Multimodal Analysis of Antipsychotic Effects on Brain

Structure and Function in First-Episode Schizophrenia. *JAMA Psychiatry* **72**, 226–

234 (2015).

24. Roiz-Santiañez, R., Suarez-Pinilla, P. & Crespo-Facorro, B. Brain Structural Effects

of Antipsychotic Treatment in Schizophrenia: A Systematic Review. *Current*

Neuropharmacology **13**, 422–434 (2015).

25. Emsley, R. *et al.* Antipsychotic treatment effects and structural MRI brain changes

in schizophrenia. *Psychological Medicine* **53**, 2050–2059 (2023).

26. Hansen, J. Y. *et al.* Local molecular and global connectomic contributions to cross-

disorder cortical abnormalities. *Nat Commun* **13**, 4682 (2022).

27. Hettwer, M. D. *et al.* Coordinated cortical thickness alterations across six

neurodevelopmental and psychiatric disorders. *Nat Commun* **13**, 6851 (2022).

28. Shafiei, G. *et al.* Spatial Patterning of Tissue Volume Loss in Schizophrenia

Reflects Brain Network Architecture. *Biological Psychiatry* **87**, 727–735 (2020).

29. Li, J. *et al.* Transcriptomic Similarity Informs Neuromorphic Deviations in

Depression Biotypes. *Biological Psychiatry* **95**, 414–425 (2024).

30. Xia, J. *et al.* Normative structural connectome constrains spreading transient brain

activity in generalized epilepsy. *BMC Medicine* **23**, 258 (2025).

31. Fan, Y.-S. *et al.* Neurodevelopmentally rooted epicenters in schizophrenia:

sensorimotor-association spatial axis of cortical thickness alterations.

2024.03.13.584752 Preprint at https://doi.org/10.1101/2024.03.13.584752 (2024).

32. Seeley, W. W., Crawford, R. K., Zhou, J., Miller, B. L. & Greicius, M. D.

Neurodegenerative diseases target large-scale human brain networks. *Neuron* **62**,

42–52 (2009).

33. Yau, Y. *et al.* Network connectivity determines cortical thinning in early

Parkinson's disease progression. *Nat Commun* **9**, 12 (2018).

34. Bullmore, E. & Sporns, O. Complex brain networks: graph theoretical analysis of

structural and functional systems. *Nat Rev Neurosci* **10**, 186–198 (2009).

35. Whitaker, K. J. *et al.* Adolescence is associated with genomically patterned

consolidation of the hubs of the human brain connectome. *Proc. Natl. Acad. Sci.*

U.S.A. **113**, 9105–9110 (2016).

36. Bethlehem, R. A. I. *et al.* Brain charts for the human lifespan. *Nature* **604**, 525–533

(2022).

37. García-San-Martín, N. *et al.* Molecular and micro-architectural mapping of gray

matter alterations in psychosis. *Mol Psychiatry* 1–10 (2024) doi:10.1038/s41380-

024-02724-0.

38. Constantinides, C. *et al.* Brain ageing in schizophrenia: evidence from 26

international cohorts via the ENIGMA Schizophrenia consortium. *Mol Psychiatry*

28, 1201–1209 (2023).

39. Walther, S. *et al.* Structural alterations of the motor cortex and higher order cortical

areas suggest early neurodevelopmental origin of catatonia in schizophrenia.

Schizophrenia Research **263**, 131–138 (2024).

40. Holmes, A. *et al.* Disruptions of Hierarchical Cortical Organization in Early

Psychosis and Schizophrenia. *Biological Psychiatry: Cognitive Neuroscience and*

Neuroimaging **8**, 1240–1250 (2023).

41. Xia, M., Sun, X., Bu, X., Li, Q. & He, Y. Big connectome imaging data in

psychiatric disorders. *Medicine Plus* **1**, 100038 (2024).

42. Hawrylycz, M. J. *et al.* An anatomically comprehensive atlas of the adult human

brain transcriptome. *Nature* **489**, 391–399 (2012).

43. Hansen, J. Y. *et al.* Mapping neurotransmitter systems to the structural and

functional organization of the human neocortex. *Nat Neurosci* **25**, 1569–1581

(2022).

44. Lotter, L. D. *et al.* Regional patterns of human cortex development correlate with

underlying neurobiology. *Nat Commun* **15**, 7987 (2024).

45. Thompson, P. M. *et al.* ENIGMA and global neuroscience: A decade of large-scale

studies of the brain in health and disease across more than 40 countries. *Transl*

Psychiatry **10**, 1–28 (2020).

46. Sydnor, V. J. *et al.* Neurodevelopment of the association cortices: Patterns,

mechanisms, and implications for psychopathology. *Neuron* **109**, 2820–2846

(2021).

47. Shafiei, G. *et al.* Neurophysiological signatures of cortical micro-architecture. *Nat*

Commun **14**, 6000 (2023).

48. Wheeler, A. L. & Voineskos, A. N. A review of structural neuroimaging in

schizophrenia: from connectivity to connectomics. *Front. Hum. Neurosci.* **8**, (2014).

49. Yao, G. *et al.* Cortical structural changes of morphometric similarity network in

early-onset schizophrenia correlate with specific transcriptional expression patterns.

BMC Med **21**, 479 (2023).

50. Yao, G. *et al.* Transcriptional patterns of the cortical Morphometric Inverse

Divergence in first-episode, treatment-naïve early-onset schizophrenia. *NeuroImage*

285, 120493 (2024).

51. Stephan, K. E., Baldeweg, T. & Friston, K. J. Synaptic Plasticity and Dysconnection

in Schizophrenia. *Biological Psychiatry* **59**, 929–939 (2006).

52. Pettersson-Yeo, W., Allen, P., Benetti, S., McGuire, P. & Mechelli, A.

Dysconnectivity in schizophrenia: Where are we now? *Neuroscience &*

Biobehavioral Reviews **35**, 1110–1124 (2011).

53. Sebenius, I. *et al.* Structural MRI of brain similarity networks. *Nat. Rev. Neurosci.*

26, 42–59 (2025).

54. Prasad, K. *et al.* Structural covariance networks in schizophrenia: A systematic

review Part II. *Schizophr Res* **239**, 176–191 (2022).

55. Ramsay, I. S., Mueller, B., Ma, Y., Shen, C. & Sponheim, S. R. Thalamocortical

connectivity and its relationship with symptoms and cognition across the psychosis

continuum. *Psychological Medicine* **53**, 5582–5591 (2023).

56. Canu, E., Agosta, F. & Filippi, M. A selective review of structural connectivity

abnormalities of schizophrenic patients at different stages of the disease.

Schizophrenia Research **161**, 19–28 (2015).

57. Jiang, Y. *et al.* Neuroimaging epicenters as potential sites of onset of the

neuroanatomical pathology in schizophrenia. *Science Advances* **10**, eadk6063

(2024).

58. Brown, J. A. *et al.* Patient-Tailored, Connectivity-Based Forecasts of Spreading

Brain Atrophy. *Neuron* **104**, 856-868.e5 (2019).

59. Shafiei, G. *et al.* Network structure and transcriptomic vulnerability shape atrophy

in frontotemporal dementia. *Brain* **146**, 321–336 (2023).

60. Chopra, S. *et al.* Network-Based Spreading of Gray Matter Changes Across

Different Stages of Psychosis. *JAMA Psychiatry* **80**, 1246–1257 (2023).

61. Seeley, W. W. Mapping Neurodegenerative Disease Onset and Progression. *Cold

Spring Harb Perspect Biol* **9**, a023622 (2017).

62. Hartz, S. M. *et al.* G72 influences longitudinal change in frontal lobe volume in

schizophrenia. *Am J Med Genet B Neuropsychiatr Genet* **153B**, 640–647 (2010).

63. Vázquez-Bourgon, J. *et al.* Variations in Disrupted-in-Schizophrenia 1 gene

modulate long-term longitudinal differences in cortical thickness in patients with a

first-episode of psychosis. *Brain Imaging and Behavior* **10**, 629–635 (2016).

64. Fornito, A. & Bullmore, E. T. Reconciling abnormalities of brain network structure

892   and function in schizophrenia. *Current Opinion in Neurobiology* **30**, 44–50 (2015).

893   65. Wagner, H. H. & Dray, S. Generating spatially constrained null models for

894   irregularly spaced data using Moran spectral randomization methods. *Methods in*

895   *Ecology and Evolution* **6**, 1169–1178 (2015).

896   66. Betzel, Richard F., B., Danielle S. Specificity and robustness of long-distance

897   connections in weighted, interareal connectomes. *Proceedings of the National*

898   *Academy of Sciences* vol. 115 4880–4889 (2018).

899   67. Williams, L. M. Voxel-based morphometry in schizophrenia: implications for

900   neurodevelopmental connectivity models, cognition and affect. *Expert Review of*

901   *Neurotherapeutics* **8**, 1049–1065 (2008).

902   68. Hammer, T. B. *et al.* Structural brain correlates of sensorimotor gating in

903   antipsychotic-naive men with first-episode schizophrenia. *Journal of Psychiatry and*

904   *Neuroscience* **38**, 34–42 (2013).

905   69. Dong, D. *et al.* Compressed sensorimotor-to-transmodal hierarchical organization in

906   schizophrenia. *Psychological Medicine* **53**, 771–784 (2023).

907   70. Dennis, E. L. *et al.* Development of brain structural connectivity between ages 12

908   and 30: A 4-Tesla diffusion imaging study in 439 adolescents and adults.

909   *NeuroImage* **64**, 671–684 (2013).

910   71. Vanes, L. D. *et al.* White matter tract myelin maturation and its association with

911   general psychopathology in adolescence and early adulthood. *Human Brain*

912   *Mapping* **41**, 827–839 (2020).

913   72. Wen, X. *et al.* First-year development of modules and hubs in infant brain

914   functional networks. *NeuroImage* **185**, 222–235 (2019).

73. Karlsgodt, K. H. *et al.* Developmental disruptions in neural connectivity in the
pathophysiology of schizophrenia. *Development and Psychopathology* **20**, 1297–
1327 (2008).

74. Karlsgodt, K. H., Jacobson, S. C., Seal, M. & Fusar-Poli, P. The Relationship of
Developmental Changes in White Matter to the Onset of Psychosis. *Curr Pharm*
Des **18**, 422–433 (2012).

75. Seshadri, S., Zeledon, M. & Sawa, A. Synapse-specific contributions in the cortical
pathology of schizophrenia. *Neurobiology of Disease* **53**, 26–35 (2013).

76. Xue, K. *et al.* Brain connectivity and transcriptomic similarity inform abnormal
morphometric similarity patterns in first-episode, treatment-naïve major depressive
disorder. *Journal of Affective Disorders* (2024) doi:10.1016/j.jad.2024.11.021.

77. Wang, X., Lin, Q., Xia, M. & He, Y. Differentially categorized structural brain hubs
are involved in different microstructural, functional, and cognitive characteristics
and contribute to individual identification. *Human Brain Mapping* **39**, 1647–1663
(2018).

78. Chételat, G. *et al.* Relationships between brain metabolism decrease in normal aging
and changes in structural and functional connectivity. *NeuroImage* **76**, 167–177
(2013).

79. Beul, S. F. & Hilgetag, C. C. Neuron density fundamentally relates to architecture
and connectivity of the primate cerebral cortex. *NeuroImage* **189**, 777–792 (2019).

80. Matejuk, A. & Ransohoff, R. M. Crosstalk Between Astrocytes and Microglia: An
Overview. *Front. Immunol.* **11**, (2020).

81. Vainchtein, I. D. & Molofsky, A. V. Astrocytes and Microglia: In Sickness and in
Health. *Trends in Neurosciences* **43**, 144–154 (2020).

82. Michell-Robinson, M. A. *et al.* Roles of microglia in brain development, tissue

maintenance and repair. *Brain* **138**, 1138–1159 (2015).

83. Reemst, K., Noctor, S. C., Lucassen, P. J. & Hol, E. M. The Indispensable Roles of

Microglia and Astrocytes during Brain Development. *Front. Hum. Neurosci.* **10**,

(2016).

84. Dietz, A. G., Goldman, S. A. & Nedergaard, M. Glial cells in schizophrenia: a

unified hypothesis. *The Lancet Psychiatry* **7**, 272–281 (2020).

85. Cabral, J., Hugues, E., Kringelbach, M. L. & Deco, G. Modeling the outcome of

structural disconnection on resting-state functional connectivity. *NeuroImage* **62**,

1342–1353 (2012).

86. Dear, R. *et al.* Cortical gene expression architecture links healthy neurodevelopment

to the imaging, transcriptomics and genetics of autism and schizophrenia. *Nat*

Neurosci **27**, 1075–1086 (2024).

87. Erkol, C. *et al.* White Matter Measures and Cognition in Schizophrenia. *Front.*

Psychiatry **11**, (2020).

88. Brambilla, P. *et al.* Assessment of cerebral blood volume in schizophrenia: A

magnetic resonance imaging study. *Journal of Psychiatric Research* **41**, 502–510

(2007).

89. Wu, X.-L., Yan, Q.-J. & Zhu, F. Abnormal synaptic plasticity and impaired

cognition in schizophrenia. *World J Psychiatry* **12**, 541–557 (2022).

90. Marek, S. *et al.* Reproducible brain-wide association studies require thousands of

individuals. *Nature* **603**, 654–660 (2022).

91. Chopra, S. *et al.* Network-based Molecular Constraints on in vivo Synaptic Density

Alterations   in   Schizophrenia.   2025.03.22.25324465   Preprint   at

https://doi.org/10.1101/2025.03.22.25324465 (2025).

92. Ng, B. *et al.* Integration across biophysical scales identifies molecular and cellular correlates of person-to-person variability in human brain connectivity. *Nat Neurosci* **27**, 2240–2252 (2024).

93. Crespo-Facorro, B. *et al.* A Practical Clinical Trial Comparing Haloperidol, Risperidone, and Olanzapine for the Acute Treatment of First-Episode Nonaffective Psychosis. *J. Clin. Psychiatry* **67**, 1511–1521 (2006).

94. Gardner, M. *et al.* ComBatLS: A Location- and Scale-Preserving Method for Multi-Site Image Harmonization. *Human Brain Mapping* **46**, e70197 (2025).

95. Strimmer, K. A unified approach to false discovery rate estimation. *BMC Bioinformatics* **9**, 303 (2008).

96. Rodríguez-Sánchez, J. M. *et al.* Ten-year course of cognition in first-episode non-affective psychosis patients: PAFIP cohort. *Psychological Medicine* **52**, 770–779 (2022).

97. Markello, R. D. *et al.* neuromaps: structural and functional interpretation of brain maps. *Nat Methods* **19**, 1472–1479 (2022).

98. Amunts, K. *et al.* BigBrain: An Ultrahigh-Resolution 3D Human Brain Model. *Science* **340**, 1472–1475 (2013).

99. Mihalik, A. *et al.* Canonical Correlation Analysis and Partial Least Squares for Identifying Brain–Behavior Associations: A Tutorial and a Comparative Study. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging* **7**, 1055–1067 (2022).

Acknowledgments

R.R.G. is funded by *Plan de Generación de Conocimiento* (PID2021-122853OA-I00), *Plan de Consolidación* (CNS2023-143647) and *ERANET Neuron JTC* 2023 (ERP-

988 2023-23684211). Both R.R.G. and N.G.S. are funded by the *EMERGIA Junta de*
989 *Andalucía* program (EMERGIA20_00139). R.A.A. is funded by *Plan de Consolidación*
990 (CNS2022-136110).

Author Contributions

N.G.S. performed data curation, methodological design, data analysis, and drafted the
manuscript; R.A.I.B., P.S., A.M., J.Se., I.S., C.A.M., L.D., G.S., S.E.M., M.R.V.,
R.A.A., J.V.B., B.M., J.Su., B.C.F., and R.R.G contributed to data acquisition, provided
advice on data analysis, and participated in writing and editing the manuscript. R.R.G.
also contributed to conceptualization and supervision of the work. All authors approved
the submitted version of the manuscript.

Competing interests

999 R.AI.B. and J.Se. hold equity in and are directors of Centile Bioscience. All other
1000 authors report no competing interests.