Anxiolytic management in eating disorder patients receiving cognitive behavioral therapy:

A quality improvement brief report

Short running title: Anxiolytic and behavioral therapy for eating disorders

Brad E.R. Smith, MD [a], Katelyn A. Greenberg, MA [b], Sreya Vadapalli, MBS [b], Sheldon R. Garrison, PhD [b],*

[a] Rogers Behavioral Health, Eating Disorder Recovery, Oconomowoc, WI, United States 53066

[b] Rogers Behavioral Health, Research Center, Oconomowoc, WI, United States 53066

* Corresponding author:

Sheldon Garrison, PhD

Rogers Behavioral Health

Research Center – Ladish Building

34700 Valley Road

Oconomowoc, WI 53066

Phone: 262-646-1627

sheldon.garrison@rogersbh.org

Abstract

Eating disorders (EDs) are serious lifelong and debilitating health conditions characterized by complex adverse eating behaviors resulting in weight loss and decreased quality of life. One evidence-based treatment is a form of cognitive-behavioral therapy (CBT) called exposure therapy where patients progress through a hierachy of exposures. This treatment approach results in concern related to CBT efficacy for patients concurrently treated with anxiolytic medications. These medications, which are being incorporated into treatment protocols, may impact the anxiety-provoking components of the exposure response, potentially limiting its efficacy. Therefore, patients were assessed in multiple ways to ensure that providers could prescribe anxiolytics without risk to diminishing the well-established response to CBT. To ensure that patients treated with anxiolytics progressed at a similar rate to those who did not receive them, a quality improvement study was conducted to compare all patients in residential, partial hospitalization and intensive outpatient eating disorder treatment programs.

Keywords: Anti-anxiety medication, psychotherapy, benzodiazepine, anorexia nervosa, bulimia nervosa, binge eating disorder.

Introduction

The impact of eating disorders (EDs) is felt broadly, with a lifetime prevalence in the general population of 0.9% (Qian et al., 2022). The onset of EDs, most commonly anorexia nervosa, bulimia nervosa and binge eating disorder, begins with behavioral and cognitive abnormalities, leading to multi-system issues. Early symptoms may manifest as difficult eating under 10 years (Pinhas et al., 2011), transitioning in adolescence to dieting, body dissatisfaction and body image concerns during critical neurodevelopmental periods (Rohde et al., 2015).

The complex neurocognitive involvement of EDs results in inadequate pharmacological options. Current evidence-based treatment for EDs is cognitive behavioral therapy (CBT), specifically with fear extinction and anti-depressants (Crone et al., 2023; Farrell et al., 2019; P. Hay, 2020). Comorbid disorders present in 56-92% of ED patients, including over half with anxiety (Wade et al., 2023). This adds an unexpected complexity to treating when anxiolytic medications are implemented to manage anxiety, introducing concerns regarding how they may affect CBT efficacy.

The goal of this retrospective quality improvement study was to analyze changes in ED symptom severity, as well as severity of co-occurring depressive and obsessive-compulsive disorder (OCD) symptoms in a high-acuity patient population undergoing ED recovery. Providers began incorporating anxiolytics for the management of patients with an ED. To ensure the anxiolytics did not interfere with other treatment modalities, patients prescribed anxiolytics were evaluated using multiple assessments during the course of their treatment. Improvement for patients receiving anxiolytic treatment (anxiolytic) with standard of care was compared to those receiving standard of care alone (non-anxiolytic) to ensure a consistent degree of improvement.

Methods

Electronic medical records were retrospectively analyzed for all patients discharged from an ED program at Rogers Behavioral Health, a behavioral healthcare system that provides mental health and addiction treatment services, between January 1, 2018 and May 31, 2024. Inclusion criteria were: (1) ≥18 years of age; (2) completion of pre- and post-assessments given at admission (baseline); and, at discharge for at least one measure; (3) admission to an ED program at any treatment level (inpatient, residential, partial hospitalization and intensive outpatient). Each patient's most intensive level of care was analyzed. All patients received intensive daily cognitive-behavioral therapy (CBT) with exposure and response prevention (ERP), a standard of care treatment option. The anxiolytics administered to patients were prescribed according to physician preference and clinical judgment, and included alprazolam, clonazepam, diazepam, gabapentin, hydroxyzine, and lorazepam. Patient data was extracted from the electronic health records and de-identified.

Outcomes

The primary outcome measure was the Eating Disorder Examination Questionnaire (EDE-Q) (Fairburn & Beglin, 1994) to measure ED behavior frequency from admission to discharge. Secondary outcomes included the Yale-Brown Cornell Eating Disorder Scale (YBC-EDS) (Mazure et al., 1994) to measure ED symptom severity, the Quick Inventory of Depression Symptomatology (QIDS) scale (Rush et al., 2003) to measure depressive symptoms and the Yale-Brown Obsessive-Compulsive Scale (Y-BOCS) scale to measure obsessive-compulsive disorder (OCD) symptom severity (Goodman, Price, Rasmussen, Mazure, Delgado, et al., 1989; Goodman, Price, Rasmussen, Mazure, Fleischmann, et al., 1989). Additional secondary outcomes included exposure hierarchy percent completion to quantify CBT treatment progress,

length of stay (LOS), and readmission rate. The demographic variables collected included age, sex, race, and ethnicity. An additional clinical variable employed in the analysis was level of care, consisting of inpatient, residential, partial hospitalization, and intensive outpatient.

Data analysis

The treatment groups for the following analyses were divided into those who were prescribed anxiolytic medication alongside standard therapy ('anxiolytic') and those not prescribed anxiolytics ('non-anxiolytic'). The non-anxiolytic group was control-matched to the anxiolytic group by randomly selecting the matched controls using the MatchIt package within R, version 4.3.3 (Ho et al., 2011; R Core Team, 2021). They were control-matched based on age, sex, race, and EDE-Q score at admission to ensure similar baseline characteristics between the anxiolytic and non-anxiolytic groups. Demographic characteristics were reported as frequencies and compared between treatment groups using Chi-square tests. Primary analyses were within- and between-group comparisons in assessment scores at admission and discharge, obtained using t tests. Assessment outcomes were reported as mean scores and standard error. Secondary analyses compared outcomes between levels of care in addition to treatment group, completed using t tests. Readmission rates were reported as frequencies and LOS was reported as number of days. For inpatient and residential, LOS is measured in calendar days, whereas partial hospitalization and intensive outpatient are measured in treatment days. Statistical analyses were completed using R.

RESULTS

Demographics and baseline clinical characteristics

Demographic data for the 242 patients who met eligibility criteria were reported at baseline (**Table 1**). Upon admission, EDEQ scores for both treatment groups fell within the clinical range denoting 'moderate' severity. YBC-EDS scores for patients taking anxiolytic medication alongside standard therapy were slightly higher than those not taking anxiolytics; the anxiolytic group fell within the clinical range indicating 'moderate' severity, while the non-anxiolytic group fell within the range indicating 'mild' severity. YBOCS-SR and QIDS scores were moderate in both groups (**Table 1**).

Clinical improvement in both groups across all measures

The anxiolytic group demonstrated a 40% reduction in ED behavior frequency and a 36% improvement in symptom severity using the EDEQ and YBC-EDS, respectively. This degree of improvement was consistent in the non-anxiolytic group, with a 46% reduction in ED behavior and a 41% improvement in symptom severity ($p < 0.05$) (**Figure 1a-b**). Both groups had similar improvement in depressive symptoms, as measured by the QIDS, with the anxiolytic group having a 36% reduction in score, and the non-anxiolytic group 37%. Co-occurring OCD symptoms, assessed with the YBOCS-SR, improved in both the anxiolytic and non-anxiolytic groups by 27% and 35%, respectively. A clinically significant change in depressive symptoms was observed for both groups, with the anxiolytic group reporting a 36% reduction in symptoms, and the non-anxiolytic group reporting a 37% reduction ($p < 0.05$) (**Figure 1c-d**). Exposure hierarchy completion was similar between the anxiolytic group (37%) and the non-anxiolytic group (36%) (**Figure 1e**). Readmission frequency did not significantly differ between treatment groups, which was 3% and 0% for the anxiolytic and the non-anxiolytic groups, respectively. Further analyses breaking down by level of care in addition to treatment status demonstrated that LOS differed at

the residential and partial hospitalization levels only, with the anxiolytic patients requiring more treatment days than the non-anxiolytic patients, with no differences in LOS for inpatient and intensive outpatient levels of care (**Table 1**).

Discussion

This retrospective analysis of anxiolytic utilization in ED patients suggests that treatment progress with CBT was unaffected. The anxiolytic and non-anxiolytic groups were observed to be in similar clinical ranges at baseline (e.g., moderate). Importantly, both treatment groups showed marked improvement in ED symptomatology, reaching mild levels at discharge, implying that anxiolytics did not impact the rate of improvement or CBT efficacy. Furthermore, both treatment groups exhibited significant reductions in OCD and depressive symptoms from admission to discharge, indicating that patient clinical progress altogether was uninhibited by administration of anxiolytic medication (**Table 1**).

The current study, investigating a large ED patient cohort, addresses longstanding concerns of anxiolytics compromising CBT efficacy and supports that anxiolytics can be administered alongside standard therapy. The hesitancy surrounding concurrent anxiolytic use for ED patients undergoing CBT can be largely attributed to a dearth of controlled trials conducted with mixed results on this treatment combination (P. J. Hay & Claudino, 2012), particularly with high-acuity patient participants being treated in more intensive levels of care (e.g., residential, partial hospitalization). The current analysis captures an integral characteristic within this population that is not well-represented in the literature, that is, the usage of anxiolytic medication for patients with severe cognitive rigidity at baseline. This quality improvement project corroborates effectiveness of anxiolytics, particularly in early stages of treatment, when an individual may have trouble initially engaging in robust CBT due to high levels of specific and/or general anxiety.

Disclosure statement

No potential conflict of interest was reported by the author(s).

ETHICS DECLARATION

This Rogers Behavioral Health Institutional Review Board of Rogers Behavioral Health determined that this quality improvement project was non-human subjects research and waived ethical approval for this work

Funding

This work was generously funded by the Rogers Behavioral Health Foundation and the Lynn S Nicholas Family Foundation.

Data availability statement

All data produced in the present study are available upon reasonable request to the authors.

REFERENCES

Crone, C., Fochtmann, L. J., Attia, E., Boland, R., Escobar, J., Fornari, V., Golden, N., Guarda, A., Jackson-Triche, M., Manzo, L., Mascolo, M., Pierce, K., Riddle, M., Seritan, A., Uniacke, B., Zucker, N., Yager, J., Craig, T. J., Hong, S. H., & Medicus, J. (2023). The American Psychiatric Association Practice Guideline for the Treatment of Patients With Eating Disorders. *Am J Psychiatry*, *180*(2), 167-171. https://doi.org/10.1176/appi.ajp.23180001

Fairburn, C. G., & Beglin, S. J. (1994). Assessment of eating disorders: interview or self-report questionnaire? *Int J Eat Disord*, *16*(4), 363-370. https://www.ncbi.nlm.nih.gov/pubmed/7866415

Farrell, N. R., Brosof, L. C., Vanzhula, I. A., Christian, C., Bowie, O. R., & Levinson, C. A. (2019). [Exploring Mechanisms of Action in Exposure-Based Cognitive Behavioral Therapy for Eating Disorders: The Role of Eating-Related Fears and Body-Related Safety Behaviors]. *Behav Ther*, *50*(6), 1125-1135. https://doi.org/10.1016/j.beth.2019.01.008

Goodman, W. K., Price, L. H., Rasmussen, S. A., Mazure, C., Delgado, P., Heninger, G. R., & Charney, D. S. (1989). The Yale-Brown Obsessive Compulsive Scale. II. Validity. *Arch Gen Psychiatry*, *46*(11), 1012-1016. https://doi.org/10.1001/archpsyc.1989.01810110054008

Goodman, W. K., Price, L. H., Rasmussen, S. A., Mazure, C., Fleischmann, R. L., Hill, C. L., Heninger, G. R., & Charney, D. S. (1989). The Yale-Brown Obsessive Compulsive Scale. I. Development, use, and reliability. *Arch Gen Psychiatry*, *46*(11), 1006-1011. https://doi.org/10.1001/archpsyc.1989.01810110048007

Hay, P. (2020). Current approach to eating disorders: a clinical update. *Intern Med J*, *50*(1), 24-29. https://doi.org/10.1111/imj.14691

Hay, P. J., & Claudino, A. M. (2012). Clinical psychopharmacology of eating disorders: a research update. *Int J Neuropsychopharmacol*, *15*(2), 209-222. https://doi.org/10.1017/S1461145711000460

Ho, D., Imai, K., King, G., & Stuart, E. A. (2011). MatchIt: Nonparametric Preprocessing for Parametric Causal Inference. *Journal of Statistical Software*, *42*(8), 1 - 28. https://doi.org/10.18637/jss.v042.i08

Mazure, C. M., Halmi, K. A., Sunday, S. R., Romano, S. J., & Einhorn, A. M. (1994). The Yale-Brown-Cornell Eating Disorder Scale: development, use, reliability and validity. *J Psychiatr Res*, *28*(5), 425-445. https://doi.org/10.1016/0022-3956(94)90002-7

Pinhas, L., Morris, A., Crosby, R. D., & Katzman, D. K. (2011). Incidence and age-specific presentation of restrictive eating disorders in children: a Canadian Paediatric Surveillance Program study. *Arch Pediatr Adolesc Med*, *165*(10), 895-899. https://doi.org/10.1001/archpediatrics.2011.145

Qian, J., Wu, Y., Liu, F., Zhu, Y., Jin, H., Zhang, H., Wan, Y., Li, C., & Yu, D. (2022). An update on the prevalence of eating disorders in the general population: a systematic review and meta-analysis. *Eat Weight Disord*, *27*(2), 415-428. https://doi.org/10.1007/s40519-021-01162-z

R Core Team, R. (2021). *A Language and Environment for Statistical Computing* (Version R version 4.3.3). https://www.R-project.org/

Rohde, P., Stice, E., & Marti, C. N. (2015). Development and predictive effects of eating disorder risk factors during adolescence: Implications for prevention efforts. *Int J Eat Disord*, *48*(2), 187-198. https://doi.org/10.1002/eat.22270

Rush, A. J., Trivedi, M. H., Ibrahim, H. M., Carmody, T. J., Arnow, B., Klein, D. N., Markowitz, J. C., Ninan, P. T., Kornstein, S., Manber, R., Thase, M. E., Kocsis, J. H., & Keller, M. B.

(2003). The 16-Item Quick Inventory of Depressive Symptomatology (QIDS), clinician rating (QIDS-C), and self-report (QIDS-SR): a psychometric evaluation in patients with chronic major depression. *Biol Psychiatry*, *54*(5), 573-583. https://doi.org/10.1016/s0006-3223(02)01866-8

Wade, T. D., Shafran, R., & Cooper, Z. (2023). Developing a protocol to address co-occurring mental health conditions in the treatment of eating disorders. *Int J Eat Disord*. https://doi.org/10.1002/eat.24008

Table 1.

Demographic characteristics

	Anxiolytic	Non-anxiolytic	*p* value
n	121	121	
Age (y) [mean (SEM)]	25.4 (0.9)	24.7 (0.8)	0.6
Sex (%)			0.509
Female	83.5	79.3	
Male	16.5	20.7	
Race and Ethnicity (%)			0.545
White	93.4	88.4	
Black	0.8	0.8	
Asian or Pacific Islander	2.5	1.7	
Other	1.7	3.3	
Ethnicity (%)			0.418
Hispanic or Latino	7.4	7.4	
Not Hispanic or Latino	91.7	85.1	

Missing values not displayed

Treatment response, comparison between groups at baseline and discharge.

	Baseline			Discharge		
Variable [mean (SEM), n]	Anxiolytic	Non-anxiolytic	*p* value	Anxiolytic	Non-anxiolytic	*p* value
EDEQ total score	3.3 (0.15), 121	3.2 (0.1), 121	0.689	2.0 (0.1), 120	1.8 (0.1), 121	0.179
YBC-EDS total score	16.5 (0.6), 120	15.0 (0.6), 115	0.698	10.5 (0.6), 120	8.9 (0.5), 121	0.098
QIDS total score	14.8 (0.5), 121	12.3 (0.5), 121	**<0.001**	9.5 (0.5), 121	7.8 (0.4), 121	**0.009**
YBOCS-SR total score	17.4 (0.9), 112	16.8 (0.8), 113	0.563	12.6 (0.8), 111	10.9 (0.7), 113	0.098
Exposure hierarchy completion (%)				36.8 (1.9), 121	36.3 (1.9), 121	0.85
Length of Stay (days) [range]						
Inpatient				33.7 (5.4), 3 (23-40)	38.5 (18.5), 2 [20-57]	0.775
Residential				63.2 (4.5), 58 [13-189]	50.1 (2.7), 47 [14-98]	**0.022**
Partial hospitalization				35.5 (1.5), 55 [11-70]	31.4 (1.4), 65 [10-72]	**0.046**
Intensive outpatient				24.4 (3.9), 5 [17-38]	25.0 (3.4), 7 [14-42]	0.911
Readmissions (%, n)				3.3, 4	0.0, 0	

Missing values not displayed

Figure 1. Non-anxiolytic- and anxiolytic-treated ED patients improved across all measures. (**a**) Anxiolytic-treated patients had a 40% improvement in EDE-Q score (p <0.001; Effect Size (ES) 0.64), and non-anxiolytic-treated patients has a 46% improvement (p <0.001; ES 0.72). (**b**) Anxiolytic-treated patients had a 36% improvement in YBC-EDS (p <0.001; ES 0.58), and non-anxiolytic-treated patients has a 41% improvement (p <0.001; ES 0.73). (**c**) Anxiolytic-treated patients had a 36% improvement in QIDS (p <0.001; ES 0.74), and non-anxiolytic-treated patients has a 37% improvement (p <0.001; ES 0.88). (**d**) Anxiolytic-treated patients had a 27% improvement in YBOCS (p <0.001; ES 0.67), and non-anxiolytic-treated patients has a 35% improvement (p <0.001; ES 0.90). (**e**) Non-anxiolytic-treated patients had a similar completion of their exposure hierarchy to the anxiolytic treated group (p <0.001).

Figure 1

A

B

C

D

E