Associations between weather and *Plasmodium vivax* malaria in an Amazonian elimination setting: a distributed lag analysis from 2017–2024.

Gabriella Barratt Heitmann,[1]* Xue Wu,[2] Anna T. Nguyen,[1] Astrid Altamirano-Quiroz,[3] Sydney R. Fine,[2] Bryan Fernandez-Camacho,[3] Antony Barja,[3] Renato Cava,[3] Verónica Soto-Calle,[3] Hugo Rodriguez,[4] Gabriel Carrasco-Escobar,[3] Adam Bennett,[2] Alejandro Llanos-Cuentas,[3] Erin A. Mordecai,[5,6] Michelle S. Hsiang,[2,7,8,9] Jade Benjamin-Chung[1,9]

1. Department of Epidemiology and Population Health, School of Medicine, Stanford University, Stanford, CA, USA
2. Malaria Elimination Initiative, Institute for Global Health Sciences, University of California San Francisco (UCSF), San Francisco, CA, USA
3. Instituto de Medicina Tropical, Alexander von Humboldt, Universidad Peruana Cayetano Heredia, Lima, Perú
4. Universidad Nacional de la Amazonía Peruana, Loreto, Perú
5. Department of Biology, Stanford University, Stanford, CA, USA
6. Woods Institute for the Environment, Stanford University, Stanford, CA, USA
7. Department of Epidemiology and Biostatistics, UCSF, San Francisco, CA, USA
8. Department of Pediatrics, UCSF, San Francisco, CA, USA
9. Chan Zuckerberg Biohub, San Francisco, CA, USA

*Corresponding author

ABSTRACT

Background

Plasmodium vivax (*Pv*) is the predominant malaria species in countries approaching elimination.

Environmental factors can guide intervention targeting.Yet, research that considers the long-term

relapse periodicity of *Pv* is limited, particularly in Latin America.

Methods

We merged *Pv* malaria incidence data from 2017–2024 from 136 communities in the Peruvian

Amazon with hourly weather data from the ERA5 dataset. Predictors included weekly minimum

and maximum temperature and total weekly precipitation. We fit non-linear distributed lag

models using a lookback period of 2–16 weeks. We conducted sub-group analyses by

community type (adjacent to river versus highway) and El Niño Southern Oscillation (ENSO)

period. Temperature models were adjusted for total precipitation; precipitation models were

adjusted for maximum temperature.

Findings

Minimum temperature at the 90th percentile (23.7ºC) was associated with 10% (95% CI 5%–

14%) higher malaria incidence compared to the lowest minimum temperature (20.4ºC) at a 7-

week lag. Maximum temperature at the 90th percentile (33.7ºC) was associated with 10% (95%

CI 8%–13%) higher malaria incidence compared to the lowest maximum temperature (29.6ºC) at

a 9-week lag. Total weekly precipitation at the 90th percentile (1000mm) was associated with

29% (95% CI 24%–33%) higher malaria incidence compared to weeks with no precipitation at

an 11-week lag. Incidence was higher and associations were stronger in communities adjacent to

rivers versus highways. Malaria incidence was lower during El Niño periods, and there was

evidence of interaction on the multiplicative scale for the association between incidence, all

weather predictors, and ENSO period.

Interpretation

Pv malaria incidence was positively associated with higher temperatures and precipitation in an

elimination setting in Peru, particularly in riverine communities during non-El Niño years, with

longer lag periods than previously reported for such associations. These findings can inform

malaria elimination interventions to combat the long-lasting effects of weather on *Pv*

transmission.

Research in context
Evidence before this study
We searched PubMed for ("weather" OR "climate" OR "temperature" OR "precipitation") AND
("malaria") AND ("Amazon"). The search yielded 76 results. The most relevant studies looked at
the influence of weather variables on all malaria incidence in neighboring Brazil and Venezuela.
The study in Brazil found positive associations with temperatures between 25–30ºC and
precipitation >4.46cm at 1-week lags, but negative associations with temperatures >25ºC at 2–3-
week lags. The study in Venezuela found positive associations with mean temperatures in the
range 20–30ºC and no association with precipitation. Both studies emphasized that malaria
transmission was highly heterogeneous.
61

Added value of this study
Our study used a distributed lag analysis to account for the long-term relapse periodicity
common in *Pv* malaria, which is predominant in the Peruvian Amazon. Our study is, to our
knowledge, the first to exclusively focus on climatic drivers of *Pv* malaria at long-term lags in an
elimination setting dominated by the *Ny. darlingi* vector. This transmission system is especially
understudied compared to *Plasmodium falciparum* malaria, yet *Pv* malaria is the predominant
species in elimination settings worldwide.
69

Implications of all the available evidence
Our study contributes to an important body of literature characterizing climatic drivers of *Pv*
malaria in the Peruvian Amazon. Our findings indicate that over 1–4-month lags, higher
temperatures and precipitation could contribute to higher malaria incidence, particularly during
neutral El Niño Southern Oscillation (ENSO) years. We also found that communities near rivers
were more sensitive to changes in temperature, while communities near highways were more
sensitive to changes in precipitation.

INTRODUCTION

As countries approach malaria elimination, the proportion of malaria infections due to *Plasmodium vivax* (*Pv*) increases.[1] *Pv* presents unique challenges for malaria elimination efforts because of its ability to survive in a wide range of environments, including temperate settings, by utilizing a dormant hypnozoite stage.[2] Relapsing infections occur weeks to months after a prior infection or relapse, and serve as a significant reservoir for persistent transmission.[3 4,5] *Pv* infections can be latent, as well asymptomatic, sub-patent, or minimally symptomatic, making diagnosis and surveillance a challenge. Further, symptomatic infections are low density and often missed through standard diagnostics.[4,6,7] These characteristics greatly obfuscate *Pv* transmission biology and make *Pv* harder to detect, treat, and ultimately eliminate.

Over 90% of Latin America's malaria burden is concentrated in the Amazon Basin, and and a recent review of malaria elimination progress highlighted the need to study risk heterogeneity within the Amazon.[8] Parasite populations and vector behavior vary significantly within this region,[9–11] and this variation is likely driven by adaptations to different environmental conditions.[10,12,13] The Amazon Basin is also undergoing rapid environmental and climatic changes[14–16], and malaria elimination will require a fine-scale understanding of how those changes are altering transmission dynamics.

There is strong biologic plausibility for temperature and precipitation to influence *Pv* transmission (Figure 1).[6,17–22] Increased precipitation can increase standing water presence, soil moisture, and humidity, which in turn increase breeding habitat for malaria vectors.[23] However, this relationship is complicated: in dry-land communities, heavy rain can flush mosquito larvae, leading to decreased mosquito populations, whereas in riverine settings such as in the Amazon, flooding from heavy rains can increase breeding ground and mosquito populations.[23] Temperature has strong effects on the capacity of the *Nyssorhynchus* vector to transmit malaria via the mosquito biting rate, mosquito abundance, parasite incubation rate, and mosquito longevity.[24,25] Finally, weather could be a direct trigger for relapse infections for *Pv* infections[26], and there is other indirect evidence that relapses may be triggered by antigens in uninfected mosquito saliva (biting rate is temperature-sensitive).[3]

Prior research on the relationship between weather and malaria incidence in Latin America has found positive associations with rainfall but mixed associations with higher temperatures, and there is significant spatial heterogeneity[27,28]. Additionally, prior studies have not focused specifically on *Pv* malaria, including those conducted in the Amazon, where *Pv* accounts for 77% of the transmission burden.[11]10/21/2025 6:09:00 PM The Loreto Region has maximum temperatures between 30–35ºC thought to be above the malaria transmission thermal optimum (25ºC), and daily temperature extremes are thought to impact malaria transmission[29]. To our knowledge, the thermal biology of *Pv* transmission by *Nyssorhynchus* (formerly *Anopheles*)

darlingi, the primary vector in this region, has not been well characterized in disease ecology
studies.
Furthermore, the Amazon region is subject to longer-term climate oscillations due to the El Niño
Southern Oscillation (ENSO). The ENSO cycle is characterized by hot, dry conditions during El
Niño years and rainy conditions during La Niña years; these periods oscillate with neutral
periods every 3–7 years[15]. One prior study in the Brazilian Amazon found that El Niño and La
Niña were associated with decreased malaria incidence, but there was significant spatial
heterogeneity[28].
124
125 Our objective was to assess the influence of rainfall and temperature on *Pv* incidence in Loreto
126 Region, Peru, one of the only remaining regions with malaria transmission in the country and
127 key to the Peruvian government's target of malaria elimination by 2030.[30] Information about
128 climatic drivers of *Pv* transmission is needed to inform effective tailoring and targeting of
129 interventions for malaria elimination.
130
131
METHODS
Study design
We analyzed malaria incidence data collected from January 2017 to December 2024 in all 136
communities in San Juan Bautista, Punchana, Alto Nanay, and Belen districts in Loreto region,
Peru, an area covering 2,569 km^2 (Figure 2). The Peruvian Ministry of Health (MINSA)
conducted a census in 2017 that was used to estimate the starting population size of each
community. The average population size of each community was 321 individuals (range: 15 to
3223).
140
Peru's malaria burden is concentrated in the Loreto Region, which includes a large swath of the
Peruvian Amazon.[7] Since the 1990s, rapid urbanization and deforestation in the area surrounding
Iquitos, the capital of the Department of Loreto, has contributed to increased malaria burden[7],
likely through increased forest edge habitat, which promotes mosquito breeding, survival, and
biting[14]. While this region is approaching elimination, sustained, endemic transmission remains,
with an annual incidence rate of 17.4 cases per 1000 in 2019,[30] and *Pv* malaria accounts for
approximately 80% of the malaria burden.[7] Transmission typically peaks in Loreto between
February and July[7] and there is a sizable burden of asymptomatic infections.[18] Asymptomatic
infections typically go undetected and uncounted in routine malaria surveillance in Peru,[18] since
only febrile individuals are tested.
151
From 2005-2010, there was a large effort towards malaria elimination in Peru via the Project for
Malaria Control in Andean Border Areas (PAMAFRO) program that improved case
management, including through deployment of community case workers and distribution of
insecticide-treated bed nets.[7] Transmission declined during the project, but the project did not

achieve elimination, and transmission has since risen.[31] Following PAMAFRO, the government
of Peru adopted Plan Malaria Cero (PMC) in 2017 to achieve malaria elimination by 2030.
Control activities over the study period involved active case detection (ACD), reactive case
detection (RACD), distribution of long-lasting insecticide-treated bed nets (ITN), and indoor
residual spraying[32].
161

Outcome data
The primary outcome was weekly community-level *Pv* malaria incidence primarily measured
through passive case detection conducted by MINSA, though PMC activities including ACD and
RACD also occurred over the study period and were recorded in surveillance data. Both physical
records stored in health posts and online records compiled by MINSA were used in this analysis.
Febrile patients presented for care at a health post and were tested for malaria via microscopy.
According to PMC policy[32], blood smears are stained with 2% Giemsa for 30 minutes. Parasite
densities are calculated from the number of asexual parasites per 200 leukocytes (or per 500, if
<10 asexual parasites/200 leukocytes), assuming a leukocyte count of 8,000/µL. A blood smear
is considered negative if examination of 100 high power fields does not reveal asexual parasites.
Thin smears are used for parasite species identification. Slides are read by two microscopists. If
there are discordant results, the results are determined by a third microscopist. Only *Pv* cases
were included in the present study, and the data did not distinguish between primary infections
versus relapse cases. Cases were matched to community census data to confirm community of
residence and then matched to geocoordinates for the community centroid. Population counts per
community were calculated from a government census conducted from 2017. The FocaL Mass
Drug Administration for Vivax Malaria Elimination (FLAME)[33] trial in Peru (NCT05690841)
conducted a census on 30 of these communities in 2023, which was used to estimate an annual
population growth rate for each year from 2017–2024. Adjusted population counts per year were
estimated by calculating the annual growth rate using the 2023 and 2017 census counts. The
same flat annual growth rate was applied to all 136 communities with a population greater than
15 people in the study area.
184

185 We obtained data for ITN distribution from MINSA with dates of delivery for communities
186 included in the study from 2018–2024. Time since most recent ITN delivery was discretized into
187 never received, 0–1 year, 1–3 years, and more than 3 years.
188

Environmental variables
We obtained temperature and precipitation data from the ERA5-Land Hourly dataset, 11
kilometer resolution, collected by the Copernicus Climate Change Service[34]. We used the air
temperature at 2 meters above the surface (K) band for all temperature variables and the total
precipitation (m) band for precipitation variables. We matched weather variables to incidence
data via community centroid.
195

196   We imputed temperature or precipitation values less than 0 as missing. We aggregated hourly
197   temperature values into the weekly minimum and maximum temperature; and total precipitation
198   observed each week. All data extraction was performed using the Python API for Google Earth
199   Engine on Google Colab servers. Aggregation over time and statistical modeling were performed
200   in R version 4.2.1.
201
202   We obtained data on the Oceanic Niño Index (ONI), a measure of the strength of the El Niño-
203   Southern Oscillation (ENSO), from the National Oceanic and Atmospheric Association National
204   Weather Service Climate Prediction Center. A 1-unit increase in ONI represents a 1ºC higher
205   average sea-surface temperature compared to a 30-year reference period[35].
206
207   We manually designated riverine versus highway communities via Google Earth hybrid satellite
208   image using QGIS version 3.32; we checked our designations with ground-truth observations
209   from 2023 for a subset of communities.
210

Statistical analysis
212   We published a pre-analysis plan at https://osf.io/f89te. Deviations from the pre-analysis plan are
213   listed in Table S1. We removed years 2020 and 2021 from our primary analysis because of
214   strong changes in incidence and case detection resulting from behavior changes and health
215   system capacity during the COVID-19 pandemic.
216
To assess the timing of the association between malaria incidence and weather variables, we fit
distributed lag models. We chose a 2–16-week lookback period (i.e., an infection during the
course of week 20 looked back to week 4, 5, …, 18) to account for variation in the potential
cumulative effect of rainfall and temperature on *Pv* transmission and long-lasting impacts on
relapse cases (Figure 1). For weather exposures (weekly minimum temperature, weekly
maximum temperature, and total precipitation), we fit non-linear distributed lag (DL) models[36]
with a log link and Poisson family using weekly malaria case counts per community as the
dependent variable with an offset for log community population size using the R package DLNM
version 2.4.7[37]. For the distributed lag cross-basis function, we specified a log function with 2
knots to allow for more flexible variation in the short term, and natural splines with 2 degrees of
freedom for the non-linear predictors, which yielded the lowest AIC and BIC in model testing.
All incidence ratios were calculated relative to reference weeks. In all models, we restricted
weather measurements in the input data to observations between the 5th and 95th percentile of the
variable's empirical distribution to remove outlier effects. For temperature predictors, we used
20.5 for minimum temperature and 28.1ºC for maximum temperature as references. These values
were selected because they correspond to the minimum value within the data range during the
study period. For total precipitation, weeks with 0 mm of precipitation were used as the
reference. Temperature variables were adjusted for the cross-basis matrix for total precipitation,
and total precipitation was adjusted for the cross-basis matrix for minimum temperature, in

accordance with AIC/BIC model testing. All DL models included fixed effects for year,
community, and years since ITN distribution, and a smooth term for ONI with 3 knots. All
positive cases were matched to weather data. Thus, we performed a complete case analysis.
Because incidence data for the weather models was aggregated at the community level, we did
not test for spatial autocorrelation or adjust for spatial clustering.

To assess the relationship between each predictor and malaria incidence, we fit generalized
additive models (GAMs) with a Poisson family and log link and an offset for community
population size. We fit models for the week with the strongest association in the DL model
(primary lag). All GAMs included fixed effects for year, community, and years since ITN
delivery and a smooth term for ONI with 3 knots. We calculated confidence intervals using
1,000 bootstrap iterations with replacement.

We conducted subgroup analyses using the primary lag for temperature and precipitation
predictors stratifying by ENSO periods and community type. We defined periods of 5 or more
consecutive months of sea surface temperature abnormality of 0.5℃ or above/-0.5℃ or below as
El Niño/La Niña periods, respectively[35]. To assess interaction effects between weather and
ENSO period or community type on the multiplicative scale, we fit generalized linear models
(GLMs) that included the same covariates from GAM models as fixed effects.

This study was approved by the Institutional Review Board of Stanford University (72291) and
by the Dirección Universitaria de Asuntos Regulatorios de la Investigación de la Universidad
Peruana Cayetano Heredia (211747).

Role of the funding source

We received funding from the National Institute of Allergy and Infectious Disease, GSK, and the
King Center on Global Development. Funding sources did not contribute to the study design,
data collection, analysis, writing, or decision to submit for publication. MH and JBC are Chan-
Zuckerberg Biohub Investigators.

RESULTS

During the study period from January 2017 to December 2024, minimum temperature varied
from 17–25℃, and maximum temperature varied from 28–41℃ (Figure 3A). Total weekly
precipitation varied from 7mm per week to 2,587mm per week (Figure 3B). There were 104 El
Niño weeks and 92 La Niña weeks over the study duration, and incidence varied by ENSO
period, with the highest incidence during neutral periods. Malaria incidence was highest from
2017–2019, then decreased dramatically in 2020 and 2021 and increased to levels below the
2019 levels in 2022–2024. Malaria incidence did not appear to follow a distinct seasonal trend
(Figure 3C). The mean incidence for all communities was 10.4 cases per 10,000 person-weeks.

Temperature

There was a positive association between minimum temperature and malaria incidence at 2–13
weeks lags, and a negative association at 14–16-week lags when comparing the 90[th] percentile of
minimum temperature (23.7ºC) to the lowest value (20.5ºC) (Figure 4A, 4C). Where this
association was highest, at a 7-week lag, the association between minimum temperature and
incidence was examined in a marginal effects plot (Figure 4B). At minimum temperatures of
20.5–22ºC, incidences were similar. At higher temperatures, there was a positive, non-linear
association with incidence of 23 cases per 10,000 at 20.5ºC compared to 37 cases per 10,000 at
23.8ºC. At a 7-week lag, compared to weeks with the lowest minimum temperature, weeks with
a minimum temperature of the 90[th] percentile were associated with up to 9% higher malaria
incidence (95% CI 5%–14%), or incidence ratio of 1.09 (Figure 4A).

The association between maximum temperature and malaria incidence was non-linear and hump-
shaped across lag time; there was a positive association between maximum temperature and
malaria incidence at 4–13-week lags when comparing the 90[th] percentile of maximum
temperature (33.7ºC) to the lowest value (28.1ºC) (Figure 4D, 4F). Where this association was
highest, at a 9-week lag, the association between maximum temperature and incidence was
examined in a marginal effects plot (Figure 4E). Incidence increased from 16 cases per 10,000 to
26 cases per 10,000 as maximum temperature increased from 29.6ºC to 34ºC. Compared to
weeks with the lowest maximum temperature, weeks with a maximum temperature of the 90[th]
percentile were weakly associated with up to 11% higher malaria incidence (95% CI 8%–14%),
or incidence ratio of 1.11 at a 9-week lag (Figure 4D).

Precipitation

The association between precipitation and malaria incidence was non-linear and positive at lags
of 5–16 weeks, when comparing the 90[th] percentile of precipitation (1000mm) to a reference of
no precipitation (Figure 4G, 4I). Where this association was highest, at a 11-week lag, the
association between maximum temperature and incidence was examined in a marginal effects
plot (Figure 4H). Incidence increased from about 17 cases per 10,000 to about 23 cases per
10,000 as precipitation increased from 50mm to 1150mm (Figure 4H). At an 11-week lag,
compared to weeks with no precipitation, weeks with precipitation at the 90th percentile were
associated with up to 28% higher malaria incidence (95 CI 24%–32%), or incidence ratio of 1.28
(Figure 4G).

Community Type Sub-group

Malaria incidence was higher in riverine versus highway-adjacent communities (Table 1, Figure
5A–C). Using GLMs, the association between malaria and minimum temperature was similar by
community type, and there was no evidence of interaction on the multiplicative scale (Table 1).
However, when using GAMs that allowed for non-linear relationships, the association with

incidence and minimum temperature was stronger in riverine versus highway-adjacent
communities at minimum temperatures above 22ºC (Figure 5A). For maximum temperature,
GLMs showed associations were stronger in riverine versus highway-adjacent communities: the
incidence ratio for a 1ºC increase in maximum temperature was 1.13 (95% CI 1.11–1.15) in
riverine communities compared to 1.03 (0.98–1.09) in highway communities (interaction p =
0.001) (Table 1). For precipitation, GLMs showed associations were weaker for riverine versus
highway-adjacent communities: the incidence ratio for a 500mm increase in precipitation was
1.06 (95% CI 1.02–1.09) in riverine communities compared to 1.58 (95% CI 1.44–1.74) in
highway communities (interaction $p < 0.001$) (Table 1).
325

ENSO Sub-group
Malaria incidence was lower during El Niño periods compared to La Niña and neutral periods
(Table 2, Figure 5D–F). Using GLMs, associations between minimum temperature and malaria
incidence were positive when stratifying by ENSO period and strongest during neutral periods.
The incidence ratio for a 1ºC increase in minimum temperature was 1.34 (95% CI 1.29–1.40)
during neutral periods, 1.28 (95% CI 1.21–1.36) during El Niño periods (interaction $p = 0.217$),
and 1.03 (95% CI 0.98–1.09) during La Niña periods (interaction $p < 0.001$) (Table 2).
333

334 Associations between maximum temperature and malaria incidence were positive when
335 stratifying by ENSO period and strongest during neutral periods. The incidence ratio for a 1ºC
336 increase in maximum temperature was 1.14 (95% CI 1.11–1.18) during neutral periods, 1.11
337 (95% CI 1.07–1.15) during El Niño periods (interaction $p = 0.195$), and 1.06 (95% CI 1.02–1.09)
338 during La Niña periods (interaction $p < 0.001$) (Table 2).
339

340 Associations between total precipitation and malaria incidence were positive when stratifying by
341 ENSO period but strongest during La Niña periods. The incidence ratio for a 500mm increase in
342 precipitation was 1.07 (95% CI 1.01–1.12) during neutral periods, 1.06 (95% CI 0.99–1.14)
343 during El Niño periods (interaction $p = 0.944$), and 1.17 (95% CI 1.10–1.24) during La Niña
344 periods (interaction $p = 0.029$) (Table 2).
345

DISCUSSION
Higher temperatures were associated with modestly higher *Pv* malaria incidence, and total
precipitation was associated with moderately higher incidence in an elimination setting in Peru.
Associations generally lasted several weeks, with more sustained, stronger associations for
precipitation. We only observed associations between malaria incidence and minimum
temperatures above 22ºC, suggesting that cooler minimum temperatures may be a limiting factor
for *Pv* malaria incidence. Incidence was lowest during El Niño periods, which are typically dry
in this region, and higher in wetter La Niña periods. The strength of associations was stronger
during neutral periods for temperature predictors and stronger during La Niña periods for

precipitation. Overall, periods of higher temperatures and cumulative rainfall were associated
with increased malaria burden in the subsequent several weeks.

Our findings suggest that weather influences malaria transmission over long, 1–4-month lags in
this region. This lasting association may reflect both initial infections and the relapse periodicity
of *Pv* malaria. The efficacy of standard of care anti-relapse treatment is approximately 75-80%,[38]
but effectiveness can be significantly lower due to challenges with adherence.[39,40] Further, many
infections go undetected and untreated due to healthcare access challenges or lack of or minimal
symptoms. As such, the vast majority of *Pv* infections are estimated to be relapse infections,[41,42]
and incidence measured in this study included both new and relapse infections. Additionally,
relapses from the tropical strains of *Pv* are estimated to range anywhere from 2 weeks[4] to 2–9
months,[21] supporting our finding. The extended association with weather over many weeks could
also reflect a high prevalence of asymptomatic and low parasitemia infections[4,5], which could
sustain new, onward transmission well past the initial meteorological trigger. Another study in
this region found that PCR-confirmed *Pv* prevalence was as high as 25%, most of which were
attributed to asymptomatic, low parasitemia infections.[43]

Precipitation

We found that higher precipitation was significantly associated with higher malaria incidence at
lags of 5–16 weeks. These results are generally in accordance with other studies of the effect of
rainfall on *Pv* infections. A meta-analysis in Mauritania found that *Pv* incidence was highest
during and after the rainy season[44] and that decreased rainfall was significantly correlated with
decreased malaria burden; a study in South Korea, a temperate area, found that increased
precipitation was associated with higher malaria incidence at a 10-week lag,[45] while a similar
study in a tropical area of China also found positive associations with precipitation at lags 2–4
weeks.[46] The lasting influence of precipitation likely reflects increased breeding ground
following rainfall, which could impact multiple transmission cycles. The 5–16 week lag may
also include the transition to the dry season, when the vector can still thrive in moist
environments but flying and biting behavior is uninhibited by falling rain.

Temperature

We found that higher minimum temperatures were associated with higher malaria incidence from
4–11-week lags. This finding has mixed support in the literature: though the thermal optimum of
the *Pv–Ny. darlingi* coupling has not been studied directly, *Pf* malaria transmission peaks at
25°C,[24,47] and minimum temperatures in our study were in the range of 20–24°C. Our finding of
the strongest associations with minimum temperature is also biologically plausible given that *Ny.*
darlingi typically bite humans at nighttime,[48] and daily temperatures are lowest at night.
However, a study of *Pv*-dominant Amazon border regions found that *Pv* incidence was
negatively associated with minimum temperatures between 17–25°C at 0- and 6-month lags, with
the stronger association at 6 months[49]. We attribute the difference at 0 months to divergent model

structures and lookback periods. We also found a negative association at 16 weeks, suggesting
there may be negative associations at longer lags that we did not explore due to our focus on the
faster time to relapse common in our study site.
Our finding that higher maximum temperatures in the range of 30–34ºC were associated with
increased incidence was surprising because it conflicts with findings in neighboring Brazil[28] and
some previous disease ecology studies. One simple explanation is that during hotter
temperatures, people may spend more time outdoors, or spend less time under bednets or in
screened areas. However, this explanation does not address the well-documented thermal limits
on vector behavior. Laboratory studies of *Pf* malaria have found that the thermal optimum for
Nyssorhynchus is at 25ºC with an upper limit of 32.6ºC,[24] and empirical studies have found that
Pf malaria transmission declines above 28ºC[47] in Africa. However, a study in *Pv*-dominant
neighboring regions of the Amazon with similar maximum temperatures (26.8–35.2ºC) found
that a 1ºC increase in maximum temperature was associated with higher *Pv* incidence at 1- and
2-month lags, consistent with our findings.[50] It is generally thought that *Anopheles gambiae* and
Anopheles stephensi biting rate, larva survival, fecundity, adult survival decline at the maximum
temperature range in our study (30–34ºC)[25], and a laboratory study of *Nyssorhynchus darlingi* in
Brazil found that larval development, adult lifespan, and body size all declined as temperatures
increased from 20ºC to 28ºC though there was some geographic variation.[51] It is possible that
microclimates, topographical variation, and forest cover tempered high temperatures in our study
site, and such fine scale phenomena would not have been accounted for given that we used
temperature data with 11km resolution. Alternatively, the vector in our study sites may be
adapted to higher maximum temperatures,[52] preventing population decline.

Another possible explanation is that associations with maximum temperature reflect relapsing *Pv*
infections, which could be triggered by temperature itself, physiological conditions, or co-
infections that are more common at higher temperatures in this setting. Extrinsic triggers of *Pv*
relapse may include untreated *P. falciparum*[53] or other co-infections that result in host
inflammation, subsequent primary *Pv* infections, seasonal changes in sunlight and temperature,
and mosquito bites and their associated immune responses.[26] In our study site, *P. falciparum* is
uncommon, though dengue is increasingly common,[54] and it is well-documented that dengue
transmission is positively associated with maximum temperatures similar to those in our study[55–
57] and has a higher thermal optimum (29ºC) and upper limit (34.5ºC) than malaria.[24] Co-
infections with dengue could thus occur at higher temperatures and lead to relapsing *Pv*
infections. It is also possible that higher temperatures lead to host inflammation and heat stress
that activate the hypnozoite and increase the reservoir of active infectious hosts, potentially
leading to onward transmission and multiple cycles of infection, explaining the delayed positive
association.

ENSO sub-group analyses

We found that incidence was generally lowest during El Niño periods compared to neutral or La
Niña periods, when incidence was similar. There is mixed support for this finding in the
literature. One study in neighboring Colombia found that caseloads were higher during El Niño
events[58], possibly because higher temperatures and lower rainfall lead to decreased river
discharge and water stagnation, creating an optimal malaria breeding ground. A study in the
Brazilian Amazon found that incidence was lower during El Niño and La Niña periods[28],
somewhat contradicting our results. The rivers in our study region tend to be shallow and small,
and some dry up completely during strong El Niño periods, such as in August 2023.
Additionally, the Amazon has been undergoing historic drought due to global climate change[16],
with 2023 being a particularly pronounced drought year. Climate change may be dramatically
altering the effects of weather independently and on a longer time scale than any individual
ENSO period studied here. Drought brought on by El Niño and climate change may decrease the
availability of stagnant pools of water necessary for mosquito breeding, explaining the overall
lowest incidence during El Niño periods.
448
Additionally, temperatures were highest during El Niño periods, potentially limiting mosquito
activity. On the other hand, because La Niña periods are associated with wetter conditions,[59] in
the context of historic drought that has dried up rivers in our study region, increases in rainfall
could potentially create temporary pools of stagnant water, explaining the higher incidence
during these periods in our study region. We also found evidence of interaction between La Niña
periods and weather predictors; associations between malaria incidence and precipitation were
strongest during La Niña periods, while associations between malaria incidence and temperature
were weakest during La Niña periods. These interaction effects suggest that variation in
precipitation drives malaria transmission during La Niña periods, while during normal and El
Niño periods, temperature largely drives transmission.
459

Community type sub-group analysis
Incidence was higher in river- versus highway-adjacent communities, and the association with
maximum temperature was also stronger in riverine communities. On the other hand, the
association with precipitation was stronger in highway-adjacent communities. It is possible that
rainfall is more likely to create stagnant pools of water in highway-adjacent communities than
riverine communities, where water bodies are larger and rainfall may result in a more modest
increase in ideal water bodies for mosquito breeding. Conversely, in river-adjacent communities
where mosquitoes can breed uninhibited, temperature may be more limiting, explaining the
stronger association with maximum temperature in these communities. In addition, higher
vegetation near rivers may create cooler microclimates that approximate the malaria thermal
optimum better than more urbanized highway communities, which may be hotter.
471

Limitations

Our study had several limitations. We were unable to distinguish between initial and relapse
infections in our incidence data; we would expect that weather would influence initial and
relapse cases over different lag periods, but our analysis was not able to investigate this.
However, using distributed lag models allowed us to investigate associations over a 4-month
period, including potential relapse cases. We were also unable to identify which cases were
identified via passive surveillance versus RACD/ACD, which were done as malaria control
efforts by the Peruvian Ministry of Health, greatly affecting case detection and potentially
altering our results. However, these were done as a reaction to higher caseloads, suggesting that
passive surveillance-detected incidence was already higher when case detection was done. A
follow-up study using genomic methods to differentiate relapse and initial infections could
elucidate how weather influences primary versus relapse infections. The study period also
included two years (2021 and early 2022) where control efforts for the COVID-19 pandemic
likely limited malaria transmission as well.
486

While we selected ERA-5 Land remote sensing data for its temporal coverage, its 11 km spatial
resolution prevented identification of small-scale microclimates, such as those created by forest
cover and topographical features, and thus did not capture small-resolution variation that may
have a large effect on mosquito breeding and survival habitat.[60] Additionally, publicly available
surface water data did not reflect our ground observations of surface water in the study site, so
we did not include it in this analysis. Further research that considers mediation by surface water
help determine the complicated dynamics between surface water, drought, and malaria incidence
in this region. While we initially considered land cover variables, they were limited in statistical
power. Further research could confirm relationships between land cover, deforestation, and
malaria incidence in this region and how that is mediated by climate change and drought. This
study was also correlational, and inferred relationships with individual weather variables may be
confounded by collinearity and correlation with other weather variables and thus made it difficult
to tease out individual direct effects.
500

Conclusions
In our study of malaria incidence in Amazonian Peru, we observed generally positive
associations with higher temperatures and higher rainfall for extended lag periods beginning 4–5
weeks after the initial weather event and enduring for 1–4 months. We also found lower
incidence during El Niño years and higher incidence in riverine communities, which may be
used to concentrate resources and time malaria interventions to achieve elimination in this
setting. Our findings indicated that the coupled transmission and relapse cycle of *Pv–Ny. darlingi*
may have more complicated associations with higher temperatures than other malaria parasite –
vector pairings, a critical finding and target for future research in the face of climate change and
global warming. These findings provide critical context to ongoing malaria elimination efforts in
the Amazon region, since apparent successes or failures of malaria interventions may be due in
part to long-lasting effects of weather on initial infection and relapse.

REFERENCES

1 Kattenberg JH, Erhart A, Truong MH, *et al.* Characterization of Plasmodium falciparum and
Plasmodium vivax recent exposure in an area of significantly decreased transmission
intensity in Central Vietnam. *Malaria Journal* 2018; **17**: 180.

2 Global technical strategy for malaria 2016–2030, 2021 update. 2021.

3 Hulden L, Hulden L. Activation of the hypnozoite: a part of Plasmodium vivax life cycle and
survival. *Malar J* 2011; **10**: 90.

4 Olliaro PL, Barnwell JW, Barry A, *et al.* Implications of Plasmodium vivax Biology for Control,
Elimination, and Research. *Am J Trop Med Hyg* 2016; **95**: 4–14.

5 da Silva-Nunes M, Moreno M, Conn JE, *et al.* Amazonian malaria: Asymptomatic human
reservoirs, diagnostic challenges, environmentally-driven changes in mosquito vector
populations, and the mandate for sustainable control strategies. *Acta Trop* 2012; **121**: 281–
91.

6 Howes RE, Battle KE, Mendis KN, *et al.* Global Epidemiology of Plasmodium vivax. *Am J*
Trop Med Hyg 2016; **95**: 15–34.

7 Rosas-Aguirre A, Gamboa D, Manrique P, *et al.* Epidemiology of Plasmodium vivax Malaria
in Peru. *Am J Trop Med Hyg* 2016; **95**: 133–44.

8 Ferreira MU, Gamboa D, Torres K, *et al.* Evidence-Based Malaria Control and Elimination in
the Amazon: Input from the International Center of Excellence in Malaria Research Network
in Peru and Brazil. *Am J Trop Med Hyg* 2022; **107**: 160–7.

9 Carrasco-Escobar G, Gamboa D, Castro MC, *et al.* Micro-epidemiology and spatial
heterogeneity of P. vivax parasitaemia in riverine communities of the Peruvian Amazon: A
multilevel analysis. *Sci Rep* 2017; **7**: 8082.

10 Rosas-Aguirre A, Guzman-Guzman M, Gamboa D, *et al.* Micro-heterogeneity of malaria
transmission in the Peruvian Amazon: a baseline assessment underlying a population-based
cohort study. *Malar J* 2017; **16**: 312.

11 da Silva-Nunes M, Moreno M, Conn JE, *et al.* Amazonian malaria: asymptomatic human
reservoirs, diagnostic challenges, environmentally driven changes in mosquito vector
populations, and the mandate for sustainable control strategies. *Acta Trop* 2012; **121**: 281–
91.

12 Rufalco-Moutinho P, Moura Kadri S, Peres Alonso D, *et al.* Ecology and larval population
dynamics of the primary malaria vector Nyssorhynchus darlingi in a high transmission setting
dominated by fish farming in western Amazonian Brazil. *PLoS One* 2021; **16**: e0246215.

13 Parker BS, Paredes Olortegui M, Peñataro Yori P, *et al.* Hyperendemic malaria transmission
in areas of occupation-related travel in the Peruvian Amazon. *Malar J* 2013; **12**: 178.

14 MacDonald AJ, Mordecai EA. Amazon deforestation drives malaria transmission, and malaria
burden reduces forest clearing. *Proceedings of the National Academy of Sciences* 2019; **116**:
22212–8.

15 Cai W, McPhaden MJ, Grimm AM, *et al.* Climate impacts of the El Niño–Southern Oscillation
on South America. *Nat Rev Earth Environ* 2020; **1**: 215–31.

16 Espinoza J-C, Jimenez JC, Marengo JA, *et al.* The new record of drought and warmth in the
Amazon in 2023 related to regional and global climatic features. *Sci Rep* 2024; **14**: 8107.

17 Gething PW, Van Boeckel TP, Smith DL, *et al.* Modelling the global constraints of
temperature on transmission of Plasmodium falciparum and P. vivax. *Parasites Vectors*
2011; **4**: 92.

18 Gilman RH, Cabrera L, Lee E, *et al.* ENDEMIC MALARIA IN THE PERUVIAN AMAZON
REGION OF IQUITOS. *The American Journal of Tropical Medicine and Hygiene* 2003; **69**:
45–52.

19 Stratman-Thomas WK. The Influence of Temperature on Plasmodium Vivax. *The American*
Journal of Tropical Medicine and Hygiene 1940; **s1-20**: 703–15.

20 Suh E, Grossman MK, Waite JL, *et al.* The influence of feeding behaviour and temperature
on the capacity of mosquitoes to transmit malaria. *Nat Ecol Evol* 2020; **4**: 940–51.

21 White NJ. Determinants of relapse periodicity in Plasmodium vivax malaria. *Malar J* 2011; **10**:
297.

22 Sadoine ML, Smargiassi A, Ridde V, Tusting LS, Zinszer K. The associations between
malaria, interventions, and the environment: a systematic review and meta-analysis. *Malar J*
2018; **17**: 73.

23 Hiwat H, Bretas G. Ecology of Anopheles darlingi Root with respect to vector importance: a
review. *Parasites Vectors* 2011; **4**: 177.

24 Mordecai EA, Caldwell JM, Grossman MK, *et al.* Thermal biology of mosquito-borne disease.
Ecol Lett 2019; **22**: 1690–708.

25 Villena OC, Ryan SJ, Murdock CC, Johnson LR. Temperature impacts the environmental
suitability for malaria transmission by Anopheles gambiae and Anopheles stephensi. *Ecology*
2022; **103**: e3685.

26 Schäfer C, Zanghi G, Vaughan AM, Kappe SHI. *Plasmodium vivax* Latent Liver Stage
Infection and Relapse: Biological Insights and New Experimental Tools. *Annu Rev Microbiol*
2021; **75**: 87–106.

27 Colonia CB, Vásquez-Rodríguez AB, Alexander N, de la Hoz Restrepo F. Malaria,
relationship with climatic variables and deforestation in Colombia, Latin America and the
Caribbean from 2000 to 2020: a systematic review. *Malar J* 2024; **23**: 347.

28 Arisco NJ, Peterka C, Schwartz J, Castro MC. The impact of weather and extreme events on
malaria transmission in the Brazilian Amazon: a case-crossover and population-based study.
The Lancet Regional Health – Americas 2025; **49**. DOI:10.1016/j.lana.2025.101189.

29 Paaijmans KP, Blanford S, Bell AS, Blanford JI, Read AF, Thomas MB. Influence of climate
on malaria transmission depends on daily temperature variation. *Proceedings of the National*
Academy of Sciences 2010; **107**: 15135–9.

30 malaria.pdf. https://www.dge.gob.pe/portal/docs/vigilancia/sala/2022/SE42/malaria.pdf
(accessed May 2, 2024).

31 Janko MM, Recalde-Coronel GC, Damasceno CP, *et al.* The impact of sustained malaria
control in the Loreto region of Peru: a retrospective, observational, spatially-varying
interrupted time series analysis of the PAMAFRO program. *The Lancet Regional Health –*
Americas 2023; **20**. DOI:10.1016/j.lana.2023.100477.

32 189846_RM_244-2017-MINSA.PDF20180823-24725-1yk3i0y.pdf.
https://cdn.www.gob.pe/uploads/document/file/190350/189846_RM_244-2017-
MINSA.PDF20180823-24725-1yk3i0y.PDF (accessed Nov 21, 2024).

33 Fine S, Quiroz AA, Calle VS, *et al.* FocaL mass drug Administration for Plasmodium vivax
Malaria Elimination (FLAME): study protocol for an open-label cluster randomized controlled
trial in Peru. *Res Sq* 2025; : rs.3.rs-5594891.

34 Muñoz-Sabater J, Dutra E, Agustí-Panareda A, *et al.* ERA5-Land: a state-of-the-art global
reanalysis dataset for land applications. *Earth System Science Data* 2021; **13**: 4349–83.

35 Center NCP. NOAA's Climate Prediction Center.
https://origin.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ONI_v5.php
(accessed Aug 12, 2025).

36 Gasparrini A, Armstrong B, Kenward MG. Distributed lag non-linear models. *Stat Med* 2010;
29: 2224–34.

37 Gasparrini A. Distributed Lag Linear and Non-Linear Models in R: The Package dlnm. *J Stat*
Softw 2011; **43**: 1–20.

38 Llanos-Cuentas A, Lacerda MVG, Hien TT, *et al.* Tafenoquine versus Primaquine to Prevent
Relapse of Plasmodium vivax Malaria. *New England Journal of Medicine* 2019; **380**: 229–41.

39 Grietens KP, Soto V, Erhart A, *et al.* Adherence to 7-day primaquine treatment for the radical
cure of P. vivax in the Peruvian Amazon. *Am J Trop Med Hyg* 2010; **82**: 1017–23.

40 Poespoprodjo JR, Burdam FH, Candrawati F, *et al.* Supervised versus unsupervised
primaquine radical cure for the treatment of falciparum and vivax malaria in Papua,
Indonesia: a cluster-randomised, controlled, open-label superiority trial. *The Lancet Infectious*
Diseases 2022; **22**: 367–76.

41 Popovici J, Tebben K, Witkowski B, Serre D. Primaquine for *Plasmodium vivax* radical cure:
What we do not know and why it matters. *International Journal for Parasitology: Drugs and*
Drug Resistance 2021; **15**: 36–42.

42 Commons RJ, Simpson JA, Watson J, White NJ, Price RN. Estimating the Proportion of
Plasmodium vivax Recurrences Caused by Relapse: A Systematic Review and Meta-
Analysis. 2020; published online June 8. DOI:10.4269/ajtmh.20-0186.

43 Carrasco-Escobar G, Miranda-Alban J, Fernandez-Miñope C, *et al.* High prevalence of very-
low Plasmodium falciparum and Plasmodium vivax parasitaemia carriers in the Peruvian
Amazon: insights into local and occupational mobility-related transmission. *Malar J* 2017; **16**:
415.

44 Moustapha IE, Moukah MO, Salem MSOA, *et al.* Malaria prevalence in Mauritania: a
systematic review and meta-analysis. *Malaria Journal* 2023; **22**: 146.

45 Kim Y-M, Park J-W, Cheong H-K. Estimated Effect of Climatic Variables on the Transmission
of Plasmodium vivax Malaria in the Republic of Korea. *Environ Health Perspect* 2012; **120**:
1314–9.

46 Bi Y, Yu W, Hu W, *et al.* Impact of climate variability on Plasmodium vivax and Plasmodium
falciparum malaria in Yunnan Province, China. *Parasit Vectors* 2013; **6**: 357.

47 Mordecai EA, Paaijmans KP, Johnson LR, *et al.* Optimal temperature for malaria
transmission is dramatically lower than previously predicted. *Ecology Letters* 2013; **16**: 22–
30.

48 Moreno M, Saavedra MP, Bickersmith SA, *et al.* Implications for changes in Anopheles
darlingi biting behaviour in three communities in the peri-Iquitos region of Amazonian Peru.
Malar J 2015; **14**: 290.

49 Wangdi K, Wetzler E, Cox H, Marchesini P, Villegas L, Canavati S. Spatial patterns and
climate drivers of malaria in three border areas of Brazil, Venezuela and Guyana, 2016-2018.
Sci Rep 2022; **12**: 10995.

50 Wangdi K, Wetzler E, Cox H, Marchesini P, Villegas L, Canavati S. Spatial patterns and
climate drivers of malaria in three border areas of Brazil, Venezuela and Guyana, 2016–
2018. *Scientific Reports* 2022; **12**. DOI:10.1038/s41598-022-14012-4.

51 Chu VM, Sallum M a. M, Moore TE, Lainhart W, Schlichting CD, Conn JE. Regional variation
in life history traits and plastic responses to temperature of the major malaria vector
Nyssorhynchus darlingi in Brazil. *Sci Rep* 2019; **9**: 5356.

52 Couper LI, Nalukwago DU, Lyberger KP, Farner JE, Mordecai EA. How Much Warming Can
Mosquito Vectors Tolerate? *Global Change Biology* 2024; **30**: e17610.

53 Commons RJ, Simpson JA, Thriemer K, *et al.* Risk of Plasmodium vivax parasitaemia after
Plasmodium falciparum infection: a systematic review and meta-analysis. *The Lancet
Infectious Diseases* 2019; **19**: 91–101.

54 Munayco CV. Notas Desde el Campo: Brote de Dengue — Perú, 2023. *MMWR Morb Mortal
Wkly Rep* 2024; **73**. https://www.cdc.gov/mmwr/volumes/73/wr/mm7304a4_ensp.htm
(accessed Oct 27, 2024).

55 López MS, Gómez AA, Müller GV, Walker E, Robert MA, Estallo EL. Relationship between
Climate Variables and Dengue Incidence in Argentina. *Environmental Health Perspectives*
2023; **131**: 057008.

56 Fan J, Wei W, Bai Z, *et al.* A Systematic Review and Meta-Analysis of Dengue Risk with
Temperature Change. *International Journal of Environmental Research and Public Health*
2015; **12**: 1–15.

57 Hii YL, Rocklöv J, Ng N, Tang CS, Pang FY, Sauerborn R. Climate variability and increase in
intensity and magnitude of dengue incidence in Singapore. *Global Health Action* 2009; **2**:
2036.

58 Poveda G, Rojas W, Quiñones ML, *et al.* Coupling between annual and ENSO timescales in
the malaria-climate association in Colombia. *Environmental Health Perspectives* 2001; **109**:
489–93.

59 Moura MM, dos Santos AR, Pezzopane JEM, *et al.* Relation of El Niño and La Niña
phenomena to precipitation, evapotranspiration and temperature in the Amazon basin.
Science of The Total Environment 2019; **651**: 1639–51.

60 Carrasco-Escobar G, Manrique E, Ruiz-Cabrejos J, *et al.* High-accuracy detection of malaria
vector larval habitats using drone-based multispectral imagery. *PLoS Negl Trop Dis* 2019; **13**:
e0007105.

| Malaria Incidence per 10,000 person−weeks | 0 | 30 | 60 | 90 | Community Type | □ Highway | ○ Riverine |

Minimum Temperature

A) 90th percentile vs. min

B) Association at 7 Weeks

C) Predictor–Lag Surface

Maximum Temperature

D) 90th percentile vs. min

E) Association at 9 Weeks

F) Predictor–Lag Surface

Total Precipitation

G) 90th percentile vs. 0

H) Association at 11 Weeks

I) Predictor–Lag Surface

Community type	Predictor range	Incidence range	Incidence Ratio (95% CI)	Interaction p-value
Minimum temperature				
Highway	20.5–23.9	4.2–8.0	1.17 (1.08–1.28)	--
Riverine	20.4–23.8	23.2–40.0	1.27 (1.23–1.31)	0.097
Maximum temperature				
Highway	29.5–33.9	2.8–5.7	1.03 (0.98–1.09)	--
Riverine	29.6–34.0	19.0–29.5	1.13 (1.11–1.15)	0.001
Precipitation				
Highway	51–1139	3.6–7.7	1.58 (1.44–1.74)	--
Riverine	57–1161	17.7–23.3	1.06 (1.02–1.09)	<0.001

ENSO period	Predictor range	Incidence range	Incidence Ratio (95% CI)	Interaction p-value
Minimum temperature				
Normal	19.9–23.7	17.4–32.4	1.34 (1.29–1.40)	--
El Niño	21.4–24.2	3.5–13.6	1.28 (1.21–1.36)	0.217
La Niña	20.3–23.5	25.3–32.0	1.03 (0.98–1.09)	<0.001
Maximum temperature				
Normal	29.5–33.7	18.3–24.2	1.14 (1.11–1.18)	--
El Niño	30.0–34.4	6.6–11.3	1.11 (1.07–1.15)	0.195
La Niña	29.4–33.6	35.0–39.9	1.06 (1.02–1.09)	<0.001
Precipitation				
Normal	32–1103	24.0–28.5	1.07 (1.01–1.12)	--
El Niño	87–1267	8.3–12.6	1.06 (0.99–1.14)	0.944
La Niña	94–1035	24.7–35.5	1.17 (1.10–1.24)	0.029

ENSO period	Predictor range	Incidence range	Incidence Ratio (95% CI)	Interaction p-value