Seasonal variation in SARS-CoV-2 transmission in the Netherlands, 2020-2022: statistical evidence for an inverse association with solar radiation and temperature.

Don Klinkenberg[1], Jantien Backer[1], Chantal Reusken[1,2], Jacco Wallinga[1,3]

1. Center for Infectious Disease Control, National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands

2. Department of Viroscience, Erasmus MC, Rotterdam, the Netherlands

3. Department of Biomedical Data Sciences, Leiden University Medical Centre, Leiden, the Netherlands

Corresponding Author: Don Klinkenberg, RIVM-EPI, Antonie van Leeuwenhoeklaan 9, 7321 MA Bilthoven, The Netherlands; +31 88 689 8989 ; don.klinkenberg@rivm.nl

ACKNOWLEDGEMENTS

We thank all collaborators in the Data Analytics, Research, and Automated Reporting (DARA) team in the Center for Infectious Disease Control for processing and curating all COVID-19 surveillance data. We thank Dr Scott McDonald for valuable discussions on time series analyses.

No additional funding was received for this study.

Data and code: all data and code are available online at github.com/rivm-syso/covid-seasonality

Keywords: Infectious Disease Transmission; Seasons; Weather; SARS-CoV-2; Respiratory Tract Infections

ABSTRACT

In temperate regions, respiratory viruses such as SARS-CoV-2 are better transmitted in winter than in summer. Understanding how the weather is associated with SARS-CoV-2 transmissibility can enhance projections of COVID-19 incidence and improve estimation of the effectiveness of control measures. During the pandemic, transmissibility was tracked by the reproduction number R_t. This study aims to determine whether information about the daily temperature, absolute humidity, and solar radiation improves predictions of R_t in the Netherlands from 2020 to 2022, and to quantify the relationship between R_t and daily weather data. We conducted a regression analysis, accounting for immunity from vaccination and previous infection, higher transmissibility of new variants, and changes in contact behaviour due to control measures. Results show a linear association between $\log R_t$ and daily solar radiation and temperature, indicating a ratio of R_t in Winter versus Summer of 1.7 (95% CI, 1.4-2.1). The possibility that this association arises from unrelated seasonal patterns was dismissed, as weather data from earlier years provided poorer fits with only small effect sizes. This suggests a causal relationship of solar radiation and temperature with SARS-CoV-2 transmissibility, enhancing confidence in using this relationship for short-term predictions and other epidemiological analyses.

BACKGROUND

Many infectious diseases display seasonal patterns in incidence caused by seasonality in exposure (e.g. Lyme disease, Q-fever) or seasonality in contact rates (e.g. measles),[1] as well as by other mechanisms. With many pathogens, associations of transmission potential with weather variables have been found, mainly temperature and humidity.[1-3] Associations with weather variables are especially pronounced for respiratory viruses, where the causal mechanism may be related to the pathogen (e.g. virus stability related to temperature, or better persistence in drier air related to aerosole size),[4-6] the individual host (amount and duration of shedding, or antiviral defense)[4, 6] or host-host interaction (e.g. people spending less time indoors during warm and sunny weather, where most transmission takes place).[5, 7, 8]

Already early in the COVID-19 pandemic a relation between weather variables and SARS-CoV-2 transmission has been hypothesized and tested, to assess the risk for a large second wave by the end of 2020 in northern temperate regions.[9-11] Many studies focused on the transmission potential, which can be expressed as the time-dependent reproduction number R_t, defined as the mean number of secondary cases infected per primary case. Large-scale studies across many geographical locations showed associations of R_t with temperature and humidity, with often small and non-linear effect sizes. A study among 409 cities in 26 countries worldwide found a maximum R_t at a temperature around 10˚C and at an absolute humidity around 6.6 g/m^3, with effect sizes of 0.087 and 0.061 compared to minimal values at 20˚C and 11 g/m^3, respectively.[12] In a multi-county analysis in mainland USA, almost 20% of variation in R_t was explained by temperature, humidity and ultraviolet radiation, with low R_t values when the weather is hot, humid, and sunny. Overall, the effects were concentrated at the extreme ends of the ranges of the variables. For instance, R_t was largest at a temperature of 20˚C, by about 5% compared to any other temperate under 30˚C; and specific humidity affected transmissibility only at very low or very high values, suggesting that the model better explained spatial than seasonal variation in R_t.[13] In a study in 47 prefectures in Japan, the reproduction number R was found to decrease with about 6% when comparing temperatures of 0˚C vs 30˚C, and with 5% when absolute humidity increased to 10g/m^3 or higher.[14] In general, across many studies, temperature was found to be inversely associated with SARS-CoV-2 transmission, whereas the association with humidity was unclear.[15] In Europe, SARS-CoV-2 transmission was most often found to be inversely related to temperature, humidity, and solar radiation.[5]

The impact of weather on SARS-CoV-2 transmission has significant implications for epidemiological analyses, such as forecasting incidence by dynamic transmission models or estimating the effects of sets of control measures. To account for a weather effect in such analyses, it should be properly and independently quantified for the target population at hand. This requires the target population to be sufficiently large, so that the epidemic is driven by transmission events within the population, affected by the local weather, instead of reflecting epidemics in other locations by many import cases. It also requires the epidemic in the target population to have a sufficiently uniform climate, because the mechanisms by which weather may impact transmissibility are not self-evident, and may be different in different climates. The importance of focussing on single target populations is further illustrated by the fact that estimated effect sizes for smaller populations tend to be large , e.g. 46% higher R_t in winter compared to summer explained by temperature and absolute humidity in France,[16] or 22% in London and Paris,[17] whereas those from larger populations are smaller or reflect spatial rather than seasonal variation.[13, 14]

In this study, we estimated the association between SARS-CoV-2 transmissibility and temperature,
absolute humidity, and solar radiation during 25 months of the pandemic in the Netherlands. We
quantified this relationship and estimated the relative change in reproduction number in winter vs
summer resulting from the association. We assessed whether the estimated association can be
attributed to the general seasonal trend of temperature, absolute humidity, and solar radiation, or to
the specific course of the weather during the pandemic.

95
96
97
98   METHODS
99
We used the log of the daily *instantaneous* reproduction number R_t, which was estimated during the
pandemic by the National Institute of Public Health and the Environment (RIVM) in the Netherlands
(Figure 1a),[18, 19] and the variance of the estimator. It was estimated by dividing the symptom onset
incidence at time t + 5 (incubation period is estimated to be 5 days) by the number of infectious
individuals, i.e. the symptom onset incidence in the days up to t + 5 weighted by the serial interval
distribution.[20] Because of autocorrelation and day-of-week effects in the time series of daily
estimated R_t, we only used the time series of $\log R_t$ estimates of Fridays (Figure 1a, see supplementary
material S1). The observation period was from 22 February 2020 until 29 March 2022, after which R_t
estimates became less reliable due to changes in case notification policy.
109
110   Weekly genomic surveillance data were obtained from RIVM's open data.[21] They were used to
111   estimate $\log R_t^{\text{var}}$, i.e. the relative mean transmissibility at time t, caused by the circulating variants
112   Alpha, Beta, Gamma, and Delta, relative to the wildtype variant (Figure 1b, see supplementary material
113   S2).[22] After the emergence of Delta, $\log R_t^{\text{var}}$ was kept constant, because the emergence of Omicron in
114   December 2021 was for an important part due to immune escape.
115
Cumulative incidence CumIncid(t) per day of the pandemic was calculated by RIVM, by combining
serological data and daily hospitalisation data, and previously used to calculate the prevalence of
infected individuals.[23] It was available until 2 July 2021, after which CumIncid(t) was kept constant,
because waning immunity and re-infections with new variants made estimation unreliable. Vaccine
coverage data were obtained from RIVM's open data.[24] We used the weekly national coverage of the
completed primary vaccination, interpolated to obtain daily coverages. To calculate the rise of vaccine-
induced immunity VaccImm(t) in the population, the coverage was multiplied by 75% vaccine
effectiveness as estimated against the Delta variant,[25] because the primary vaccination series was
quickly followed by the rise of the Delta variant. Cumulative incidence and vaccine-induced immunity
were combined to calculate the times series of the susceptible fraction of the population as
$s_t = \left(1 - \text{CumIncid}(t)\right) \cdot \left(1 - \text{VaccImm}(t)\right)$ (Figure 1b). Note that s_t only reflects the decrease in
susceptibilty until the Summer of 2021, after which waning immunity and the rise of the Omicron
variant made reliable estimation impossible.
129
130

Figure 1. The time course of all datasets. (A) the response variable $\log R_t$, with the 95% confidence interval indicating the uncertainty of the estimator, the inverse of which is used to weigh the observations; (B) the offsets due to increased transmissibility from new variants, and decreased transmissibility from immunity due to infection and vaccination; (C) the mean weekly temperature during the pandemic and in the years 2000-2019; (D) the mean weekly absolute humidity during the pandemic and in the years 2000-2019; (E) the weekly mean of daily solar radiation during the pandemic and in the years 2000-2019.

The time series (109 weeks) of weekly reproduction numbers from 22 February 2020 to 29 March 2022
was partitioned into 41 intervention periods (see supplementary material S3), separated by changes in
COVID-19 control measures or the start or end of school holidays. After a preliminary analysis for outlier
detection based on Cook's distance (supplementary material S1),[26] the period between 26 June 2021

and 9 July 2021 was split into two separate weeks for the main analyses. This was a period when bars
and nightclubs were allowed to open, causing a short sudden increase in R_t (Figure 1a) followed by a
rapid surge in new cases and closure of bars and nightclubs in the period thereafter.

We used the weekly averaged (Tuesday to Monday) temperature T_t (degrees Celsius), absolute humidity
ah_t (g/m³), and solar radiation sr_t (kJ/cm²) as weather variables (Figure 1c,d). We used observations from
the "De Bilt" weather station, located centrally in the Netherlands,[27] and calculated absolute humidity
from relative humidity and temperature as

$ah = 2.1674 \cdot 6.112 \cdot \exp\left(17.67 \cdot T / (T + 243.5)\right) \cdot rh / (T + 273.15)$.[28]

We used the Oxford Stringency Index for the Netherlands as a compound measure for the stringency of
control measures (Oxford COVID-19 Government Response Tracker).[29] It was used to indicate changes
in control measures and compare these to changes in the estimated regression coefficients.

We conducted weighted time series regression analyses to estimate the association of the SARS-CoV-2
reproduction number R_t and weather variables between 22 February 2020 and 29 March 2022 in the
Netherlands. The full regression equation for the main analysis was

$\log R_t \sim offset\left(\log R_t^{var} + \log s_t\right) + period_t + s(T_t) * s(ah_t) * s(sr_t) + s(T_{t-7}) * s(ah_{t-7}) * s(sr_{t-7})$,   (1)

in which the lags are 7 days because of the use of weekly data, and weekly averaged weather variables.
The observations were weighted by $1/var(\log R_t)$ to reflect uncertainty in the estimated logR_t. Offsets
were included to account for a known increase in transmissibility because of the emergence of new
variants (up to and including Delta), and for a known decrease in the susceptible fraction resulting from
immunity due to natural infection (until 2 July 2021) and primary vaccination. The factor *period* was
added to correct for the effect of different contact rates due to changing intervention measures or
school holidays and to absorb other mechanisms for gradual changes in transmissibility, such as
compliance to measures, behaviour unrelated to measures, and population immunity not included in
the offsets, by booster vaccination, natural waning of immunity and immune escape due to the
emergence of the various Omicron variants.

We fitted 57 models with the last six terms of equation (1), i.e. temperature, absolute humidity, and/or
solar radiation, with or without interaction terms, in the same week or lagged by one week, with a linear
relation or with thin-plate regression splines. For this, we used generalised additive models (R package
mgcv,[30] version 1.9.0), with the REML optimisation method. We compared the models by AIC$_c$, Akaike
Information Criterion with an additional correction[31] because of the high number of parameters
(including 41 periods) relative to observations (109 weeks): $AIC_c = AIC + 2 \times df \times (df + 1) / (n - df - 1)$, in
which AIC and *df* are calculated output from the mgcv package (*df* is estimated with spline models). A
lower AIC$_c$ by at least 2 points difference indicates a better model. All AIC$_c$ values are reported as ΔAIC$_c$,
relative to the model in equation (1) without the weather terms. We translated estimated model
parameters to a mean R_t in winter vs summer, by first calculating 7-day rolling averages of temperature,
absolute humidity, and solar radiation per day-of-year with the mean weather of 2000-2019, then
predicting a daily R_t, and finally dividing the maximum weekly R_t by the minimum.

We challenged our results against the possibility that they are due to day-of-year as a confounder, i.e.
that there was an association between logR_t and the weather only because both are seasonal. We did

this by fitting models with mean daily temperature, absolute humidity, and solar radiation of the years
2000-2019, and with the five time series with the exact weather data of earlier years, all starting with a
leap year: 2000-2002, 2004-2006, 2008-2010, 2012-2014, and 2016-2018. This addressed the question if
$\log R_t$ can be explained by the general seasonal trend of the weather rather than the specific fluctuations
in temperature,absolute humidity, and solar radiation in the two years of the pandemic.
187
We did sensitivity analyses to address some specific choices we made. First, we repeated the main
analysis by using averages of weather variables across 28 weather stations in the Netherlands, weighted
by local population size. Second, we repeated the main analysis with estimates of reproduction numbers
for different days than Friday. Third, we repeated the main analysis with a single two-week period from
26 June 2021 to 9 July 2021, instead of two separate one-week periods. Fourth, we repeated the main
analyses without offsets, to assess if the *period* variable will compensate for this; if so, we can be more
confident that *period* will also absorb other gradually changing effects, in particular loss of immunity and
the rise of the Omicron variant.
196
197
198
199　RESULTS
200
201　We see a clear preference (lower AIC_c) for two linear models, both with an inverse association with solar
202　radiation: one with temperature and the other with absolute humidity (Table 1). Splines and delay terms
203　never improved model fits (lower AIC_c); including interaction terms improved model fits only for models
204　with both temperature and absolute humidity (supplementary material S4).
205
206　According to the model with temperature and solar radiation, R_t is expected to decrease by 2.0%
207　(standard error 0.5%) with every degree Celsius increase, and by 12% (s.e. 5%) with every kJ/cm^2
208　increase of solar radiation, which results in a ratio of R_t in winter (around 1 February) vs summer
209　(around 1 August) of 1.7 (95% CI: 1.4 ; 2.1). The model with absolute humidity and solar radiation
210　predicts this ratio to be 1.9 (1.5 ; 2.4) (Table 1, Figure 2).
211
212　*Period* and offsets explain 68% of the variance in $\log R_t$ (Table 1). The estimates of the period effects turn
213　out to reflect the stringency of control measures very well, as illustrated by the qualitative agreement
214　with the Oxford Stringency Index (Figures 3, S2): if measures become more stringent, the reproduction
215　number decreases. This qualitative agreement ends in Autumn 2021, after which the period-associated
216　reproduction number tends to be higher, especially when Omicron arises in 2022.
217
The effect of solar radiation and temperature (or absolute humidity) in determining overall
transmissibility explained another 8% (or 7%) of variance in $\log R_t$ (Table 1). This is clearly visible when
comparing the observed R_t and the predictions from the models with and without weather variables
(Figures 4, S3): especially during longer periods without changes in control measures (June to October
2020, and February to April 2021), fluctuations in R_t are much better explained in the model with
weather variables.
224
To assess robustness of these results, we addressed the question if the association of R_t with weather
variables was confounded by day of the year, i.e. if the association is only there because both are
seasonal. We did this by fitting the $\log R_t$ data to alternative time series of temperature, absolute
humidity, and/or solar radiation: the means of the years 2000-2019, and five periods of exact
measurements in earlier years, with all models presented in the main results. There were only eight

combinations of weather variables and models showing better fits than the model without weather
variables (most with data from 2016-2018), suggesting some association to a general seasonal pattern,
but none of the 54 models explained $\log R_t$ better than the exact weather data from 2020-2022 (Table
2). The estimated effect sizes (R_t in winter vs summer) do indicate a small general seasonal effect
explained by the weather variables, but never as large and significant as with the original 2020-2022
data (Figure 5).

236
237

238 Table 1. Supports and estimated seasonal effects of models in the main analysis, including offsets and
239 *period*. The selected model with lowest ΔAIC_c is bold-faced.

	Model coefficients (SE) [a]				R_t winter vs summer (95% CI)	Performance	
	Temperature	Absolute humidity	Solar insolation	Interaction T and ah [b]		ΔAIC_c [c]	R^{2} [d]
0	-	-	-	-	1	0	0.68
1	-0.023 (0.006)	-	-	-	1.5 (1.2 ; 1.7)	-19.2	0.74
2	-	-0.049 (0.015)	-	-	1.5 (1.2 ; 1.8)	-10.5	0.72
3			-0.16 (0.05)	-	1.3 (1.1 ; 1.6)	-8.3	0.71
4	-0.032 (0.013)	0.027 (0.034)	-	-	1.4 (1.2 ; 1.7)	-14.6	0.74
5	-0.060 (0.016)	0.059 (0.034)	-	-0.0046 (0.0017)	1.8 (1.4 ; 2.4)	-20.6	0.76
6	**-0.020 (0.005)**	**-**	**-0.12 (0.05)**	**-**	**1.7 (1.4 ; 2.1)**	**-23.3**	**0.76**
7	**-**	**-0.051 (0.014)**	**-0.16 (0.05)**	**-**	**1.9 (1.5 ; 2.4)**	**-22.4**	**0.75**
8	-0.013 (0.015)	-0.019 (0.038)	-0.14 (0.06)	-	1.8 (1.4 ; 2.3)	-17.9	0.75
9	-0.041 (0.020)	0.021 (0.042)	-0.090 (0.060)	-0.0036 (0.0018)	2.0 (1.5 ; 2.7)	-18.4	0.76

240 [a] Measurement units: temperature (˚C), absolute humidity (g/m^3), solar radiation (kJ/cm^2); [b] Interaction
241 between temperature and absolute humidity; other interaction terms never improved model fit as
242 measured by AIC_c; [c] Corrected Akaike's Information Criterion, relative to that of the least complex model
243 (first line); [d] R-squared, proportion of variance explained

244
245

Sensitivity analyses show that our decision to use the data from a single weather station did not affect
our results (supplementary material S5). We repeated our main analyses with $\log R_t$ time series of other
days than Fridays (supplementary material S5), and found that our final model with temperature and
solar radiation as linear terms had highest or considerable support (AIC_c within two points of the
minimum) with data from all days except Wednesday and Thursday. Across all weekday datasets, the
model with only temperature and no solar radiation was also supported very well. The datasets of
Tuesdays and especially Wednesdays showed lowest support overall for associations of R_t with the
weather, which supports our decision for using the Friday data, because the Tuesday and Wednesday
data were estimated with incidences of Sundays and Mondays, which were most affected by day-of-the-
week effects in reporting. The estimated ratio of reproduction number R_t in winter vs summer varied
between 1.2 and 1.9, with the estimates of the main analysis at the higher end of this range.
257

Reproduction number relative to annual mean

Figure 2. The change in reproduction number due to the associations with temperature and solar radiation, and with absolute humidity and solar radiation, expressed relative to the annual mean. (A) The direct relation with temperature; (B) the direct relation with absolute humidity; (C) the direct relation with daily solar radiation; (C) the annual cycle using the relation between reproduction number and the mean daily weather variables of the years 2000-2019.

258
259
We also repeated the main analysis with a single two-week period between 26 June 2021 and 9 July
2021 (supplementary material S5), which resulted in more complex models with interactions receiving
most support: the model with temperature and solar radiation with an estimated R_t in winter vs summer
of 2.5, and the model absolute humidity and solar radiation with an estimated R_t in Winter vs Summer of
2.9. This finding confirms that results were sensitive to the two data points indicated by evaluating
Cook's distance in the preliminary analysis, thus justifying our choice of periods to use two separate
one-week periods and reduce the influence of these data points.
267
268 Finally, we repeated the main analysis without the offset term, to check if the *period* variable indeed
269 absorbs gradual changes in R_t due to effects not included in the model. Leaving out the offset did not
270 change the estimated effects of the weather variables, but did indeed result in *period* effects
271 incorporating the offsets (supplementary material S5).
272

Figure 3. The reproduction number during the pandemic with mean annual weather and without offsets, as predicted by the model with linear temperature and solar radiation effects. It shows the contribution of the *period* estimates, mainly reflecting the effectiveness of interventions and adherence to interventions, but after the summer of 2021 also waning of immunity and the emergence of the Omicron variants. The coloured bar indicates the Oxford Stringency index, a compound measure (scale 0-100) of overall stringency of government interventions[29].

273
274
275   DISCUSSION
276
We estimated the association between transmissibility of SARS-CoV-2 during the pandemic in the
Netherlands, and the mean daily temperature, absolute humidity, and solar radiation. We found an
inverse linear association with temperature and solar radiation to explain 8% of variance in $\log R_t$ during
the pandemic, with reproduction number R_t decreasing by 2.0% for every degree Celsius increase, and
by 12% for every kJ/cm^2 daily solar energy. Extrapolating to mean weekly temperatures during the year,
this means that R_t early February (Winter) is 70% higher than R_t early August (Summer) in the
Netherlands. A model receiving similar support suggests an association with absolute humidity and solar
radiation, translated to a seasonal effect of a 90% higher R_t in winter vs summer.
285

Effect of including weather variables on the prediction of R_t

A) Model without weather variables

B) Model with temperature and solar radiation

Figure 4. The reproduction number during the pandemic, predicted by (A) the model with only offsets and *period*, and (B) the model with linear temperature and solar radiation effects.

295 Table 2. Additional analysis adressing confounding of weather variables and the reproduction number R_t
296 by day-of-year (co-seasonality). The table shows ΔAIC_c values of model fits with the original weather
297 time series during the pandemic and alternative time series of other years (lowest in bold). ΔAIC_c is
298 calculated relative to that of the least complex model, i.e. without weather variables.

Model	Dataset (years of weather data)						
variables [a]	**2020-2022**	average [b]	2016-2018	2012-2014	2008-2010	2004-2006	2000-2002
T	**-19.2**	2.2	-4.9	4.5	4.7	1.9	5.2
ah	**-10.5**	4.7	1.1	5.2	5.5	-0.2	4.7
sr	**-8.3**	0.0	4.1	-8.9	2.9	4.7	4.2
T + ah	**-14.6**	3.9	-10.9	8.8	0.6	5.1	9.7
T * ah	**-20.6**	9.7	-9.6	12	3.7	10.8	15.5
T + sr	**-23.3**	5.6	-0.3	-3.3	8.2	7.5	9.4
ah + sr	**-22.4**	5.3	4.8	-3.3	8.5	5.3	9.2
T + ah + sr	**-17.9**	9.7	-5.6	1.7	6.3	9.6	14.9
T * ah + sr	**-18.4**	15.7	-5.2	7.4	8.7	15.5	20.5

299 [a] Model variables are temperature (T), absolute humidity (ah), and solar radiation (sr), multiplication
300 indicates interaction; [b] daily average weather of the years 2000-2019

301
302
303
304

Figure 5. Additional analysis adressing confounding of weather variables and the reproduction number R_t by day-of-year (co-seasonality). The figure shows the magnitude of the estimated seasonal effect (R_t in winter vs summer, 95% CI) associated with the original weather time series during the pandemic, the average weather between 2000-2019, and alternative time series of other years. Weather variables in the models are temperature (T), absolute humidity (ah), and solar radiation (sr); multiplication indicates inclusion of an interaction term.

305
306

To estimate the effect of weather variables, we had to separate this effect from the much larger effect
on SARS-CoV transmission during the pandemic by several other factors. The effects of variants,
immunity from infection, and vaccination could be included as offsets up to Autumn 2021, after which
waning immunity and immune escape due to the Omicron variant obscured the level of immunity in the
population. However, the varying sets of sometimes rapidly changing social distancing measures had
even greater impacts which could not be estimated independently. To account for these, the stepwise
constant *period* was added to the model with changepoints at the times of changing control measures
and school holidays. These could explain large changes in $\log R_t$ (68% of variance) due to changing
control policies, but also allowed for more gradual changes due to other effects such as changing
behaviour due to epidemic fatigue, and changes in immunity or variant transmissibility not accounted
for in the offsets. The fact that the estimated effect of the weather variables does not change when
leaving out the offsets (supplementary material S5) shows that such gradual changes are indeed
absorbed by the *period* variable. Including the offsets is useful however, because it allows us to interpret
the coefficients for *period* as the effects of control measures, which could therefore be compared with
the Oxford Stringency Index as a qualitative validation (Figure 2a).[29]
322
We addressed the possibility that the association was only a result of two seasonally co-oscillating
variables without any direct or indirectly causal link, by fitting $\log R_t$ to alternative datasets of
temperature, absolute humidity, and solar radiation in the Netherlands on the same dates but in
different years, and a 20-year average. A small seasonal effect was observed with all datasets (Figure 5),
but with most datasets and models (46/54) there was no statistical evidence for an association between
$\log R_t$ and weather variables at all, and with each model $\log R_t$ was best explained by the data from 2020-
2022, the years of the pandemic. This result is a strong indication that the actual weather was causally
related with SARS-CoV-2 transmission during the pandemic. Regardless of the causal nature of the
relationship, changes in the actual solar radiation and temperature (or absolute humidity) are thus good
predictors of changes in transmissibility. Prediction models can take advantage of this by using actual
weather data to anticipate changes in incidence that will result from such changes in transmissibility.
334
A strength of our study was that we had a high-quality R_t time series of more than two years. However,
the R_t estimates were autocorrelated and R_t was highly affected by changing behaviour due to control
measures and holidays, but also related to other factors such as pandemic.[32] Changes in R_t due to
variants and immunity by infection or vaccination could be accounted for, but only up to the summer of
2021, after which waning immunity and the rise of the Omicron variant made population immunity
untrackable. We accounted for these issues by using only weekly data and including a *period* variable to
avoid assumptions on changing behaviour and immunity. However, this resulted in a considerable loss of
degrees of freedom and therefore statistical power. We used AIC_c for model selection, which strongly
favoured simpler model with linear terms rather than splines, without delays, and with no or limited
interactions, whereas many earlier studies have shown complex relations between R_t and temperature,
humidity and solar or UV radiation. The limitation in statistical power also made it difficult to correct for
more possibly relevant variables, such as mobility data, which we considered but resulted in unstable
models due to high collinearity with especially solar radiation (results not shown).
348
Earlier studies on associations between weather variables and SARS-CoV-2 transmission generally
agreed on an inverse effect of temperature. Results on the role of humidity were less consistent. Our
study shows evidence of an inverse association with solar radiation, in combination with temperature or
absolute humidity, resulting in a larger seasonal amplitude than estimated before in similar climates.[16,
17] Because most SARS-CoV-2 transmission took place indoors,[8] it is likely that this association is for a
large part due to people going outside in warm and sunny weather. However, even then the true

mechanism is still unknown: people may have fewer or shorter close contacts outside, but it does not
rule out the possibility of host or virus factors, e.g. sunlight affecting virus stability. The fact that our
results are inconclusive about the role of temperature or absolute humidity may have to do with the
limited statistical power for more complex models, as discussed above. Given that the existing literature
is more conclusive about temperature,[15] and that our sensitivity analyses with datasets from other
weekdays generally favour models with temperature over those with absolute humidity (supplementary
material S5), we have more confidence in a role for temperature rather than absolute humidity.
362
Unique to our study is that we aimed to quantify changes in SARS-CoV-2 transmissibility attributable to
changes in weather variables, to assess the magnitude of seasonal variation in transmissibility in the
Netherlands. For a reliable quantification it is important to restrict estimation to a limited geographic
area, because the relationship may differ in different geographic locations, making a single estimate
from multiple locations difficult to interpret. Keeping the model simple also adds to reliability of the
estimation, because if weather variables are included with many other explanatory variables, there is a
risk that the different variables are dependent or display spurious associations, resulting in potentially
biased estimates. Our results suggest that using solar radiation and temperature as covariates will
increase the performance of prediction models of SARS-CoV-2 transmissibility and resulting COVID-19
incidence.
373
374
375
376   CONCLUSIONS
377
378   In the Netherlands during the COVID-19 pandemic, SARS-CoV-2 transmissibility was higher with less
379   solar radiation and with lower temperature. This resulted in a 70% higher reproduction number in
380   winter compared to summer. Sensitivity analyses confirmed a robust association with weather variables,
381   suggesting a valuable role of weather data data for prediction models of COVID-19 incidence.
382
383
384
385

REFERENCES

[1] Martinez, M.E. 2018 The calendar of epidemics: Seasonal cycles of infectious diseases. *PLoS Pathog* **14**, e1007327. (doi:10.1371/journal.ppat.1007327).

[2] Metcalf, C.J.E., Walter, K.S., Wesolowski, A., Buckee, C.O., Shevliakova, E., Tatem, A.J., Boos, W.R., Weinberger, D.M. & Pitzer, V.E. 2017 Identifying climate drivers of infectious disease dynamics: recent advances and challenges ahead. *Proc Biol Sci* **284**. (doi:10.1098/rspb.2017.0901).

[3] te Beest, D.E., van Boven, M., Hooiveld, M., van den Dool, C. & Wallinga, J. 2013 Driving factors of influenza transmission in the Netherlands. *Am J Epidemiol* **178**, 1469-1477. (doi:10.1093/aje/kwt132).

[4] Lowen, A.C., Mubareka, S., Steel, J. & Palese, P. 2007 Influenza virus transmission is dependent on relative humidity and temperature. *PLoS Pathog* **3**, 1470-1476. (doi:10.1371/journal.ppat.0030151).

[5] Limaheluw, J., Dollmann, S., Folpmers, S., Beltran Beut, L., Lazarakou, A., Vermeulen, L.C. & de Roda Husman, A.M. 2024 Associations between meteorological factors and COVID-19: a global scoping review. *Front Public Health* **12**, 1183706. (doi:10.3389/fpubh.2024.1183706).

[6] Moriyama, M., Hugentobler, W.J. & Iwasaki, A. 2020 Seasonality of Respiratory Viral Infections. *Annu Rev Virol* **7**, 83-101. (doi:10.1146/annurev-virology-012420-022445).

[7] Atchison, C.J., Tam, C.C., Hajat, S., van Pelt, W., Cowden, J.M. & Lopman, B.A. 2010 Temperature-dependent transmission of rotavirus in Great Britain and The Netherlands. *Proc Biol Sci* **277**, 933-942. (doi:10.1098/rspb.2009.1755).

[8] Bulfone, T.C., Malekinejad, M., Rutherford, G.W. & Razani, N. 2021 Outdoor Transmission of SARS-CoV-2 and Other Respiratory Viruses: A Systematic Review. *J Infect Dis* **223**, 550-561. (doi:10.1093/infdis/jiaa742).

[9] Babin, S. 2020 Use of Weather Variables in SARS-CoV-2 Transmission Studies. *Int J Infect Dis* **100**, 333-336. (doi:10.1016/j.ijid.2020.09.032).

[10] Sajadi, M.M., Habibzadeh, P., Vintzileos, A., Shokouhi, S., Miralles-Wilhelm, F. & Amoroso, A. 2020 Temperature, Humidity, and Latitude Analysis to Estimate Potential Spread and Seasonality of Coronavirus Disease 2019 (COVID-19). *JAMA Netw Open* **3**, e2011834. (doi:10.1001/jamanetworkopen.2020.11834).

[11] Kissler, S.M., Tedijanto, C., Goldstein, E., Grad, Y.H. & Lipsitch, M. 2020 Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period. *Science* **368**, 860-868. (doi:10.1126/science.abb5793).

[12] Sera, F., Armstrong, B., Abbott, S., Meakin, S., O'Reilly, K., von Borries, R., Schneider, R., Roye, D., Hashizume, M., Pascal, M., et al. 2021 A cross-sectional analysis of meteorological factors and SARS-CoV-2 transmission in 409 cities across 26 countries. *Nat Commun* **12**, 5968. (doi:10.1038/s41467-021-25914-8).

[13] Ma, Y., Pei, S., Shaman, J., Dubrow, R. & Chen, K. 2021 Role of meteorological factors in the transmission of SARS-CoV-2 in the United States. *Nat Commun* **12**, 3602. (doi:10.1038/s41467-021-23866-7).

[14] Wagatsuma, K. 2023 Association of Ambient Temperature and Absolute Humidity with the Effective Reproduction Number of COVID-19 in Japan. *Pathogens* **12**. (doi:10.3390/pathogens12111307).

[15] Wei, Y., Dong, Z., Fan, W., Xu, K., Tang, S., Wang, Y. & Wu, F. 2022 A narrative review on the role of temperature and humidity in COVID-19: Transmission, persistence, and epidemiological evidence. *Eco Environ Health* **1**, 73-85. (doi:10.1016/j.eehl.2022.04.006).

[16] Paireau, J., Charpignon, M.L., Larrieu, S., Calba, C., Hoze, N., Boelle, P.Y., Thiebaut, R., Prague, M. & Cauchemez, S. 2023 Impact of non-pharmaceutical interventions, weather, vaccination, and variants on COVID-19 transmission across departments in France. *BMC Infect Dis* **23**, 190. (doi:10.1186/s12879-023-08106-1).

[17] Xu, R., Rahmandad, H., Gupta, M., DiGennaro, C., Ghaffarzadegan, N., Amini, H. & Jalali, M.S. 2021

Weather, air pollution, and SARS-CoV-2 transmission: a global analysis. *Lancet Planet Health* **5**, e671-

e680. (doi:10.1016/S2542-5196(21)00202-3).

[18] 2023 Covid-19 reproductiegetal. (2023 ed, National Institute for Public Health and the Environment

(RIVM).

[19] Wallinga, J. & Teunis, P. 2004 Different epidemic curves for severe acute respiratory syndrome

reveal similar impacts of control measures. *Am J Epidemiol* **160**, 509-516. (doi:10.1093/aje/kwh255).

[20] Gostic, K.M., McGough, L., Baskerville, E.B., Abbott, S., Joshi, K., Tedijanto, C., Kahn, R., Niehus, R.,

Hay, J.A., De Salazar, P.M., et al. 2020 Practical considerations for measuring the effective reproductive

number, Rt. *PLoS Comput Biol* **16**, e1008409. (doi:10.1371/journal.pcbi.1008409).

[21] 2024 Covid-19 rapportage van SARS-CoV-2 varianten in Nederland via de aselecte steekproef van

RT-PCR positieve monsters in de nationale kiemsurveillance. (ed. N.I.f.P.H.a.t.E. (RIVM)), 2024 ed.

Bilthoven, The Netherlands, National Institute for Public Health and the Environment (RIVM).

[22] Campbell, F., Archer, B., Laurenson-Schafer, H., Jinnai, Y., Konings, F., Batra, N., Pavlin, B.,

Vandemaele, K., Van Kerkhove, M.D., Jombart, T., et al. 2021 Increased transmissibility and global

spread of SARS-CoV-2 variants of concern as at June 2021. *Euro Surveill* **26**. (doi:10.2807/1560-

7917.ES.2021.26.24.2100509).

[23] 2021 COVID-19 besmettelijke personen per dag. (2021 ed, National Institute for Public Health and

the Environment (RIVM).

[24] 2022 Covid-19 Cumulatieve opkomst voor tenminste één COVID-19 vaccinatie, vaccinatiegraad

basisserie, vaccinatiegraad herhaalprik tegen corona binnen de najaarsronde 2022 naar gemeente en

veiligheidsregio per geboortecohort per week. (ed. N.I.f.P.H.a.t.E. (RIVM)), 2022 ed. Bilthoven, The

Netherlands, National Institute for Public Health and the Environment (RIVM).

[25] Huiberts, A.J., de Gier, B., Hoeve, C.E., de Melker, H.E., Hahne, S.J.M., den Hartog, G., Grobbee, D.E.,

van de Wijgert, J., van den Hof, S. & Knol, M.J. 2023 Vaccine effectiveness of primary and booster

COVID-19 vaccinations against SARS-CoV-2 infection in the Netherlands from July 12, 2021 to June 6,

2022: A prospective cohort study. *Int J Infect Dis* **133**, 36-42. (doi:10.1016/j.ijid.2023.04.401).

[26] Cook, R.D. 1979 Influential observations in linear regression. *Journal of the American Statistical

Association* **74**, 169-174. (doi:10.1080/01621459.1979.10481634).

[27] 2022 Daggegevens van het weer in Nederland. (ed. R.N.M.I. (KNMI)), 2022 ed, Royal Netherlands

Meteorological Institute (KNMI).

[28] Wu, P., Goldstein, E., Ho, L.M., Yang, L., Nishiura, H., Wu, J.T., Ip, D.K., Chuang, S.K., Tsang, T. &

Cowling, B.J. 2012 Excess mortality associated with influenza A and B virus in Hong Kong, 1998-2009. *J

Infect Dis* **206**, 1862-1871. (doi:10.1093/infdis/jis628).

[29] Hale, T., Angrist, N., Goldszmidt, R., Kira, B., Petherick, A., Phillips, T., Webster, S., Cameron-Blake,

E., Hallas, L., Majumdar, S., et al. 2021 A global panel database of pandemic policies (Oxford COVID-19

Government Response Tracker). *Nat Hum Behav* **5**, 529-538. (doi:10.1038/s41562-021-01079-8).

[30] Wood, S.N. 2017 *Generalized additive models : an introduction with R*. Second edition. ed. Boca

Raton, CRC Press/Taylor & Francis Group; xx, 476 pages p.

[31] Burnham, K.P., Anderson, D.R. & Burnham, K.P. 2002 *Model selection and multimodel inference : a

practical information-theoretic approach*. 2nd ed. New York, Springer; xxvi, 488 p. p.

[32] Bussemakers, C., Stappers, N., Kroese, F., van den Putte, B. & de Bruin, M. 2025 Psychosocial

determinants of handwashing and physical distancing behaviour during the COVID-19 pandemic in the

Netherlands: A longitudinal analysis. *Br J Health Psychol* **30**, e12755. (doi:10.1111/bjhp.12755).