Transparency in epidemiological analyses of cohort data

A case study of the Norwegian Mother, Father, and Child cohort study (MoBa)

Timo Roettger[3]
Adrian Dahl Askelund[1,2]
Viktoria Birkenæs[5]
Ludvig Daae Bjørndal[3]
Agata Bochynska[3]
Bernt Damian Glaser[2,3]
Tamara Kalandadze[6]
Max Korbmacher[4,7,8] *(corresponding author)*
Ivana Malovic[3]
Julien Mayor[3]
Pravesh Parekh[5]
Daniel S. Quintana[3,9,10]
Laurie J. Hannigan[1,2,11]

[1] Psychiatric Genetic Epidemiology group, Research Department, Lovisenberg Diaconal Hospital, Norway
[2] PsychGen Centre for Genetic Epidemiology and Mental Health, Norwegian Institute of Public Health, Norway
[3] University of Oslo, Norway
[4] Department of Neurology, Haukeland University Hospital, University of Bergen, Norway
[5] Centre for Precision Psychiatry, Division of Mental Health and Addiction, University of Oslo and Oslo University Hospital, Oslo, Norway
[6] Østfold University College, Norway
[7] Department of Health and Functioning, Western Norway University of Applied Sciences, Norway
[8] Mohn Medical Imaging and Visualization Centre (MMIV), Norway
[9] KG Jebsen Center for Neurodevelopmental Disorders, University of Oslo, Norway
[10] Oslo University Hospital, Norway
[11] Population Health Sciences, Bristol Medical School, University of Bristol, UK

Abstract

Background: Epidemiological research is central to our understanding of health and disease. Secondary analysis of cohort data is an important tool in epidemiological research but is vulnerable to practices that can reduce the validity and robustness of results. As such, adopting measures to increase the transparency and reproducibility of secondary data analysis is paramount to ensuring the robustness and usefulness of findings. The uptake of such practices has not yet been systematically assessed.

Methods: Using the Norwegian Mother, Father and Child Cohort study (MoBa; Magnus et al., 2006, 2016) as a case study, we assessed the prevalence of the following reproducible practices in publications between 2007-2023: preregistering secondary analyses, sharing of synthetic data, additional materials, and analysis scripts, conducting robustness checks, directly replicating previously published studies, declaring conflicts of interest and publishing publicly available versions of the paper.

Results: Preregistering secondary data analysis was only found in 0.4% of articles. No articles used synthetic data sets. Sharing practices of additional data (2.3%), additional materials (3.4%) and analysis scripts (4.2%) were rare. Several practices, including data and analysis sharing, preregistration and robustness checks became more frequent over time. Based on these assessments, we present a practical example for how researchers might improve transparency and reproducibility of their research.

Conclusions: The present assessment demonstrates that some reproducible practices are more common than others, with some practices being virtually absent. In line with a broader shift towards open science, we observed an increasing use of reproducible research practices in recent years. Nonetheless, the large amount of analytical flexibility offered by cohorts such as MoBa places additional responsibility on researchers to adopt such practices with urgency, to both ensure the robustness of their findings and earn the confidence of those using them. A particular focus in future efforts should be put on practices that help mitigating bias due to researcher degrees of freedom – namely, preregistration, transparent sharing of analysis scripts, and robustness checks. We demonstrate by example that challenges in implementing reproducible research practices in analysis of secondary cohort data - even including those associated with data sharing - can be meaningfully overcome.

1. Background

Epidemiological research is central to our understanding of why and how diseases and disorders originate and spread over time. For example, the Norwegian Mother, Father and Child Cohort Study (MoBa; Magnus et al., 2006) was created to investigate diseases and disorders, collecting longitudinal information about almost 100,000 mothers alongside their children and their children's fathers over the last 25 years. This dataset has allowed researchers to successfully answer questions about fertility, cognitive development and the development of the human brain, among many other research areas (Magnus et al., 2016).

1.1 Epidemiological research and the garden of forking paths

Epidemiological data sets, such as MoBa, are rich and complex, containing hundreds of variables, and offer a myriad of ways to learn from them. To analyze these datasets appropriately, many different methodological and analytical choices need to be made, all of which may influence the final interpretation of the data (Wicherts et al., 2016). This range of analytical decisions introduces so-called researcher degrees of freedom. On the one hand, they represent opportunities to look at the data set from many different angles, which, in turn, allows researchers to make important discoveries and generate novel hypotheses (e.g., Box, 1976; De Groot, 2014; Tukey, 1977). On the other hand, idiosyncratic choices can lead to categorically different interpretations (Simmons et al., 2011). Recent studies have shown that independent data analysts analyze the same data set in vastly different ways, leading them to arrive at different conclusions when trying to answer the same research question (e.g., Silberzahn et al., 2018; Dutilh et al., 2019; Starns et al., 2019; Coretta et al., 2023).

 For example, Silberzahn et al. (2018) invited 29 independent research teams to test the same hypothesis: whether soccer referees are more likely to issue red cards to players with darker skin tones compared to those with lighter skin tones. The teams employed a wide range of analytical approaches, leading to varying results - 20 teams (69%) found evidence supporting the hypothesis, while 9 teams (31%) did not. Among the 29 analyses, there were

21 distinct combinations of predictors used. Notably, the diversity in outcomes could not be

explained by the researchers' prior beliefs about the topic or by peer evaluations of the

quality of their analyses. The findings highlight that variability in analytic results may be an

inherent feature of the scientific process, rather than a reflection of bias or differences in

expertise. Follow up studies further showed that even seemingly inconsequential choices

regarding data cleaning, variable operationalisation, and model structure can lead to

categorically different conclusions (e.g. Coretta et al., 2023).

With rich cohort data such as MoBa, the "garden of forking paths" of possible analytical

pipelines (Gelman & Loken, 2014) is so vast - and the amount of researcher degrees of

freedom is so large - that arguably any hypothesis can be supported by the data if the data

are interrogated often enough. Indeed, some recent empirical efforts to either replicate,

reproduce, or generalise epidemiological findings have been unsuccessful (Border et al.,

2019; Seibold et al., 2021; Orben et al., 2019). This underlines the urgency of addressing

these issues in research using secondary data analysis.

105   1.2 Transparent scientific practices

Under the broad umbrella of "Open Science", a number of reform efforts have recently been

proposed across disciplines (e.g. Kidwell et al., 2016; Klein et al., 2018; Zwaan et al., 2018;

Korbmacher et al., 2023). The main focus of these efforts has been to increase the

transparency of the research process and its outputs by, for example, establishing and

archiving a public version of a preregistered research plan (Nosek et al., 2018), and sharing

of research materials and procedures, raw and processed data, and analysis scripts

(Gilmore et al., 2018; Lindsay, 2017). The central goal of these efforts is to allow the

scientific community and often also the broader public to access relevant information so that

they can critically evaluate scientific claims (Munafò et al., 2017; Vazire, 2017). Some of

these practices directly tackle challenges related to analytical flexibility. Sharing statistical

protocols and data enables transparent and unambiguous communication of all relevant

analytical decisions and, in turn, allows other researchers to assess the robustness of an

analysis against other possible analyses. Preregistration draws a transparent line between

initially planned analyses and post hoc exploratory analyses (Nosek et al., 2018), makes

questionable research practices such as selective reporting and hypothesizing after results

are known detectable (John et al., 2012), and helps reduce researcher degrees of freedom

(Mertzen et al., 2021; Simmons et al., 2011; Wicherts et al., 2016), making it a particularly

useful tool in epidemiological research.

Beyond directly addressing concerns related to researcher degrees of freedom, transparent

practices promise general benefits for both individual researchers and science at large. More

transparent research can accelerate discovery by allowing re-use of research materials,

procedures and data, which can in turn facilitate collaborations (Boland et al., 2017), and

increase efficiency and sustainability (Lowndes et al., 2017). Transparent sharing of

research materials, data and scripts makes it easier to detect and correct errors (Nuijten et

al., 2016). It can also aid in facilitating evidence synthesis (e.g. meta-analyses; Pigott &

Polanin, 2020) and enable independent reproductions and replications of results (Munafò et

al., 2017). Given this propagation of the effects of transparent and reproducible practices

across all levels of the research process – and that the goal of all epidemiological research

is to benefit patients and their families – their implementation can ultimately be seen as a

means of achieving this goal by enhancing and accelerating the clinical impact of findings.

1.3 The present paper

Recent assessments of biomedicine (Wallach et al., 2018), social sciences (Hardwicke et al.,

2020), psychological sciences (Hardwicke et al., 2022) and language sciences (Bochynska

et al., 2023) suggest that rates of transparent practices are still quite low. In epidemiological

analyses of cohort data, researchers may perceive specific challenges to implementing

transparent practices (Baldwin et al., 2022; van den Akker et al., 2019; Weston et al., 2019).

For example, epidemiological cohorts often contain sensitive and/or personal data that

cannot be shared openly. Even sharing derivatives of the original data can contain

identifying information (Rocher et al., 2019). Thus, the aim of the current paper is to assess

the current level of transparent practices found in secondary analysis of cohort data and to

offer concrete and actionable advice on how to be as transparent as possible given the

existing limitations and challenges. First, we aim to assess and quantify the adoption of

transparent practices in secondary data analysis of cohort data using MoBa as a case study.

Second, we will identify and demonstrate solutions that can facilitate the adoption of

transparency practices in cohort studies, with a particular focus on practices that tackle

issues related to researcher degrees of freedom.

152   2. Methods

153   2.1 Assessing transparent research practices in MoBa

As a case study, we assessed reproducible and transparent practices in secondary data

analysis using the MoBa dataset. We chose MoBa for several reasons: First, the MoBa

cohort has resulted in more than thousand research articles, thus providing a representative

snapshot of scientific practices while the sample size remains feasible for a systematic meta

scientific assessment. Second, MoBa-based research has been published across a long

enough time span (>17 years) to meaningfully investigate trends over time. Third, data

collection for MoBa is ongoing and thus likely to result in many more future publications.

Assessing research on MoBa, thus, gives us the opportunity to encourage researchers

engaging with MoBa to adopt more open and reproducible practices.

Specifically, we looked at the implementation of the following practices which are currently

considered among best practices for open and credible research: preregistering secondary

analyses, sharing of synthetic data, sharing of meta-data and summary statistics, sharing of

additional materials, including analysis scripts, conducting robustness checks, directly

replicating previously published studies, declaring conflicts of interest and publishing of

publicly available versions of the paper. Our study's design and data collection plan was

uploaded as a preregistration prior to data collection (https://osf.io/ef4t5/). All materials, data,

and analysis scripts related to this study are publicly available on Open Science Framework:

https://osf.io/2jqxv/.

2.1.1 Sample

173   As the primary data source, we retrieved the list of publications that have used the MoBa

174   dataset from the MoBa website[1] and obtained 1005 articles. After a pilot evaluation of 60

175   randomly sampled entries, the remaining 945 articles were coded by six raters.

2.1.2 Procedure

Measured variables and their operationalisations are shown in Table 1. We coded the year

of publication and the type of study by examining title, abstract, and if necessary, the

methods section, to establish the study characteristics. We coded whether authors claimed

the presence of preregistrations, synthetic data, other non-primary data such as metadata

and summary statistics, analysis protocols or additional materials, by examining relevant

parts of the method and result section of each article. If one of these aspects was present,

we noted how this information was accessible, and whether we could indeed access it or

not. We moreover assessed whether the authors claimed to have replicated a previous study

or not and whether authors declared the absence or presence of a conflict of interest. Data

extraction was executed via a Google Form consisting of questions to the coder and

response options, see https://osf.io/47xmh.

Table 1: Main measured variables.

Coding Category	Coding Area
Open access	Publication year
	Study type

---

[1] https://www.fhi.no/en/ch/studies/moba/for-forskere-artikler/publications/

Preregistration	Preregistration
	Preregistration method
	Preregistration accessibility
	Preregistration content
Data	Synthetic data sharing
	Synthetic data sharing method
	Synthetic data accessibility
	Synthetic data documentation
	Other data sharing
	Other data sharing method
	Other data accessibility
Analysis script	Analysis script sharing
	Analysis script sharing method
	Analysis script accessibility
Materials	Materials sharing
	Materials sharing method
	Materials accessibility
Replication	Replication
Conflict of interest	Conflict-of-Interest statement
Open publishing	Open publishing status

191 Each coder assessed individual articles for each measured variable, recording outcomes in

192 a structured form (https://osf.io/2qdzb). In total, 80% of the 945 target articles were randomly

193 drawn and coded by single coders. The remaining subset (20%) was coded by two coders to

194 assess inter-rater reliability. See the preregistration for more details on the coding

195 procedure.

2.2 How to be as transparent as possible - a toy example

We also aim to demonstrate a worked example of how researchers analyzing MoBa data can use current tools and standards for transparent practices in conducting and reporting their work. To that end, we simulated data based on the characteristics of real MoBa data in order to present a realistic example of a reproducibly planned and reported MoBa analysis. The key features that we emphasise in this example are preregistration, detailed reporting of deviations and unregistered steps, robustness checks, sharing of well-annotated analytic code, and sharing of synthetic data to facilitate computational reproducibility. All materials are available (https://osf.io/nm9ej/) and can be used as points of departure for future efforts.

In our hypothetical research project, researchers were interested in testing the independent effects of age (M = 3.46 years, SD = 2.76) and breastfeeding duration (M = 5.47 months, SD = 3.73) on height measured on six occasions during childhood in MoBa and assessing whether these effects vary according to sex (a categorical variable with levels: "Female" and "Male").

3. Results

3.1 Assessment of MoBa

Coders agreed for 91% of all items. Discrepancies between the first and the second coder were not obviously linked to any particular coding category and were easily resolved through discussion. The following overview is based on the consensus ratings. Of the 943 articles, 799 (84.7%) were coded as in-principle accessible. Only those articles were further analysed. 58 articles could not be accessed by the coder (6.2%); 24 were coded as not being a full article, e.g. only a conference abstract or an erratum (2.5%); two were coded as not being written in English (0.2%); and 60 articles were coded as having other issues, including papers that turned out to actually not analyze MoBa (6.4%).

Of those articles that were in principle accessible, 733 articles were coded as presenting

empirical data and using the MoBa corpus (91.7%). All rates reported below refer to these

733 articles. Figure 1 summarises the results in relation to the articles' year of publication.

Green represents transparent practices, and gray represents non-transparent practices.

Individual assessment categories are discussed below. A more detailed presentation of data

is accessible here (https://osf.io/ec6tf).

Figure 1: Percentage of transparent practices for articles as a function of publication year (2007-

2024). Year 2023 and 2024 were collapsed due to a small number of *n*.

3.1.1 Preregistration, data and analysis sharing

Despite its relevance for secondary data analysis, there were only 3 articles in the entire sample (0.4%) that reported having preregistered aspects of their analysis. None of the articles reported having used synthetic data.

In our corpus, we found 17 articles to share resources that could be considered some form of non-primary data (2.3%), which mostly referred to additional metadata, validation data, or summary statistics. However, only 31 of assessed articles were coded as reporting to share some form of analysis protocols either in form of step-by-step instructions or code (4.2%).

Despite these low rates, there is a trend towards more transparent practices: both additional data sharing, and analysis sharing indicate a trend upwards with 7.1% of articles sharing additional data and 21.4% of articles sharing their analysis in 2023/2024. Both practices were non-existent in the first half of the publication range (see Fig. 1B and 1D).

3.1.2 Robustness checks

Given the analytical flexibility in secondary data analysis, it is useful to evaluate how robust a statistical outcome is under different analytical assumptions. Such sensitivity analyses were indeed not uncommon in our sample: one third of articles stated that some form of a sensitivity analysis was performed (239 articles, 33%). 44 of assessed articles reported a power analysis (6%). Given the lack of sharing practices of scripts and non-primary data, the exact nature of both these sensitivity and power analyses often remained unclear. While the overall rates for robustness checks were low, general upward trends were apparent with 42.9% of articles performing a sensitivity analysis and 14.3% a power analysis in 2023/2024, both of which were almost non-existent at the beginning of the publication range (see Fig. 1E).

In our assessment, 24 articles (3.4%) reported to sharing some form of additional materials

such as algorithmic description of quality control procedures (e.g., Isungset et al., 2022) or

insights into the selection of items and results from measurement models (Øksendal et al.,

2022). 11 articles reported to have directly replicated a previous study (1.5%). Compared to

other fields, MoBa articles have a high rate of declaring conflicts of interest, approaching

ceiling effects for recently published articles. Only 142 of assessed articles were coded as

not reporting a conflict-of-interest statement (19.3%); 488 reported no conflict of interest

(66.2%); and 107 reported an existing conflict of interest (14.5%). It is important to note that

these rates likely do not reflect voluntary practices of researchers but reflect how many

journals require this form of disclosure. When it comes to publicly sharing a version of the

manuscript, 661 of assessed articles were publicly available according to our definition

(89.7%) which is a much higher rate than similar assessments in other fields (e.g.,

biomedicine: 25%; Wallach et al., 2018).

3.2 Demonstrating transparent practices

3.2.1 Preregistering the analysis and document deviations

Here we present a full toy example that demonstrates workflows and tools that can be used

to increase transparency for secondary data analyses of cohort data. The preregistration of

the toy example is uploaded to the Open Science Framework Registry prior to accessing the

data for analyses and can be accessed here: https://osf.io/eczpq. The hypothetical

researchers investigate relationships between age, breastfeeding duration, and childhood

height in MoBa and aim to limit researcher degrees of freedom in the specific context of

analysing pre-existing data through a thorough preregistration. In particular, this includes

specification of inclusion/exclusion criteria, operationalisation of relevant variables that will

be included in analytic models, and specification of hypotheses that will be tested alongside

the applied inference criteria. The researchers also disclose the nature and extent of their

knowledge of MoBa data from prior work.

The specific variables of our toy example were chosen as they illustrate variables that are

typically used in MoBa. For instance, questionnaire data may look and act differently than

what was expected in the preregistration phase. We emulated this here by simulating

irregular distributions for mother-reported breastfeeding duration, with large idiosyncratic

asymmetries in its distribution (as well as missing data), suggesting at least three underlying

data generating patterns for the variable. Given these irregularities, in the analysis phase,

the researchers decide to not treat the variable as a continuous predictor as preregistered

but rather as a categorical predictor that bins breastfeeding into short (1-2 months), medium

(3-5 months), and long duration (>5 months). Moreover, when cleaning the height variables,

the hypothetical researchers realise that the preregistration was insufficiently precise about

how "valid responses" should be defined and therefore implement a +/- 3 SD threshold for

outlier removal. These issues will likely be familiar to readers accustomed to performing

secondary data analyses and may even be part of the explanation for the limited uptake of

preregistration for this kind of work in epidemiological studies. However, there are easy-to-

implement solutions for such situations.

In our example, the researchers' note their deviations and unregistered steps in a specially

formatted table (Tab. 2, adapted from Willroth et al. 2024), designed to ensure transparency

for readers. In implementing their deviation (categorising the breastfeeding variable), they

are mindful of maintaining their Type 1 error rate at the prespecified level (introducing a

single likelihood ratio test to assess the effect of breastfeeding, which would otherwise - with

the exposure now a categorical variable - correspond to multiple model parameters). To

interrogate the impact of their modified approach on their results, they additionally perform a

sensitivity analysis with a slightly different categorisation of the breastfeeding duration

variable (0-5 months as "early cessation", 6 months as "recommended", and over 6 months

as "late cessation"). Likewise, when implementing their unregistered step (operationalising

"valid responses" on the measure of height as those within 3 standard deviations of the

mean), they again perform an additional sensitivity analysis (using a +/- 2 SD threshold) to

interrogate the impact of this decision on the results and communicate this to readers.

Table 2: *Preregistration Deviations Table for the Toy Example (the template can be accessed [here](#))*

Deviations

Details		Original Wording	Deviation Description	Reader Impact
Type	Data Preparation	Section 3.1 of preregistration: "Breastfeeding duration: a continuous variable measured in months, derived based on mothers' reports of breastfeeding in the six and 18 month questionnaires."	Based on the distribution of the breastfeeding variable, this variable was categorised into "None/early cessation", "Up to 6 months", and "Extended duration". The variable is not treated as ordinal due to the possibility of qualitative differences between non-/early stopping breastfeeders and others in the sample.	The hypothesis tests regarding breastfeeding duration (H2 and H4) are now evaluated based on a likelihood ratio test to compare the fit of the model with and without this variable. Adopting this approach, rather than drawing inferences based on beta coefficients for contrasts, ensures that the alpha for these hypothesis tests remains at 0.05. This deviation should have a limited impact on readers' interpretation of results. The impact, on the results, of the specific implementation of this deviation are examined in sensitivity analyses (wherein an alternative categorisation of the breastfeeding variable is included).
Reason	New knowledge			
Timing	After data access			

Unregistered Steps

Details		Original Wording	Unregistered Step Description	Reader Impact
Type	Data Preparation	In the preregistration, we stated that we would "retain all valid responses" in section 3.5 Statistical outliers.	In order to operationalise "valid responses", we set height values ≥ 3 standard deviations from the mean to missing.	This unregistered step was an omission of detail in the preregistration and should have a limited impact on readers' interpretation of the results. The impact, on the results, of the specific
Timing	After results known			

				implementation of this step are examined in sensitivity analyses (wherein a threshold of ≥ 2 standard deviations is used).

308   3.2.2 Sharing of analytic code and metadata

After completing the analyses, the researchers report all the details of their data preparation

and analyses in the methods section of their manuscript – with much of the detail imported

from their preregistration. However, they are aware that such a report, no matter how

diligently compiled, is unlikely to be sufficiently detailed to facilitate reproducibility without the

statistical protocol or code used to implement these steps. They provide analysis scripts in

an online repository (https://osf.io/nm9ej/), organised so that its contents are easy to

navigate and understand to possible users. A readme file in the repository itemises and

describes the files it contains, and analytic scripts include comments to allow readers to

follow them and relate their contents back to the manuscript, and the results obtained by the

researchers.

319   3.2.3 Create synthetic data to allow reproducibility

The researchers wish to share their data to allow others to directly reproduce their findings.

However, the consent given by MoBa participants does not allow data to be shared openly.

Therefore, they create a synthetic dataset which preserves the quantitative relationships

between variables without including any information corresponding to real individuals from

the sample. This dataset, available in a designated data repository (https://osf.io/nm9ej/),

can be downloaded for use with the analytic code shared, allowing the code to be run and

the results reproduced. It is important to note however that synthetic data do not allow for an

exact reproduction of the original data. General quantitative relationships are maintained but

exact values are slightly different since the involved data simulation process is stochastic.

While sharing of synthetic data based on secondary analysis of cohorts like MoBa is

relatively new and subject to some specific considerations (Jordon et al., 2022; Major-Smith

et al., 2024), guidance exists to help researchers make use of this important tool for

reproducibility (e.g., Quintana, 2020). For our example, the online repository includes code

used to (a) generate synthetic data using the *synthpop* package in R (Nowok et al., 2016),

(b) verify that it does not disclose information about individuals in the real dataset, and (c)

validates that it produces similar results to the original (simulated) data. Admittedly, currently

available materials such as our own use rather simple example data. Depending on the

analysis and the data, creating accurate and appropriate synthetic data might be challenging

for complex data sets and might require particular statistical expertise. However, we believe

it is important to emphasise that even an imperfect attempt to make research more

reproducible is worthwhile, commendable and more useful than not attempting it at all.

4. Discussion

Our assessment of transparent practices in MoBa showed that some transparent practices in

MoBa are generally low. In line with assessments from other fields (Hardwicke et al., 2020;

2022; Rowhani-Farid & Barnett, 2018; Wallach et al., 2018; Bochynska et al., 2023),

preregistration, sharing of additional data, additional materials and analysis script were rare,

although some of them indicated a positive trend towards higher rates in recent years.

Preregistration is still not a common practice in secondary data analysis of the MoBa cohort.

Preregistration can be perceived as challenging due to researchers' prior knowledge of the

data, or the perception that preregistrations are inappropriate because analytical decisions

need to be made contingent on properties of the data. As illustrated by our demonstration, all

of these perceived challenges have straightforward and actionable solutions (Baldwin et al.,

2022): Researchers can preregister their plan (alongside declarations of their prior

knowledge about the data) either by using templates specific to secondary data analysis

provided on websites, such as the Open Science Framework Registry

(https://osf.io/registries) and AsPredicted (https://aspredicted.org/), or by publicly archiving

their preregistration in other formats. Necessary deviations can be accommodated and

recorded in ways that preserve false discovery rates at pre-registered levels and prioritise

analytical transparency. Additionally, more and more journals offer specific article types

called Registered Reports, which are peer-reviewed preregistrations (Nosek & Lakens,

2014)[2]. Even though the uptake of this format remains still rather limited for medical and

epidemiological journals, there are exceptions (e.g., *BMC Medicine*, *Nature Human*

Behavior).

Sharing practices of additional data, additional materials, and analysis scripts were also low

in our assessment. Raw data of MoBa can categorically not be shared because it is outside

of participants' consent and any form of derived data tables might also not easily be shared

due to legal restrictions and privacy concerns. A possible way to circumvent these issues is

the generation of synthetic data (Quintana, 2020, Major-Smith et al. 2024), i.e. simulated

data that retain statistical properties of the original data but remove all identifying

information. None of the articles reported using synthetic data, which in principle would have

allowed the reader to reproduce their analysis without violating data protection laws.

However, it is important to note that this is a relatively new approach for sharing sensitive

data and thus not widely known. To raise awareness and reassure both administrators and

participants of cohort studies about the appropriateness of this approach, we offered a

concrete example of synthetic data simulation in our demonstration. Sharing synthetic

datasets for analyses such as those performed in our example can be transformative for the

field. The availability of tailored R packages (such as *synthpop*) means that, in many cases,

the creation of synthetic data can be both a computationally and practically simple step to

implement at the end of an analytic pipeline. For more complex designs that exceed the

restrictions of available software, manual simulations assuming simple distributions can

often offer an acceptable approximation. Even in these "higher cost scenarios", we argue

that the benefits outweigh the investments, as shared synthetic data has the potential to

---

[2] Since conducting our MoBa assessment, at least one registered report on MoBa data was published (Askelund et al. 2024), albeit by authors of the present paper only. The authors used a new wave of data collection in MoBa for the analyses to ensure that no prior knowledge of the data could influence the study design, a way to strictly control bias in registered reports based on cohort studies.

improve the quality and efficiency of the peer review process, empower readers to

understand and make secondary use of analytic code, and facilitate the first phase of

attempts to reproduce or interrogate analytic findings.

Complementary to synthetic data, sharing additional data such as metadata (i.e., information

about the data) increases the findability, the transparency, and the usability of aspects of an

analysis. Certain descriptive statistics allow researchers to reverse-engineer at least some

statistical properties of numerical distributions. Despite these benefits, sharing these forms

of information has not been widely adopted for MoBa research. The repository associated

with our demonstration can be used as a model for ways to enrich shared resources with

appropriate metadata.

Whether additional or synthetic data is available or not, a step-by-step description of the

analysis in the form of a script is the minimal requirement for other researchers to reproduce

the results and evaluate how researchers arrived at their conclusions. While this practice

was by no means a widespread standard throughout the earlier years of the period from

which we assessed studies, it has become an expectation - and often mandate - of many

publishers and funders in recent years. Yet, our assessment indicates that sharing practices

of data analysis protocols are still not common. Sharing such resources, however, is

important because humans are error-prone (Nuijten et al., 2016), and methodological

descriptions often lack the detail to recreate or critically evaluate the data analysis pipeline

(Hardwicke et al., 2018). The latter is a problem, because the outcome and interpretation of

analyses has been shown to be dependent on analytical details (e.g., Silberzahn et al.,

2018; Dutilh et al., 2019; Starns et al., 2019; Coretta et al., 2023).

The robustness of the outcome and interpretation of an analysis can be assessed by probing

the sensitivity of the results to researcher degrees of freedom. Our demonstration offered

simple sensitivity checks which can straightforwardly be scaled up. Particularly

comprehensive versions of sensitivity analyses are so-called multiverse analyses (Steegen

et al., 2016), which involves identifying and implementing all potential analytic choices that

could justifiably be made to address the research question. These analyses often involve

thousands of analytical pipelines, which may be impractical for some computationally

intensive analyses of large-scale datasets. Nonetheless, sensitivity analyses at some scale

are an important element of the toolbox for conducting secondary data analysis transparently

and robustly.

Not surprisingly, power analyses are not very common for MoBa studies. Given that most

inferential analyses in research on cohort data are based on versions of the null hypothesis

significance testing framework, decision procedures are usually binary (i.e. is a relationship

significant or not). A non-significant result can either reflect a true negative, i.e. the true

absence of a relationship, or a false negative because the analysis lacked sufficient

statistical power. To interpret a non-significant result in light of its false negative rate,

analysts can a priori estimate the statistical power of a statistical test. However, power

analysis is challenging for secondary data analysis and arguably of limited use because

researchers have no control over the sample size or the measurement process.

Researchers performing secondary data analysis can give readers of their research the best

information to interpret their results in light of statistical power by ensuring that test results

are supplemented with the presentation of estimates of effect sizes and corresponding

indices of precision.

5. Conclusion

We demonstrated that some transparent practices in the Norwegian Mother, Father and

Child Cohort study (MoBa), are more common than others, with some practices being

virtually absent. It is important to acknowledge that our review assessed research published

across a span of 17 years against criteria based on *present day* best practices, not all of

which have been widely popularised for long. Nonetheless, even recent upward trends would

– if they continue – be insufficient to see a majority of researchers analyzing MoBa data as

reproducibly as possible within the next decade. The complexity of cohort corpora such as

MoBa invites a large amount of analytical flexibility. If left unchecked, these researcher

degrees of freedom, which can be either implicit or explicit, can facilitate bias and a

proliferation of overconfident claims (Simmons et al., 2011; Gelman & Loken, 2014). A

particular focus in future efforts in the field should be put on practices that help mitigating

bias due to researcher degrees of freedom – namely, preregistration, transparent sharing of

analysis scripts, and robustness checks.

With our toy example of a transparently reported MoBa analysis, we demonstrate that

researchers performing secondary analyses of cohort data have no cause to be reluctant to

adopt transparent and reproducible practices going forward. While our demonstration and its

online materials illustrate just one possible way to document and transparently share

analysis protocols, researchers without inclination to develop their own alternatives may

consider using them as templates in future work.

Despite the general low rate of transparent and reproducible practices in our assessment, it

is important to note that our assessment shows several positive trends over time. To us,

these trends indicate that researchers are becoming more aware of the importance,

usefulness, and feasibility of transparent practices. All in all, the field of secondary data

analyses on cohorts in particular, and epidemiology in general seems to be moving towards

a more transparent and reproducible future.

The adoption of transparent principles in the field of epidemiology can be further facilitated

by making it possible and easy to register, share, and publish research outputs.

Stakeholders such as funders, journals, and institutions should invest in and create

infrastructures for these practices, and either incentivise them as a currency that increases

the competitiveness of researchers or make them obligations when publishing, acquire

funding, and compete for professional positions. We hope the present paper helps in

tracking progress over time in secondary analysis of cohort data and epidemiology more

broadly, enabling cross-disciplinary comparisons, and, through our accompanying materials,

represent a useful point of departure for future efforts.

Acknowledgment

463   Not applicable.

Clinical trial number

465   Not applicable.

Funding

TR was supported by the Research Council of Norway (ICONIC; #350026); PP was
supported by the European Union's Horizon 2020 research and innovation programme
under the Marie Skłodowska-Curie grant (#801133) and by Research Council of Norway
(#324252); ADA was supported by the South-Eastern Norway Regional Health Authority
(#2020023) and the European Union's Horizon Europe Research and Innovation programme
(FAMILY; #101057529); BG was supported by the Research Council of Norway (#336085);
DQ was supported by the Research Council of Norway (#324783) and the Kavli Trust; LH
was supported by the South Eastern Regional Health Authority of Norway (Helse Sør-Øst;
#2022083, #2019097).

Author contributions

477   Authors contributed with the following roles to the present research article (following
478   Contributor Role Taxonomy ([CRediT](#)):
479
TR: Conceptualtion, Data Curation, Software, Formal Analysis, Project Administration (lead),
Supervision, Funding acquisition, Validation, Visualization, Methodology, Writing - original
draft (lead), Writing - review and editing
ADA.: Conceptualization, Data Curation, Software, Formal Analysis, Funding acquisition,
Methodology, Writing - original draft (support), Writing - review and editing
VB: Conceptualization, Methodology, Writing - original draft (support), Writing - review and
editing
LDB: Conceptualization, Formal Analysis, Methodology, Writing - original draft (support),
Writing - review and editing
AB: Conceptualization, Data Curation, Methodology, Writing - original draft (support), Writing
- review and editing
BDG: Conceptualization, Software, Formal Analysis, Validation, Methodology, Writing -
original draft (support), Writing - review and editing
TK: Conceptualization, Data Curation, Methodology, Writing - original draft (support), Writing
- review and editing
MK: Conceptualization, Data Curation, Software, Methodology, Writing - original draft
(support), Writing - review and editing

IM: Conceptualization, Data Curation, Methodology, Writing - original draft (support), Writing
- review and editing
JM: Conceptualization, Data Curation, Methodology, Writing - original draft (support), Writing
- review and editing
PP: Conceptualization, Data Curation, Software, Formal Analysis, Funding acquisition,
Validation, Methodology, Writing - original draft (support), Writing - review and editing
DSQ: Conceptualization, Formal Analysis, Methodology, Writing - original draft (support),
Writing - review and editing
LJH: Conceptualization, Software, Formal Analysis, Project Administration (support),
Funding acquisition, Validation, Methodology, Writing - original draft (support), Writing -
review and editing

Corresponding author

Correspondence to Max Korbmacher (Max.Korbmacher@hvl.nov)

Ethic approval and consent to participate

Not applicable.

Consent for publication

We confirm that all authors of the manuscript have read and agreed to its content and are
accountable for all aspects of the accuracy and integrity of the manuscript. No further
consent requirement is applicable.

Competing interests

AB, DQ, IM, JM, PP, TK and TR have engaged with meta-scientific research and have been
actively involved in methodological debates about scientific practices, including transparency
across disciplines. ADA, LDB, LH, and PP have published secondary analyses of the MoBa
cohort. All other authors have no competing interests.

Data availability statement

All dataset(s) and materials supporting the conclusions of this article are available in the
OSF repository: https://osf.io/2jqxv/.

6. References

1. Askelund AD, Wootton RE, Torvik FA, Lawn RB, Ask H, Corfield EC, et al. Assessing

causal links between age at menarche and adolescent mental health: a Mendelian

randomisation study. BMC Med. 2024;22:155.

2. Baldwin JR, Pingault J-B, Schoeler T, Sallis HM, Munafò MR. Protecting against

researcher bias in secondary data analysis: challenges and potential solutions. Eur J

Epidemiol. 2022;37:1–10.

3. Bochynska A, Keeble L, Halfacre C, Casillas JV, Champagne I-A, Chen K, et al.

Reproducible research practices and transparency across linguistics. Glossa

Psycholinguistics. 2023;2.

4. Boland MR, Karczewski KJ, Tatonetti NP. Ten Simple Rules to Enable Multi-site

Collaborations through Data Sharing. PLOS Computational Biology. 2017;13:e1005278.

5. Border R, Johnson EC, Evans LM, Smolen A, Berley N, Sullivan PF, et al. No Support for

Historical Candidate Gene or Candidate Gene-by-Interaction Hypotheses for Major

Depression Across Multiple Large Samples. AJP. 2019;176:376–87.

6. Box GE. Science and statistics. Journal of the American Statistical Association.

1976;71:791–9.

7. Brandlistuen RE, Ystrom E, Nulman I, Koren G, Nordeng H. Prenatal paracetamol

exposure and child neurodevelopment: a sibling-controlled cohort study. International

Journal of Epidemiology. 2013;42:1702–13.

8. Brugger P. From haunted brain to haunted science: A cognitive neuroscience view of

paranormal and pseudoscientific thought. In: Houran J, Lange R, editors. Hauntings and

poltergeists: Multidisciplinary perspectives. Jefferson, NC: McFarland; 2001. p. 195–213.

9. Coretta S, Casillas JV, Roessig S, Franke M, Ahn B, Al-Hoorie AH, et al. Multidimensional

Signals and Analytic Flexibility: Estimating Degrees of Freedom in Human-Speech Analyses.

Advances in Methods and Practices in Psychological Science. 2023;6:25152459231162567.

10. De Groot AD. The meaning of "significance" for different types of research. Acta

Psychologica. 1956;148:188–94.

11. Dutilh G, Annis J, Brown SD, Cassey P, Evans NJ, Hawkins GE, et al. The Quality of

Response Time Data Inference: A Blinded, Collaborative Assessment of the Validity of

Cognitive Models. Psychon Bull Rev. 2019;:19.

12. Fischhoff B. Hindsight is not equal to foresight: The effect of outcome knowledge on

judgment under uncertainty. Journal of Experimental Psychology: Human perception and

performance. 1975;1:288.

13. Gelman A, Loken E. The garden of forking paths: Why multiple comparisons can be a

problem, even when there is no "fishing expedition" or "p-hacking" and the research

hypothesis was posited ahead of time. Department of Statistics, Columbia University. 348:1–

17.

14. Gilmore RO, Kennedy JL, Adolph KE. Practical Solutions for Sharing Data and Materials

From Psychological Research. Advances in Methods and Practices in Psychological

Science. 2018;1:121–30.

15. Hardwicke TE, Thibault RT, Kosie JE, Wallach JD, Kidwell MC, Ioannidis JPA.

Estimating the prevalence of transparency and reproducibility-related research practices in

psychology (2014–2017). Perspectives on Psychological Science. 2022;17:239–51.

16. Hardwicke TE, Wallach JD, Kidwell MC, Bendixen T, Crüwell S, Ioannidis JPA. An

empirical assessment of transparency and reproducibility-related research practices in the

social sciences (2014–2017). Royal Society Open Science. 2020;7:190806.

17. Isungset MA, Conley D, Zachrisson HD, Ystrom E, Havdahl A, Njølstad PR, et al. Social

and genetic associations with educational performance in a Scandinavian welfare state.

Proceedings of the National Academy of Sciences. 2022;119:e2201869119.

18. John LK, Loewenstein G, Prelec D. Measuring the Prevalence of Questionable Research

Practices With Incentives for Truth Telling. Psychol Sci. 2012;23:524–32.

19. Jordon J, Szpruch L, Houssiau F, Bottarelli M, Cherubin G, Maple C, et al. Synthetic

Data -- what, why and how? 2022.

20. Kidwell MC, Lazarević LB, Baranski E, Hardwicke TE, Piechowski S, Falkenberg L-S, et

al. Badges to Acknowledge Open Practices: A Simple, Low-Cost, Effective Method for

Increasing Transparency. PLOS Biology. 2016;14:e1002456.

21. Klein O, Hardwicke TE, Aust F, Breuer J, Danielsson H, Mohr AH, et al. A practical guide

for transparency in psychological science. Collabra: Psychology. 2018;4.

22. Koole SL, Lakens D. Rewarding Replications: A Sure and Simple Way to Improve

Psychological Science. Perspect Psychol Sci. 2012;7:608–14.

23. Korbmacher M, Azevedo F, Pennington CR, Hartmann H, Pownall M, Schmidt K, et al.

The replication crisis has led to positive structural, procedural, and community changes.

Commun Psychol. 2023;1:1–13.

24. Lindsay DS. Sharing Data and Materials in Psychological Science. Psychol Sci.

2017;28:699–702.

25. Lowndes JSS, Best BD, Scarborough C, Afflerbach JC, Frazier MR, O'Hara CC, et al.

Our path to better science in less time using open data science tools. Nat Ecol Evol.

2017;1:1–7.

26. Magnus P, Birke C, Vejrup K, Haugan A, Alsaker E, Daltveit AK, et al. Cohort Profile

Update: The Norwegian Mother and Child Cohort Study (MoBa). International Journal of

Epidemiology. 2016;45:382–8.

27. Magnus P, Irgens LM, Haug K, Nystad W, Skjærven R, Stoltenberg C, et al. Cohort

profile: The Norwegian Mother and Child Cohort Study (MoBa). International Journal of

Epidemiology. 2006;35:1146–50.

Major-Smith, D, Kwong, AS, Timpson, NJ, Heron, J, Northstone, K. Releasing synthetic data from the Avon Longitudinal Study of Parents and Children (ALSPAC): Guidelines and applied examples. Wellcome Open Res. 2024;9:57.

28. Mertzen D, Lago S, Vasishth S. The benefits of preregistration for hypothesis-driven bilingualism research. Bilingualism: Language and Cognition. 2021;24:807–12.

29. Munafò MR, Nosek BA, Bishop DVM, Button KS, Chambers CD, Percie du Sert N, et al. A manifesto for reproducible science. Nat Hum Behav. 2017;1:1–9.

30. Nguyen RHN, Wilcox AJ, Skjaerven R, Baird DD. Men's body mass index and infertility. Human Reproduction. 2007;22:2488–93.

31. Nickerson RS. Confirmation bias: A ubiquitous phenomenon in many guises. Review of general psychology. 1998;2:175–220.

32. Nosek BA, Ebersole CR, DeHaven AC, Mellor DT. The preregistration revolution. PNAS. 2018;115:2600–6.

33. Nosek BA, Lakens D. Registered Reports. Social Psychology. 2014;45:137–41.

34. Nowok B, Raab GM, Dibben C. synthpop: Bespoke Creation of Synthetic Data in R. Journal of Statistical Software. 2016;74:1–26.

35. Nuijten MB, Hartgerink CHJ, van Assen MALM, Epskamp S, Wicherts JM. The prevalence of statistical reporting errors in psychology (1985–2013). Behav Res. 2016;48:1205–26.

36. Øksendal E, Brandlistuen RE, Holte A, Wang MV. Associations between poor gross and fine motor skills in pre-school and peer victimization concurrently and longitudinally with follow-up in school age – results from a population-based study. Brit J of Edu Psychol. 2022;92:557–75.

37. Orben A, Przybylski AK. The association between adolescent well-being and digital

technology use. Nat Hum Behav. 2019;3:173–82.

38. Pigott TD, Polanin JR. Methodological Guidance Paper: High-Quality Meta-Analysis in a

Systematic Review. Review of Educational Research. 2020;90:24–46.

39. Quintana DS. A synthetic dataset primer for the biobehavioural sciences to promote

reproducibility and hypothesis generation. eLife. 2020;9:e53275.

40. Rocher L, Hendrickx JM, de Montjoye Y-A. Estimating the success of re-identifications in

incomplete datasets using generative models. Nat Commun. 2019;10:3069.

41. Roth C, Magnus P, Schjølberg S, Stoltenberg C, Surén P, McKeague IW, et al. Folic

Acid Supplements in Pregnancy and Severe Language Delay in Children. JAMA.

2011;306:1566–73.

42. Rowhani-Farid A, Barnett AG. Badges for sharing data and code at Biostatistics: an

observational study. F1000Research. 2018;7:90.

43. Seibold H, Czerny S, Decke S, Dieterle R, Eder T, Fohr S, et al. A computational

reproducibility study of PLOS ONE articles featuring longitudinal data analyses. PLOS ONE.

2021;16:e0251194.

44. Silberzahn R, Uhlmann EL, Martin DP, Anselmi P, Ullrich J, Aust F, et al. Many analysts,

one dataset: Making transparent how variations in analytical choices affect results.

Advances in Methods and Practices in Psychological Science. 2017.

https://doi.org/10.5167/uzh-148470.

45. Simmons JP, Nelson LD, Simonsohn U. False-positive psychology: Undisclosed

flexibility in data collection and analysis allows presenting anything as significant.

Psychological Science. 2011;22:1359–66.

46. Starns JJ, Cataldo AM, Rotello CM, Annis J, Aschenbrenner A, Bröder A, et al.

Assessing Theoretical Conclusions With Blinded Inference to Investigate a Potential

Inference Crisis. Advances in Methods and Practices in Psychological Science. 2019;2:335–

49.

47. Steegen S, Tuerlinckx F, Gelman A, Vanpaemel W. Increasing Transparency Through a

Multiverse Analysis. Perspect Psychol Sci. 2016;11:702–12.

48. Sterling TD. Publication Decisions and their Possible Effects on Inferences Drawn from

Tests of Significance–or Vice Versa. Journal of the American Statistical Association.

1959;54:30–4.

49. Tukey JW. Exploratory Data Analysis. Reading, MA: Addison-Wesley; 1977.

50. Tversky A, Kahneman D. Judgment under uncertainty: Heuristics and biases. science.

1974;185:1124–31.

51. van den Akker O, Weston S, Campbell L, Chopik B, Damian R, Davis-Kean P, et al.

Preregistration of secondary data analysis: A template and tutorial. Meta-Psychology.

2021;5.

52. Vazire S. Quality Uncertainty Erodes Trust in Science. Collabra: Psychology. 2017;3:1.

53. Wallach JD, Boyack KW, Ioannidis JPA. Reproducible research practices, transparency,

and open access data in the biomedical literature, 2015–2017. PLOS Biology.

2018;16:e2006930.

54. Weston SJ, Ritchie SJ, Rohrer JM, Przybylski AK. Recommendations for Increasing the

Transparency of Analysis of Preexisting Data Sets. Advances in Methods and Practices in

Psychological Science. 2019;2:214–27.

55. Wicherts JM, Veldkamp CLS, Augusteijn HEM, Bakker M, Van Aert R, Van Assen

MALM. Degrees of freedom in planning, running, analyzing, and reporting psychological

studies: A checklist to avoid p-hacking. Frontiers in psychology. 2016;7:1832.

56. Willroth EC, Atherton OE. Best Laid Plans: A Guide to Reporting Preregistration

Deviations. Advances in Methods and Practices in Psychological Science. 2024;7(1).

57. Zwaan RA, Etz A, Lucas RE, Donnellan MB. Making replication mainstream. Behavioral

and Brain Sciences. 2018;41:e120.