Quality Improvement (QI) Project: Enhancing Surgical Safety Checklist Utilization at a Rural Primary Hospital

Authors:

- Melaku Teshale Gemechu, M.D., General Surgeon[1] (Corresponding author)
  - Email- melakuteshale0916@gmail.com
- Anteneh Cheru Adinew, M.D., Obstetrician and Gynecologist[1]
  - Email- antenehcheru34@gmail.com
- Elias Ermias Ledamo, M.D., Health Service Quality Director[1]
  - Email- eliasermias05@gmail.com

Affiliation:

1. Wondo Genet Primary Hospital, Sidama, Ethiopia

Abstract

Background: Despite evidence demonstrating the benefits of the World Health Organization (WHO) Surgical Safety Checklist (SSC), its utilization is less frequent in countries with a lower Human Development Index (HDI). This quality improvement project aimed to enhance SSC utilization in a rural primary hospital to address critical issues, including inconsistent administration of prophylactic antibiotics, frequent instances of lost or mislabeled surgical specimens, and inadequate preparedness for essential surgical instruments and blood products.

Methods: This quality improvement project was conducted at a Primary Hospital, a rural facility in Ethiopia, to address suboptimal utilization of the WHO Surgical Safety Checklist (SSC). The surgical team included 2 specialists, 2 anesthetists, and 5 nurses, with an average SSC utilization rate of 50% and a completion rate of 80% over the prior 18 months. A Plan-Do-Study-Act (PDSA) approach was employed to enhance adherence. Key interventions included translating the SSC into Amharic, conducting a two-day training session, introducing pre- and post-operative briefings, and requiring surgeons to remain in the operating room until checklist completion. Bi-weekly audits and regular supervision were conducted to monitor progress, with feedback loops established to guide adjustments. The effectiveness of the intervention was assessed through quantitative and qualitative methods. Key performance indicators included SSC utilization and completion rates, with secondary metrics such as prophylactic antibiotic administration and surgical site infection rates. Trend analysis using run charts and Interrupted Time Series (ITS) analysis evaluated changes over time, while staff interviews provided insights into behavioral and attitudinal shifts. Sustained improvements were reinforced through ongoing training, monthly supervision, and staff recognition initiatives.

Results: The quality improvement project demonstrated a significant and sustained improvement in the utilization and completion rates of the WHO Surgical Safety Checklist (SSC). Baseline measurements in October 2023 showed a utilization rate of 53% and a completion rate of 65%. Following targeted interventions, both metrics steadily improved, reaching 100% by June 2024 and maintaining this level through September 2024.

Conclusion: The QI project demonstrated that achieving 100% utilization and completion of the WHO Surgical Safety Checklist (SSC) is feasible even in rural hospitals of low- and middle-income countries (LMICs). Formal training proved crucial for improving adherence, addressing gaps seen with informal approaches, and fostering better team communication and culture. Challenges such as incomplete Sign-out sections and language barriers were mitigated by adapting the SSC to local contexts, including translation. These findings highlight the importance of structured interventions and localized solutions in enhancing surgical safety practices and inspire broader implementation in similar resource-limited settings.

Introduction

The World Health Organization (WHO) Surgical Safety Checklist (SSC) has proven effective in reducing surgical morbidity across various settings, and substantial evidence supports its role in lowering both surgical mortality and morbidity rates (1–4). Its adoption in Ethiopian hospitals is a key aspect of the Ethiopian Hospital Reform Implementation Guidelines (EHRIG). Moreover, One of the Ethiopian Ministry of Health's perioperative quality improvement targets for 2025 is achieving 100% utilization of the SSC (5).

The utilization of the SSC is generally high in various countries; however, there is significant diversity across resource contexts and patient groups, even within countries with similar resources. While most facilities are reported to use the SSC, it is only done with reasonable consistency. On average, the SSC is used less frequently in countries with a lower Human Development Index and in those where the common or official languages are not among the six WHO official languages. Significant gaps exist in the usage of SSC for patients undergoing less sophisticated or more urgent surgeries, as well as for obstetric and gynecological procedures (6). Various studies have identified barriers to effective SSC implementation, including the need for changes in the workflow of healthcare professionals, as well as in their perception of the checklist and patient safety in general. The factors that hinder or promote the necessary changes are focused on the checklist itself, the implementation process, and the local context, including language, staffing levels, types of surgeries, patient load, and presence of healthy

teamwork (6,7). The absence of positive role models, unenthusiastic team members, hierarchical obstacles, limited knowledge of proper usage, and inadequate implementation procedures have also been reported as barriers to effective SSC adoption (8).

At Wondo Genet Primary Hospital (WGPH), a newly functioning rural facility with an inexperienced operation team, adherence to the WHO Surgical Safety Checklist (SSC) was initially poor. The checklist, designed to ensure critical safety steps are followed, was often underused or inadequately completed. This lack of compliance led to issues such as inconsistent administration of prophylactic antibiotics, which are crucial for preventing surgical site infections. There were also frequent cases of lost or mislabeled surgical specimens, compromising post-operative care. Moreover, the inconsistent use of the SSC hindered preparedness for essential surgical instruments and blood products, both critical for managing complex surgeries.

Recognizing these gaps, it became evident that a quality improvement (QI) project was necessary to enhance SSC adherence. Implementing a structured QI initiative would standardize procedures and ensure all team members consistently followed the checklist. This would improve antibiotic administration, reduce specimen handling errors, and ensure preparedness for surgical instruments and blood products, ultimately minimizing delays and complications. Focusing on the SSC as part of the QI efforts would build a safety culture, leading to better patient outcomes and greater operational efficiency (9,10).

Methods

Context

Wondo Genet Primary Hospital, established in September 2022, is 22 km from Hawassa, Ethiopia. The hospital's surgical unit is staffed by a team that includes one general surgeon, one Obstetrician and Gynecologist, one integrated emergency surgical officer, two anesthetists, five scrub nurses, and two cleaners. The hospital provided major surgical service for 226 patients over the 18 month period.

Despite utilization of the WHO Surgical Safety Checklist (SSC) since the start of surgical services, the hospital has not met the national target of 100% SSC completion over the past 18 months. This gap in compliance presents a significant challenge in meeting national surgical safety standards. After identifying four areas for improvement in the surgical unit, poor utilization of the Surgical Safety Checklist (SSC) was ranked as the top priority based on a prioritization matrix. Over the past six months, the average utilization of the SSC in the WGPH operating room case team was only 50%, with a completion rate of 80%. This low adherence is likely to lead to increased surgical morbidity. Hence, the project aimed to achieve 100% utilization and completion of the SSC after 3 months of intervention, by the end of June 2024.

Baseline measurement

Data were initially gathered from patient charts identified through the OR registry over the past year. Each chart was reviewed for the presence and completeness of the WHO Surgical Safety Checklist (SSC), and the findings were recorded using

an audit tool. The results are displayed in Figure 1. The analysis revealed an average SSC utilization rate of 50% and a

completion rate of 80%. Non-utilization was particularly common in emergency and late-night procedures, with the sign-out

section being the most frequently incomplete part of the checklist.

Figure 1: Surgical safety checklist utilization and completeness rate from Oct/2023- Mar/2024.

Figure 2: Fishbone diagram depicting factors contributing to poor SSC utilization.

Intervention

In our project aimed to improve the utilization and completion of the WHO SSC, we used a driver diagram to systematically map out the key factors influencing adherence. We aimed to increase both the SSC utilization rate of 50 % and completion rate of 80 % to 100 % in operating rooms within three months. We identified several primary drivers, including enhanced staff training, leadership commitment, and effective communication between surgical teams. These were further broken down into secondary drivers, such as conducting regular SSC briefings, fostering a culture of safety, and ensuring accountability through monitoring. Change ideas, such as implementing daily reminders for staff, revising the SSC form to make it more user-friendly translating it to the local language, and conducting refresher training sessions, were derived from these drivers. The driver diagram, shown below (Figure 3), helped our team visualize the relationships between these factors, enabling us to focus our interventions on areas most likely to influence improvement, and guiding our efforts through targeted PDSA cycles.

Figure 2: A driver diagram illustrating the primary and secondary drivers, along with the planned change ideas to achieve the target.

Strategy

The Plan-Do-Study-Act (PDSA) QI model served as the framework for this project, drawing on the success of similar past initiatives (9,10). After implementing the intervention strategy, the hospital's QI director and one operating room (OR) team member audited WHO Surgical Safety Checklist (SSC) compliance and completeness every two weeks. Daily monitoring was conducted throughout the three month period. Here's how the PDSA cycle was applied to our QI project with the change ideas listed above:

Plan:

In the planning phase, the team identified key areas for improvement in the compliance and completeness of the WHO Surgical Safety Checklist (SSC) within the Operating Room (OR). Based on previous audits, low utilization and incomplete sections, particularly during emergency and late-night surgeries, were major issues. To improve the utilization of the Surgical Safety Checklist (SSC) at Wondo Genet Primary Hospital, several targeted interventions will be implemented. The SSC will be translated into Amharic to enhance understanding and facilitate its consistent use among staff. Additionally, a two-day onsite training led by a General Surgeon and an Obstetrician and Gynecologist will be conducted to improve staff awareness and skills in using the SSC effectively. To ensure proper checklist completion, surgeons will be required to remain in the operating room until the third part of the SSC (sign-out section) is completed. The head nurse will oversee printing 1.5 times the SSC formats used in the previous month to ensure availability and maintain accurate records by discarding incomplete or vague forms.

Furthermore, pre- and post-operative briefings will be initiated to foster open communication and reduce hierarchical barriers within the surgical team.

To sustain these efforts, shared accountability and regular monitoring will be prioritized. Staff will take turns conducting audits to promote shared responsibility for compliance, and a focal supervisor will perform bi-monthly audits to identify and address gaps. Additionally, the QI director will conduct monthly supervision to review SSC adherence and guide improvements. To motivate the team, bi-annual staff recognition days will be introduced, celebrating individuals who excel in SSC compliance. These measures aim to create a sustainable culture of safety and accountability, ultimately improving surgical outcomes and patient care at the hospital.

Do:

Over a three-month period, the team implemented a series of planned interventions to improve the utilization of the Surgical Safety Checklist (SSC). An Amharic-translated version of the SSC was introduced to enhance understanding and facilitate usage among staff. To build capacity, a two-day training session was conducted on February 28 and 29 by a general surgeon and an Obstetrician and Gynecologist, focusing on the importance and proper application of the SSC. Workflows were also adjusted to ensure surgeons remained in the operating room until the SSC was fully completed, emphasizing accountability and thoroughness. Pre- and post-operative briefings were systematically introduced for every surgical case, fostering open communication and addressing hierarchical barriers.

To monitor the effectiveness of these interventions, bi-weekly audits and regular supervision were conducted. These audits aimed to track compliance with the new processes, identify gaps, and provide timely feedback to staff. Supervision ensured consistent adherence to the checklist and allowed for the resolution of challenges as they arose. This structured approach helped establish a culture of accountability and collaboration, with the ultimate goal of improving patient safety and surgical outcomes.

Study:

During the implementation period, data was systematically collected through bi-weekly audits to evaluate the effectiveness of the interventions. These audits focused on SSC compliance, utilization, and completeness. A notable improvement in SSC utilization was observed, driven largely by the training sessions and the introduction of pre- and post-operative briefings. Staff interviews confirmed that these changes had an immediate impact, enhancing their understanding and commitment to the checklist. The adjustments made to workflows also contributed to higher completion rates, particularly during emergency and late-night surgeries, as staff availability and processes became more streamlined.

The audits also revealed enhanced adherence to the sign-out process, previously identified as the most neglected section of the SSC. This improvement was attributed to the requirement that surgeons remain in the operating room until the checklist was fully completed. Additionally, the pre- and post-operative briefings received overwhelmingly positive feedback from staff, who noted that these sessions reinforced the importance of teamwork and the SSC. Collectively, these findings demonstrate the effectiveness of the interventions in fostering a culture of safety and accountability within the surgical team.

Act:

Based on the results from the study phase, several actions were taken to sustain and build upon the improvements achieved. Regular training sessions were scheduled every three months to address areas where compliance remained low, ensuring continuous staff development. The successful Amharic translation of the SSC was extended for an additional three months to further enhance understanding and completion rates. Monthly supervision by the QI director and bi-annual staff recognition were formally integrated into the hospital's standard practices to maintain accountability and motivation. Additionally, the practice of rotating audit responsibilities among all operating room staff was continued, fostering shared responsibility and sustained compliance with the SSC.

Study of the intervention

Assessing the Impact of the Intervention

1. Key Performance Indicators (KPIs):

    a. Primary Indicators: The rate of SSC compliance (% of surgeries where the SSC is utilized) and rate of completed SSC (% of fully completed SSC among the utilized checklists).

    b. Secondary Indicators: The rate of prophylactic antibiotics administration when its applicable, SSI rates, and rate of lost or unlabeled surgical specimens.

2. Trend analysis was done using run charts to visualize trends in SSC utilization and completion rate. Interrupted Time Series (ITS) analysis depicting the collect data at multiple time points before and after the intervention will also be used.

3. Interviews with staff were conducted to gather qualitative insights on changes in attitudes, behaviors, and perceived outcomes after the interventions. The staffs were also interviewed to assess whether they attribute observed improvements to the interventions.

Ethical considerations

- Patient consent for publication: Not required.

Results

The QI project showed gradual improvement in the utilization and completeness of the WHO Surgical Safety Checklist (SSC) over three months. By June, the project achieved a 100% rate for both metrics, highlighting the effectiveness of the implemented changes and the team's commitment to enhancing patient safety in the operating room. Figures 4 and 5 illustrate the results. No missed or unlabeled surgical specimens were reported, and prophylactic antibiotics were administered to all eligible cases during the intervention period. There were instances of missed necessary equipment during surgery, primarily due to resource limitations. These issues were addressed by using substitute or similar items.

Figure 3: Audit result of utilization rate of the WHO Surgical Safety Checklist from October 2023 – September 2024

Figure 4: Audit result of completion rate of the WHO Surgical Safety Checklist from October 2023 – September 2024.

Discussion

The QI project aimed to achieve a 100% utilization and completion rate of the WHO Surgical Safety Checklist (SSC), based on the assumption of an overall low surgical case burden in the facility. Formal training was essential, as the initial informal approach proved ineffective in improving WHO Surgical Safety Checklist (SSC) adherence. This finding aligns with

various studies that highlight the importance of structured education in driving compliance with clinical protocols (8,11–14). The formal training not only increased staff understanding but also reinforced the consistent use of the SSC, contributing to the overall success of the QI project.

In low- and middle-income countries, especially in rural hospitals, the utilization of the WHO Surgical Safety Checklist (SSC) is often low (6,15,16). Language barriers have been identified as a significant factor contributing to this issue, as noted in other studies(6,13). Addressing these barriers, such as by translating the SSC into local languages, could facilitate easier implementation and improve adherence among surgical staff.

The "Sign-out" section of the three-part checklist was frequently left incomplete compared to others, primarily due to uncertainty about when it should be filled and because surgeons frequently departed the operating room before the checklist was fully completed. Other studies showed similar results (6,13).

While good communication is essential for effective SSC utilization, the reverse also proved true in our OR, as team communication significantly improved following the full implementation of the SSC. Similar studies support this idea, although there is uncertainty about the direction of the effect. Proper SSC utilization not only enhances communication but also fosters a stronger culture of teamwork within the surgical environment (13,17).

Conclusion

This initiative intended to demonstrate that such high compliance rates were attainable even in rural hospitals of low- and middle-income countries (LMICs), where completion rates are typically lower. By successfully achieving 100% for both utilization and completeness, the project provided a compelling example that rigorous adherence to safety protocols can be implemented in these settings. Adopting the checklist in the local context and language, and addressing context-associated barriers played a major role. The results aimed to inspire other rural hospitals to adopt similar practices, ultimately enhancing patient safety and surgical outcomes in regions often underserved in terms of healthcare quality.

References

1. Haynes AB, Weiser TG, Berry WR, Lipsitz SR, Breizat AHS, Dellinger EP, et al. A Surgical Safety Checklist to Reduce Morbidity and Mortality in a Global Population. N Engl J Med. 2009 Jan 29;360(5):491–9.

2. Bergs J, Hellings J, Cleemput I, Zurel Ö, De Troyer V, Van Hiel M, et al. Systematic review and meta-analysis of the effect of the World Health Organization surgical safety checklist on postoperative complications. Br J Surg. 2014 Jan 27;101(3):150–8.

3. Haugen AS, Sevdalis N, Søfteland E. Impact of the World Health Organization Surgical Safety Checklist on Patient Safety. Anesthesiology. 2019 Aug 1;131(2):420–5.

4. Martis WR, Hannam JA, Lee T, Merry AF, Mitchell SJ. Improved compliance with the World Health Organization Surgical Safety Checklist is associated with reduced surgical specimen labelling errors. N Z Med J Online. 2016;129(1441):63.

5. Kifle F, Iverson KR, Belay E, Buno Teko E, Dawit A, Deneke A, et al. Towards Establishing a National Perioperative Quality Improvement Network in LMICs: Implementation Experiences From Ethiopia. Ann Surg Open. 2024 Aug 19;5(3):e480.

6. Delisle M, Pradarelli JC, Panda N, Koritsanszky L, Sonnay Y, Lipsitz S, et al. Variation in global uptake of the Surgical Safety Checklist. Br J Surg. 2020 Jan 5;107(2):e151–60.

7. Bergs J, Lambrechts F, Simons P, Vlayen A, Marneffe W, Hellings J, et al. Barriers and facilitators related to the implementation of surgical safety checklists: a systematic review of the qualitative evidence. BMJ Qual Saf. 2015 Dec;24(12):776–86.

8. Sendlhofer G, Mosbacher N, Karina L, Kober B, Jantscher L, Berghold A, et al. Implementation of a Surgical Safety Checklist: Interventions to Optimize the Process and Hints to Increase Compliance. Quach C, editor. PLOS ONE. 2015 Feb 6;10(2):e0116926.

9. Bete DY, Sibhatu MK, Godebo MG, Abdulahi IJ, Liyew TW, Minas SM, et al. Improving surgical safety checklist utilisation at 23 public health facilities in Ethiopia: a collaborative quality improvement project. BMJ Open Qual. 2023 Nov 1;12(4):e002406.

10. Röhsig V, Maestri RN, Parrini Mutlaq MF, Brenner De Souza A, Seabra A, Farias ER, et al. Quality improvement strategy to enhance compliance with the World Health Organization Surgical Safety Checklist in a large hospital: Quality improvement study. Ann Med Surg. 2020 Jul;55:19–23.

11.   Sutton-Smith L. A quality improvement project to improve the identification and management of delirium. Nurs Crit Care. 2021 May;26(3):183–9.

12.   Melekie TB, Getahun GM. Compliance with Surgical Safety Checklist completion in the operating room of University of Gondar Hospital, Northwest Ethiopia. BMC Res Notes. 2015 Dec;8(1):361.

13.   Bashford T, Reshamwalla S, McAuley J, Allen NH, McNatt Z, Gebremedhen YD. Implementation of the WHO Surgical Safety Checklist in an Ethiopian Referral Hospital. Patient Saf Surg. 2014;8(1):16.

14.   Alidina S, Chatterjee P, Zanial N, Alreja SS, Balira R, Barash D, et al. Improving surgical quality in low-income and middle-income countries: why do some health facilities perform better than others? BMJ Qual Saf. 2021 Dec;30(12):937–49.

15.   Hellar A, Tibyehabwa L, Ernest E, Varallo J, Betram MM, Fitzgerald L, et al. A Team-Based Approach to Introduce and Sustain the Use of the WHO Surgical Safety Checklist in Tanzania. World J Surg. 2020 Mar;44(3):689–95.

16.   White MC, Baxter LS, Close KL, Ravelojaona VA, Rakotoarison HN, Bruno E, et al. Evaluation of a countrywide implementation of the world health organisation surgical safety checklist in Madagascar. PLOS ONE. 2018 Feb 5;13(2):e0191849.

17.   Pugel AE, Simianu VV, Flum DR, Patchen Dellinger E. Use of the surgical safety checklist to improve communication and reduce complications. J Infect Public Health. 2015 May 1;8(3):219–25.