Estimates of epidemiological parameters for H5N1 influenza in humans: a rapid review

Authors: Jack Ward[1], Joshua W. Lambert[1,*], Timothy W. Russell[1,2,*], James M. Azam[1], Adam J. Kucharski[1], Sebastian Funk[1], Billy J. Quilty[3], Oswaldo Gressani[4], Niel Hens[4,5], W. John Edmunds[1]

Affiliations
1. Centre for the Mathematical Modelling of Infectious Diseases, Department of Infectious Disease Epidemiology, London School of Hygiene & Tropical Medicine, London, UK.
2. The Francis Crick Institute, 1 Midland Road, London NW1 1AT.
3. Charité Center for Global Health, Charité Universitätsmedizin Berlin, Berlin, Germany.
4. Data Science Institute, I-BioStat,, Hasselt University, Hasselt, Belgium.
5. Centre for Health Economic Research and Modelling Infectious Diseases, Vaccine and Infectious Disease Institute, University of Antwerp, Antwerp, Belgium

* These authors contributed equally and share second authorship

Abstract

Background

The ongoing H5N1 panzootic in mammals has amplified zoonotic pathways to facilitate human infection. Characterising key epidemiological parameters for H5N1 is critical should it become widespread.

Aim

To identify and estimate critical epidemiological parameters for H5N1 from past and current outbreaks, and to compare their characteristics with human influenza subtypes and the 2003 Netherlands H7N7 outbreak.

Methods

We searched PubMed, Embase, and Cochrane Library for systematic reviews reporting parameter estimates from primary data or meta-analyses. To address gaps, we searched PubMed and Google Scholar for studies of any design providing relevant estimates. We estimated the basic reproduction number for the outbreak in the US and the 2003 Netherlands H7N7 outbreak. In addition we estimated the serial interval for H5N1 using data from previous household clusters in Indonesia. We also applied a branching process model to simulate transmission chain size and duration to assess if simulated transmission patterns align with observed dynamics.

Results

From 46 articles, we identified H5N1's epidemiological profile as having lower transmissibility ($R_0 < 0.2$) but higher severity compared to human subtypes. Evidence suggests H5N1 has a longer incubation (~4 days vs ~2 days) and serial intervals (~6 days vs ~3 days) than human subtypes, impacting transmission dynamics. The epidemiology of the US H5 outbreak is

similar to the 2003 Netherlands H7N7 outbreak. Key gaps remain regarding latent and

infectious periods.

Conclusions

We characterised critical epidemiological parameters for H5N1 infection. The current U.S.

outbreak shows lower pathogenicity, but similar transmissibility compared to prior outbreaks.

Longer incubation and serial intervals may enhance contact tracing feasibility. These

estimates offer a baseline for monitoring changes in H5N1 epidemiology.

Introduction

Highly Pathogenic Avian Influenza A (H5N1) was first isolated in 1997 in Hong Kong [1]. The

primary reservoir of H5N1 is aquatic birds [2], with occasional human outbreaks occurring in

the last two decades [3]. Following reassortment with low pathogenic avian influenza viruses

in the Western hemisphere in 2020, new reassortant H5N1 genotypes (clade 2.3.4.4b) have

spread globally in wild bird populations, and have caused devastating outbreaks in domestic

flocks [4]. In addition, there is evidence of sustained mammal-to-mammal transmission of

H5N1 in European fur farms (genotype BB), marine mammals in South America (B3.2) and

dairy cattle in the United States (US) (B3.13) [4]. Subsequently, a second clade 2.3.4.4b

genotype, D1.1, has been detected in U.S. dairy cattle [5] .

The outbreak in US dairy cattle was first detected in February 2024, in Texas [6], before

spreading to other states [7]. As of 1st April 2025, there have been 70 documented human

cases. 41 of which have been linked to exposure to dairy herds, 24 to poultry farming and

culling, two linked to other animal exposure and three without any known exposure to sick or

infected animals [8] (**Supplementary Table S3**). All cases where genetic sequencing has

been reported (*N*=51) have been attributed to infection from clade 2.3.4.4b viruses

(**Supplementary Table S3**).

H5N1 continues to pose a threat to global biosecurity. Given the risk of viral reassortment

within a dually infected human or other mammalian species infected with a human influenza

virus, there is a possibility that the resulting variant could be capable of sustained human-to-

human transmission [9]. Longini et al. 2005 [9], emphasise the importance of key

epidemiological parameters required at the start of an outbreak; the reproduction number,

incubation, latent and infectious periods, the serial interval and the case fatality risk (CFR).

Outside of CFR and the reproduction number [10,11], the key parameters for H5N1 have not

been well characterised in relation to human influenza. Given increased mammalian H5N1

infection and the increasing level of human exposure at the human-mammal interface, these

parameters become key to understanding how H5N1 can be controlled compared to

traditional human influenza outbreaks.

This rapid review examines these critical epidemiological parameters for H5N1,

incorporating seropositivity data, and provides a comparative analysis with other influenza

subtypes. To address gaps in existing estimates, we utilised mathematical models to

estimate the reproduction number for the current U.S. outbreak and calculated the serial

interval using data from previous household clusters in Indonesia. Additionally, we applied a

branching process model to simulate transmission chain size and duration, enabling us to

assess whether simulated patterns align with observed outbreak dynamics.

____

Methods

Given the emerging situation of H5N1 in the US as well as globally there is a need to rapidly assess evidence of its epidemiology and compared to human influenzas. Rapid reviews are valuable in accelerating this process to provide actionable and relevant evidence to make informed decisions [12]. To this end, we conducted a rapid review to identify existing estimates of critical epidemiological parameters for H5N1 and to compare these with other human influenza subtypes.

Search strategy

This review incorporated two separate but related key questions, we prepared separate search strategies to answer each question. Key question 1: What are the epidemiological parameters of Highly Pathogenic Avian Influenza A (H5N1) in humans and how does it compare to the human epidemiology of pandemic and seasonal influenza strains? Key question 2: How does the epidemiology of H5N1 compare to H7N7 in the Netherlands?

Key question 1

In order to gather information on the epidemiological parameters of H5N1 in humans and pandemic and seasonal influenza strains, we undertook a staged search to first identify existing systematic reviews and another search to identify missing or more recent studies with other designs [13].

We conducted an initial search for systematic reviews in PubMed, Embase, and Cochrane Library. We included systematic reviews that provided parameter estimates for the reproduction number (R_0), dispersion parameter (k), incubation period, latent and infectious periods, CFR, or seroprevalence estimates (for H5N1 only) in humans. In addition to CFR we expanded the search to include the Infection fatality risk (IFR). CFR may be an

overestimate in some settings due to mild and subclinical infections not being detected [14].

IFR estimates help categorise the true risk of death from infection [14].

The search terms used are listed in **Supplementary Table S1**. Studies were included if they

provided quantitative estimates based on primary epidemiological data or meta-analyses of

such data. Studies were excluded if they lacked quantitative estimates, focused solely on

animal models, or did not specify influenza subtype.

Following the initial search, a tailored search was conducted in PubMed and Google Scholar

(**Supplementary Table S1**) to address gaps and identify missing parameters. This search

targeted studies reporting estimates for the dispersion parameter, incubation period, latent

and infectious periods, and CFR in humans, as these were not adequately covered by the

initial search. Search terms were narrowed (**Supplementary Table S1)** to capture specific

epidemiological parameters. We included studies of any design that provided relevant

estimates using epidemiological data. We excluded studies which reported assumed values

(parameters had been assumed for modelling purposes and had been based on opinion or

other data sources). The Google Scholar search was limited to the first 30 results per query

(sorted by relevance) to further streamline the review process. In addition we included

pertinent studies and grey literature which had been identified without a focused search

strategy.

For each included study, the study design, the study description, parameter estimate,

subtype were extracted. Where individual data points were obtained from systematic

reviews, details of the original primary studies referenced in those reviews were also

extracted. This information was consolidated into a central database, which is provided as a

supplementary resource. Ethical approval was not required as this was a review of published

literature.

A meta-analysis was not conducted due to the large heterogeneity across the included

studies. Additionally, the limited availability of comparable data for certain parameters, such

as the dispersion parameter and infectious periods, constrained the practicality of pooling

results. Given the evolving situation, a narrative approach was deemed more appropriate for

summarising existing evidence and identifying gaps in the literature.

Key question 2

Given the high rates of conjunctivitis reported among US H5N1 cases [15], it was deemed

useful to provide an epidemiological comparison to the 2003 H7N7 outbreak in the

Netherlands where cases displayed a similar phenotype [16]. We conducted a search to

identify related studies in PubMed using the following search terms (H7N7 OR "influenza A

virus H7N7") AND (Netherlands) AND (2003) AND (outbreak OR epidemiology). We

included studies of any design that provided relevant estimates using epidemiological data.

Data extraction was conducted in the same method as mentioned previously.

Reproduction number estimation

We estimated the basic reproduction number for the current outbreak in the US using the

package {epichains} [17]. We estimated the R_0 distribution under two scenarios, making an

assumption that cases with unknown exposure were infected from an undetected case of

H5N1. Scenario 1: assuming 67 independent spillover cases and three clusters of two

cases: the Missouri case of unknown exposure, and the two Californian cases of unknown

exposure each with a hypothetical source case. Scenario 2: assuming 67 independent

spillover cases, a cluster of 3 (the Missouri case, their probable household contact, and a

hypothetical source case), and two clusters of two (both Californian cases). Full details are

given in the Supplementary Information. List of H5N1 cases and their exposures are listed in

Supplementary table S3.

For H7N7 we estimated the R_0 for the 2003 outbreak in the Netherlands using the same

method. Assuming 84 independent spillover cases and a cluster of 5 (2 spillover cases and 3

household members) [18].

Serial interval estimation

We did not identify any primary estimates for the serial interval of H5N1 in the literature. To

address this, we estimated the serial interval using data collected by Aditama et al. 2012 [19]

from 22 human cases not exposed to zoonotic sources during H5N1 outbreaks in Indonesia

between 2005 and 2009 (**Supplementary Figure S4**). We fitted lognormal and gamma

distributions to the number of onsets for a given day from this data to estimate the serial

interval. Full details are given in the Supplementary Information.

Outbreak size distribution

To assess how varying transmission parameters influence H5N1 outbreak size and duration,

we used a branching process model to explore the transmission chain size and length,

across scenarios of varying reproduction numbers, dispersion (k), and offspring distributions.

This was across all possible combinations of parameters. We used a reproduction number

between 0 and 1.1 at 0.1 intervals, and for the Negative Binomial offspring distribution we

varied the dispersion parameter (k) at 0.1, 0.5, 5, 1,000. Outbreaks with no secondary

transmission were categorised as size and length 0. Full details for the branching process

are given in the Supplementary Information.

Computational details

220 The branching process simulation is implemented in the {epichains} R package [17], plots

221 use the {ggplot2} R package [20]. All analyses were run using R version 4.4.1 (2024-06-14)

222 [21]. All code to reproduce this report (including serial interval estimation) is available on

223 GitHub at https://github.com/cmmid/h5n1_uk_scenario_modelling.

226 ____

Results

228 We included 42 articles in this review (**Supplementary Figure S1**), comprising 7 systematic

229 reviews, 35 studies of other types and 2 sources of grey literature.

H5N1 transmission dynamics

H5N1 R_0 estimates are usually estimated well below 1, four estimates were reported in a

systematic review by Biggerstaff et al. 2014 [10], ranging from 0—0.25 [19,22], with one

outlier: Yang et al. reported an R_0 of 1.14 (95% CI: 0.61—2.14) during a 2006 household

outbreak in Indonesia [23]. Two additional studies reported estimates of 0.06 (95% CI: 0.01–

0.2) and 0.2 [24,25]. Biggerstaff et al. 2014 [10] summarised R_0 estimates for previous

influenza pandemics and seasonal influenza, reporting median values between 1.27 and

1.80 (**Figure 1**). Values were generally higher in confined settings compared to community

settings [10]. We identified two studies not included in this systematic review; both estimates

were from the 2009 pandemic, reporting values of 1.6 in Korea [26] and 1.32 (95% CI:1.29–

1.36) in Spain [27].

In our analysis, we estimated the median R_0 for the US outbreak as 0.04 (95% CrI: 0.02–

0.09) for scenario 1, and 0.05 (95% CrI: 0.02–0.10) for scenario 2 (**Supplementary Figure

S2a**). For scenario 1, the posterior estimate for *k* was estimated as a median of 1.45 (95%

CrI: 0.05–4.26) and 4.0 (95% CrI: 0.14–10.9) for scenario 2. One study provided a *k*

estimate for H5N1 of 0.75 [25], while Fraser et al. estimated *k* for H1N1 at 0.94 (95% CI:

0.59—1.72) [28].

Figure 1. Reproduction number estimates for influenza

Estimates for H5N1 from the current US outbreak and previous outbreaks where a reproduction number was reported. Compared to estimates from previous influenza pandemics (Median point estimate of R in the community setting for all waves of illness). Solid coloured bars represent the uncertainty around the central estimate, which was reported as IQR, and 95% CI.

We used a branching process model to further illustrate H5N1 transmission dynamics. With

a Poisson offspring distribution, most outbreaks result in no secondary transmission when R

< 0.7, and clusters of more than 20 are rare but can occur when R > 0.4 but outbreaks of at

least 20 cases only happen in at least 10% of outbreaks once R > 0.9 (**Figure 2A**).

Introducing moderate heterogeneity (k = 0.5) on average reduces secondary transmission

and even for R = 1.1 less than 10% of outbreaks exceed 50 human-to-human cases (**Figure**

2). Variations in k, from 0.1 (highly heterogeneous) to 1,000 (approximately Poisson), show

that greater heterogeneity leads to smaller and shorter outbreaks on average [29], typically

lasting fewer than five transmission generations (**Figure 2B & Supplementary Figure S5)**.

These patterns align with observed dynamics, suggesting limited potential for large

outbreaks under current transmission parameters.

Figure 2. Simulated H5N1 outbreak size

Simulated outbreak size (secondary cases excluding the index case) using a single-type branching process model across values of the reproduction number (R), with (A) a Poisson offspring distribution and (B) a Negative binomial offspring distribution. The latter with varied values of k between 0.1 and 1,000.

Incubation period

A systematic review by Bui et al. 2015 [30] reported three incubation period estimates for

H5N1 ranging from 3.3–5.0 days [31–33]. Two additional studies reported estimates ranging

from 2–9.5 days [23,34]. 11 studies (including one systematic review) reported 12 incubation

period estimates for human influenza subtypes. Estimates for Influenza A subtypes ranged

from 1.34 to 2.33 [35–45]. One estimate was identified for Influenza B, with a median

incubation period of 0.6 days [40] (**Figure 3**).

Figure 3. Incubation period estimates across influenza subtypes

Incubation period estimates for H5N1 from previous outbreaks, compared to estimates for other influenza subtypes. Mean estimates are represented by triangle points, and median estimates are shown using circle points. Uncertainty is represented 95% CI, IQR and range.

Latent & Infectious period

We could not identify any estimates of the latent period distribution for human H5N1

infections. The latent period for human influenza subtypes were reported in four studies with

mean values ranging from 0.4 to 2.62 days [41,46–48] (**Supplementary Figure S6**). There

was limited literature regarding the infectious period for H5N1. One study reported a

probable range of 5–13 days for a household outbreak in Indonesia 2006 [23]. Six studies

reported eight estimates for the infectious period for human influenza subtypes, the mean

infectious period ranged from 1 to 3.9 days [26,41,46–49] (**Supplementary Figure S6**).

Serial interval

From the outbreak in Indonesia we estimated the serial interval distribution to have a median

of 6.8 days (95% CrI: 0.3–13.3) when fit to a gamma distribution, and a median of 6.4 days

(95% CrI: 0.3–12.7) when fit to a lognormal distribution (**Supplementary Figure S4**). The

results from the Leave-One-Out (LOO) analysis suggested that the gamma distribution

would predict unseen data more accurately, and is therefore the preferable parametric

distribution in this case (**Supplementary Table S2**). We did not identify other studies

providing serial interval estimates for H5N1 specifically.

A systematic review by Vink et al. 2014 [50], listed estimates for human influenza subtypes.

Estimates ranged from 1.9 to 4.66 days for H1N1, 1.9 to 5 days for H1N1pdm09, 3.1 to 3.5

for H3N2 and 3.4 to 4.9 for Influenza B respectively [50] (**Figure 4**). Additionally the authors

report seven estimates which were estimated from index case–to–case intervals, these

range from 1.7–3.7 days [50] (**Figure 4**). We found two articles not included in this

systematic review. These estimates ranged from 2.1–3.4 [41,48] (**Figure 4**).

Figure 4. Serial interval estimates across influenza subtypes

Serial interval estimates for H5N1 from previous outbreaks, compared to estimates for other influenza subtypes. Mean estimates are represented by triangle points, and median estimates are shown using circle points. Uncertainty is represented by 95% CI, 95% CrI and range. Data source studies for the Vink et al. estimates are referenced in the data collection document. The upper 95% CrIs for the two estimated values extend to 13.3 (gamma) and 12.6 days (log normal)

Serological evidence of H5N1 human infections

We identified a systematic review by Chen et al. [51], which provides a comprehensive estimate of serological evidence of human infections with H5N1, across populations with different occupational and behavioural exposures [51].   Studies adhering to the WHO seropositivity criteria (neutralising antibody titer ≥1:80 confirmed by a second assay, such as

hemagglutination inhibition test [HAI, titer ≥1:160], enzyme-linked immunosorbent assay, or

western blot [51]) reported higher seroprevalence among high-risk occupational groups,

particularly poultry cullers and workers [51]. In contrast, no seropositive results were

detected among close contacts of cases in healthcare, household, or social settings [51]

(**Figure 5**). Chen et al. also reported estimates of seroprevalence utilising non-standardised

antibody titer criteria (different antibody titer threshold defined by each study [51]) (**Figure 5**).

Since this review was published, there have been three further serological studies in high-

risk occupational settings specific to the 2.3.4.4b clade. Gomaa et al. reported an estimated

seroprevalence of 4.6% (95% CI: 3.3—6.2) in workers exposed to poultry infected with clade

2.3.4.4b H5N1 in five live bird markets in Egypt [52]. The criteria for seropositivity was not

clearly defined in this study. In the US, Shittu et al. conducted microneutralization assays

(MN) on sera samples taken from 14 recently symptomatic farm workers at two Texas dairy

farms. Two (14.3% 95% CI: 4.0—40) showed evidence of having neutralising antibodies to a

recombinant influenza A H5N1 virus [53], although only MN assay results were reported. A

larger study by Mellis et al. (2024) analysed sera samples from 115 dairy workers from dairy

farms in Michigan and Colorado [54]. Workers were deemed eligible for sampling if they had

worked on dairies with herds with laboratory-confirmed infection with HPAI A(H5) viruses

within 90 days prior to sampling and had reported no illness on the day of specimen

collection [54]. Out of the 115 workers, 8 (7%, 95% CI: 3.6—13.1) had serological evidence

of recent infection with H5N1. All positive cases reported milking cows or cleaning the

milking parlour [54]. This study reported neutralising antibody titers and HAI antibody titers

≥1:40 which does not meet the WHO criteria.

Figure 5. Serological evidence of H5N1 human infections

Relevant serological evidence of H5N1 human infections across estimates from 1997—2020, Egypt 2022—2023 and the current outbreak in the US. WHO seropositivity criteria refers to neutralising antibody titer ≥1:80 with a positive result using a 2nd confirmatory assay [i.e., hemagglutination inhibition test (HAI) (HAI antibody titer ≥1:160), enzyme-linked immunosorbent assay, or western blot assay] [51]. Non-standard (NS) criteria refers to different antibody titer threshold defined by each study rather than a neutralising (NT) antibody titer ≥1:80 with a positive result confirmed by a 2nd assay (i.e. HAI antibody titer ≥1:40, ELISA or western blot assay) [51]

348 ───

Severity profile of Infections

We identified a systematic review by Lai et al. which reported the CFR of H5N1 from 1997 to 2015 [11]. The overall CFR was estimated to be 53.5% with clade-specific CFRs ranging from 33.3% to 100% [11] (**Figure 6)**. CFR also ranged between age groups among pediatric cases with 12-17 year olds recording the highest CFR (80.4%; 95% CI:68–90%) compared to 0-5 year olds (27.5%; 95% CI: 19–38%) [55]. One study, conducted by Li et al, estimated the IFR for H5N1, using surveillance and seroprevalence studies [56]. The authors analyse the 1997 Hong Kong and 2006 Turkey outbreaks and estimate IFRs of 14% (95% CI: 7—29) and 33% (95% CI: 14—61) (respectively) [56].

In comparison to H5N1 estimates from historical influenza pandemics were lower, with the 1918 pandemic being the most severe with a CFR of 2-3% [57] (**Figure 6**). 9 other studies reported 17 estimates for the CFR or IFR for seasonal and pandemic influenza [27,58–65]

(**Figure 6**). Estimates range depending on how the denominator was defined, with the

fatality rate increasing when only including laboratory-confirmed cases compared to

infections [59]. Fatality rates also varied depending on demographic with risk being affected

by severity of infection (type of hospitalisation and symptoms) and age [59–65].

As of 31st of March 2025 the CFR in the US is 1.43% (95% CI: 0.04–7.7%) with 1 death

being recorded out of 70 cases with recorded outcomes (**Supplementary table S3**). This

decreases to 1.25% (95% CI: 0.03–6.8%) when including cases identified via serological

studies (an additional 10 cases) [54,66]. CFR from cases where genotypes have been

reported is 8.33% (1/14; 95% CI: 0.21–38.5%) for D1.1, 0% (0/1; 95% CI: 0–97.5%) for D1.2

and 0% (0/19; 95% CI: 0–18%) for B3.13 (**Supplementary table S3**).

Figure 6. Severity profile of previous H5N1 outbreaks compared across influenza subtypes

Fatality risk estimates by outbreak. Case fatality risk (CFR) estimates are represented by circles and infection fatality risk (IFR) estimates by triangles. Uncertainty around these estimates are represented by 95% CI, IQR and range.

Comparisons to the 2003 Netherlands Avian Flu Epidemic 2003

A total of 89 human confirmed cases were reported for this outbreak [67]. We estimate R_0 to

be a median of 0.05 (95%CI: 0.02–0.12) when only including confirmed cases

(**Supplementary Figure S3a**). The posterior estimate for *k* was estimated as a median of

0.94 (95%: CrI: 0.78–1.1).

Regarding symptoms 78 cases (88%) presented with conjunctivitis only, 5 (5.6%) presented

with conjunctivitis and influenza-like illness and 2 cases (2.2%) presented with influenza-like

illness only and 4 (4.5%) did not fit the case definitions [67]. In comparison, among the 70

cases confirmed in the US (as of 30th March 2025) 61 cases (87%) have been recorded as

having conjunctivitis or ocular symptoms (**Supplementary table S3**).

In terms of severity the CFR for the outbreak in the Netherlands was 1.1% (95% CI: 0.03–

6.1%, with 1 death being recorded out of 89 cases [67]. When including infections identified

via serology, the IFR can be estimated as 0.1% (95% CI: 0–0.56%), with approximately 1000

people experiencing an infection with H7N7 [68]. The individual who died of H7N7,

experienced pneumonia followed by acute respiratory distress syndrome; the only individual

who developed severe respiratory symptoms during this outbreak [67] (CFR for severe

cases; 100%). Compared to the US, three cases have developed severe respiratory

symptoms (defined as symptoms that required hospitalisation), with 1 fatality (severe case

CFR; 33.3%)

A high percentage of positive H7 serology was also recorded in this outbreak, with a study of

56 household contacts of PCR-positive poultry workers reporting that 33 (58.9%) had

detectable antibodies against H7 [69]. In the Netherlands 86 (97%) cases had been exposed

via infected poultry and 3 (3%) had no known exposure to H7N7 infected poultry but were

household contacts of confirmed cases [69]. As of April 2025, 41 of (59%) US cases have

been exposed via dairy herds, 24 (34%) via poultry farms and culling, 2 (3%) via other other

animal exposures and 3 (4%) with unknown exposure source (**Supplementary table S3**).

Discussion

We reviewed the critical parameters for H5N1 and human influenza subtypes, and

additionally we estimated the R_0 for the current H5N1 outbreak in the US and the serial

interval for H5N1. To our knowledge this is the first estimate for the serial interval distribution

for H5N1 using H5N1 epidemiological data. We show that H5N1 has a different

epidemiological profile when compared to human influenza subtypes. Currently, H5N1 has a

much lower transmission potential than previous pandemic or seasonal human influenza

subtypes, with R < 0.2. H5N1 also appears to have a longer incubation period (~4 days vs

~2 days) and likely has a longer serial interval than is typical of human influenza (~6 days vs

~3 days). Data on latent and infectious periods are limited. Previous outbreaks of H5N1

have been typified by a high CFR and IFR, however the current IFR for the US outbreak is

currently low, and more similar to the CFR observed for the H7N7 outbreak in the

Netherlands in 2003, which was also characterised by high rates of conjunctivities among

cases. Serological studies may point to poultry workers and cullers being the groups at most

risk of infection. There may be weak evidence indicating that workers exposed to infection

with Clade 2.3.4.4b in similar settings might be more likely to be seropositive than workers

exposed to other strains of H5N1(**Figure 5**) [52–54].

Our results reaffirm the low transmissibility of H5N1 to or between humans. This could be

attributed to its replication preference for α2,3-linked sialic acid receptors predominantly

located in the lower respiratory tract (LRT) and the eye [70], specifically lung alveoli and

conjunctiva [71] in humans. In comparison human influenza viruses preferentially bind to

α2,6-linked sialic acid receptors (SAα2,6) [72], which are found in higher levels in the upper

respiratory tract (URT) [73].

We found the incubation period of H5N1 to be longer compared to human subtypes (**Figure**

3). This may be partially explained by H5N1's potential use of the eye as a portal of entry

before subsequent transmission to the respiratory tract [70]. Initial ocular infection may be

mild or subclinical, prolonging the incubation period before respiratory symptoms appear

[70]. Relevant for contact tracing purposes, we found limited evidence of a longer infectious

period of 5–13 days [23] though further studies are clearly required to confirm or refute these

data (**Supplementary Figure S6**). For the serial interval we estimated a median of ~6 days.

This is nearly double the length of typical human influenza subtypes [50] (**Figure 4**). There is

limited literature to provide a direct comparison for our estimate for H5N1 however avian

influenza A (H7N9) was estimated to have a median serial interval of 9 days in humans [74],

providing further evidence that avian influenza may exhibit longer serial intervals in humans.

The prolonged incubation period and serial interval of H5N1 may also be influenced by

reduced host susceptibility. It has been suggested that the expression of initial symptoms

after H5N1 infection stems from cellular damage compared to human subtypes where initial

symptoms may arise earlier due to the adaptive immune response triggered by previous

exposure [75].

A systematic review by Chen et al. indicated that individuals involved in poultry culling and

processing are at the highest risk of H5N1 infection [51]. Recent US studies additionally

highlight the risks of occupational exposure, suggesting that milking cows or cleaning milking

machinery pose significant risks [54]. The seroprevalence rates among both the dairy

workers in the US and the poultry workers in Egypt exceed those of poultry workers

reviewed by Chen et al. [51,53,54] (**Figure 5**), perhaps pointing to higher levels of

asymptomatic infection in workers exposed to Clade 2.3.4.4b viruses compared with other

H5N1 viruses. However, the seroprevalence estimates in the US and potentially the

Egyptian study did not follow WHO criteria for defining thresholds for seropositivity. It is also

not possible to standardise exposure across these studies and as such, it is not clear

whether there is higher seropositivity in workers exposed to clade 2.3.4.4b than other H5N1

clades. Nevertheless, this finding does suggest that the level of asymptomatic infection

needs to be monitored closely.

The reported CFR and IFR for H5N1, based on previous outbreaks, were much higher than

previous seasonal and pandemic influenza strains [57]. The historical severity of H5N1 is

well noted, however the current (as of April 2025) CFR stands at 1.43%, significantly lower

than what would be expected from historic experience. It has been argued that the high

mortality of H5N1 might be due to H5N1 virus pneumonia [71], indeed the CFR for US cases

which have reported severe respiratory symptoms (*N*=3) is 33%. This would be more in line

with what is seen in historical outbreaks. With the vast majority of cases reporting mild ocular

symptoms (**Supplementary table S3**), this indicates an unusual phenotype of H5N1

infection, which was previously infrequently associated with ocular disease [76].

This phenotype has however been documented in H7 subtype viruses, with 80% of

documented human infections being associated with ocular complications and an influenza-

virus positive eye swab [77]. This was seen during the 2003 H7N7 outbreak in the

Netherlands with 97% of cases presenting with conjunctivitis. The CFR during this outbreak

was 1.1% which is similar to what is currently recorded in the US, with the one fatality being

recorded in a patient who developed pneumonia [67]. We estimated the R_0 for this outbreak

(among confirmed cases) as 0.05, which is similar to what we estimated for the US outbreak

(if cases with unknown exposure have indeed been infected from another undetected case).

The R_0 for this outbreak may be higher given the high percentage of household contacts to

infected poultry workers having detectable antibodies against H7 [69]. It is plausible that

patients with conjunctivitis may have the potential to expose household members when

sharing items such as towels and washcloths [69]. This would suggest there is a risk of this

type of transmission in the US.

Despite the severity of this outbreak currently being low, given the high prevalence of H5N1

among cattle [78] and the level of human exposure, there is a risk of reassortment leading to

a subtype capable of sustaining human to human transmission. Therefore, proactive

containment measures are crucial to mitigate the potential for a global health crisis.

Our study is subject to several limitations. The interpretation of transmissibility in the US

based case exposure source is uncertain. These cases may have been exposed via

zoonotic sources or alternatively may have been at the end of longer undetected

transmission chains.; leading to caution when interpreting our estimates. Additionally there is

substantial heterogeneity regarding some parameters, which may result from differences in

study designs, populations, and methodologies as well as differences in surveillance and

reporting practices.

Conclusion

We have assessed and estimated critical epidemiological parameters for human H5N1 based on past and current outbreaks. The outbreak in the US appears to have lower pathogenicity than observed in previous H5N1 outbreaks, and its clinical and epidemiological picture appears more similar to the H7N7 outbreak in the Netherlands in 2003. H5N1 may have a longer incubation period, and serial interval compared to human influenza subtypes. These characteristics may allow for more effective contact tracing more than is typically the case for influenza (depending on the degree of presymptomatic transmission). Despite these insights, data on H5N1 infections remain sparse and critical gaps remain in our understanding. Addressing these gaps and continually monitoring the epidemiology is imperative to enhance our preparedness and assess whether the risk from these viruses is potentially escalating.

References

[1] Claas EC, Osterhaus AD, van Beek R, De Jong JC, Rimmelzwaan GF, Senne DA, et al. Human influenza A H5N1 virus related to a highly pathogenic avian influenza virus. Lancet 1998;351:472–7.

[2] Charostad J, Rezaei Zadeh Rukerd M, Mahmoudvand S, Bashash D, Hashemi SMA, Nakhaie M, et al. A comprehensive review of highly pathogenic avian influenza (HPAI) H5N1: An imminent threat at doorstep. Travel Med Infect Dis 2023;55:102638.

[3] CDC. Current Bird Flu Situation in Wild Birds. Avian Influenza (Bird Flu) 2024. https://www.cdc.gov/bird-flu/situation-summary/wildbirds.html (accessed September 26, 2024).

[4] Peacock T, Moncla L, Dudas G, VanInsberghe D, Sukhova K, Lloyd-Smith JO, et al.

The global H5N1 influenza panzootic in mammals. Nature 2024:1–2.

[5] Pekar JE, Peacock TP, Moncla L, Gangavarapu P, Venkatesh D, Goldhill DH, et al.

Timing and molecular characterisation of the transmission to cattle of H5N1 influenza A

virus genotype D1.1, clade 2.3.4.4b 2025.

[6] Burrough ER, Magstadt DR, Petersen B, Timmermans SJ, Gauger PC, Zhang J, et al.

Highly pathogenic avian influenza A(H5N1) clade 2.3.4.4b virus infection in domestic

dairy cattle and cats, United States, 2024. Emerg Infect Dis 2024;30:1335–43.

[7] Sah R, Srivastava S, Kumar S, Mehta R, Donovan S, Sierra-Carrero L, et al. Concerns

on H5N1 avian influenza given the outbreak in U.S. dairy cattle. Lancet Reg Health Am

2024;35:100785.

[8] CDC. H5 Bird Flu: Current Situation. Avian Influenza (Bird Flu) 2024.

https://www.cdc.gov/bird-flu/situation-summary/index.html (accessed December 6,

2024).

[9] Longini IM Jr, Nizam A, Xu S, Ungchusak K, Hanshaoworakul W, Cummings DAT, et al.

Containing pandemic influenza at the source. Science 2005;309:1083–7.

[10] Biggerstaff M, Cauchemez S, Reed C, Gambhir M, Finelli L. Estimates of the

reproduction number for seasonal, pandemic, and zoonotic influenza: a systematic

review of the literature. BMC Infect Dis 2014;14:480.

[11] Lai S, Qin Y, Cowling BJ, Ren X, Wardrop NA, Gilbert M, et al. Global epidemiology of

avian influenza A H5N1 virus infection in humans, 1997-2015: a systematic review of

individual case data. Lancet Infect Dis 2016;16:e108–18.

[12] Tricco AC, Langlois EV, Straus SE. Rapid reviews to strengthen health policy and

systems: A Practical Guide. Geneva: World Health Organization; 2017.

[13] King VJ, Stevens A, Nussbaumer-Streit B, Kamel C, Garritty C. Paper 2: Performing

rapid reviews. Syst Rev 2022;11:151.

[14] Lessler J, Salje H, Van Kerkhove MD, Ferguson NM, Cauchemez S, Rodriquez-

Barraquer I, et al. Estimating the severity and subclinical burden of Middle East

respiratory syndrome Coronavirus infection in the Kingdom of Saudi Arabia. Am J

556 Epidemiol 2016;183:657–63.

557 [15] Garg S, Reinhart K, Couture A, Kniss K, Davis CT, Kirby MK, et al. Highly pathogenic

558 avian influenza A(H5N1) virus infections in humans. N Engl J Med 2025;392:843–54.

559 [16] Meijer A, Wilbrink B, du Ry van Beest Holle M, Fouchier RAM, Natrop G, Bosman A, et

560 al. Highly pathogenic avian influenza virus A(H7N7) infection of humans and human-to-

561 human transmission during avian influenza outbreak in the Netherlands. Int Congr Ser

562 2004;1263:65–8.

563 [17] Azam JM, Funk S, Finger F. epichains: Simulating and Analysing Transmission Chain

564 Statistics Using #> Branching Process Models 2024.

565 [18] CDC. Past Examples of Probable Limited, Non-Sustained, Person-to-Person Spread of

566 Avian Influenza A Viruses. Avian Influenza (Bird Flu) 2024. https://www.cdc.gov/bird-

567 flu/php/avian-flu-summary/h5n1-human-infections.html (accessed March 31, 2025).

568 [19] Aditama TY, Samaan G, Kusriastuti R, Sampurno OD, Purba W, Misriyah, et al. Avian

569 influenza H5N1 transmission in households, Indonesia. PLoS One 2012;7:e29971.

570 [20] Wickham H. ggplot2: Elegant Graphics for Data Analysis 2016.

571 [21] R Core Team. R: A Language and Environment for Statistical Computing 2021.

572 [22] Bettencourt LMA, Ribeiro RM. Real time bayesian estimation of the epidemic potential

573 of emerging infectious diseases. PLoS One 2008;3:e2185.

574 [23] Yang Y, Halloran ME, Sugimoto JD, Longini IM Jr. Detecting human-to-human

575 transmission of avian influenza A (H5N1). Emerg Infect Dis 2007;13:1348–53.

576 [24] Ferguson NM, Fraser C, Donnelly CA, Ghani AC, Anderson RM. Public health. Public

577 health risk from the avian H5N1 influenza epidemic. Science 2004;304:968–9.

578 [25] Saucedo O, Martcheva M, Annor A. Computing human to human Avian influenza □□

579 via transmission chains and parameter estimation. Math Biosci Eng 2019;16:3465–87.

580 [26] Lee Y, Lee DH, Kwon H-D, Kim C, Lee J. Estimation of the reproduction number of

581 influenza A(H1N1)pdm09 in South Korea using heterogeneous models. BMC Infect Dis

582 2021;21:658.

583 [27] Larrauri Cámara A, Jiménez-Jorge S, Mateo Ontañón S de, Pozo Sánchez F, Ledesma

Moreno J, Casas Flecha I, et al. Epidemiology of the 2009 influenza pandemic in Spain.

The Spanish Influenza Surveillance System. Enferm Infecc Microbiol Clin 2012;30 Suppl

4:2–9.

[28] Fraser C, Cummings DAT, Klinkenberg D, Burke DS, Ferguson NM. Influenza

transmission in households during the 1918 pandemic. Am J Epidemiol 2011;174:505–

14.

[29] De Serres G, Gay NJ, Farrington CP. Epidemiology of transmissible diseases after

elimination. Am J Epidemiol 2000;151:1039–48; discussion 1049–52.

[30] Bui C, Bethmont A, Chughtai AA, Gardner L, Sarkar S, Hassan S, et al. A Systematic

Review of the Comparative Epidemiology of Avian and Human Influenza A H5N1 and

H7N9 - Lessons and Unanswered Questions. Transbound Emerg Dis 2016;63:602–20.

[31] Beigel JH, Farrar J, Han AM, Hayden FG, Hyer R, de Jong MD, et al. Avian Influenza A

(H5N1) Infection in Humans. N Engl J Med 2005;353:1374–85.

[32] Cowling BJ, Jin L, Lau EHY, Liao Q, Wu P, Jiang H, et al. Comparative epidemiology of

human infections with avian influenza A H7N9 and H5N1 viruses in China: a population-

based study of laboratory-confirmed cases. Lancet 2013;382:129–37.

[33] Oner AF, Bay A, Arslan S, Akdeniz H, Sahin HA, Cesur Y, et al. Avian influenza A

(H5N1) infection in eastern Turkey in 2006. N Engl J Med 2006;355:2179–85.

[34] Huai Y, Xiang N, Zhou L, Feng L, Peng Z, Chapman RS, et al. Incubation period for

human cases of avian influenza A (H5N1) infection, China. Emerg Infect Dis

2008;14:1819–21.

[35] Tom BDM, Van Hoek AJ, Pebody R, McMenamin J, Robertson C, Catchpole M, et al.

Estimating time to onset of swine influenza symptoms after initial novel A(H1N1v) viral

infection. Epidemiol Infect 2011;139:1418–24.

[36] Carrat F, Vergu E, Ferguson NM, Lemaitre M, Cauchemez S, Leach S, et al. Time lines

of infection and disease in human influenza: a review of volunteer challenge studies. Am

J Epidemiol 2008;167:775–85.

[37] Ferguson NM, Cummings DAT, Cauchemez S, Fraser C, Riley S, Meeyai A, et al.

Strategies for containing an emerging influenza pandemic in Southeast Asia. Nature

2005;437:209–14.

[38] Nishiura H. Early efforts in modeling the incubation period of infectious diseases with an

acute course of illness. Emerg Themes Epidemiol 2007;4:2.

[39] Nishiura H, Inaba H. Estimation of the incubation period of influenza A (H1N1-2009)

among imported cases: addressing censoring using outbreak data at the origin of

importation. J Theor Biol 2011;272:123–30.

[40] Lessler J, Reich NG, Brookmeyer R, Perl TM, Nelson KE, Cummings DAT. Incubation

periods of acute respiratory viral infections: a systematic review. Lancet Infect Dis

2009;9:291–300.

[41] Canini L, Carrat F. Population modeling of influenza A/H1N1 virus kinetics and symptom

dynamics. J Virol 2011;85:2764–70.

[42] Lessler J, Reich NG, Cummings DAT, New York City Department of Health and Mental

Hygiene Swine Influenza Investigation Team, Nair HP, Jordan HT, et al. Outbreak of

2009 pandemic influenza A (H1N1) at a New York City school. N Engl J Med

2009;361:2628–36.

[43] Wang C, Yu E, Xu B, Wang W, Li L, Zhang W, et al. Epidemiological and clinical

characteristics of the outbreak of 2009 pandemic influenza A (H1N1) at a middle school

in Luoyang, China. Public Health 2012;126:289–94.

[44] Shen J, Niu J. Epidemiologic parameters and evaluation of control measure for 2009

novel influenza a (H1N1) in Xiamen, Fujian Province, China. Virol J 2012;9.

https://doi.org/10.1186/1743-422x-9-20.

[45] Cao B, Li X-W, Mao Y, Wang J, Lu H-Z, Chen Y-S, et al. Clinical features of the initial

cases of 2009 pandemic influenza A (H1N1) virus infection in China. N Engl J Med

2009;361:2507–17.

[46] Tuite AR, Greer AL, Whelan M, Winter A-L, Lee B, Yan P, et al. Estimated

epidemiologic parameters and morbidity associated with pandemic H1N1 influenza.

CMAJ 2010;182:131–6.

[47] Cori A, Valleron AJ, Carrat F, Scalia Tomba G, Thomas G, Boëlle PY. Estimating

influenza latency and infectious period durations using viral excretion data. Epidemics

2012;4:132–8.

[48] Chan LYH, Morris SE, Stockwell MS, Bowman NM, Asturias E, Rao S, et al. Estimating

the generation time for influenza transmission using household data in the United

States. medRxiv 2024:2024.08.17.24312064.

[49] Cauchemez S, Carrat F, Viboud C, Valleron AJ, Boëlle PY. A Bayesian MCMC

approach to study transmission of influenza: application to household longitudinal data.

Stat Med 2004;23:3469–87.

[50] Vink MA, Bootsma MCJ, Wallinga J. Serial intervals of respiratory infectious diseases: a

systematic review and analysis. Am J Epidemiol 2014;180:865–75.

[51] Chen X, Wang W, Wang Y, Lai S, Yang J, Cowling BJ, et al. Serological evidence of

human infections with highly pathogenic avian influenza A(H5N1) virus: a systematic

review and meta-analysis. BMC Med 2020;18:377.

[52] Gomaa M, Moatasim Y, El Taweel A, Mahmoud SH, El Rifay AS, Kandeil A, et al. We

are underestimating, again, the true burden of H5N1 in humans. BMJ Glob Health

2023;8:e013146.

[53] Shittu I, Silva D, Oguzie JU, Marushchak LV, Olinger GG, Lednicky JA, et al. A one

health investigation into H5N1 avian influenza virus epizootics on two dairy farms.

medRxiv 2024:2024.07.27.24310982. https://doi.org/10.1101/2024.07.27.24310982.

[54] Mellis AM, Coyle J, Marshall KE, Frutos AM, Singleton J, Drehoff C, et al. Serologic

evidence of recent infection with highly pathogenic avian influenza A(H5) virus among

dairy workers — Michigan and Colorado, June–august 2024. MMWR Morb Mortal Wkly

Rep 2024;73:1004–9.

[55] Oner AF, Dogan N, Gasimov V, Adisasmito W, Coker R, Chan PKS, et al. H5N1 avian

influenza in children. Clin Infect Dis 2012;55:26–32.

[56] Li FCK, Choi BCK, Sly T, Pak AWP. Evidence-based public health policy and practice:

Finding the real case-fatality rate of H5N1 avian influenza. Journal of Epidemiology and

Community Health (1979-) 2008;62:555–9.

[57] WHO. Pandemic Influenza Risk Management. 2017.

[58] McDonald SA, Teirlinck AC, Hooiveld M, van Asten L, Meijer A, de Lange M, et al.

Inference of age-dependent case-fatality ratios for seasonal influenza virus subtypes

A(H3N2) and A(H1N1)pdm09 and B lineages using data from the Netherlands.

Influenza Other Respi Viruses 2023;17:e13146.

[59] Wong JY, Kelly H, Ip DKM, Wu JT, Leung GM, Cowling BJ. Case fatality risk of

influenza A (H1N1pdm09): a systematic review: A systematic review. Epidemiology

2013;24:830–41.

[60] Abdalla O, Mohammed M, Hakawi AM, Aljifri A, Abdalla M, Eltigani S, et al. Hospital-

based surveillance of influenza A(H1N1)pdm09 virus in Saudi Arabia, 2010-2016. Ann

Saudi Med 2020;40:1–6.

[61] Fujikura Y, Kawano S, Kouzaki Y, Shinoda M, Hara Y, Shinkai M, et al. The (H1N1)

2009 pandemic influenza pneumonia among adult patients in Japan. Jpn J Infect Dis

2014;67:100–4.

[62] Petrović V, Seguljev Z, Cosić G, Ristić M, Nedeljković J, Dragnić N, et al. Overview of

the winter wave of 2009 pandemic influenza A(H1N1)v in Vojvodina, Serbia. Croat Med

J 2011;52:141–50.

[63] Kim P, Coleman B, Kwong JC, Plevneshi A, Hassan K, Green K, et al. Burden of severe

illness associated with laboratory-confirmed influenza in adults aged 50-64 years, 2010-

2011 to 2016-2017. Open Forum Infect Dis 2023;10:ofac664.

[64] Presanis AM, De Angelis D, New York City Swine Flu Investigation Team, Hagy A,

Reed C, Riley S, et al. The severity of pandemic H1N1 influenza in the United States,

from April to July 2009: a Bayesian analysis. PLoS Med 2009;6:e1000207.

[65] Altmann M, Fiebig L, Soyka J, von Kries R, Dehnert M, Haas W. Severe cases of

pandemic (H1N1) 2009 in children, Germany. Emerg Infect Dis 2011;17:186–92.

[66] Shittu I, Silva D, Oguzie JU, Marushchak LV, Olinger GG, Lednicky JA, et al. A one

health investigation into H5N1 avian influenza virus epizootics on two dairy farms. Clin

Infect Dis 2025;80:331–8.

[67] Fouchier RAM, Schneeberger PM, Rozendaal FW, Broekman JM, Kemink SAG,

Munster V, et al. Avian influenza A virus (H7N7) associated with human conjunctivitis

and a fatal case of acute respiratory distress syndrome. Proc Natl Acad Sci U S A

2004;101:1356–61.

[68] Milieu RRVV, National Institute of Public Health and the Environment. Avian Flu

Epidemic 2003: Public health consequences. Executive summary 630940004 2010.

https://www.rivm.nl/bibliotheek/rapporten/630940004.html (accessed March 31, 2025).

[69] Du Ry van Beest Holle M, Meijer A, Koopmans M, de Jager CM. Human-to-human

transmission of avian influenza A/H7N7, The Netherlands, 2003. Euro Surveill

2005;10:264–8.

[70] Kumlin U, Olofsson S, Dimock K, Arnberg N. Sialic acid tissue distribution and influenza

virus tropism. Influenza Other Respi Viruses 2008;2:147–54.

[71] Granata G, Simonsen L, Petrosillo N, Petersen E. Mortality of H5N1 human infections

might be due to H5N1 virus pneumonia and could decrease by switching receptor.

Lancet Infect Dis 2024;24:e544–5.

[72] Zeng H, Goldsmith CS, Maines TR, Belser JA, Gustin KM, Pekosz A, et al. Tropism and

infectivity of influenza virus, including highly pathogenic avian H5N1 virus, in ferret

tracheal differentiated primary epithelial cell cultures. J Virol 2013;87:2597–607.

[73] Couceiro JNSS, Paulson JC, Baum LG. Influenza virus strains selectively recognize

sialyloligosaccharides on human respiratory epithelium; the role of the host cell in

selection of hemagglutinin receptor specificity. Virus Res 1993;29:155–65.

[74] Zhou L, Li Q, Uyeki TM. Estimated Incubation Period and Serial Interval for Human-to-

Human Influenza A(H7N9) Virus Transmission. Emerg Infect Dis 2019;25:1982–3.

[75] Hermesh T, Moltedo B, López CB, Moran TM. Buying time-the immune system

determinants of the incubation period to respiratory viruses. Viruses 2010;2:2541–58.

[76] Belser JA, Zeng H, Katz JM, Tumpey TM. Ocular tropism of influenza A viruses:

identification of H7 subtype-specific host responses in human respiratory and ocular

cells. J Virol 2011;85:10117–25.

[77] Belser JA, Lash RR, Garg S, Tumpey TM, Maines TR. The eyes have it: influenza virus

infection beyond the respiratory tract. Lancet Infect Dis 2018;18:e220–7.

[78] CDC. Current H5N1 Bird Flu Situation in Dairy Cows. Avian Influenza (Bird Flu) 2024.

https://www.cdc.gov/bird-flu/situation-summary/mammals.html (accessed November 26,

2024).