CSF pharmacokinetics-pharmacodynamics of linezolid in critically brain injured patients, with or without central nervous system healthcare-associated infection. The PK-Pop-LCR Study: A Multicenter Pharmacokinetics and Pharmacodynamics Population Study.

Claire DAHYOT-FIZELIER[1,2], Alexia CHAUZY[1], Kévin CHALARD[3], Fanny BERNARD[1,2], Hugues de COURSON[4], Pierre-Etienne LEBLANC[5], Gilles FRANCONY[6], Russel CHABANNE[7], Karim LAKHAL[8], Raphaël CINOTTI[8], Charles GREGOIRE[9], Julien POTTECHER[10], Belaid BOUHEMAD[11], Assil MERLAUD[1], Christophe ADIER[1], Jean-Claude LECRON[13] Ombeline REMY[2], William COUET[1,14], Nicolas GREGOIRE[1,14], Sandrine MARCHAND[1,14], For the PK-Pop-LCR Study Group (Members and affiliations listed in Acknowledgements, linked to pubmed record).

1. Université de Poitiers, INSERM U1070, PHAR2, Poitiers, France
2. Service d'anesthésie-réanimation médecine périopératoire, CHU de Poitiers, Poitiers, France
3. Service d'anesthésie réanimation médecine périopératoire, Chu Montpellier – Montpellier, France
4.   Service d'anesthésie réanimation médecine périopératoire, Chu Bordeaux- Chu Bordeaux – Bordeaux, France
5. Service d'anesthésie réanimation médecine périopératoire, APHP Kremlin Bicêtre – Paris, France
6. Service d'anesthésie réanimation médecine périopératoire, Chu Grenoble – Grenoble, France
7. Service d'anesthésie réanimation médecine périopératoire, Chu Clermont Ferrand – Clermont Ferrand, France
8. Service d'anesthésie réanimation médecine périopératoire, Chu Nantes – Nantes, France
9. Service d'anesthésie réanimation médecine périopératoire, Hôpital Fondation Adolphe de Rothschild, APHP – Paris, France
10. Service d'anesthésie réanimation médecine périopératoire, Chu Strasbourg – Strasbourg, France
11. Service d'anesthésie réanimation médecine périopératoire, Chu Dijon – Dijon, France
12. Centre Hospitalier Universitaire de Poitiers, Direction de la recherche clinique et innovation, Poitiers, France
13. Université de Poitiers, LITEC, Poitiers, France
14. Service de Toxicologie-Pharmacocinétique, CHU de Poitiers, Poitiers, France

CORRESPONDING AUTHOR:

Claire Dahyot-Fizelier, MD, PhD

Université of Poitiers College of Medicine and Pharmacy, INSERM U1070, 6 rue de la milétrie TSA 51115, 86073 POITIERS Cedex 9, France

claire.dahyot@gmail.com

+33 612 91 25 31/ +33 549 44 48 40

Abstract

Background

Linezolid is an alternative to vancomycin for treating Gram-positive central nervous system (CNS) healthcare-

associated infections. The recommended dosing regimen remains debated.

Methods

PK-Pop-LCR, a prospective population pharmacokinetic-pharmacodynamic multicenter study, included brain

injured patients with an external ventricular drainage receiving linezolid at different dosing regimens. The

cerebrospinal fluid (CSF) penetration of linezolid was investigated and a population pharmacokinetic model

developed using plasma and CSF data. Monte Carlo simulations were conducted to calculate probability of

target attainment (PTA) and cumulative fraction of response (CFR) in CSF against methicillin-resistant *Staph-*

ylococcus aureus (MRSA) and methicillin-resistant *Staphylococcus epidermidis* (MRSE), for different dosing

regimens. The plasma pharmacodynamic target, AUC/MIC > 100, was used in CSF.

Findings

Over 25 patients included, blind adjudication confirmed 14 cases of CNS infections. Mean $AUC_{CSF}/fAUC_{plasma}$

ratio was close to 80% and not different between patients with and without CNS infection, despite higher CSF

cytokines levels in CNS-infected patients. The recommended dose of 1200 mg/24h allowed to reach PTAs $\geq$

90% for MICs $\leq$ 0.5 mg/L, and CFR of 3.2% and 40% for MRSA and MRSE, respectively. 2 700 mg/24h

would allow to achieve PTA > 90% for MIC up to 1 mg/L and CFR of 90% for MRSE. None of dosing regi-

mens tested was appropriate for MRSA infections.

Interpretation

We confirmed the extensive CSF distribution of linezolid. Higher doses than those recommended should be

considered to treat CNS infection in critically brain injured patients. However, pharmacodynamic target for

CNS infections should be further investigated to confirm these findings.

Fundings: French Ministry of Health**.**

Keywords: linezolid, external ventricular drainage, healthcare associated meningitis, critical care, *Staphylo-*

coccus

INTRODUCTION

Central nervous system (CNS) healthcare-associated infections (CNS-HAI) are associated with high morbi-mortality. Recently, a mortality rate of 9% was reported, with an unfavorable prognosis in 50% of surviving patients. One year mortality rises to almost 40% in ventriculitis[1,2]. In literature, incidences varies according to the clinical situation, ranging from less than 1% in post-operative cases of scheduled surgery to 10% in average in external ventricular drain (EVD) associated infections in ICU[3].

In two-thirds of cases, causative organisms are gram-positive bacteria, with a majority of *Staphylococcus aureus* and coagulase-negative bacteria[4]. To date, high-dose of vancomycin is still recommended as a first-line for CNS-HAI, despite its limited CNS distribution and narrow therapeutic index with its renal toxicity[5]. Indeed, blood-brain barrier (BBB) and blood-cerebrospinal fluid barrier (BCSFB) limit penetration of hydrophilic antibiotics such as vancomycin, exposing patients to the risk of therapeutic failure. Recently, vancomycin diffusion of 9 to 13% in patients with CNS infections was reported, with median CSF concentrations of 2.9 +/- 2.2 mg/L[6], failing to reach the EUCAST minimum inhibitory concentration (MIC) threshold for resistant *Staphylococcus*[7].

Linezolid is indicated against multidrug resistant Gram-positive bacteria for the treatment of community- and hospital-acquired pneumonia and skin structure infections[8]. It is also increasingly prescribed for CNS infections in ICU patients as an alternative when methicillin-resistant *S. aureus* (MRSA) strains have a vancomycin minimum inhibitory concentration (MIC) higher than 1 mg/L, as linezolid characteristics enable it crossing barriers more easily [5]. Recently, linezolid was proposed in tuberculous meningitis, especially for drug-resistant ones [9–11]. Indeed, high linezolid CSF penetration from 60 to 90% was reported, but mainly in non-infected situations or extra-cerebral infections[12]. However, those few studies have limitations: few patients, no proven CNS infection for most of them, comparison of CSF linezolid concentrations to total plasma concentrations, no inflammatory biomarkers and EVD flow rates reported, and only one administration scheme studied[9,13–18]. Despite high CNS concentrations of linezolid, several studies reported that pharmacodynamics (PD) targets were not achieved, suggesting the need to increase the studied recommended doses of 600mg q12h[13,14].

The aim of the present study was to explore the linezolid CSF pharmacokinetics (PK) and efficacy in ICU patients with an EVD, at different dosing regimens, with or without infection.

METHODS

Study design and participants

PK-Pop-LCR study, a national prospective population PK/PD multicenter study (*CPP17-016a/2017-002993-37*), involving 14 French ICU, was approved by the Committee for the Protection of Persons of Sud-Ouest and Outre-Mer IV (EudraCT: 2017-002993-37) and the "Agence Nationale de Sécurité du Médicament et des Produits de Santé". It was also registered at the Clinical Trials web site: NCT03481569. Linezolid was one of the 9 broad-spectrum antibiotics studied. Patients, after obtaining informed consent from legal surrogate, were included in 11 of participating centers from January 2019 to February 2020.

Brain injured patients (≥18 years of age), requiring intensive care management with EVD, treated for a CNS or extra-CNS infection by linezolid were eligible. Declared cases of CNS infection were confirmed by a masked adjudication committee according to IDSA definition[5]. Exclusion criteria were acute renal failure (creatinine clearance (Clcr) < 50 mL/min and/or continuous hemodialysis), pregnancy, contraindication to linezolid, patient or family refusal to be involved in the study, patient with reinforced protection or deprived of freedom subsequent to a legal or administrative decision. All clinical and biological data, recorded on a data collection template, are detailed in the supplemental materials.

Intervention and sample collection

All patients received linezolid (infusion solution at 2mg/mL) by intravenous (IV) infusion over 30 minutes, 600 mg twice a day (b.i.d., n=17 patients) or 600 mg thrice a day (t.i.d., n=8). The PK study was performed within a dosing interval at steady state.

Blood samples were collected at the start of IV injection, at the end of infusion, 2 hours and 6 hours after the administration, and just before the next administration, i.e. 8 or 12 hours after administration (t.i.d. or b.i.d. respectively). CSF samples were collected over one–hour intervals *via* the EVD between two infusions, as follows: H-1/H0, H0/H1, H1/H2, H2/H3, H3/H4, H5/H6, with the last sample depending on the regimen, i.e. H7/H8 or H11/H12 (t.i.d. or b.i.d. respectively). One additional blood and CSF samples were collected for albumin, Interleukin (IL) 6, IL8, IL10, IL17 and TNF-alpha measurements.

Blood samples were centrifuged at 3000rpm (4°C; 10min). Plasma and CSF samples were stored at -80°C until analysis. Samples' exact time of collection was recorded and dead volume of EVD tubing considered for PK analysis.

Linezolid, cytokine and albumin measurements

Total plasma and unbound CSF concentrations of linezolid were measured using an appropriate validated LC–MS/MS method (limit of quantification of 0.01 mg/L for both matrices). Concentrations of albumin, IL6, IL8, IL10, IL17A and TNF-alpha were quantified in plasma and CSF. Linezolid, cytokines and albumin assays are detailed in Supplemental Materials. CSF-to-plasma albumin concentrations ratios (Q_{alb}) were calculated to assess the BCSFB function.

PK-PD analysis

Population PK analysis

Linezolid total plasma and unbound CSF concentrations were analyzed simultaneously using the non-linear

mixed-effect modelling approach in NONMEM 7.4 (ICON Development Solutions, Ellicott City, MD, USA).

Description of the development, evaluation and covariates selection are detailed in Supplemental materials.

The final structural model is shown on Figure S1. Free maximal concentrations (C_{max}), free areas under the

concentration-time curves over 24h ($fAUC_{0-24}$) in plasma and CSF were estimated at steady-state from 5 000

concentration-time profiles simulated with the final model.

Probability of Target Attainment (PTA) and Cumulative Fraction of Response (CFR)

The PTA of linezolid in plasma and CSF ($fAUC_{0-24}$/MIC > 100)[20] was estimated from Monte-Carlo

simulations for 1 200, 1 800 and 2 700 mg daily doses, delivered in 2 or 3 injections (over 30 or 180 min) or

by continuous infusion. To evaluate linezolid dosing regimens efficacy against MRSA and methicillin resistant

S. epidermidis (MRSE), PTAs were compared with linezolid MIC frequency distributions from EUCAST[21].

CFRs were calculated for each dosing regimen[22].

Statistical analysis

All statistical analysis was performed using R statistical software (v4.2.3; R Core Team 2023). Descriptive

statistics were presented as median and interquartiles. Cytokines concentrations between plasma and CSF, Q_{alb}

and cytokine concentrations in CSF between CNS and extra-CNS infection groups were compared with a

nonparametric Mann Whitney U test (p < 0.05). Cytokines concentrations below the detection limit were

considered as half of this threshold.

RESULTS

Patients and data

Among 25 patients included, 16 CNS and 9 extra-CNS infections were declared. Among the 16 CNS

infections, 14 were confirmed after adjudication. Patient demographics, clinical, biological characteristics and

pathogens identified are summarized in Table S1. The underlying diseases were mostly hemorrhagic stroke,

with subarachnoid or intracranial hemorrhage. A total of 87 plasma and 159 CSF samples were considered for

PK analysis.

Inflammatory markers

In the "CNS Infection" group, median concentrations of all cytokines except TNF-alpha were significantly

higher in CSF than in plasma (Figure 1). In CSF of patients with CNS infections, cytokines' median

concentrations were 4-fold higher for IL6 (726 pg/mL [522-4823] vs 161 pg/mL [59-1195]), 6-fold higher for

IL8 (1016 pg/mL [441-1553] vs 172 pg/mL [10-266]) and significantly higher for IL10 and IL17A (Figure 2).

The median Q_{alb} was not significantly different between "CNS Infection" (7.3×10^{-3} [5.5×10^{-3} - 18.4×10^{-3}])

and "Extra-CNS Infection" (8.9×10^{-3} [6.3×10^{-3} - 14.2×10^{-3}]) groups, with a larger variability in the "CNS

Infection" group (Figure S2).

Plasma and CSF concentrations and population PK model parameters

The final PK model described observed data well in plasma and CSF, as indicated by the visual predictive

check, individual predictions vs. observed data and goodness of fit plots (Figures S3, S4, and S5). The final

model parameters were all estimated with good precision (Table S2).

Predicted plasma and CSF concentrations, and secondary PK parameters for the two dosing regimens are

shown on Figure 3 and Table 1.

Over all relevant covariates, only the effect of body weight on the central plasma compartment volume of

distribution was found to significantly improve the model. Neither CNS infection nor inflammation

biomarkers and Q_{alb} had a significant impact on linezolid CSF distribution and were therefore not included in

the final model. However, linezolid elimination rate from CSF via the EVD (CL_{EVD}) was not negligible (Table

S1), corresponding to 27.8% of $Q_{CSF,out}$ on average, which tended to decrease the mean CSF linezolid

exposure, resulting in a lower CSF-to-plasma fAUC$_{0-24}$ ratio as CL_{EVD} increased (Figure 4).

PTA and CFR

The PTA and CFRs for various dosing regimens showed that recommended daily dose (1 200 mg/24h) would

allow reaching PTA $\geq$ 90% only for MIC $\leq$ 0.5 mg/L, and CFR of 3.2% and 40% for MRSA and MRSE,

respectively. Only a daily dose of 2 700 mg would allow to achieve PTA > 90% against pathogens with MIC

up to 1 mg/L and thus cover the MICs frequency distribution for MRSE with CFR = 90% (Figure 5 and Table

2). None of the dosing regimens tested appeared to be appropriate to treat MRSA infections. None of the

tested modalities of administration of linezolid had any impact on PTA and CFR.

DISCUSSION

To our knowledge, this is the most comprehensive population PK study of linezolid in plasma and CSF in a large number of critically ill patients with proven CNS infection, which explored several linezolid dosing regimens.

Few studies conducted a robust PK analysis to describe linezolid disposition in CNS, with heterogeneous methodological approaches and small sample sizes, preventing accurate characterization of the wide interindividual variability in the ICU population[23]. Traditional compartmental modelling was only occasionally conducted[13]. In the present study, estimated mean steady-state PK parameters were similar to those reported in the previous studies, with a total Vd of 36.4 L and total CL of linezolid of 8.4.L/h[14–18]. Only one study reported higher findings without any clear explanation (Vd=101.3 L; CL=16.6 L/h)[13].

Considering linezolid CSF distribution, flat concentrations profiles were observed with around 3 hours delay in peak concentrations compared to plasma, in accordance with the amphiphilic properties of linezolid and with previous reported observations in ICU patients[13–17]. Our results confirmed a good central penetration of linezolid around 80%, despite significant inter-studies variability with CSF/plasma AUC ratio ranging from 41.3% to 77% in patients without CNS infection[13,15–18], and from 80% to 98% in patients with CNS infections[14,24]. However, all these previous studies compared AUCs in CSF with total plasma AUCs, without taking into account the protein binding of linezolid, probably leading to a slight underestimation of ratios.

In our study, we observed significantly higher CSF than plasma concentrations of all cytokines in patients with CNS infection, except for TNF-alpha, which suggest a local neuroinflammation of BCSFB in these patients. This difference of response in serum and CSF between ICU patients with or without CNS infection was not previously reported. In bacterial meningitis, an increase of all cytokines in CNS was already described, but only transiently for TNF-alpha and IL-1[25]. Despite these significant differences, cytokine levels were not covariates influencing linezolid diffusion, unlike meropenem for which high CSF IL-6 levels were associated with increased CSF penetration in ventriculitis patients[26]. Besides, the brain barriers integrity marker, CSF/plasma albumin ratio (Q_{alb}), did not influence linezolid diffusion either, contrasting with the impact described for meropenem and vancomycin, known for their limited CNS diffusion[27,28]. In our population, median Q_{alb} below 10 x 10^{-3} in both groups suggested only a mild BCSFB dysfunction[19], which could explain the lack of influence of this covariate on linezolid diffusion, whose penetration is particularly good. Nevertheless, caution must be taken when interpreting this parameter and its influence in CNS infections studies as it may be influenced by several factors, including the rostrocaudal gradient within the CSF, with Q_{alb} twice as low in cerebral ventricles samples as in lumbar ones[29,30]. Interestingly, the EVD rate influenced the CSF exposure of linezolid, as also recently described for vancomycin and meropenem in literature[31,32]. The higher the EVD rate, the lower the AUC ratio, reflecting the increased linezolid clearance. Thus, in case of CNS infection, it might

be advisable to limit CSF drainage, under optimal cerebral hemodynamics, to limit the additional clearance
due to EVD, reducing CSF linezolid concentrations.

From a pharmacodynamics perspective, plasma PK/PD index related to linezolid efficacy is not clearly
defined and varies among studies, utilizing AUC/CMI or %T>CMI[13–18]. In the first study describing *in vivo*
linezolid PD from animal models of acute infections, the more relevant of the two PK/PD parameters was
difficult to determine[33]. In seriously ill patients, a linear relationship between these two parameters was
characterized for linezolid at AUC/MIC values < 120, and higher clinical success rates occurred at AUC/MIC
ratios of 80-120 or T>MIC of 100%[20]. However, careful interpretation of these results is essential as only 50%
of patients received linezolid intravenously and only 2 patients were treated for a CNS infection. In literature,
plasma PK/PD targets are currently used to optimize the treatment of cerebral infections, with AUC/MIC >
100 most often retained, as no target based on CSF concentrations exist[13,16,17]. Thus, depending on studies,
indexes and targets used varied, leading to different conclusions. Luque *et al.,* targeting 24-h AUC/MIC $\geq$ 80
in CSF, considered that higher than standard doses were necessary (AUC/MIC of 21 for pathogens with a MIC
of 2 mg/L)[13]. In contrast, Beer *et al.* suggested that 600 mg linezolid twice daily provided satisfactory PK/PD
results in ventriculitis patients (T>MIC of 99.8% for pathogens with a MIC of 2 mg/L)[14].

According to our results, selecting as PK/PD target a 24-h CSF AUC/MIC ratio > 100, the recommended dose
of 1200 mg/day achieved PTAs > 90% only for Staphylococci with MICs < 0.5 mg/L. Increasing daily dose to
2700 mg/day would allow reaching PTA > 90% for MICs up to 1 mg/L only, and would allow covering
linezolid MIC distribution for MRSE (CFR = 90%) but not for MRSA (CFR = 32%). Whatever doses, none of
the administration modalities tested showed any impact on PTA and CFR values, and hence any benefit in
terms of efficacy for MRSA-infected patients. These results could challenge linezolid indication, considering
that linezolid is recommended as a second-line treatment or for MRSA with vancomycin MIC higher than 1
mg/L[5]. Nevertheless, clinical efficacy of 600mg twice daily was reported in cases of failure of vancomycin
treatment for CNS MRSA infections, whereas most PK/PD studies in CSF recommend increasing dosages[34].

If we had adopted 100% *f*T>MIC as CSF PK/PD target, results would have diverged significantly. While 1800
mg daily, whatever the administration scheme, seemed insufficient for pathogens with MICs > 0.5 mg/L when
targeting CSF AUC/CMI > 100 (PTAs = 98%, 73% and 21% for MICs of 0.5, 1 and 2 mg/L, respectively),
with 100% CSF *f*T>MIC, continuous infusion or splitting the dose into 600 mg t.i.d would have resulted in
higher PTA for MICs up to 2 mg/L ($\geq$ 98% and $\geq$ 85%, respectively) (Figure S6). Thus, for MRSE, 1200 mg
daily by continuous infusion would have been sufficient (CFR > 95%), while 2400 mg daily by continuous
infusion would have been appropriate for MRSA (CFR = 95%) (Table S3).

Increasing dosing regimens raises concerns about the risk of linezolid hematological toxicity. However, an
increased risk was exclusively reported in cases of renal failure, with the need to adapt dosing regimen when

Clcr < 60 mL/min[35,36]. Brain-injured patients, on the other hand, are frequently exposed to augmented renal
clearance (ARC)[38], as demonstrated here with 18 of our 25 patients exhibiting ARC. Thus, none reached the
toxicity threshold (i.e. residual concentration > 10 mg/L), even with 1800 mg daily[37]. Recent studies involving
ICU patients highlighted risks of underdosing in cases of ARC, indicating the usefulness of dose adaptation.
An ICU prospective multicenter study recommended increasing linezolid up to 2400 daily for patients with
ARC[39]. An early therapeutic drug monitoring is perhaps the first step to optimize linezolid dosages in ICU
patients, even if linezolid does not require adjustment based on renal clearance theoretically.

The first strength of our study is its design. Its multicenter framework allowed to include the highest number
of patients with cerebral infections, all confirmed by a blinded adjudication committee (n=14), and to evaluate
linezolid cerebral penetration at two dosing regimens (600mg b.i.d for 17 patients; 600mg t.i.d for 8 patients).
Second, it is the largest ICU PK/PD study based on Monte Carlo simulations reporting PTA and for the first
time CFR of linezolid against MRSE and MRSA, coupled with CSF and plasma cytokine concentrations in
CNS and extra-CNS infected patients. For the first time, the EVD flow rate was described to influence linezol-
id diffusion in CSF.

The first limitation of our study is the lack of patients with renal clearance lower than 80 mL/min, making it
impossible to demonstrate any link between linezolid and creatinine clearances, and to propose a dosing regi-
men adjustment based on renal function. Secondly, the absence of follow-up of clinical recovery and bacterial
eradication made impossible to determine a PD threshold adapted to CNS infections. The need for targets
based on CSF concentrations is urgent, but the complexity of gathering a large number of infected patients
undergoing monotherapy treatment as clinicians often resort to antibiotic combinations, make PK/PD analysis
difficult and uncertain[40]. An approach based on a hollow-fiber infection model that enables continuous culture
of bacteria and mimics the clinical drugs CSF concentration profiles, could overcome these limitations.

In conclusion, study findings support extensive linezolid diffusion into CSF of severely brain-injured patients,
influenced by the EVD flow rate, and highlight that currently recommended doses (600mg b.i.d) would not be
appropriate for covering infections due to staphylococci in ICU patients with normal or ARC, whatever
PK/PD index chosen. Considering AUC/MIC > 100 in CSF, 2700mg daily would cover only MRSE, whereas
with 100% fT>MIC, daily continuous infusions of 1200mg and 2400mg would be appropriate for MRSE and
MRSA, respectively. However, these results should be interpreted with caution without any well-defined PD
target for cerebro-meningeal infections. Further PK/PD studies including clinical outcomes in larger popula-
tion are necessary to identify such linezolid PD index in CSF of severe brain-injured patients.

 Figures and Tables

292 Figure 1: Boxplots of concentrations of IL6, IL8, IL10, IL17A and TNF-alpha in CSF (in blue) and plasma (in

293 red) for "Extra-CNS Infection" group (Top line boxplots) and for "CNS Infection" group (Bottom line

294 boxplots). * p<0.05.

309 Figure 2: Boxplots of CSF concentrations of IL6, IL8, IL10, IL17A and TNF-alpha in patients with extra-

310 CNS infections (in blue) and patients with CNS infections (in yellow). * p<0.05.

324   Figure 3. Simulations of the mean free concentration-time profiles of linezolid at steady-state in plasma (red

325   solid lines), cranial CSF (blue solid lines) and EVD (grey dotted lines) after administration of 600mg q12h and

326   600mg q8h.

342   Figure 4: Relationship between CSF/free plasma AUC ratio and EVD flow rate. Circles represent the

343   individual predictions of CSF/free plasma AUC ratio for the mean EVD flow rate measured on the PK day for

344   each patient. The line represents the fit of a linear function and the grey shaded area the 95% confidence

345   interval.

350   Figure 5. PTA (fAUC$_{0-24}$ > 100) in plasma (dashed lines) and CSF (solid lines) in simulated patients receiving
351   different daily doses of linezolid (1 200, 1 800 and 2 700 mg/day). The distribution of linezolid MICs for
352   methicillin resistant *S. aureus* (MRSA) and *S. epidermidis* (MRSE) is shown.

372 Table 1. Median free plasma and CSF PK parameters at steady-state after administration of 600mg q12h (n=17
373 patients) and 600mg q8h (n=8 patients) of linezolid.
374

	600 mg q12h		600 mg q8h	
	Plasma	CSF	Plasma	CSF
Free C_{max} (mg/L)	12.2 ± 3.4	5.3 ± 2.0	13.7 ± 3.7	6.7 ± 2.7
T_{max} (h)	0.5 ± 0.0	3.4 ± 0.6	0.5 ± 0.0	2.9 ± 0.4
Free AUC_{0-24} (h.mg/L)	102.4 ± 42.6	85.3 ± 37.6	150.7 ± 61.3	123.3 ± 53.6
$AUC_{CSF}/fAUC_{pl}$	0.838 ± 0.151		0.823 ± 0.147	

375
376
377

Table 2. Cumulative fraction of response (CFR) of linezolid against methicillin resistant *S. aureus*
(MRSA) and *S. epidermidis* (MRSE) in simulated patients receiving different daily doses

		1 200 mg/day	1 800 mg/day	2 700 mg/day
MRSA	Plasma	3.9%	16%	38%
	CSF	3.2%	13%	32%
MRSE	Plasma	48%	78%	94%
	CSF	40%	70%	90%

378
379

Acknowledgments

We thank the patients and their relatives and caregivers, local site clinical and research staff, regional

coordinating centre staff of the university hospital of Poitiers, and all the members of the PK-Pop-LCR Study

for their active participation.

Financial support

This work was supported by the French Ministry of Social Affairs and Health as source of funding for this

study (Grant numbers: PHRCN-16-0501).

Authors contributions

CDF, AC, NG, WC and SM conceived the study, participated in protocol development, study design, data

interpretation, and co-wrote the first draft of the report. CD-F coordinated the study. CDF, FB, KC, HdC, PEL,

GF, RCh, RCi, JP, BB, AM participated in inclusion of patients and data collection. DCF and FB participated

in the masked central adjudication committee. OR verified and CA, JCL analyzed samples. CDF, OR, AC,

JCL, FB and AM accessed and verified the data. AC, AM, NG and SM provided statistical and

pharmacokinetic expertise. All authors attest to have had full access to all the data in the study, contributed to

manuscript revision and to have approved the final manuscript prior to submission. All collaborators for the

PK-Pop-LCR study study group, full names and academic degrees, are reported in the supplementary

appendix.

Declaration of interests

CDF received honoraria for lectures from Pulsion and support for attending meetings and/or travel from

Codman, SOPHYSA and Pfizer. JP received honoraria for lectures from Masimo, Edwards Lifesciences,

support for attending meetings and/or travel from Masimo and consulting fees from AOP Orphan, Takeda,

Masimo and Acticor Biotech. All other authors report no potential conflicts.All authors have submitted the

ICMJE for Disclosure of potential Conflicts of Interest.

Further Members of the PK-Pop-LCR study group with linked authorship in Pubmed

Fanny BERNARD, M.D (Centre Hospitalier Universitaire de Poitiers, Service d'Anesthésie-Réanimation et

Médecine Péri-Opératoire, Poitiers, France.), Carine DOMINIQUE, M.D (Centre Hospitalier Universitaire de

Poitiers, Service d'Anesthésie-Réanimation et Médecine Péri-Opératoire, Poitiers, France.), Tony

TRIQUARD, M.D (Centre Hospitalier Universitaire de Poitiers, Service d'Anesthésie-Réanimation et

Médecine Péri-Opératoire, Poitiers, France.), Amélie RUBIN, M.D (Centre Hospitalier Universitaire de

Poitiers, Service d'Anesthésie-Réanimation et Médecine Péri-Opératoire, Poitiers, France.), Jean-Claude

LECRON, M.D., PhD (Université de Poitiers, LITEC, France), Adriana DELWAIL (Université de Poitiers,

LITEC, France), Rémy BELLIER, M.D. (Centre Hospitalier Universitaire de Poitiers, Service d'Anesthésie-
Réanimation et Médecine Péri-Opératoire, Poitiers, France.), Thierry BENARD, M.D. (Centre Hospitalier
Universitaire de Poitiers, Service d'Anesthésie-Réanimation et Médecine Péri-Opératoire, Poitiers, France.),
Nadia IMZI, C.R.A. (Centre Hospitalier Universitaire de Poitiers, Direction de la Recherche Clinique, Poitiers,
France.), Angela KOSTENCOVSKA, C.R.A (Centre Hospitalier Universitaire de Poitiers, Direction de la
Recherche Clinique, Poitiers, France.), Sabrina SEGUIN, C.R.A (Centre Hospitalier Universitaire de Poitiers,
Direction de la Recherche Clinique, Poitiers, France.), Véronique FERRAND-RIGALLAUD, C.R.A. (Centre
Hospitalier Universitaire de Poitiers, Direction de la Recherche Clinique, Poitiers, France.), Séverine
CLERJAUD, (Centre Hospitalier Universitaire de Poitiers, Direction de la Recherche Clinique, Poitiers,
France.), Flora DJANIKIAN, M.D. (Centre Hospitalier Universitaire de Montpellier, Anaesthesia and
Intensive Care Department, Montpellier, France.), Pierre-François PERRIGAULT, M.D. (Centre Hospitalier
Universitaire de Montpellier, Anaesthesia and Intensive Care Department, Montpellier, France.), Karim
LAKKHAL, M.D. (Nantes Université, Centre Hospitalier Universitaire de Nantes, Service d'Anesthésie
Réanimation, F-44000, Nantes, France.), Matthieu BIAIS, M.D., Ph.D. (Centre Hospitalier Universitaire de
Bordeaux, Anaesthesia and Intensive Care Unit, Bordeaux, France), Grégoire CHADEFAUX, M.D. (Centre
Hospitalier Universitaire de Bordeaux, Anaesthesia and Intensive Care Unit, Bordeaux, France) Aurore
RODRIGUES, M.D. (Centre Hospitalier Universitaire du Kremlin-Bicêtre, Paris, France), Jean-François
PAYEN, M.D., PhD (Centre Hospitalier Universitaire de Grenoble, Anaesthesia and Intensive Care
Department, Grenoble, France), Magdalena SZCZOT M.D. (Centre Hospitalier Universitaire de Strasbourg,
Anaesthesia and Intensive Care Department, Strasbourg, France), Stéphane HECKETSWEILER C.R.A.
(Centre Hospitalier Universitaire de Strasbourg, Direction de la Recherche Clinique, Strasbourg, France.),
Nicolas ENGRAND M.D.(Hôpital Fondation Adolphe de Rotchild, Anaesthesia and Intensive Care Unit,
Paris, France), Michel WOLFF, M.D, PhD. (Hôpital Sainte Anne, Service de Neuroréanimation, Paris,
France), Gérard AUDIBERT, M.D, PhD. (Centre Hospitalier de Nancy, Service d'Anesthésie Réanimation,
Nancy, France), Frédéric DAILLER, M.D. (Hospices civils de Lyon, Groupement Est, Service d'Anesthésie
Réanimation, Lyon, France).

References

1. Luque-Paz D, Revest M, Eugène F. Ventriculitis: A Severe Complication of Central Nervous System Infections. Open Forum Infect Dis 2021;8(6).

2. Srihawan C, Castelblanco RL, Salazar L. Clinical Characteristics and Predictors of Adverse Outcome in Adult and Pediatric Patients With Healthcare-Associated Ventriculitis and Meningitis. Open Forum Infect Dis 2016;3(2).

3. Martin RM, Zimmermann LL, Huynh M, Polage CR. Diagnostic Approach to Health Care- and Device-Associated Central Nervous System Infections. J Clin Microbiol 2018;56(11).

4. Ramanan M, Lipman J, Shorr A, Shankar A. A meta-analysis of ventriculostomy-associated cerebrospinal fluid infections. BMC Infect Dis 2015;15(3).

5. Tunkel AR, Hasbun R, Bhimraj A. Infectious Diseases Society of America's Clinical Practice Guidelines for Healthcare-Associated Ventriculitis and Meningitis. Clin Infect Publ Infect Soc Am 2017;2017;64(6):e34–65.

6. Jalusic KO, Hempel G, Arnemann P-H. Population pharmacokinetics of vancomycin in patients with external ventricular drain-associated ventriculitis. Br J Clin Pharmacol 2021;87(6):2502–10.

7. The European Committee on Antimicrobial Susceptibility Testing. Breakpoint tables for interpretation of MICs and zone diameters [Internet]. 2024;Available from: http://www.eucast.org

8. Hashemian SMR, Farhadi T, Ganjparvar M. Linezolid: a review of its properties, function, and use in critical care. Drug Devel Ther 2018;12:1759–67.

9. Sun F, Ruan Q, Wang J. Linezolid manifests a rapid and dramatic therapeutic effect for patients with life-threatening tuberculous meningitis. Antimicrob Agents Chemother 2014;58(10):6297–301.

10. Smith AGC, Gujabidze M, Avaliani T. Clinical outcomes among patients with tuberculous meningitis receiving intensified treatment regimens. Int J Tuberc Lung J Int Union Tuberc Lung Dis 2021;25(8):632–9.

11. Kempker RR, Smith AGC, Avaliani T. Cycloserine and Linezolid for Tuberculosis Meningitis: Pharmacokinetic Evidence of Potential Usefulness. Clin Infect Dis 2022;75(4):682–9.

12. Viaggi B, Cangialosi A, Langer M. Tissue Penetration of Antimicrobials in Intensive Care Unit Patients: A Systematic Review-Part II. Antibiot Basel Switz 2022;11(9).

13. Luque S, Grau S, Alvarez-Lerma F. Plasma and cerebrospinal fluid concentrations of linezolid in neurosurgical critically ill patients with proven or suspected central nervous system infections. Int J Antimicrob Agents 2014;44(5):409–15.

14. Beer R, Engelhardt KW, Pfausler B. Pharmacokinetics of Intravenous Linezolid in Cerebrospinal Fluid and Plasma in Neurointensive Care Patients with Staphylococcal Ventriculitis Associated with External Ventricular Drains. Antimicrob Agents Chemother 2007;51(1):379–82.

15. Myrianthefs P, Markantonis SL, Vlachos K. Serum and cerebrospinal fluid concentrations of linezolid in neurosurgical patients. Antimicrob Agents Chemother 2006;50(12):3971–6.

16. Wu X, Tang Y, Zhang X, Wu C, Kong L. Pharmacokinetics and pharmacodynamics of linezolid in
plasma/cerebrospinal fluid in patients with cerebral hemorrhage after lateral ventricular drainage by
Monte Carlo simulation. Drug Devel Ther 2018;12:1679–84.

17. Zhao W, Kong L, Wu C, Wu X. Prolonged infusion of linezolid is associated with improved
pharmacokinetic/pharmacodynamic (PK/PD) profiles in patients with external ventricular drains. Eur J
Clin Pharmacol 2020;

18. Viaggi B, Paolo AD, Danesi R. Linezolid in the central nervous system: comparison between
cerebrospinal fluid and plasma pharmacokinetics. Scand J Infect Dis 2011;43(9):721–7.

19. Reiber H, Felgenhauer K. Protein transfer at the blood cerebrospinal fluid barrier and the quantitation of
the humoral immune response within the central nervous system. Clin Chim Acta 1987;163(3):319–28.

20. Rayner CR, Forrest A, Meagher AK, Birmingham MC, Schentag JJ. Clinical Pharmacodynamics of
Linezolid in Seriously Ill Patients Treated in a Compassionate Use Programme. Clin Pharmacokinet
2003;42(15):1411–23.

21. The European Committee on Antimicrobial Susceptibility Testing. Antimicrobial wild type distributions
of microorganisms [Internet]. 2024;Available from: https://mic.eucast.org/search/

22. Mouton JW, Dudley MN, Cars O, Derendorf H, Drusano GL. Standardization of
pharmacokinetic/pharmacodynamic (PK/PD) terminology for anti-infective drugs: an update. J
Antimicrob Chemother 2005;55(5):601–7.

23. Kumta N, Roberts JA, Lipman J, Wong WT, Joynt GM, Cotta MO. A Systematic Review of Studies
Reporting Antibiotic Pharmacokinetic Data in the Cerebrospinal Fluid of Critically Ill Patients with
Uninflamed Meninges. Antimicrob Agents Chemother 2020;65(1):e01998-20.

24. Yogev R, Damle B, Levy G, Nachman S. Pharmacokinetics and Distribution of Linezolid in
Cerebrospinal Fluid in Children and Adolescents. Pediatr Infect J 2010;29(9):827–30.

25. Yekani M, Memar MY. Immunologic biomarkers for bacterial meningitis. Clin Chim Acta
2023;548:117470.

26. König C, Grensemann J, Czorlich P, Schlemm E, Kluge S, Wicha SG. A dosing nomograph for
cerebrospinal fluid penetration of meropenem applied by continuous infusion in patients with
nosocomial ventriculitis. Clin Microbiol Infect Off Publ Eur Soc Clin Microbiol Infect Dis
2022;28(7):1022.e9-1022.e16.

27. Nau R, Lassek C, Kinzig-Schippers M, Thiel A, Prange HW, Sörgel F. Disposition and elimination of
meropenem in cerebrospinal fluid of hydrocephalic patients with external ventriculostomy. Antimicrob
Agents Chemother 1998;42(8):2012–6.

28. Ishikawa M, Yamazaki S, Suzuki T, Uchida M, Iwadate Y, Ishii I. Correlation between vancomycin
penetration into cerebrospinal fluid and protein concentration in cerebrospinal fluid/serum albumin ratio.
J Infect Chemother J Jpn Soc Chemother 2019;25(2):124–8.

29. Tarnaris A, Toma AK, Chapman MD. Rostrocaudal dynamics of CSF biomarkers. Neurochem Res
2011;36(3):528–32.

30. Weisner B, Bernhardt W. Protein fractions of lumbar, cisternal, and ventricular cerebrospinal fluid. Sep
Areas Ref J Neurol Sci 1978;37(3):205–14.

31.   Chen C, Xu P, Xu T, Zhou K, Zhu S. Influence of cerebrospinal fluid drainage and other variables on the
plasma vancomycin trough levels in postoperative neurosurgical patients. Br J Neurosurg
2021;35(2):133–8.

32.   Li X, Wang X, Wu Y, et al. Plasma and cerebrospinal fluid population pharmacokinetic modeling and
simulation of meropenem after intravenous and intrathecal administration in postoperative neurosurgical
patients. Diagn Microbiol Infect Dis 2019;93(4):386–92.

33.   Andes D, van Ogtrop ML, Peng J, Craig WA. In Vivo Pharmacodynamics of a New Oxazolidinone
(Linezolid). Antimicrob Agents Chemother 2002;46(11):3484–9.

34.   Fu X, Lin Z, Chen S, Hong L, Yu X, Wu S. Treatment of Intracranial Infection Caused by Methicillin-
Resistant Staphylococcus epidermidis with Linezolid Following Poor Outcome of Vancomycin Therapy:
A Case Report and Literature Review. Infect Drug Resist 2021;14:2533–42.

35.   Shi C, Xia J, Ye J. Effect of renal function on the risk of thrombocytopaenia in patients receiving
linezolid therapy: A systematic review and meta□analysis. Br J Clin Pharmacol 2022;88(2):464–75.

36.   Crass RL, Cojutti PG, Pai MP, Pea F. Reappraisal of Linezolid Dosing in Renal Impairment To Improve
Safety. Antimicrob Agents Chemother 2019;63(8).

37.   Cattaneo D, Orlando G, Cozzi V, et al. Linezolid plasma concentrations and occurrence of drug-related
haematological toxicity in patients with Gram-positive infections. Int J Antimicrob Agents
2013;41(6):586–9.

38.   Bilbao-Meseguer I, Rodríguez-Gascón A, Barrasa H, Isla A, Solinís MÁ. Augmented Renal Clearance in
Critically Ill Patients: A Systematic Review. Clin Pharmacokinet 2018;57(9):1107–21.

39.   Wang X, Wang Y, Yao F. Pharmacokinetics of Linezolid Dose Adjustment for Creatinine Clearance in
Critically Ill Patients: A Multicenter, Prospective, Open-Label, Observational Study. Drug Devel Ther
2021;15:2129–41.

40.   Delattre IK, Taccone FS, Jacobs F, et al. Optimizing β-lactams treatment in critically-ill patients using
pharmacokinetics/pharmacodynamics targets: are first conventional doses effective? Expert Rev Anti
Infect Ther 2017;15(7):677–88.

541
542