Postpartum Psychosis: could genetic vulnerability to insomnia or short sleep duration be protective?

Chiara Petrosellini MRCOG[1,2,3*], Sofia H Eriksson FRCP PhD[4,5], Nicholas Meyer MRCPsych PhD[6,7], Olivia Protti MRCOG MRCPsych[1,8], Nicholas Bass MD MRCPsych[2], Karoline Kuchenbaecker PhD[2], Dimitrios Siassakos MDRes FRCOG[1,3] & Andrew McQuillin PhD[2]

[1] Elizabeth Garrett Anderson Institute for Women's Health, University College London
[2] Division of Psychiatry, University College London
[3] Elizabeth Garrett Anderson Wing, University College London Hospitals NHS Foundation Trust
[4] Department of Clinical and Experimental Epilepsy, Institute of Neurology, University College London
[5] National Hospital for Neurology and Neurosurgery, University College London Hospitals NHS Foundation Trust
[6] Insomnia and Behavioural Sleep Medicine Clinic, University College London Hospitals NHS Foundation Trust
[7] Department of Psychosis Studies, Institute of Psychology, Psychiatry and Neuroscience, King's College London
[8] East London Mother and Baby Unit, East London NHS Foundation Trust

*Correspondence to:
Dr Chiara Petrosellini
chiara.petrosellini.22@ucl.ac.uk
UCL EGA Institute for Women's Health
84-86 Chenies Mews, London WC13 6HU
Tel: +44(0)20 3108 2035

Abstract

Postpartum Psychosis (PP) is a severe and understudied perinatal mental illness which disproportionately affects women with bipolar disorder (BD). A relationship between sleep disturbance and PP is often assumed, but is poorly understood. From a cohort of 2099 individuals with BD, 343 parous women were identified and screened for perinatal psychiatric complications. We compared 117 women who developed PP with 226 who did not. Polygenic Risk Scores (PRS) for BD, schizophrenia, insomnia, short sleep, long sleep, sleep efficiency and sleep duration were computed using PRS-CS. Logistic regression was used to model the effect of each PRS on PP. Higher PRS for insomnia and short sleep were associated with reduced risk of PP. Individuals in the lowest decile for insomnia PRS (RR 1.96, 95% CI 1.25-3.07, $p = 3.50 \times 10^{-3}$) and short sleep PRS (RR 2.23, 95% CI 1.40-3.54, $p = 7.94 \times 10^{-4}$) had approximately double the risk of PP than individuals in the highest decile. The other PRS were not associated with PP. Mendelian Randomisation analyses did not support a causal relationship between sleep traits and PP. However, we demonstrate that the integration of PRS with bipolar subtype can improve prediction accuracy. Individuals with genetic vulnerability to insomnia or short sleep may develop a heightened tolerance to sleep disruption earlier in life, mitigating the impact of childbirth on mood. These findings suggest that genetic susceptibility to sleep disturbance may be important in the aetiology of PP, offering a new potential avenue for risk stratification and targeted prevention.

Introduction

Postpartum psychosis (PP) is a severe but treatable mental illness which affects 1-2 in 1000 women who give birth. Most episodes present with mania or a mixed affective state, often accompanied by the presence of psychotic or cognitive symptoms(1). PP is a psychiatric emergency due to its rapid onset and escalation, coupled with the associated risks of suicide and infanticide(2).

Whilst the disease classification of PP is a source of ongoing debate(3), it is widely conceptualised as a form of bipolar disorder (BD) with a vulnerability to childbirth. Women with BD carry a 20% risk of developing a psychotic or manic episode postpartum(4), and a third of women who experience first-onset PP will be diagnosed with BD long-term(5). Despite the close relationship between the two conditions, a previous psychiatric history does not explain most cases of PP and only a proportion of women with BD experience a postnatal relapse. Improved prediction of PP therefore requires better understanding of the role of other perinatal triggers.

Sleep is a fundamental physiological necessity which is almost universally disrupted around childbirth. Pregnancy and the postpartum period are times of extreme change in sleep quantity, quality and pattern(6). There is increasing recognition that these changes can represent both symptoms and causes of maternal mood and anxiety disorders(7). Outside the perinatal period, sleep loss is a well-established trigger of psychosis(8) and a recognised risk factor for suicidality(9). Experimentally-induced sleep deprivation has been shown to induce mania in women with postnatal psychosis(10) and cause a switch from depression to mania(11). Reduced need or ability to sleep is one of the earliest and most common presentations of PP(12) and sleep has been hypothesized to be the final common pathway through which other factors produce psychosis in susceptible women(13, 14).

There is increasing evidence that sleep may play a role in perinatal episodes for women with BD. Parous women with BD who report sleep loss as a lifetime trigger of manic episodes have been shown to be twice as likely to have had PP as women with BD who have had children and do not report sleep loss as a personal trigger(15). A prospective study has also demonstrated that women with BD who lose one or more nights of sleep over the intrapartum period have a significantly higher risk of developing PP compared to those who do not(16). It is plausible that there is a subset of women with BD who are particularly vulnerable to psychiatric deterioration following sleep disruption, whilst others may be more resilient when sleep patterns are disturbed.

Both BD(17) and sleep traits(18) are thought to have polygenic inheritance, whereby the combined effects of multiple common genetic variants contribute to disease risk. The data from genome-wide association studies (GWAS) of complex traits can be used to derive Polygenic Risk Scores (PRS), which provide an estimate of an individual's relative susceptibility to disease. Whilst a PRS alone has

limited practical application at present, the combination of PRS with other risk factors has been

increasingly shown to improve outcome prediction and clinical decision-making in psychiatry(19). A

PRS approach can also be used to elucidate disease mechanisms and has provided support to the

view that BD and first-onset PP may be biologically distinct(20).

BD demonstrates significant heterogeneity in its pathogenesis, clinical presentation and

outcomes(21). At present, prediction of PP in women with BD is imprecise. As a result, resource

allocation for pregnant women with pre-existing mental illness is also inefficient. In a direct

comparison of women with BD who did and who did not develop PP following childbirth, PRS for BD

and schizophrenia were found not to differ significantly between the two groups(20). Our initial

objective was to replicate this finding. We then sought to establish whether genetic vulnerability to

sleep disorders can be used to improve prediction of PP in women with BD. We chose to investigate

the following sleep traits, which we hypothesized would be associated with differing genetic risk

profiles in women with BD who develop PP and those who do not:

1. **Insomnia:** difficulty in initiating or maintaining sleep, despite adequate opportunity to sleep(22).

Observational(12, 23) and qualitative(24, 25) studies have consistently identified insomnia as one

of the most common symptoms of PP. In prospective studies of high-risk cohorts, insomnia has

been shown to predict the emergence of BD(26) and has been associated with an increased

likelihood of developing delusions and hallucinations(27).

2. **Short Sleep:** commonly defined as habitual sleep duration of less than 6 hours per night(28).

Reduced sleep duration is often a prodrome of mania(29) and has been shown to predict

deterioration in psychosis symptoms(30).

3. **Long Sleep:** commonly defined as habitual sleep duration of more than 9 hours per night(31).

Individuals with BD have been shown to have significantly longer sleep duration than healthy

controls, both in periods of euthymia(32) and during depressive relapses(33).

4. **Sleep Efficiency:** the proportion of time in bed which is spent asleep(34). There is consistent

evidence that individuals with BD have lower sleep efficiency than healthy controls(32) and a

recent meta-analysis has supported an association between reduced sleep efficiency and both

early and chronic psychosis(35).

5. **Sleep Duration:** the total amount of sleep obtained per nocturnal sleep episode or across the 24

hour period(36). It is consistently reported as a clinically relevant trait across mood episodes and

is linked with both onset and recurrence of BD(37). While short and long sleep have more clearly

established associations with affective and psychotic disorders, we also examined sleep duration

as a continuous trait given its importance in the interplay between sleep and mood(38-40).

Methods

Participants

Individuals with a diagnosis of BD based on the International Classification of Diseases, 10[th]

Revision(41) were recruited from National Health Service (NHS) general practices, hospitals and

community mental health teams in England. Participants were also recruited through the Bipolar UK

charity. This included individuals with Bipolar I Disorder (BDI), Bipolar II Disorder (BDII) and

Schizoaffective Bipolar Disorder (SABD). Ancestrally matched healthy controls were recruited from

the same study sites in which BD cases were recruited and from the NHS blood transfusion

service(42). All controls were screened for the absence of a lifetime history of psychiatric illness

including all major psychotic and affective disorders. All BD cases and healthy controls were of White

British descent and had at least three out of four grandparents of the same ancestry. Written informed

consent was obtained from all participants. The study received NHS Research Ethics Committee and

Health Research Authority approval (MREC/03/11/090).

Study Measures and Definitions

All participants with BD completed semi-structured interviews using the lifetime version of the

Schizophrenia and Affective Disorder Schedule (43) and the Operational Criteria Checklist for

Psychotic Illness (44). Female participants who have given birth were screened for a history of

perinatal psychiatric complications using the International Cohort Collection for Bipolar Disorder

(ICCBD) variables checklist(45). In this cohort, Postpartum Psychosis (PP) was defined as an

episode of mania, psychosis or psychotic depression following childbirth.

There is no universal consensus on the timeframe which defines the postnatal period, and there is no

widely accepted definition for what qualifies as a 'perinatal' episode in women with pre-existing mental

illness. Whilst the Diagnostic and Statistical Manual of Mental Disorders (DSM) defines this time

period as four weeks(46) and the International Classification of Diseases (ICD) defines this as six

weeks following delivery(47), perinatal services in the United Kingdom are designed to care for

women until at least a year following childbirth(48). This reflects an increasing recognition that the

biopsychosocial changes associated with motherhood have profound personal and public health

implications which extend well beyond the early postnatal period(49). In research studies, definitions

of PP have varied widely, with some restricting onset to within two weeks of delivery and others

including all episodes occurring within the first postpartum year(50). Whilst most episodes of PP

emerge within the first two weeks postpartum(51), onset beyond three months is well documented.

Such late-onset presentations have been described in both historical and contemporary literature,

and have been linked to physiological transitions such as the resumption of menstruation or the

weaning process(52). To reflect the full breadth of clinical vulnerability, we included all episodes

occurring within the first twelve months postpartum, consistent with standard definitions of the

perinatal period(48). We conducted an additional sensitivity analysis in which we only included
individuals who met strict ICD definitions for a perinatal episode. The data were coded in a binary
format, to separate parous women with BD into two groups: those who had and those who had not
experienced PP following childbirth.
197
198   ## Discovery datasets for sleep phenotypes
199   Summary statistics of GWAS were obtained for insomnia(53), short and long sleep(54), sleep
200   efficiency and sleep duration(55) to conduct our analyses.
201
The insomnia GWAS(53) was conducted in participants recruited to the UK Biobank and 23andMe.
Insomnia was defined as self-reported difficulty in falling asleep or staying asleep, which was
assessed using a single question: 'Do you have trouble falling asleep at night or do you wake up in
the middle of the night?'. Answers to this question were then dichotomised to create a binary
phenotype. A total of 593,724 cases and 1,771,286 controls were included in the GWAS. We used
summary statistics from a subset of 386,988 European individuals from the UK Biobank (109,548
cases and 277,440 controls) to generate our PRS.
209
Austin-Zimmerman et al. performed a multi-ancestry GWAS of self-reported short and long sleep
duration in adults recruited to the UK Biobank and the Million Veteran Programme(54). Self-reported
sleep duration was measured as a continuous trait and reported in hourly increments for a total of
493,142 individuals. Short sleep was defined as habitual sleep duration ≤5 hours, whilst long sleep
was defined as habitual sleep duration ≥10 hours. Normal sleep duration was defined as 7-8 hours
per night. Their GWAS meta-analyses compared short versus normal sleep duration and long versus
normal sleep duration. We used summary statistics from a subset of 445,966 individuals of European
descent to calculate PRS for short sleep (47,054 cases and 382,950 controls) and long sleep (15,962
cases and 382,950 controls).
219
Jones et al. used accelerometer data from 85,670 UK Biobank participants to perform a GWAS of
eight objective sleep traits(55). Sleep efficiency was defined as sleep duration divided by the time
period of nocturnal inactivity.
This measure was reported as a percentage ratio in 84,810 individuals of European ancestry. We
used summary statistics from this subset to generate PRS for sleep efficiency in our sample. The
same summary statistics also included data for accelerometer-derived sleep duration in 85,449
individuals, reported as a continuous measure. We used these to generate PRS for objective sleep
duration in our sample as a follow-on analysis, to assess whether these findings differed to those
based on binary, self-reported measures of sleep duration reported by Austin-Zimmerman et al(54).
229
The remaining accelerometer-derived sleep traits reported by Jones et al., were not included in our
analyses as current evidence does not strongly support their association with BD or psychosis. While
other objective sleep parameters such as M10 timing(56, 57), sleep midpoint(58) and sleep

fragmentation(59, 60) may play a role in affective or psychotic disorders, the associations between
these traits and illness onset or relapse risk are either limited or inconsistent. They were therefore not
included in our hypothesis-driven analyses.
236

Genotyping

Participants provided blood or saliva samples for DNA extraction and genome-wide single nucleotide
polymorphism (SNP) genotyping was performed. The data was generated at the Broad Institute
(Boston, USA) using the Illumina PsychArray or the Illumina Global Screening Array (GSA)
genotyping platforms. Strict quality control (QC) was performed separately for each platform using
PLINK1.9(61). Imputation was undertaken against the Haplotype Reference Consortium panel using
the Sanger Imputation Server(62). Details of the genotyping, QC and imputation methods used have
been described elsewhere(63).
245

Polygenic Risk Score (PRS) Calculation

Summary statistics for BD(17) and schizophrenia(64) were obtained from the Psychiatric Genetic
Consortium. Summary statistics for sleep phenotypes were obtained from the GWAS described
above. PRS were computed for BD, schizophrenia and our sleep traits of interest in both subjects
with BD and healthy controls. These were generated using the PRS-CS auto method(65), which
provides a single score for each individual without the need for p value selection thresholds. PRS-CS
requires GWAS summary statistics and a linkage disequilibrium reference panel, and we chose to use
the European reference panel from the 1000 Genomes Project(66) for the latter.
254

255 For each trait of interest, the raw PRS values were normalised into Z scores centred around a mean
256 of 0 and a standard deviation of 1 in the healthy control subjects, allowing direct comparison across
257 our cohorts (GSA, PsychArray and healthy controls). The normally distributed PRS for the
258 PsychArray and GSA datasets were then merged for downstream analysis.
259

Statistical Analysis

Data analysis was conducted in R using version 4.3.2(67). Logistic regression analyses were
performed to examine the effect of each PRS on PP following childbirth in women with BD. The
genotyping platform used and the first three population principal components were added as
covariates in all analyses to account for chip and ancestry confounding. To account for multiple
testing, we used the p.adjust function in R to apply the Bonferroni correction method(68) to each PRS
analysis. The pROC package in R(69) was used to compute the Receiver Operating Characteristic
(ROC) curves and Area Under the Curve (AUC) for significant PRS and for bipolar subtype. For each
ROC analysis, 95% confidence intervals for AUCs were obtained using 2000 stratified bootstrap
replicates, a standard approach for obtaining stable ROC confidence intervals(70). Differences
between AUCs were assessed using DeLong's test for two correlated ROC curves(71).
271

For sleep traits in which we observed a significant association between PRS and PP diagnosis, we

also added BD PRS and schizophrenia PRS Z scores as covariates in the regression model. In doing

so, we attempted to account for possible confounding from differences in relative genetic

susceptibility to psychotic illness. Individuals with BDI or SABD are thought to have a higher risk of

PP following childbirth than individuals with BDII(72). We therefore also adjusted for bipolar subtype

in our regression analyses. To further test the validity of our significant findings, we performed an

additional sensitivity analysis in which we only included individuals who developed PP within 6 weeks

of childbirth in line with the strict ICD definition of a 'perinatal episode' and compared their PRS with

individuals who did not develop PP following childbirth.

We performed power calculations using the Additive Variance Explained and Number of Genetic

Effects Method of Estimation (AVENGEME) package(73) to estimate whether our analyses were

sufficiently powered to detect a significant effect. Of note, AVENGEME is designed to perform power

calculations for PRS analyses using older p-value thresholding methods, which limits its application to

analyses performed with the newer PRS-CS auto method.

Mendelian Randomisation (MR) Analyses

For sleep phenotypes where we identified significant associations between PRS and PP, we

proceeded with two-sample Mendelian Randomisation (MR) studies to examine whether exposure to

specific sleep traits may be causally related to our outcome of interest: PP in women with BD. MR is

an epidemiological method which uses genetic variants as instrumental variables to assess causal

relationships between potential risk factors and an outcome of interest(74). By using genetic variants

as natural experiments, this approach reduces bias from confounding and reverse causation,

providing more robust evidence for causal inference. The genetic variants used in MR analyses must

satisfy three key assumptions: (i) that they are strongly associated with the exposure of interest, (ii)

that they are independent of confounders that could influence the outcome, and (iii) that they only

influence the outcome through their effect on the exposure(75). While the genetic association with the

exposure can be evaluated, the absence of confounding and exclusion restriction cannot be tested

directly.

We extracted genome-wide significant, linkage-disequilibrium independent SNPs from sleep GWAS

summary statistics and used these as genetic instruments for insomnia(53) and short sleep(54). We

then estimated SNP-outcome associations by using individual-level data from our cohort of parous

women with BD. All MR analyses were conducted using the TwoSampleMR package in R(76), which

uses the Inverse Variance Weighted (IVW), MR Egger, Weighted Median, Simple Mode and Weighted

Mode MR methods to test the relationship between the genetic instruments and outcomes of interest.

We conducted heterogeneity tests using both MR Egger and IVW methods, and assessed directional

horizontal pleiotropy via the MR Egger intercept.

Our findings are reported in accordance with the Strengthening the reporting of observational studies

in epidemiology using MR (STROBE-MR) checklist(77). In line with STROBE-MR guidance(78), we

limited MR testing to hypotheses supported by our own findings, to minimise the risk of false-positive

findings from exploratory analyses.

Results

PRS were initially generated for 2099 individuals with BD and 1818 healthy controls in our wider
cohort.
343 female participants with BD who had given birth at the point of recruitment were included in our
final analyses. Of these individuals, 213 (62.1%) met criteria for BDI, 37 (10.8%) met criteria for BDII
and 28 (8.2%) met criteria for SABD. In 65 (19.0%) participants, the BD subtype could not be
specified. Of all parous women with BD, 117 had experienced PP following childbirth at the point of
recruitment and 226 had not. Those who developed PP had symptom onset between one day and 8
months postpartum. Participants with an underlying diagnosis of a high-risk Bipolar subtype (BPI or
SABD) were more likely to develop PP than those without($p=1.387\times10^{-4}$). Women who did and who
did not develop PP did not significantly differ in their age or parity at the point of recruitment.

A demographic summary of individuals included in our analyses is shown in Table 1. There was no
significant difference in the distribution of genotyping platform used in participants who did and who
did not develop PP following childbirth (Chi-squared= 2.717, df= 1, p= 0.099).

Individuals with lower insomnia PRS were found to be at increased risk of PP than individuals with
higher PRS. Each one standard deviation increase in insomnia PRS was associated with an odds
ratio of 0.718 for PP (95% CI 0.575-0.897; $p=3.5 \times 10^{-3}$). This finding remained significant despite
stringent correction for multiple testing (Table 2). Each one standard deviation increase in insomnia
PRS was therefore associated with an almost 30% decrease in the odds of developing PP. This
association remained significant when BD PRS, schizophrenia PRS and bipolar subtype were added
as a covariates in the logistic regression model. Women in the bottom decile for insomnia PRS had
approximately twice the risk of PP than those in the top decile (RR 1.96, 95% CI 1.25-3.07; Table 3).
The distribution of insomnia PRS in women who did and who did not develop PP is shown in Figure
1A.
We performed a sensitivity analysis comparing individuals who did not develop PP with a subset of
individuals who met strict ICD-11 criteria for PP (onset within six weeks postpartum). Whilst this
sample size was underpowered to detect a statistically significant effect, the direction of association
remained unchanged, with insomnia PRS being lower in individuals who developed PP (mean Z-
Score -0.37, SD 0.84) compared to those who did not (mean Z-Score 0.04, SD 1.01).

Individuals with lower short sleep PRS were also found to be at increased risk of PP compared to
individuals with higher PRS. Each one standard deviation increase in short sleep PRS was
associated with an odds ratio of 0.674 for PP (95% CI 0.535-0.849; $p=7.94\times10^{-4}$). This result
withstood Bonferroni correction for multiple testing (Table 2). Each one standard deviation increase in
short sleep PRS was therefore associated with a decreased odds of PP by approximately one third.
As with insomnia, the association between short sleep PRS and PP remained significant when BD

PRS, schizophrenia PRS and bipolar subtype were added as covariates in the logistic regression
analysis. Women in the bottom decile for short sleep PRS had over twice the risk of PP than those in
the top decile (RR 2.23, 95% CI 1.40-3.54; Table 3). The distribution of short sleep PRS in individuals
who did and who did not develop PP following childbirth is shown in Figure 1B. A sensitivity analysis
in which only individuals meeting ICD-11 criteria for PP were included as cases was underpowered to
detect a statistically significant effect, however the direction of effect was unchanged. Short sleep
PRS was found to be lower in individuals who developed PP (mean Z-Score -0.58, SD 0.86)
compared to those who did not (mean Z Score 0.17, SD 1.02).

Combining the data for insomnia and short sleep PRS improved prediction of PP in our cohort
compared to using bipolar subtype alone, as demonstrated by the increased area under the curve
(AUC) in the joint analysis (Figure 2). A model including bipolar subtype alone yielded an AUC of
0.602 (95% CI: 0.555-0.648), whereas adding insomnia and short sleep PRS increased the AUC to
0.697 (95% CI: 0.637-0.753). DeLong's test confirmed that this improvement in discrimination was
statistically significant (Z= –3.91, p = 9.29 × 10□□). This suggests improved predictive accuracy
when polygenic risk information is included alongside clinical factors.

Logistic regression comparing individuals with BD who did and who did not develop PP following
childbirth did not identify a significant association between BD PRS and PP or between schizophrenia
PRS and PP (Table 2). We did not identify significant associations between long sleep PRS and PP
(Figure 1C), between sleep efficiency PRS and PP (Figure 1D) or between accelerometer-derived
sleep duration PRS and PP (Figure 1E) in our cohort. AVENGEME calculations suggested that all our
analyses had >80% power to detect a significant effect (p<0.05), even when stringent thresholds are
used in the model. The exception to this was long sleep, where our study may have been
underpowered to a detect a significant association (p<0.05) between PRS and PP.

Across all MR methods, we found no evidence to support a causal association between insomnia and
PP or short sleep and PP (Table 4). The MR analysis for insomnia showed significant heterogeneity in
both MR-Egger (Q = 27.50, df = 9, p = 1.16 × 10□³) and IVW (Q = 29.61, df = 10, p = 9.92 × 10□□)
tests, indicating potential variability in the causal estimates of the different genetic variants. We found
no evidence to suggest significant horizontal pleiotropy (MR-Egger intercept=0.10, SE = 0.12, p =
0.43).
For short sleep, we found no evidence of significant heterogeneity using MR-Egger (Q = 5.78, df =
15, p = 0.98) or IVW (Q= 6.19, df = 16, p = 0.99) tests. Additionally, the MR-Egger intercept was 0.03
(SE = 0.05, p = 0.53), indicating no evidence of directional horizontal pleiotropy.

Discussion

To the best of our knowledge, this study is the first to test the hypothesis that sleep PRS may

contribute to perinatal risk stratification in women with BD. We identified that individuals with lower

PRS for insomnia and short sleep were at significantly higher risk of developing PP. Women in the

bottom decile for each trait had approximately double the risk of PP compared to those in the top

decile. Robust statistical corrections were applied to account for potential confounders including

bipolar subtype, other PRS, and multiple testing, strengthening the validity of these results. Our

findings support previous work indicating that PRS of sleep phenotypes may contribute to risk

stratification in BD outside the perinatal period(79). These results suggest that incorporating sleep-

related PRS could enhance clinical risk stratification models and inform future preventative

approaches.

A genetic correlation (r_g) of 0.624 has been reported between insomnia and short sleep(54),

suggesting that genetic variants associated with one trait are likely to also influence the other to a

considerable extent. Notably, the strongest associations in the short sleep GWAS were identified in

SNPs in close proximity to *TCF4* and *FOXP2*, both encoding transcription factors which have also

been implicated in insomnia(53). If genetic vulnerabilities to self-reported insomnia and short sleep

play a role in PP risk, they may therefore do so through shared biological mechanisms. Interestingly,

emerging evidence suggests that both *TCF4* and *FOXP2* regulate neurodevelopmental pathways

relevant to psychosis. *TCF4* is essential for normal brain cortical development(80) and has been

implicated in schizophrenia(81) and major depression(82). *TCF4* was identified as the only gene

mediating the bidirectional causal relationship between insomnia and major depression in a MR

study(83), and forebrain overexpression of *TCF4* in mice has been shown to increase arousal and

shorten sleep duration(84). Similarly, *FOXP2* regulates genes involved in neuronal differentiation,

cortical development and synaptic plasticity(85). *FOXP2* polymorphisms have been implicated in

dopaminergic signalling, language processing and cognitive function in schizophrenia(86). These

findings suggest that genetic influences on arousal, cortical maturation and cognitive regulation may

represent a mechanistic framework through which sleep-related vulnerability may influence

postpartum psychiatric risk.

Insomnia and short sleep are clinically distinct, with the former representing an internally generated

state of hyperarousal leading to subjective perception of poorer sleep quality. This is not reliably

associated with objective short sleep duration(87), which can also be due to choice or external

circumstances. Nevertheless, the GWAS we used to generate PRS for these phenotypes are both

based on self-report, thereby capturing individuals' subjective experiences of their sleep. Individuals

who are biologically more prone to experience insomnia or short sleep may develop a heighted

tolerance to sleep disruption over time. It is plausible that the impact of acute postnatal sleep loss on

mood stability may be mitigated in women with BD who subjectively experience chronic sleep

restriction prior to childbirth. In the pathogenesis of PP, this would suggest that psychiatric

deterioration is not determined by sleep loss per se, but rather by the novelty associated with sleep

disruption. This interpretation is consistent with the observation that primiparity is a known risk factor

for PP(88), as first-time mothers may experience a heighted response to sleep loss compared to

multiparous women who would have previously been exposed to perinatal sleep disruption.

Perceived adjustment to chronic sleep loss is often a misperception, and subjective adaptation may

occur despite cumulative physiological impairment and sustained sleep pressure(89-91).

Nevertheless, this psychological adaptation may play a role in affective resilience. There is growing

interest in the concept of sleep resilience, which describes the capacity to maintain emotional,

cognitive and physical functioning when sleep or circadian patterns are disturbed(92). Significant

individual differences in vulnerability are observed and resilience in one functional domain does not

necessarily extend to others(93). While resilience is known to moderate the cognitive consequences

of poor sleep(94, 95) and mediate the relationship between sleep and mental illness(96-98), it is

uncertain whether this adaptability can be acquired through chronic sleep disturbance. Vulnerability to

sleep loss has been described as trait-like, as individual behavioural responses appear to be

relatively stable across both acute and more prolonged sleep restriction(99-101). However, the

existing evidence comes from small studies with brief observation periods which do not capture the

cumulative impact of lifelong sleep behaviours or inherited vulnerability. A computational modelling

study simulated the long-term effects of habitual sleep patterns on responses to acute sleep

deprivation, identifying that short sleepers were more resilient and had longer duration of waking

episodes following recovery sleep(102). This is consistent with evidence that habitual short sleepers

may be more tolerant of homeostatic sleep pressure(103). While there is no longitudinal evidence that

genetic vulnerability to insomnia or short sleep mitigates the effects of sleep loss on mood, our

findings raise the possibility that these traits may confer resilience in the context of PP.

Our analyses suggest that incorporating PRS for insomnia and short sleep alongside clinical variables

may improve the accuracy of PP risk prediction. Bipolar subtype(72, 104), use of prophylactic

medication(4, 105), perinatal psychiatric history(72), and family history(106, 107) are important

clinical predictors of PP. Whilst data was not available for all of these factors in our cohort, future

predictive models should aim to integrate these clinical variables with PRS information. This

combined approach could improve risk stratification and represents a promising step toward more

personalised preventive strategies for women at high risk of PP. At present, however, PRS are not yet

suitable for clinical implementation due to their well-recognised limitations(108). Therefore, the

translational implication of this work lies in advancing mechanistic understanding and generating

hypotheses, rather than suggesting imminent clinical adoption.

Our MR analyses do not support a causal relationship between insomnia and PP or short sleep and

PP, indicating that the observed associations may reflect correlated genetic or environmental

vulnerabilities rather than direct causation. However, these null findings may also have been

influenced by the weak genetic instruments available and the heterogeneity among them. The

relatively low estimated SNP heritability of our traits of interest (8.2% for insomnia(53) and 11.9% for

short sleep(54)) also somewhat violates the first assumption of MR. Furthermore, it is important to

note that these sleep traits are highly polygenic, and PRS will therefore capture genetic susceptibility

much more accurately than an MR model with relatively few variants. Whilst pre-existing insomnia or

short sleep may represent a useful clinical marker in perinatal risk assessment, there is insufficient

evidence to support a causal association between genetic predisposition to these traits and PP at

present.

We did not find evidence to suggest a significant association between long sleep PRS and PP risk.

Women who experienced PP had higher PRS for long sleep than participants who did not, albeit

below the threshold for statistical significance. This is in keeping with the hypothesis that individuals

who are chronically exposed to sleep restriction prior to childbirth may be less vulnerable to PP.

Conversely, women who are biologically predisposed to habitual long sleep may be more affected by

acute perinatal sleep loss. Our study lacked sufficient power to test this theory, in part because long

sleep has a lower SNP heritability compared to the other sleep phenotypes examined(54), resulting in

reduced predictive power of its PRS. Improved GWAS of long sleep may offer the opportunity to

robustly test this notion in future studies.

Our study did not identify an association between sleep efficiency PRS and PP outcome. Reduced

sleep efficiency is often observed in actigraphy studies of individuals with BD(32) and has been

implicated in psychosis relapse(35). Recent evidence also suggests that higher sleep efficiency at

baseline predicts lower subjective sleepiness during prolonged sleep deprivation, indicating a

potential role in mood regulation(109). It is plausible that women who are able to effectively capitalise

on available sleep opportunities may experience less mood disturbance from perinatal sleep

disruption, whilst those with lower sleep efficiency may be more susceptible to psychiatric

deterioration. Study designs which integrate subjective and objective assessments of sleep efficiency

may be helpful to test this theory further in perinatal cohorts.

While we found a significant association between PRS for self-reported short sleep and PP, we did

not identify a significant association between accelerometer-derived sleep duration PRS and PP. This

may in part reflect the weak genetic correlation between self-reported and objectively measured sleep

duration observed in the discovery GWAS (r_g=0.43) (55). Observational data also show that objective

and subjective sleep measures can differ substantially in the same individual, with studies frequently

reporting low concordance between self-reported and actigraphy-derived estimates of sleep

duration(110, 111). In perinatal populations, subjective perceptions of sleep may be more closely

linked to mental health outcomes than objective measures. Several studies have found self-reported

sleep measures to be more predictive of postpartum depression than any actigraphy-derived

metric(112-115). Taken together with our results, these findings suggest that subjective perceptions

of sleep (and the genetic liability underlying them) may be more clinically relevant than objective

sleep measures in predicting vulnerability to perinatal mood disorders.

Despite our study having adequate power to detect a significant association, we found no significant

difference in BD PRS or schizophrenia PRS in women who did and who did not develop PP following

childbirth. Our results are consistent with previous work in which BD and schizophrenia PRS were

compared in parous women with BD who did and who did not develop PP(20). Psychotic symptoms

are highly prevalent in BD(116) and women with BD have a significantly increased risk of affective

psychoses in the perinatal period(117). The estimated heritability of BD lies in the region of 60-85%

and common genome-wide variants are thought to account for a significant proportion of this(17).

Given the substantial genetic contribution to phenotypic variance in BD and the high genetic

correlation between BD and schizophrenia(118), it would be plausible that PP may represent a more

severe BD phenotype and that PRS associated with psychotic illnesses could be used to improve

outcome prediction. Our negative finding may partly reflect the limited explanatory power of current

PRS for BD and schizophrenia, or it may indicate that the hypothesis that PP represents a more

severe bipolar phenotype is incorrect. Irrespective of the underlying cause, our results and existing

evidence indicate that PP cannot simply be explained by individual genetic susceptibility to psychotic

illness.

Our findings should be considered within the context of some important limitations. The retrospective

cross-sectional design of our study renders it susceptible to recall bias. Data collection primarily relied

on participant reports and clinical details could not always be validated through case notes.

Additionally, the interviews captured case-control status at a specific point in time, potentially missing

PP in subsequent pregnancies among individuals labelled as controls. Despite our evidence-based

rationale for including cases occurring up to 12 months postpartum, this broader definition inevitably

increases heterogeneity within the PP group. However, this is not a limitation unique to our

methodology, as PP is increasingly recognised as a highly heterogeneous clinical entity(119), and the

absence of standardised diagnostic criteria continues to hinder research in this area.

While our psychiatric phenotyping was comprehensive, the lack of detailed sleep data limits the

conclusions that can be drawn on the relationship between PRS, sleep phenotype and PP outcome.

Moreover, sleep-related PRS are relatively weak genetic instruments, and the extent to which they

reflect sleep behaviour, particularly in individuals with BD, remains to be clarified. In addition, all

available data were based on lifetime, retrospective measures that do not provide sufficient

granularity to evaluate potential perinatal-specific differences. Important clinical variables such as

medication use and treatment response were recorded without reference to timing in relation to

pregnancy or childbirth. The absence of longitudinal data limited our ability to assess whether specific

exposures during the perinatal period influenced risk of PP. This represents a limitation of our dataset

and highlights the need for prospective studies that capture clinical information specific to the

perinatal period. Several important potential confounding factors were also not available in our

dataset, including socioeconomic status, smoking, substance misuse and physical health

comorbidities, which are known to influence both sleep and mental health and could contribute to the

observed associations(120). Future studies incorporating richer clinical and sociodemographic data

will be essential to better disentangle these relationships.

Finally, our sample characteristics may also limit the generalisability of our findings. Women with high-

risk bipolar subtypes (BPI or SABD) were overrepresented in our study cohort compared to their

prevalence the general population(121) and our findings may not be epidemiologically representative

of all women with BD. It is important to note that our findings pertain specifically to women with BD

and should not be used to draw conclusions on the relationship between sleep PRS and PP risk in

low-risk individuals. Furthermore, the exclusive inclusion of individuals of European ancestry restricts

the generalisability of findings to other populations. Future work should investigate the role of sleep

PRS in women experiencing new-onset PP without a pre-existing history of mental illness to elucidate

this relationship further, and should include more diverse populations to improve generalisability.

PP is a complex, multifactorial illness with potentially devastating consequences. Improved prediction

of PP is essential to facilitate early intervention and mitigate its impact on maternal, child and public

health. In women with BD, prediction of perinatal episodes is particularly challenging due to clinical

heterogeneity and controversies in disease classification. We demonstrate that individuals with BD

who develop PP may have a lower genetic propensity for insomnia and short sleep. These findings

suggest that genetic vulnerability to sleep disturbance may contribute to the aetiology of PP,

highlighting the potential value of assessing individual sleep history in perinatal psychiatric risk. In

particular, our results prompt further inquiry into whether individuals with a genetic predisposition to

chronically short or restricted sleep may be more resilient to the acute sleep loss associated with

childbirth. In view of the limitations described above, however, our findings should be considered

exploratory and will require replication in larger, well-characterised perinatal cohorts. Future work

should prioritise multi-ancestry recruitment, unified definitions of PP and the use of both objective and

subjective sleep measures. Prospective studies in women with and without pre-existing mental illness

are essential to further understand the role of PRS in perinatal psychiatric risk. These efforts will not

only contribute to early intervention and improved management of conditions like PP, but will also

drive the ongoing progress of precision medicine in mental healthcare.

Acknowledgments

The authors would like to gratefully acknowledge all DNA Polymorphisms In Mental illness (DPIM) study participants for their contribution to this work. Participant recruitment was funded by the Medical Research Council (MRC grant code G1000708), the Neuroscience Research Charitable Trust, the Camden and Islington NHS Foundation Trust, a research lectureship from the Priory Hospitals and the National Institute for Health and Care Research (NIHR) Mental Health Research Network (MHRN). The study also received support from Bipolar UK (formerly the UK Manic Depression Fellowship). Genotyping was funded by the Stanley Center for Psychiatric Research at the Broad Institute. Parts of the analyses were supported by a project that has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (Grant agreement No. 948561). CP is supported by the Rosetrees Trust and the Elizabeth Garrett Anderson Hospital Charity. DS is supported by the NIHR. AM, NB and SE are supported by the NIHR University College London Hospitals Biomedical Research Centre funding scheme.

Conflicts of Interest

NM has received speaker fees from Idorsia pharmaceuticals. SE has received honoraria for educational activities from Eisai, Fidia, Lincoln, and UCB pharma. CP, OP, KK, NB, DS and AM have no conflicts of interest to declare.

Data availability

The GWAS summary statistics used for these analyses is publicly available through the original publications for the different phenotypes of interest: bipolar disorder(17), schizophrenia(64), insomnia(53), short and long sleep(54), sleep efficiency and sleep duration(55). The participant data set used for this study as well as the code for the analyses conducted is available upon request.

References

1.   Perry A, Gordon-Smith K, Jones L, Jones I. Phenomenology, Epidemiology and Aetiology of Postpartum
Psychosis: A Review. Brain Sci. 2021;11(1).
2.   Luykx JJ, Di Florio A, Bergink V. Prevention of Infanticide and Suicide in the Postpartum Period-the
Importance of Emergency Care. JAMA Psychiatry. 2019;76(12):1221-2.
3.   Spinelli M. Postpartum psychosis: a diagnosis for the DSMV. Arch Womens Ment Health.
2021;24(5):817-22.
4.   Wesseloo R, Kamperman AM, Munk-Olsen T, Pop VJ, Kushner SA, Bergink V. Risk of Postpartum
Relapse in Bipolar Disorder and Postpartum Psychosis: A Systematic Review and Meta-Analysis. Am J
Psychiatry. 2016;173(2):117-27.
5.   Rommel AS, Molenaar NM, Gilden J, Kushner SA, Westerbeek NJ, Kamperman AM, et al. Long-term
outcome of postpartum psychosis: a prospective clinical cohort study in 106 women. Int J Bipolar Disord.
2021;9(1):31.
6.   Ross LE, B.J M, M S. Sleep and perinatal mood disorders: a critical review. J Psychiatry Neurosci.
2005;30(4).
7.   Lawson A, Murphy KE, Sloan E, Uleryk E, Dalfen A. The relationship between sleep and postpartum
mental disorders: A systematic review. J Affect Disord. 2015;176:65-77.
8.   Zanini M, Castro J, Coelho FM, Bittencourt L, Bressan RA, Tufik S, et al. Do sleep abnormalities and
misaligned sleep/circadian rhythm patterns represent early clinical characteristics for developing psychosis in
high risk populations? Neurosci Biobehav Rev. 2013;37(10 Pt 2):2631-7.
9.   Harris LM, Huang X, Linthicum KP, Bryen CP, Ribeiro JD. Sleep disturbances as risk factors for suicidal
thoughts and behaviours: a meta-analysis of longitudinal studies. Sci Rep. 2020;10(1):13888.
10.   Strouse TB, Szuba MP, Baxter LR, Jr. Response to sleep deprivation in three women with postpartum
psychosis. J Clin Psychiatry. 1992;53(6):204-6.
11.   Wehr TA, Goodwin FK, Wirz-Justice A, Breitmaier J, Craig C. 48-hour sleep-wake cycles in manic-
depressive illness: naturalistic observations and sleep deprivation experiments. Arch Gen Psychiatry.
1982;39(5):559-65.
12.   Heron J, McGuinness M, Blackmore ER, Craddock N, Jones I. Early postpartum symptoms in puerperal
psychosis. BJOG. 2008;115(3):348-53.
13.   Sharma V, Mazmanian D. Sleep loss and postpartum psychosis. Bipolar Disord. 2003;5(2):98-105.
14.   Bilszta JL, Meyer D, Buist AE. Bipolar affective disorder in the postnatal period: investigating the role
of sleep. Bipolar Disord. 2010;12(5):568-78.
15.   Lewis KJS, Di Florio A, Forty L, Gordon-Smith K, Perry A, Craddock N, et al. Mania triggered by sleep
loss and risk of postpartum psychosis in women with bipolar disorder. J Affect Disord. 2018;225:624-9.
16.   Perry A, Gordon-Smith K, Lewis KJS, Di Florio A, Craddock N, Jones L, et al. Perinatal sleep disruption
and postpartum psychosis in bipolar disorder: Findings from the UK BDRN Pregnancy Study. J Affect Disord.
2024;346:21-7.
17.   Mullins N, Forstner AJ, O'Connell KS, Coombes B, Coleman JRI, Qiao Z, et al. Genome-wide association
study of more than 40,000 bipolar disorder cases provides new insights into the underlying biology. Nat Genet.
2021;53(6):817-29.
18.   Madrid-Valero JJ, Gregory AM. Behaviour genetics and sleep: A narrative review of the last decade of
quantitative and molecular genetic research in humans. Sleep Med Rev. 2023;69:101769.
19.   Murray GK, Lin T, Austin J, McGrath JJ, Hickie IB, Wray NR. Could Polygenic Risk Scores Be Useful in
Psychiatry?: A Review. JAMA Psychiatry. 2021;78(2):210-9.
20.   Di Florio A, Mei Kay Yang J, Crawford K, Bergink V, Leonenko G, Pardiñas AF, et al. Post-partum
psychosis and its association with bipolar disorder in the UK: a case-control study using polygenic risk scores.
The Lancet Psychiatry. 2021;8(12):1045-52.
21.   Vieta E, Berk M, Schulze TG, Carvalho AF, Suppes T, Calabrese JR, et al. Bipolar disorders. Nat Rev Dis
Primers. 2018;4:18008.
22.   Wilson S, Anderson K, Baldwin D, Dijk DJ, Espie A, Espie C, et al. British Association for
Psychopharmacology consensus statement on evidence-based treatment of insomnia, parasomnias and
circadian rhythm disorders: An update. J Psychopharmacol. 2019;33(8):923-47.
23.   Sharma V, Smith A, Khan M. The relationship between duration of labour, time of delivery, and
puerperal psychosis. J Affect Disord. 2004;83(2-3):215-20.

24.   Jefferies D, Schmied V, Sheehan A, Duff M. The river of postnatal psychosis: A qualitative study of
women's experiences and meanings. Midwifery. 2021;103:103165.
25.   Engqvist I, Ferszt G, Ahlin A, Nilsson K. Psychiatric nurses' descriptions of women with postpartum
psychosis and nurses' responses--an exploratory study in Sweden. Issues Ment Health Nurs. 2009;30(1):23-30.
26.   Ritter PS, Hofler M, Wittchen HU, Lieb R, Bauer M, Pfennig A, et al. Disturbed sleep as risk factor for
the subsequent onset of bipolar disorder--Data from a 10-year prospective-longitudinal study among
adolescents and young adults. J Psychiatr Res. 2015;68:76-82.
27.   Zaks N, Velikonja T, Parvaz MA, Zinberg J, Done M, Mathalon DH, et al. Sleep Disturbance in
Individuals at Clinical High Risk for Psychosis. Schizophr Bull. 2022;48(1):111-21.
28.   Cappuccio FP, D'Elia L, Strazzullo P, Miller MA. Sleep duration and all-cause mortality: a systematic
review and meta-analysis of prospective studies. Sleep. 2010;33(5):585-92.
29.   Lewis KS, Gordon-Smith K, Forty L, Di Florio A, Craddock N, Jones L, et al. Sleep loss as a trigger of
mood episodes in bipolar disorder: individual differences based on diagnostic subtype and gender. Br J
Psychiatry. 2017;211(3):169-74.
30.   Meyer N, Joyce DW, Karr C, de Vos M, Dijk DJ, Jacobson NC, et al. The temporal dynamics of sleep
disturbance and psychopathology in psychosis: a digital sampling study. Psychol Med. 2021;52(13):1-10.
31.   Bin YS, Marshall NS, Glozier N. Sleeping at the Limits: The Changing Prevalence of Short and Long
Sleep Durations in 10 Countries. American Journal of Epidemiology. 2013;177(8):826-33.
32.   De Crescenzo F, Economou A, Sharpley AL, Gormez A, Quested DJ. Actigraphic features of bipolar
disorder: A systematic review and meta-analysis. Sleep Med Rev. 2017;33:58-69.
33.   Kaplan KA, McGlinchey EL, Soehner A, Gershon A, Talbot LS, Eidelman P, et al. Hypersomnia subtypes,
sleep and relapse in bipolar disorder. Psychol Med. 2015;45(8):1751-63.
34.   Reed DL, Sacco WP. Measuring Sleep Efficiency: What Should the Denominator Be? J Clin Sleep Med.
2016;12(2):263-6.
35.   Bagautdinova J, Mayeli A, Wilson JD, Donati FL, Colacot RM, Meyer N, et al. Sleep Abnormalities in
Different Clinical Stages of Psychosis: A Systematic Review and Meta-analysis. JAMA Psychiatry.
2023;80(3):202-10.
36.   Kline C. Sleep Duration. In: Gellman MD, Turner JR, editors. Encyclopedia of Behavioral Medicine. New
York, NY: Springer New York; 2013. p. 1808-10.
37.   Chung J, Kim Y-C, Jeong J-H. Bipolar Disorder, Circadian Rhythm and Clock Genes. Clin
Psychopharmacol Neurosci. 2024;22(2):211-21.
38.   Short MA, Booth SA, Omar O, Ostlundh L, Arora T. The relationship between sleep duration and mood
in adolescents: A systematic review and meta-analysis. Sleep Med Rev. 2020;52:101311.
39.   Barber KE, Rackoff GN, Newman MG. Day-to-day directional relationships between sleep duration and
negative affect. Journal of Psychosomatic Research. 2023;172:111437.
40.   Lee SA, Mukherjee D, Rush J, Lee S, Almeida DM. Too little or too much: nonlinear relationship
between sleep duration and daily affective well-being in depressed adults. BMC Psychiatry. 2024;24(1):323.
41.   World Health Organization. ICD-10 : international statistical classification of diseases and related
health problems : tenth revision. 2nd ed. Geneva: World Health Organization; 2004.
42.   NHS Blood and Transplant. Clinical and research: Innovation focused on improving patient outcomes
2024 [Available from: https://www.nhsbt.nhs.uk/research-and-development/.
43.   Spitzer RL, Endicott J, Robins E. Research Diagnostic Criteria: Rationale and Reliability. Archives of
General Psychiatry. 1978;35(6):773-82.
44.   McGuffin P, Farmer A, Harvey I. A Polydiagnostic Application of Operational Criteria in Studies of
Psychotic Illness: Development and Reliability of the OPCRIT System. Archives of General Psychiatry.
1991;48(8):764-70.
45.   Castro VM, Minnier J, Murphy SN, Kohane I, Churchill SE, Gainer V, et al. Validation of electronic
health record phenotyping of bipolar disorder cases and controls. Am J Psychiatry. 2015;172(4):363-72.
46.   American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition,
Text Revision (DSM-5-TR) 2022.
47.   World Health Organization. ICD-11: International Statistical Classification of Diseases and Related
Health Problems (11th ed.). Geneva: World Health Organization; 2019.
48.   National Collaborating Centre for Mental Health. The Perinatal Mental Health Care Pathways. 2018.
49.   Royal College of Psychiatrists. College Report CR238- – Infant and early childhood mental health: the
case for action. 2023.
50.   Jones I, Smith S. Puerperal psychosis: identifying and caring for women at risk. Advances in Psychiatric
Treatment. 2009;15(6):411-8.

51.   Langan Martin J, McLean G, Cantwell R, Smith DJ. Admission to psychiatric hospital in the early and
late postpartum periods: Scottish national linkage study. BMJ Open. 2016;6(1):e008758.

52.   Brockington I. Late onset postpartum psychoses. Arch Womens Ment Health. 2017;20(1):87-92.

53.   Watanabe K, Jansen PR, Savage JE, Nandakumar P, Wang X, Hinds DA, et al. Genome-wide meta-
analysis of insomnia prioritizes genes associated with metabolic and psychiatric pathways. Nat Genet.
2022;54(8):1125-32.

54.   Austin-Zimmerman I, Levey DF, Giannakopoulou O, Deak JD, Galimberti M, Adhikari K, et al. Genome-
wide association studies and cross-population meta-analyses investigating short and long sleep duration. Nat
Commun. 2023;14(1):6059.

55.   Jones SE, van Hees VT, Mazzotti DR, Marques-Vidal P, Sabia S, van der Spek A, et al. Genetic studies of
accelerometer-based sleep measures yield new insights into human sleep behaviour. Nat Commun.
2019;10(1):1585.

56.   McGowan NM, Goodwin GM, Bilderbeck AC, Saunders KEA. Circadian rest-activity patterns in bipolar
disorder and borderline personality disorder. Transl Psychiatry. 2019;9(1):195.

57.   Esaki Y, Obayashi K, Saeki K, Fujita K, Iwata N, Kitajima T. Association between circadian activity
rhythms and mood episode relapse in bipolar disorder: a 12-month prospective cohort study. Translational
Psychiatry. 2021;11(1):525.

58.   Carpenter JS, Crouse JJ, Park SH, Shin M, Tonini E, Guo W, et al. Actigraphy-derived circadian rhythms,
sleep-wake patterns, and physical activity across clinical stages and pathophysiological subgroups in young
people presenting for mental health care. Journal of Psychiatric Research. 2025;186:396-406.

59.   McCarthy MJ, Gottlieb JF, Gonzalez R, McClung CA, Alloy LB, Cain S, et al. Neurobiological and
behavioral mechanisms of circadian rhythm disruption in bipolar disorder: A critical multi-disciplinary literature
review and agenda for future research from the ISBD task force on chronobiology. Bipolar Disord.
2022;24(3):232-63.

60.   Formica MJC, Fuller-Tyszkiewicz M, Reininghaus U, Kempton M, Delespaul P, de Haan L, et al.
Associations between disturbed sleep and attenuated psychotic experiences in people at clinical high risk for
psychosis. Psychol Med. 2024;54(9):2254-63.

61.   Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-generation PLINK: rising to the
challenge of larger and richer datasets. GigaScience. 2015;4(1).

62.   McCarthy S, Das S, Kretzschmar W, Delaneau O, Wood AR, Teumer A, et al. A reference panel of
64,976 haplotypes for genotype imputation. Nat Genet. 2016;48(10):1279-83.

63.   Grigoroiu-Serbanescu M, van der Veen T, Bigdeli T, Herms S, Diaconu CC, Neagu AI, et al.
Schizophrenia polygenic risk scores, clinical variables and genetic pathways as predictors of phenotypic traits of
bipolar I disorder. J Affect Disord. 2024;356:507-18.

64.   Trubetskoy V, Pardiñas AF, Qi T, Panagiotaropoulou G, Awasthi S, Bigdeli TB, et al. Mapping genomic
loci implicates genes and synaptic biology in schizophrenia. Nature. 2022;604(7906):502-8.

65.   Ge T, Chen C-Y, Ni Y, Feng Y-CA, Smoller JW. Polygenic prediction via Bayesian regression and
continuous shrinkage priors. Nature Communications. 2019;10(1):1776.

66.   Durbin RM, Altshuler D, Durbin RM, Abecasis GR, Bentley DR, Chakravarti A, et al. A map of human
genome variation from population-scale sequencing. Nature. 2010;467(7319):1061-73.

67.   R Core Team. RStudio: Integrated Development for R. RStudio, PBC, Boston, MA2024.

68.   Dunn OJ. Multiple Comparisons among Means. Journal of the American Statistical Association.
1961;56(293):52-64.

69.   Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez JC, et al. pROC: an open-source package for R
and S+ to analyze and compare ROC curves. BMC Bioinformatics. 2011;12:77.

70.   Carpenter J, Bithell J. Bootstrap confidence intervals: when, which, what? A practical guide for
medical statisticians. Stat Med. 2000;19(9):1141-64.

71.   DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated
receiver operating characteristic curves: a nonparametric approach. Biometrics. 1988;44(3):837-45.

72.   Di Florio A, Gordon-Smith K, Forty L, Kosorok MR, Fraser C, Perry A, et al. Stratification of the risk of
bipolar disorder recurrences in pregnancy and postpartum. Br J Psychiatry. 2018;213(3):542-7.

73.   Palla L, Dudbridge F. A Fast Method that Uses Polygenic Scores to Estimate the Variance Explained by
Genome-wide Marker Panels and the Proportion of Variants Affecting a Trait. Am J Hum Genet.
2015;97(2):250-9.

74.   Davey Smith G, Hemani G. Mendelian randomization: genetic anchors for causal inference in
epidemiological studies. Hum Mol Genet. 2014;23(R1):R89-98.

75.   Davies NM, Holmes MV, Davey Smith G. Reading Mendelian randomisation studies: a guide, glossary,
and checklist for clinicians. BMJ. 2018;362:k601.
76.   Haycock PC, Borges MC, Burrows K, Lemaitre RN, Harrison S, Burgess S, et al. Design and quality
control of large-scale two-sample Mendelian randomization studies. Int J Epidemiol. 2023;52(5):1498-521.
77.   Skrivankova VW, Richmond RC, Woolf BAR, Davies NM, Swanson SA, VanderWeele TJ, et al.
Strengthening the reporting of observational studies in epidemiology using mendelian randomisation (STROBE-
MR): explanation and elaboration. BMJ. 2021;375:n2233.
78.   Skrivankova VW, Richmond RC, Woolf BAR, Yarmolinsky J, Davies NM, Swanson SA, et al.
Strengthening the Reporting of Observational Studies in Epidemiology Using Mendelian Randomization: The
STROBE-MR Statement. Jama. 2021;326(16):1614-21.
79.   Lewis KJS, Richards A, Karlsson R, Leonenko G, Jones SE, Jones HJ, et al. Comparison of Genetic
Liability for Sleep Traits Among Individuals With Bipolar Disorder I or II and Control Participants. JAMA
Psychiatry. 2020;77(3):303-10.
80.   Flora A, Garcia JJ, Thaller C, Zoghbi HY. The E-protein Tcf4 interacts with Math1 to regulate
differentiation of a specific subset of neuronal progenitors. Proc Natl Acad Sci U S A. 2007;104(39):15382-7.
81.   Stefansson H, Ophoff RA, Steinberg S, Andreassen OA, Cichon S, Rujescu D, et al. Common variants
conferring risk of schizophrenia. Nature. 2009;460(7256):744-7.
82.   Wray NR, Ripke S, Mattheisen M, Trzaskowski M, Byrne EM, Abdellaoui A, et al. Genome-wide
association analyses identify 44 risk variants and refine the genetic architecture of major depression. Nature
Genetics. 2018;50(5):668-81.
83.   Cai L, Bao Y, Fu X, Cao H, Baranova A, Zhang X, et al. Causal links between major depressive disorder
and insomnia: A Mendelian randomisation study. Gene. 2021;768:145271.
84.   Zhou R, Zhang C, Gan R, Yin X, Wang M, Shi B, et al. Transcriptional regulation of daily sleep amount
by TCF4-HDAC4-CREB complex in mice. Sleep. 2025;48(5).
85.   Hickey SL, Berto S, Konopka G. Chromatin Decondensation by FOXP2 Promotes Human Neuron
Maturation and Expression of Neurodevelopmental Disease Genes. Cell Reports. 2019;27(6):1699-711.e9.
86.   Salmón-Gómez G, Suárez-Pinilla P, Setién-Suero E, Martínez-Asensi C, Ayesa-Arriola R. FoxP2 and
Schizophrenia: a systematic review. J Psychiatr Res. 2025;190:205-15.
87.   Vgontzas AN, Fernandez-Mendoza J, Liao D, Bixler EO. Insomnia with objective short sleep duration:
the most biologically severe phenotype of the disorder. Sleep Med Rev. 2013;17(4):241-54.
88.   Di Florio A, Jones L, Forty L, Gordon-Smith K, Blackmore ER, Heron J, et al. Mood disorders and parity -
a clue to the aetiology of the postpartum trigger. J Affect Disord. 2014;152-154:334-9.
89.   Banks S, Dinges D. Chapter 6 - Chronic Sleep Deprivation. Fifth Edition ed: Elsevier Inc; 2011. p. 67-75.
90.   Skorucak J, Arbon EL, Dijk DJ, Achermann P. Response to chronic sleep restriction, extension, and
subsequent total sleep deprivation in humans: adaptation or preserved sleep homeostasis? Sleep. 2018;41(7).
91.   Simpson NS, Diolombi M, Scott-Sutherland J, Yang H, Bhatt V, Gautam S, et al. Repeating patterns of
sleep restriction and recovery: Do we get used to it? Brain Behav Immun. 2016;58:142-51.
92.   Kay DB, Sherriff D. How are sleep and resilience related and how can sleep resilience be harnessed to
improve psychological, biological, and social outcomes? Research Directions: Sleep Psychology. 2024;1:e12.
93.   Tkachenko O, Dinges DF. Interindividual variability in neurobehavioral response to sleep loss: A
comprehensive review. Neurosci Biobehav Rev. 2018;89:29-48.
94.   Mantua J, Brager AJ, Alger SE, Adewale F, Skeiky L, Balkin TJ, et al. Self-Reported Sleep Need,
Subjective Resilience, and Cognitive Performance Following Sleep Loss and Recovery Sleep. Psychological
Reports. 2021;124(1):210-26.
95.   Hughes JM, Ulmer CS, Hastings SN, Gierisch JM, Howard MO. Sleep, resilience, and psychological
distress in United States military Veterans. Mil Psychol. 2018;30(5):404-14.
96.   Arora T, Grey I, Östlundh L, Alamoodi A, Omar OM, Hubert Lam KB, et al. A systematic review and
meta-analysis to assess the relationship between sleep duration/quality, mental toughness and resilience
amongst healthy individuals. Sleep Med Rev. 2022;62:101593.
97.   Allan AC, Gamaldo AA, Gamaldo CE, Gunia BC, Razzak I, Ighodaro E, et al. The promotion of sleep
wellness: Resilience as a protective factor. Front Sleep. 2023;2.
98.   Cai H, Guo J, Zhou J, Lai Y. Psychological resilience mediates the association between sleep quality and
anxiety symptoms: A repeated measures study in college students. PLoS One. 2023;18(10):e0292887.
99.   Van Dongen HP, Baynard MD, Maislin G, Dinges DF. Systematic interindividual differences in
neurobehavioral impairment from sleep loss: evidence of trait-like differential vulnerability. Sleep.
2004;27(3):423-33.

100.   Rupp TL, Wesensten NJ, Balkin TJ. Trait-like vulnerability to total and partial sleep loss. Sleep.
2012;35(8):1163-72.
101.   Dennis LE, Wohl RJ, Selame LA, Goel N. Healthy Adults Display Long-Term Trait-Like Neurobehavioral
Resilience and Vulnerability to Sleep Loss. Sci Rep. 2017;7(1):14889.
102.   Piltz SH, Diniz Behn CG, Booth V. Habitual sleep duration affects recovery from acute sleep
deprivation: A modeling study. Journal of Theoretical Biology. 2020;504:110401.
103.   Aeschbach D, Postolache TT, Sher L, Matthews JR, Jackson MA, Wehr TA. Evidence from the waking
electroencephalogram that short sleepers live under higher homeostatic sleep pressure than long sleepers.
Neuroscience. 2001;102(3):493-502.
104.   Di Florio A, Forty L, Gordon-Smith K, Heron J, Jones L, Craddock N, et al. Perinatal Episodes Across the
Mood Disorder Spectrum. JAMA Psychiatry. 2013;70(2):168-75.
105.   Jairaj C, Seneviratne G, Bergink V, Sommer IE, Dazzan P. Postpartum psychosis: A proposed treatment
algorithm. J Psychopharmacol. 2023:2698811231181573.
106.   Jones I, Craddock N. Familiality of the puerperal trigger in bipolar disorder: results of a family study.
Am J Psychiatry. 2001;158(6):913-7.
107.   Jones I, Craddock N. Do puerperal psychotic episodes identify a more familial subtype of bipolar
disorder? Results of a family history study. Psychiatr Genet. 2002;12(3):177-80.
108.   Abu-El-Haija A, Reddi HV, Wand H, Rose NC, Mori M, Qian E, et al. The clinical application of polygenic
risk scores: A points to consider statement of the American College of Medical Genetics and Genomics
(ACMG). Genetics in Medicine. 2023;25(5).
109.   Pasetes LN, Goel N. Bidirectional Predictors Between Neurobehavioural Measures During Total Sleep
Deprivation and Baseline and Recovery Sleep Measures. J Sleep Res. 2025:e70031.
110.   Matthews KA, Patel SR, Pantesco EJ, Buysse DJ, Kamarck TW, Lee L, et al. Similarities and differences in
estimates of sleep duration by polysomnography, actigraphy, diary, and self-reported habitual sleep in a
community sample. Sleep Health. 2018;4(1):96-103.
111.   Benz F, Riemann D, Domschke K, Spiegelhalder K, Johann AF, Marshall NS, et al. How many hours do
you sleep? A comparison of subjective and objective sleep duration measures in a sample of insomnia patients
and good sleepers. J Sleep Res. 2023;32(2):e13802.
112.   Dorheim SK, Bondevik GT, Eberhard-Gran M, Bjorvatn B. Subjective and objective sleep among
depressed and non-depressed postnatal women. Acta Psychiatr Scand. 2009;119(2):128-36.
113.   Bei B, Milgrom J, Ericksen J, Trinder J. Subjective perception of sleep, but not its objective quality, is
associated with immediate postpartum mood disturbances in healthy women. Sleep. 2010;33(4):531-8.
114.   Park EM, Meltzer-Brody S, Stickgold R. Poor sleep maintenance and subjective sleep quality are
associated with postpartum maternal depression symptom severity. Arch Womens Ment Health.
2013;16(6):539-47.
115.   Stremler R, McMurray J, Brennenstuhl S. Self-Reported Sleep Quality and Actigraphic Measures of
Sleep in New Mothers and the Relationship to Postpartum Depressive Symptoms. Behav Sleep Med.
2020;18(3):396-405.
116.   Dunayevich E, Keck PE, Jr. Prevalence and description of psychotic features in bipolar mania. Curr
Psychiatry Rep. 2000;2(4):286-90.
117.   Jones I, Chandra PS, Dazzan P, Howard LM. Bipolar disorder, affective psychosis, and schizophrenia in
pregnancy and the post-partum period. Lancet. 2014;384(9956):1789-99.
118.   Song J, Pasman JA, Johansson V, Kuja-Halkola R, Harder A, Karlsson R, et al. Polygenic Risk Scores and
Twin Concordance for Schizophrenia and Bipolar Disorder. JAMA Psychiatry. 2024.
119.   Cohen LS, Arakelian M, Church TR, Dunk MM, Gaw ML, Yoon HE, et al. The phenomenology of
postpartum psychosis: preliminary findings from the Massachusetts General Hospital Postpartum Psychosis
Project. Molecular Psychiatry. 2025;30(6):2537-44.
120.   Papadopoulos D, Etindele Sosso FA. Socioeconomic status and sleep health: a narrative synthesis of 3
decades of empirical research. J Clin Sleep Med. 2023;19(3):605-20.
121.   Merikangas KR, Jin R, He JP, Kessler RC, Lee S, Sampson NA, et al. Prevalence and correlates of bipolar
spectrum disorder in the world mental health survey initiative. Arch Gen Psychiatry. 2011;68(3):241-51.
895

898 ## Tables and table legends

900 **Table 1: Demographic summary of parous female participants with Bipolar Disorder**

Demographics	All (n=343)	PP following childbirth (n=117)	No PP following childbirth (n=226)	p value
Age[1]	51(16)	53(14)	50(16)	0.684[3]
Bipolar Subtype:				
BPI	213	86	127	**1.387×10^{-4}**[4]
SABD	28	12	16	
BPII	37	3	34	
BPNOS	65	16	49	
Parity[2]:				
1	124	44	80	0.776[5]
≥2	219	73	146	

Notes: BPI= Bipolar I Disorder; BPII= Bipolar II Disorder; SABD= Schizoaffective Bipolar Disorder; BPNOS= Bipolar Disorder

not otherwise specified

[1] Median age (interquartile range) at recruitment

[2] Defined as the total number of pregnancies carried to a viable gestation at the point of recruitment

Significance set at p < 0.05 (significant findings shown in bold):

[3] Wilcoxon rank sum test comparing age between the two groups

[4] Pearson's Chi-squared test comparing the prevalence of high-risk Bipolar subtypes (BPI or SABD) between the two

groups

[5] Pearson's Chi-squared test of independence comparing parity between the two groups

Table 2: Logistic regressions between PRS and Postpartum Psychosis in parous women with

Bipolar Disorder

PRS	Odds Ratio (95% CI)	p value[1]	Bonferroni_p[1]
Bipolar Disorder	1.255 (0.988-1.594)	0.063	0.441
Schizophrenia	1.158 (0.922-1.454)	0.207	1.000
Insomnia	0.718 (0.575-0.897)	$\mathbf{3.50 \times 10^{-3}}$	$\mathbf{2.45 \times 10^{-2}}$
Short Sleep	0.674 (0.535-0.849)	$\mathbf{7.94 \times 10^{-4}}$	$\mathbf{5.56 \times 10^{-3}}$
Long Sleep	1.100 (0.887-1.365)	0.384	1.000
Sleep Efficiency	1.074 (0.851-1.356)	0.546	1.000
Sleep Duration	1.056 (0.846-1.317)	0.630	1.000

918 *Notes:* Logistic regression analyses performed with Postpartum Psychosis diagnosis following childbirth as the outcome

919 variable and adjusted for 3 population principal components and genotyping platform.

920 PRS= Polygenic Risk Score; Bonferroni_p= p value adjusted using the Bonferroni method to correct for multiple testing

921 [1]Significance set at $p < 0.05$ (significant findings shown in bold)

923 **Table 3: Risk of Postpartum Psychosis according to sleep PRS decile**

Variable	Risk in lowest decile (%)	Risk in highest decile (%)	Relative Risk[1] (95% CI)
Insomnia PRS	27.6	14.1	**1.96 (1.25-3.07)**
Short Sleep PRS	29.3	13.1	**2.23 (1.40-3.54)**

Notes: Risk of Postpartum Psychosis according to insomnia/ short sleep PRS decile, assuming a background risk of 20% in

individuals with bipolar disorder(4, 72)

[1] Relative risk of Postpartum Psychosis in bottom decile compared to top decile for sleep trait PRS (significant findings shown

in bold)

Table 4: Two-Sample Mendelian Randomisation studies testing causal association between
sleep traits and Postpartum Psychosis
933

Mendelian Randomisation Method	Beta	SE	P value
Insomnia[1]:			
Inverse Variance Weighted	0.384	0.849	0.651
MR Egger	-2.364	3.414	0.506
Weighted median	0.416	0.731	0.569
Simple mode	0.620	1.239	0.628
Weighted mode	0.432	0.822	0.611
Short Sleep[2]:			
Inverse Variance Weighted	-0.262	0.270	0.330
MR Egger	-0.800	0.880	0.377
Weighted median	-0.320	0.373	0.391
Simple mode	-0.095	0.580	0.872
Weighted mode	-0.393	0.491	0.435

935 [1] 11 Single Nucleotide Polymorphisms

936 [2] 17 Single Nucleotide Polymorphisms

939 ## Figures and figure legends

941 **Figure 1: Sleep trait PRS distributions in parous women with bipolar disorder**

942
943
Notes:

Comparisons of Polygenic Risk Scores (PRS) across sleep traits in individuals who did and who did not develop postpartum

psychosis following childbirth. Each plot depicts the distribution of PRS Z-scores for a sleep phenotype for each of the two

groups. Higher Z-scores indicate greater genetic predisposition for the respective sleep trait. The PRS mean value for each

group is shown in the corresponding box plot.

ns= not significant with significance set at p < 0.05

**p ≤ 0.01 with significance set at p < 0.05

***p ≤ 0.001 with significance set at p < 0.05

Figure 2: Prediction of Postpartum Psychosis using bipolar subtype alone and in combination

with PRS

Notes: Receiver Operating Characteristic (ROC) curves comparing the predictive ability of bipolar subtype alone versus bipolar

subtype combined with insomnia and short sleep PRS for Postpartum Psychosis.

The ROC curve illustrates the trade-off between sensitivity and 1-specificity, while the Area Under the Curve (AUC) quantifies

overall predictive accuracy. The grey diagonal line represents a model with no predictive power (random chance), where AUC =

0.5.

An AUC closer to 1 indicates better model performance.

970