Title

Obesity-related brain atrophy is independent of Alzheimer's disease protein pathways

Authors

Filip Morys[1,2], Lang Liu[1,2,3], Konstantin Senkevich[1,2,3,4], Ziv Gan-Or[1,2,3], Alain Dagher[1,2]

[1] The Neuro (Montreal Neurological Institute-Hospital), McGill University, Montreal, QC,
Canada

[2] Department of Neurology and neurosurgery, McGill University, Montréal, QC, Canada

[3] Department of Human Genetics, McGill University, Montréal, QC, Canada

[4] Department of Specialized Medicine, Division of Medical Genetics, McGill University Health
Centre, Montreal, QC, Canada.

Corresponding author

Filip Morys

filip.morys@mcgill.ca

3801 University Street

QC H3A 2B4, Montreal

Canada

Abstract

Background: Obesity increases the risk for Alzheimer's disease (AD) and other dementias.

Obesity causes structural brain injury, and it has been suggested that this may contribute to

the development of AD pathology. Neurodegeneration in AD results from the aggregation of

misfolded and dysfunctional tau and amyloid-β. However, it remains unknown whether

adiposity-related brain injury acts through tau and amyloid deposition or as an independent

cause of neurodegeneration.

Objective: Here, we tested whether obesity, cerebrovascular disease, and obesity-related

metabolic risk score were associated with structural brain and cognitive changes via the

same mechanisms as AD or independent of them.

Methods: We used the UK Biobank sample of over 33,000 individuals aged 64 years on

average. We tested the influence of the microtubule-associated protein tau (*MAPT)* and

apolipoprotein E (*APOE)* risk alleles involved in tau and amyloid-β synthesis, folding, and

clearance, as well as AD polygenic risk score (PRS) to define genetic risk of AD.

Specifically, we investigated whether these genetic risk factors moderated the relationship

between obesity and brain structure and cognition.

Results: We found that *MAPT* and *APOE* status and AD PRS did not moderate the

relationship between obesity and brain atrophy. We also found limited evidence for the

moderation of *MAPT* and *APOE* of the cerebrovascular disease-brain structure relationship

as well as the metabolic risk score-brain structure relationship.

Conclusions: We conclude that the mechanisms linking obesity with brain atrophy are most

likely independent of the ones governing AD-related protein deposition.

Keywords

Obesity, Alzheimer's disease, APOE, MAPT, brain structure

Introduction

Among the most significant risk factors for Alzheimer's Disease (AD) are age, family history, and obesity and its associated metabolic dysfunctions [1,2]. However, how genetic risk for AD and obesity interact to cause neurodegeneration with advancing age is not known.

AD is caused by the progressive accumulation and propagation of abnormal isoforms of amyloid-β and tau, which aggregate as plaques and neurofibrillary tangles [3]. These proteins are neurotoxic and their accumulation is associated with loss of neurons, synapses, and neuropil, which can be detected as cortical volume loss and white matter damage with magnetic resonance imaging (MRI) [4]. Longitudinal MRI studies in AD have demonstrated a spatially stereotyped and progressive pattern of cortical thinning [5].

The expression, function, and deposition of tau and amyloid-β are regulated by the genes encoding microtubule-associated protein tau (*MAPT*) and apolipoprotein E (*APOE*). The *MAPT* gene regulates the expression and function of tau protein and is represented by H1 and H2 haplotypes in European populations [6]. The H1 haplotype is an ancestral inversion at chromosome 17q21 that includes several genes besides *MAPT* [7]. H1 is a risk factor for a number of neurodegenerative diseases, including Parkinson's disease, progressive supranuclear palsy, frontotemporal dementia and corticobasal degeneration [8]. The H1 haplotype has also been associated with AD, and with a higher risk of dementia in PD [8–10].

The *APOE* gene has three alleles - ε2, ε3, and ε4 - and its product apolipoprotein E is associated with cholesterol transport and regulation of amyloid-β seeding, aggregation, and deposition [11]. The ε4 allele confers an increased risk of AD [12,13]. Additionally, the risk of AD is linked to other genes involved in neuroinflammation, endocytosis, or lipid metabolism [14]. These genes can be incorporated into a polygenic risk score (PRS), which is associated with earlier age of onset and higher annual incidence of AD.

Obesity in middle-age is associated with cerebrovascular dysfunction, grey and white matter atrophy, and cognitive decline [15,16]. Animal and human studies show that adiposity-related factors leading to brain injury are hypertension, diabetes and insulin resistance,

inflammation, and dyslipidemia, labelled together as metabolic syndrome [15–20]. In addition,

the cortical atrophy pattern associated with higher BMI in middle-age overlaps spatially with

that described in AD [21]. This suggests that chronic obesity may lead to brain injury that

predisposes people to AD.

However, whether adiposity-related brain atrophy is associated with AD-related protein

deposition remains unknown, although some evidence points towards an association. For

instance, insulin resistance can directly cause cerebrovascular dysfunction, but it also

contributes to the accumulation of amyloid-β and related neurotoxicity [22,23]. Similar dual

effects on cerebral vasculature and protein accumulation are observed in relation to

hypertension, inflammation, and dyslipidemia [15,24–30]. Thus, while some studies suggest that

obesity and AD share protein-related mechanisms [31–35], obesity might also operate through

independent pathways.

One possibility is that prolonged adiposity and metabolic syndrome damage the brain

through the amyloid/tau cascade of AD in vulnerable individuals (e.g., those carrying high-

risk variants such as *APOE4*, *MAPT* H1, or a high PRS for AD[36]. If this is the case, then

genetic risk for AD and associated protein deposition should moderate the relationship

between obesity and brain atrophy [37]. Conversely, metabolic injury and genetics may act

independently [21,38,39]. For example, adiposity-related brain damage may lower the threshold

to dementia by reducing cognitive reserve independently of amyloid or tau deposition. In this

case, obesity and genetic risk would have independent effects on brain atrophy.

There are now emerging treatments for both obesity and AD-related proteinopathy, making it

important to better understand the interaction of these two processes. Here, we examine the

role of genetic risk in modulating the relationship between obesity and brain atrophy and

cognitive function. Specifically, we look for interactions between obesity measures and

MAPT and *APOE* genotypes and AD PRS. We use the UK Biobank (UKBB) dataset with its

extensive phenotypic and genetic information to test the two mentioned mechanisms in a

sample of over 33,000 individuals. Overall, our findings mostly support an independent role

for genetic risk and adiposity.

Materials and methods

Participants

The UKBB sample is a large-scale, multi-site study conducted in the UK [40,41]. The current

research was performed under UKBB study ID 45551. Prior to all analyses we excluded

individuals with neurological disorders diagnosed before the neuroimaging timepoint and we

only included individuals who participated in the neuroimaging session. Participants gave

written informed consent and the study was approved by the North-West Multi-Centre

Research Ethics Committee. All ethical regulations relevant to human research participants

were followed. The sample size for this study was 33,769 (mean age=64 years; age range:

45-83 years, SD=9; mean BMI=26.21kg/m^2, SD=3.97; 17,210 women).

Neuroimaging data

Neuroimaging data were collected using 3T Siemens Skyra scanners at 3 sites in the UK.

Imaging   protocol   details   are   described   online   at

https://biobank.ctsu.ox.ac.uk/crystal/refer.cgi?id=2367 and in previous publications [41]. We

used T1-weighted structural images with a 0.8mm3 isotropic voxel size. Here, we used

cortical thickness and cortical surface area imaging-derived phenotypes for each parcel of

the Desikan-Killiany-Tourville atlas provided by the UKBB [42–44] as well as subcortical volume

measurements for the following regions of interest: thalamus, caudate, putamen, pallidum,

hippocampus,   amygdala,   and   nucleus   accumbens.   To   investigate   white   matter

microstructure, we used mean diffusivity and fractional anisotropy values for 48 standard

parcels from the Johns Hopkins University white matter atlas provided by UKBB [45,46]. Finally,

we used total volume of white matter hyperintensities (WMH) as a better proxy of

cerebrovascular disease than BMI. Cortical data were obtained using FreeSurfer 6.0.0,

subcortical volumes were derived using Functional Magnetic Resonance Imaging of the

Brain (FMRIB)'s Integrated Registration and Segmentation Tool (FIRST), FA and MD were

obtained using custom-made scripts, while WMH was obtained using Brain Intensity

AbNormality Classification Algorithm (BIANCA) [47–50]. Automated quality control was

performed by the UKBB initiative.

Genotyping data quality control and haplotype analysis

We analysed 487,410 samples from the 2019 release of the UK Biobank (UKB), applying

standard genotyping quality control measures as described by Bycroft et al. [51]. We excluded

individuals of non-European ancestry, determined through both self-reported data and

genetic principal component analysis. This was done to remove confounding effects due to

population stratification. Further exclusions were made for individuals with relatedness closer

than that of first cousins. Data from participants whose self-reported sex did not correspond

to their genetic sex were removed from the dataset. In addition, SNPs with minor allele

frequency (MAF) < 1%, missing rate > 1%, imputation quality INFO score < 0.3, and

significant deviation from Hardy-Weinberg equilibrium (HWE) with $P < 1 \times 10^{-10}$ were

excluded. MAPT haplotypes were based on genotypes of rs1052553 (H1: A; H2: G). *APOE*

alleles were based on combinations of genotypes derived from rs429358 and rs7412 (ε2:

rs429358 T/ rs7412 T; ε3: T/C; ε4: C/C). There were 8,666 *APOE ε4* carriers (heterozygous

and homozygous) and 20,337 *MAPT* H1 carriers (heterozygous and homozygous).

Alzheimer's disease polygenic risk score calculation

The AD PRS was calculated using the posterior effect size of SNPs inferred from the latest

GWAS summary statistics [14]. The posterior effect size was calculated by PRS-CS, a

software that applies a Bayesian regression framework, continuous shrinkage (CS) priors

and the linkage disequilibrium reference panel from 1000 Genome Projects European

samples on SNP effect sizes [52].

Obesity-related variables

BMI was used as a main indicator of obesity. In addition, we investigated the following

measures of obesity-related metabolic dysfunction: serum C-reactive protein, non-fasting

plasma glucose, plasma glycated haemoglobin A1c (HbA1c), systolic and diastolic blood

pressure, plasma triglycerides, total cholesterol, high density lipoprotein and low density

lipoprotein, as well as waist-to-hip ratio and body fat percentage. Blood sampling and

handling are described in [53]. All blood parameters were collected at a baseline timepoint,

which took place on average 8 years prior to the brain imaging timepoint. Weight and body

fat percentage were measured with the Tanita BC418ma bioimpedance device (Tanita,

Tokyo, Japan) during the imaging visit. Metabolic variables were selected based on our

previous work [16].

Cognitive measures

To test the influence of AD genetic risk and obesity on cognition, we used the following

cognitive measures provided by the UK Biobank [54,55]: 1) working memory (digit span task); 2)

fluid intelligence (set of reasoning tasks); 3) executive function (tower rearranging test); 4)

prospective memory (number of times an intention was forgotten on a prospective memory

task); 5) visuospatial memory (pairs matching task); 6) reaction time (button press task).

Because tests of visuospatial memory and reaction time were administered more than once,

we used an average value of all trials. For prospective memory, visuospatial memory, and

reaction time, higher scores indicate worse performance. Cognitive measures were collected

during the imaging visit.

Statistical analyses

Statistical analysis was performed in R (v. 4.3.1). We began by investigating the

associations between genotypes of interest (*APOE ε4*, *MAPT* H1, AD PRS) and brain and

cognitive measures of interest - cortical thickness, cortical surface area, subcortical volumes,

white matter FA and MD, and cognition. We used general linear model analysis with genetic

measures as predictors and brain or cognitive measures as outcome variables. Next, to

investigate the relationship between obesity and neurocognitive phenotypes, we used the

GLM analysis with BMI as a predictor variable and brain/cognitive measures as outcome

variables. To investigate whether the relationship between BMI, brain structure and cognition

differed as a function of *MAPT/APOE* genotype or AD PRS, we introduced an interaction

term between BMI and genotype/PRS in the GLM. Next, we performed similar GLM

analyses with WMH volume as a predictor variable instead of BMI. This was only done for

grey matter measures and cognition as outcome variables due to high collinearity between

WMH and white matter FA and MD. Finally, to capture the combined effect of multiple

obesity-related variables, we first ran a principal component analysis with *varimax* rotation to

identify a latent variable that encompasses all obesity-related variables in this study. We

used scores of the first rotated component (explaining 24% of total variance in all obesity-

related variables; Supplemental Table 1, Supplemental Figure 1) representing overall

cardiometabolic risk as a predictor variable in GLMs analogous to the previous analyses with

BMI and WMH volume. According to the best available practices, GLM analyses included

the following covariates: age, sex, age^2, age*sex, age^2*sex, imaging site, time difference

between imaging acquisition of the first participant and each remaining participant, time

difference squared (not for cognitive analysis), as well as genotype measurement batch and

first 15 genetic principal components [56]. These covariates were included to control for

potential confounding effects of demographics, imaging protocols, and genetic ancestry. Sex

differences exist in how *APOE* can affect the brain and AD risk [57,58]. We therefore repeated

the analyses investigating the interaction of genotype and obesity-related variables on brain

phenotypes after sex disaggregation (i.e. separately for females and males). We corrected

for multiple comparisons using false discovery rate separately for each analysis modality

(e.g. cognition or cortical thickness) [59]. Scripts used to analyse the data for this study are

available at https://github.com/FilipMorys/MAPT_ApoE.

Results

APOE ε4 status, MAPT H1 haplotype and AD PRS associations with neurocognitive

phenotypes

We found significant widespread differences in multiple brain and cognitive measures

between *APOE* ε4 carriers and non-carriers, as well as individuals with H1 vs. H2

haplotypes. *APOE ε4* carriers showed higher cortical surface area in frontal and occipital

brain areas and lower cortical thickness predominantly in the temporal and frontal brain

areas, as well as lower volumes of the bilateral thalamus and hippocampus (Figure 1a-b,

Supplemental Tables 2-3). They also showed lower FA in posterior brain regions and higher

MD in anterior brain regions (Figure 2a-b, Supplemental Table 4). *APOE* ε4 carriers also had

poorer visuospatial memory, prospective memory, fluid intelligence, working memory, and

executive function (Supplemental Table 5).

Individuals with *MAPT* H1 haplotype showed higher surface area globally as well as lower

cortical thickness in the frontal, temporal, and occipital lobes (Figure 1c, Supplemental Table

6). H1 haplotype was also associated with lower volumes of the bilateral putamen and

nucleus accumbens (Figure 1d, Supplemental Table 7), as well as higher FA and lower MD

throughout the white matter (Supplemental Table 8, Figure 2c-d). Individuals with *MAPT* H1

haplotype had better working memory (Supplemental Table 9). Finally, AD PRS was not

significantly associated with any brain phenotypes (Supplemental Tables 10-12), but was

associated with worse prospective memory, fluid intelligence, and executive function

(Supplemental Table 13).

Figure 1 Associations between genotypes and grey matter phenotypes; **a/b** - *APOE* ε4 carriers vs. non-carriers;
c/d - MAPT H1 vs H2 haplotype.

Figure 2 Associations between genotypes and grey matter phenotypes; **a** - fractional anisotropy in *APOE* ε4 carriers vs. non-carriers; **b** - mean diffusivity in *APOE* ε4 carriers vs. non-carriers; **c** - fractional anisotropy in *MAPT* H1 vs H2 haplotype; **d** - mean diffusivity in *MAPT* H1 vs H2 haplotype.

Associations between BMI, genotype, and grey matter

We found significant negative relationships between BMI and cortical thickness in the bilateral fronto-temporal and cingulate brain regions, as well as positive relationships in the occipital, parietal, and dorsal frontal regions (Figure 3a, Supplemental Table 14). In terms of

cortical surface area, BMI showed mainly negative associations in the temporal, occipital,

parietal, and frontal brain regions (Figure 3a, Supplemental Table 14). Subcortical volumes

were negatively associated with BMI with the exception of the left amygdala, which showed

a positive association with BMI and right amygdala, which showed no association (Figure 3b,

Supplemental Table 15). *MAPT* genotype, *APOE* genotype, and AD PRS did not moderate

the relationship between BMI and grey matter phenotypes (Figure 3c-f, Supplemental Tables

16-21), suggesting that the effects of BMI and AD genetic risk on the brain are independent.

Figure 3 Associations between body mass index (BMI) and grey matter phenotypes; **a** Relationship between BMI

and cortical thickness (CT) and surface area (SA); **b** relationship between BMI and subcortical volumes; **c-f**

effects of genotypes on cortical thickness and surface area in regions significantly related to BMI.

Associations between WMH volume, genotype, and grey matter

WMH volume was associated with widespread greater surface area and reduced cortical

thickness (Supplemental Figure 2a, Supplemental Table 22). WMH volume was also

significantly negatively associated with subcortical volumes except the caudate nucleus,

which showed a positive association (Supplemental Figure 2b, Supplemental Table 23) [60].

We found that *APOE* genotype moderated the relationship between WMH volume and

bilateral hippocampal volume. Here, the negative slope of the relationship between WMH

volume and hippocampal volume was steeper for *APOE* ε4 carriers (Supplemental Figure

2c). We also found a moderating effect of *MAPT* genotype on the relationship between WMH

volume and pallidum volume. The relationship between WMH load and pallidum volume was

positive for individuals with H1 haplotype, and negative for individuals with H2 haplotype

(Supplemental Figure 2d). No other significant moderating effects were found (Supplemental

Tables 24-29).

Associations between metabolic risk score, genotype, and grey matter

The first rotated component from PCA analysis representing an obesity metabolic risk score

was negatively associated with cortical surface area in most brain regions, negatively related

to cortical thickness in fronto-temporal and cingulate brain regions, and positively related to

cortical thickness in occipital, parietal, and dorsal frontal regions (Supplemental Figure 3a,

Supplemental Table 30). Metabolic risk score was also associated with lower volume of the

bilateral thalamus, left putamen, bilateral pallidum, and bilateral nucleus accumbens, as well

as higher volume of the left amygdala (Supplemental Figure 3b, Supplemental Table 31).

We found significant moderating effects of the *APOE* genotype on the relationship between

this component and cortical surface area, in which the slope of the relationship between

surface area and metabolic risk score was positive for *APOE* ε4 carriers and negative for

APOE ε4 non-carriers (Supplemental Figure 3c-d). No other effects were found

(Supplemental Tables 32-37).

Associations between BMI, genotype, and white matter

BMI showed widespread negative associations with FA in most brain regions - e.g., fornix,

corona radiata, corpus callosum or cerebellar peduncles - with only a few positive

associations with FA within the corticospinal tracts, pontine crossing tract, middle cerebellar

peduncle, right posterior limb of internal capsule, and right cingulum hippocampus (Figure

4a, Supplemental Table 38). BMI showed positive associations with MD, most notably within

the superior fronto-occipital fasciculi, and anterior corona radiata, as well as negative

associations within the middle cerebellar peduncle, pontine crossing tract, and corticospinal

tracts (Figure 4b, Supplemental Table 38). Overall, BMI-associated FA and MD changes

overlapped, but showed opposite directions of associations. We found no moderating effects

of genotype on the relationship between BMI and white matter microstructure measures

(Figure 4c-4d; Supplemental Tables 39-41).

Figure 4 Associations between body mass index (BMI) and white matter microstructure; **a** Relationship between

BMI and fractional anisotropy (FA); **b** relationship between BMI and mean diffusivity (MD); **c-f** effects of

genotypes on fractional anisotropy and mean diffusivity in regions significantly related to BMI.

Associations between metabolic risk score, genotype, and white matter

Associations between metabolic risk score and white matter diffusivity measures mimicked

the ones between BMI and white matter. Overall, more widespread negative associations

were observed between the metabolic risk score and FA, with more widespread positive

associations with MD (Supplemental Figure 4, Supplemental Table 42). No moderating

effects of genotype were observed (Supplemental Tables 43-45).

Associations between BMI, genotype, and cognition

Higher BMI was associated with longer reaction time, poorer prospective memory, lower fluid

intelligence, and lower working memory (Supplemental Table 46). We did not observe any

moderating effects of genotype on the associations between BMI and cognitive functions

(Supplemental Tables 47-49).

Associations between WMH, genotype, and cognition

Individuals with higher WMH volume had a longer reaction time, poorer prospective memory,

lower fluid intelligence, poorer working memory, as well as lower executive functions

(Supplemental Table 50). The relationship between WMH and prospective memory was

moderated by *APOE* genotype. Here, ε4 carriers show a steeper negative relationship

between WMH load and prospective memory (Supplemental Figure 5). No other significant

associations were observed (Supplemental Tables 51-53).

Associations between metabolic risk score, genotype, and cognition

Overall metabolic risk score was associated with lower fluid intelligence and working

memory (Supplemental Table 54). In individuals with H1 haplotype we did not observe a

significant relationship between the first principal component and visuospatial memory, while

in individuals with H2 haplotype this relationship was negative (Supplemental Figure 6).

Remaining associations were not statistically significant (Supplemental Tables 55-57).

Sex-disaggregated analyses

After sex disaggregation, we did not find any significant interactions between obesity-related

variables and any investigated genetic factors in males (Supplemental Tables 58-81). In

females, we found a significant interaction between APOE genotype and WMH on cortical

thickness and volume of the right caudate (Supplemental Figure 7), where APOE ε4 carriers

with higher WMH load exhibited lower cortical thickness and caudate volume. We also

observed an interaction effect indicating that females with higher BMI and APOE ε4 allele

showed lower volume of the left nucleus accumbens (Supplemental Tables 82-105).

Discussion

Obesity causes brain atrophy and increases the risk for AD [16,61]. To investigate whether the

mechanisms governing AD and obesity-related brain changes act in synergy, we tested

whether genetic risk for AD moderates the relationship between obesity and brain atrophy.

We utilised three measures of obesity: BMI, a comprehensive metabolic risk score, and

WMH volume as a direct measure of cerebrovascular disease. We first investigated how

genes involved in tau and amyloid-β expression and clearance (*MAPT* and *APOE*) as well as

AD PRS affect brain structure and cognition. We then examined whether these genetic

factors moderate the effect of BMI, metabolic risk score, and WMH volume on brain and

cognition. We showed strong effects of *MAPT* and *APOE* genes as well as WMH, obesity

and metabolic syndrome   measures on all investigated neurocognitive phenotypes.

However, there was limited evidence for an influence of AD-related genetic risk on obesity-

brain structure/cognition associations. This argues that obesity itself, as measured by BMI,

might cause brain injury independent of AD pathology.

On the other hand, we did find some evidence of AD genetic risk moderation on the

associations of WMH and metabolic risk scores with brain structure. Specifically, we found

that *APOE* ε4 gene carriers displayed a more negative relationship between WMH volume

and hippocampal volume. We also found that *APOE* ε4 status moderated the relationship

between overall metabolic risk score and cortical surface area, as well as WMH volume and

prospective memory. Finally, *MAPT* haplotype only moderated the relationship between

overall metabolic risk score and visuospatial memory.

Effects of APOE ε4, MAPT H1, and AD PRS on brain structure and cognition

In line with previous studies, we found that *MAPT* H1 haplotype and *APOE* ε4 genotype

were strongly associated with all investigated domains of brain structure and cognition. Most

notably, *APOE* ε4 was associated with lower cortical thickness and hippocampal and

thalamic volume, which are hallmarks of AD [62,63]. At the same time, *MAPT* H1 haplotype was

associated with higher surface area globally, as well as decreased cortical thickness. The

positive association with cortical surface area is in line with recent literature on *MAPT*

inversion Given that cortical surface area is determined genetically at an early age, and

MAPT expression seems to be greatest during the prenatal period [64,65], it has been

suggested that the increase in cortical surface area reflects a neurodevelopmental effect of

the H1 haplotype. *MAPT* H1 was also associated with denser white matter microstructure

and better working memory in our healthy sample. Overall, these effects of *MAPT* H1 could

represent a neurodevelopmental influence that may account for one of the mechanisms by

which it has been selected in the course of evolution. Conversely, *MAPT* H1 was associated

with reduced cortical thickness, which perhaps accounts for the neurodegenerative risk.

Finally, AD PRS was associated with slight reductions in cognitive function but not any of the

investigated brain measures, suggesting that any neuroanatomical effects might be too small

to be found in this population.

Effects of obesity, metabolism, and cerebrovascular disease on brain structure and cognition

Obesity was associated with widespread cortical and subcortical atrophy and disruptions in

white matter integrity. We found a typical fronto-temporal cortical thinning signature of

obesity and metabolic risk score. This is in line with previous studies in this and other

samples showing that obesity causes brain atrophy in areas similar to the ones affected in

AD [16,21,66,67]. Lower subcortical volumes with BMI are also in line with previous reports [16]. Our

findings of altered white matter microstructure are also consistent with the literature [68–70].

The widespread lower cortical surface area with obesity is somewhat novel. We also show

that WMH volume, a measure of chronic ischemic damage, is associated with lower cortical

thickness and volume of subcortical structures, supporting the fact that cerebrovascular

dysfunction in obesity leads to grey matter injury [71].

Relative absence of moderating effects of AD-related genes

Our investigation of moderating effects of AD-related genes on BMI-related brain atrophy

revealed that obesity mostly affects brain structure independent of AD pathophysiology. In

other words, genes affecting the deposition of tau and amyloid-β, as well as AD PRS, did not

alter how obesity is associated with all tested grey and white matter phenotypes, as well as

cognition. Overall, our findings suggest that obesity-related brain atrophy and cognitive

decline are not the result of AD-like pathology.

While some studies using positron emission tomography have reported associations

between amyloid-β pathology and obesity measures [34,35,72], these findings may be influenced

by sample characteristics (e.g., individuals with mild cognitive impairment) or limited sample

sizes. Our results align with our previous work showing no correlation between obesity-

related cortical thickness changes and whole brain patterns of amyloid-β or tau deposition [21].

The apparently contradictory findings might reflect different temporal stages in disease

progression. While MAPT and APOE regulate protein homeostasis continuously, their

variants would only moderate obesity's effects on brain structure if obesity was acting

through protein-related mechanisms. The lack of such moderation in our study suggests that

obesity initially causes brain changes through independent pathways, even though obesity-

related damage might eventually create conditions that promote protein accumulation in later

disease stages. This explains why studies conducted at different disease timepoints (i.e. in

individuals with mild cognitive impairment or AD) might show varying relationships between

obesity and protein pathology - later associations could reflect downstream consequences

rather than the initial mechanism of action of obesity.

Nonetheless, we did find some limited moderating effects of *APOE* and *MAPT* genotypes on

some obesity-brain relationships. For example, all *APOE* ε4 carriers have a more negative

relationship between WMH volume and hippocampal volume. This is in line with literature

showing lower hippocampal volumes in *APOE* ε4 carriers and suggests that cerebrovascular

disease might potentiate this relationship [62,63]. Similarly, females with higher WMH volume

and APOE ε4 allele exhibited lower cortical thickness throughout the brain, which

corroborates previous findings reporting that this allele might be more influential in brain

pathology in females vs males [73]. Interestingly and opposed to the relationship with BMI, we

found that *APOE* ε4 status moderates the relationship between metabolic risk score and

cortical surface area. Conceivably, metabolic derangements associated with obesity affect

the brain differently than BMI and this relationship seems to be moderated by *APOE*

genotype. The *APOE* gene controls not only the processing and clearance of amyloid-β, but

also cholesterol transport or neuroinflammation, amongst others [13,74]. *APOE* ε4 can also lead

to vascular dysfunction and neuroinflammation through non-amyloid dependent pathways [13].

As such, this could mean that the moderation of *APOE* on the metabolic risk score-brain

structure relationship is due to the functions of apolipoprotein E unrelated to amyloid-β

deposition, especially given that the metabolic risk score captures factors such as

hypercholesterolemia and systemic inflammation. There is also a possibility that obesity-

associated metabolic factors, such as e.g. dyslipidemia, can lead to amyloid pathology [29,30].

Conclusions

Given that most analyses performed in this study failed to show a moderating effect of AD

genetic risk on the relationship between BMI, metabolic syndrome, or WMH volume and the

brain, we conclude that it is likely that the effects of obesity on the brain are mostly

cerebrovascular in nature rather than directly involving tau and amyloid accumulation. How

can obesity lead to AD in this scenario? It is possible that some brain regions (such as the

medial temporal lobe) are inherently more vulnerable to damage. These brain regions are

affected by obesity, and in individuals with a high risk for AD, this could be enough to

facilitate or potentiate the AD-related pathology and lead to the disease. Indeed, the one

mediating effect of genetics here was one of *APOE* ε4 genotype on the relationship between

cerebrovascular damage and hippocampal atrophy.

The findings of this study should be considered within the context of its limitations. The

cross-sectional nature of the dataset prevents us from making definitive conclusions about

causality and directionality of associations. Further, the genes investigated here are not

direct measures of protein deposition. As such, some of our results are not necessarily

indicative of the associations of obesity-related variables with amyloid-β deposition. Finally,

we used BMI as a measure of obesity; this approach has sometimes been criticised as BMI

does not directly measure the percentage or distribution of body fat and is a suboptimal

biomarker of obesity risk on an individual level [75,76]. However, it performs well at population

levels [77–79]. Nevertheless, by calculating metabolic risk score, we used more direct indicators

of obesity risk to validate our findings. Future studies should explore links between obesity

and AD using comprehensive longitudinal designs with wider age ranges, different imaging

modalities (e.g., positron emission tomography), as well as a broader range of genetic,

behavioural, lifestyle, and environmental risk factors for AD. In addition, given different

mechanisms and associations between AD-risk genotypes and brain phenotypes for

different ethnicities [80,81], it is crucial that future research addresses those aspects [82,83]. Prior

to exclusions, our sample included only 2.6% percent of individuals of other ethnicities,

which was not enough to conduct desired, disaggregated analyses.

Overall, this study fails to support the hypothesis that obesity causes neurodegeneration

through the deposition of dysfunctional AD proteins. This opens avenues to investigate

alternative pathways, potentially leading to new therapeutic targets. Clinically, the findings

highlight the need for interventions that focus on weight management and correcting

metabolic abnormalities, rather than solely targeting protein deposition, to better preserve

brain health in obese individuals at risk for AD and other dementias.

Author contributions

FM, ZGO, and AD conceptualised the study, FM, LL, KS analysed the data, AD and ZGO supervised the work, FM and AD drafted the manuscript, all authors edited the final version of the manuscript.

Acknowledgments

This research used the NeuroHub infrastructure and was undertaken thanks in part to funding from the Canada First Research Excellence Fund, awarded through the Healthy Brains, Healthy Lives initiative at McGill University. This research was enabled in part by support provided by Calcul Québec and the Digital Research Alliance of Canada. This research has been conducted using the UK Biobank Resource under Application Number 45551.

Statements and declarations

Funding statement

AD was funded by the Canadian Institutes of Health Research. Z.G.O. is supported by the Fonds de recherche du Québec—Santé (FRQS) Chercheurs-boursiers award and is a William Dawson Scholar.

Ethical considerations

The UK Biobank study was approved by the North-West Multi-Centre Research Ethics Committee. All ethical regulations relevant to human research participants were followed.

Consent to participate

All participants gave their written informed consent

Consent for publication

Not applicable

Declaration of conflicting interests

470   *Data and code availability*

471   Data used in this study are available upon request from the UK Biobank. Code used to

472   analyse the dataset is deposited at https://github.com/FilipMorys/MAPT_ApoE.

References

1. Silva MVF, Loures C de MG, Alves LCV, et al. Alzheimer's disease: risk factors and
potentially protective measures. *J Biomed Sci* 2019; 26: 33.

2. Yu J-T, Xu W, Tan C-C, et al. Evidence-based prevention of Alzheimer's disease:
systematic review and meta-analysis of 243 observational prospective studies and 153
randomised controlled trials. *J Neurol Neurosurg Psychiatry* 2020; 91: 1201–1209.

3. Busche MA, Hyman BT. Synergy between amyloid-β and tau in Alzheimer's disease.
Nat Neurosci 2020; 23: 1183–1193.

4. Bayram E, Caldwell JZK, Banks SJ. Current understanding of magnetic resonance
imaging biomarkers and memory in Alzheimer's disease. *Alzheimers Dement (N Y)*
2018; 4: 395–413.

5. Planche V, Manjon JV, Mansencal B, et al. Structural progression of Alzheimer's
disease over decades: the MRI staging scheme. *Brain Commun* 2022; 4: fcac109.

6. Spillantini MG, Goedert M. Tau pathology and neurodegeneration. *Lancet Neurol* 2013;
12: 609–622.

7. Zody MC, Jiang Z, Fung H-C, et al. Evolutionary toggling of the MAPT 17q21.31
inversion region. *Nat Genet* 2008; 40: 1076–1083.

8. Sánchez-Juan P, Moreno S, de Rojas I, et al. The MAPT H1 haplotype is a risk factor
for Alzheimer's disease in APOE ε4 non-carriers. *Front Aging Neurosci* 2019; 11: 327.

9. Setó-Salvia N, Clarimón J, Pagonabarraga J, et al. Dementia risk in Parkinson disease:
disentangling the role of MAPT haplotypes: Disentangling the role of MAPT haplotypes.
Arch Neurol 2011; 68: 359–364.

10. Pastor P, Moreno F, Clarimón J, et al. MAPT H1 haplotype is associated with late-onset
Alzheimer's disease risk in APOE □4 noncarriers: Results from the dementia genetics
Spanish consortium. *J Alzheimers Dis* 2016; 49: 343–352.

11. Kanekiyo T, Xu H, Bu G. ApoE and Aβ in Alzheimer's disease: accidental encounters or
partners? *Neuron* 2014; 81: 740–754.

12. Sun Y-Y, Wang Z, Huang H-C. Roles of ApoE4 on the Pathogenesis in Alzheimer's
Disease and the Potential Therapeutic Approaches. *Cell Mol Neurobiol* 2023; 43: 3115–
3136.

13. Martens YA, Zhao N, Liu C-C, et al. ApoE Cascade Hypothesis in the pathogenesis of
Alzheimer's disease and related dementias. *Neuron* 2022; 110: 1304–1317.

14. Bellenguez C, Küçükali F, Jansen IE, et al. New insights into the genetic etiology of
Alzheimer's disease and related dementias. *Nat Genet* 2022; 54: 412–436.

15. García-García I, Michaud A, María ·, et al. Mechanisms linking obesity and its metabolic
comorbidities with cerebral grey and white matter changes. *Rev Endocr Metab Disord*
2022; 1: 1–11.

16. Morys F, Dadar M, Dagher A. Association Between Midlife Obesity and Its Metabolic
Consequences, Cerebrovascular Disease, and Cognitive Decline. *J Clin Endocrinol*
Metab 2021; 106: e4260–e4274.

17. Pedditizi E, Peters R, Beckett N. The risk of overweight/obesity in mid-life and late life
for the development of dementia: A systematic review and meta-analysis of longitudinal
studies. *Age Ageing* 2016; 45: 14–21.

18. Lloret A, Monllor P, Esteve D, et al. Obesity as a risk factor for Alzheimer's disease:
Implication of leptin and glutamate. *Front Neurosci* 2019; 13: 508.

19. Moroz N, Tong M, Longato L, et al. Limited Alzheimer-type neurodegeneration in
experimental obesity and type 2 diabetes mellitus. *J Alzheimers Dis* 2008; 15: 29–44.

20. Walker JM, Dixit S, Saulsberry AC, et al. Reversal of high fat diet-induced obesity
improves glucose tolerance, inflammatory response, β-amyloid accumulation and
cognitive decline in the APP/PSEN1 mouse model of Alzheimer's disease. *Neurobiol*
Dis 2017; 100: 87–98.

21. Morys F, Potvin O, Zeighami Y, et al. Obesity-Associated Neurodegeneration Pattern
Mimics Alzheimer's Disease in an Observational Cohort Study. *J Alzheimers Dis* 2023;
91: 1059–1071.

22. de la Monte SM. Contributions of brain insulin resistance and deficiency in amyloid-
related neurodegeneration in Alzheimer's disease. *Drugs* 2012; 72: 49–66.

23. Mullins RJ, Diehl TC, Chia CW, et al. Insulin Resistance as a Link between Amyloid-
Beta and Tau Pathologies in Alzheimer's Disease. *Front Aging Neurosci* 2017; 9: 118.

24. van den Kerkhof M, van der Thiel MM, Postma AA, et al. Hypertension Correlates With
Stronger Blood Flow Pulsatility in Small Perforating Cerebral Arteries Assessed With 7
Tesla Magnetic Resonance Imaging. *Hypertension* 2023; 80: 802–810.

25. Iadecola C, Gottesman RF. Neurovascular and Cognitive Dysfunction in Hypertension.
Circ Res 2019; 124: 1025–1044.

26. Kinney JW, Bemiller SM, Murtishaw AS, et al. Inflammation as a central mechanism in
Alzheimer's disease. *Alzheimers Dement* 2018; 4: 575–590.

27. Mun KT, Hinman JD. Inflammation and the Link to Vascular Brain Health: Timing Is
Brain. *Stroke* 2022; 53: 427–436.

28. Reitz C. Dyslipidemia and the risk of alzheimer's disease. *Current Atherosclerosis*
Reports 2013; 15: 307.

29. Kang SH, Yoo H, Cheon BK, et al. Distinct effects of cholesterol profile components on
amyloid and vascular burdens. *Alzheimers Res Ther* 2023; 15: 197.

30. Rudajev V, Novotny J. Cholesterol as a key player in amyloid β-mediated toxicity in
Alzheimer's disease. *Front Mol Neurosci* 2022; 15: 937056.

31. Meakin PJ, Coull BM, Tuharska Z, et al. Elevated circulating amyloid concentrations in
obesity and diabetes promote vascular dysfunction. *J Clin Invest* 2020; 130: 4104–4117.

32. Platt TL, Beckett TL, Kohler K, et al. Obesity, Diabetes, and Leptin Resistance Promote
Tau Pathology in a Mouse Model of Disease. *Neuroscience* 2016; 315: 162.

33. So SW, Fleming KM, Nixon JP, et al. Early Life Obesity Increases Neuroinflammation,
Amyloid Beta Deposition, and Cognitive Decline in a Mouse Model of Alzheimer's
Disease. *Nutrients*; 15. Epub ahead of print 27 May 2023. DOI: 10.3390/nu15112494.

34. Dolatshahi M, Commean PK, Rahmani F, et al. Alzheimer disease pathology and
neurodegeneration in midlife obesity: A pilot study. *Aging Dis* 2024; 15: 1843–1854.

35. Kim S, Yi H-A, Won KS, et al. Association between visceral adipose tissue metabolism
and Alzheimer's disease pathology. *Metabolites* 2022; 12: 258.

36. Edwards FA. A Unifying Hypothesis for Alzheimer's Disease: From Plaques to
Neurodegeneration. *Trends Neurosci* 2019; 42: 310–322.

37. Mole JP, Fasano F, Evans J, et al. Genetic risk of dementia modifies obesity effects on
white matter myelin in cognitively healthy adults. *Neurobiol Aging* 2020; 94: 298–310.

38. Pegueroles J, Pané A, Vilaplana E, et al. Obesity impacts brain metabolism and
structure independently of amyloid and tau pathology in healthy elderly. *Alzheimers*
Dement 2020; 12: e12052.

39. Cogswell PM, Lundt ES, Therneau TM, et al. Evidence against a temporal association
between cerebrovascular disease and Alzheimer's disease imaging biomarkers. *Nat*
Commun 2023; 14: 3097.

40. Sudlow C, Gallacher J, Allen N, et al. UK Biobank: An Open Access Resource for
Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old Age.
PLoS Med 2015; 12: e1001779.

41. Miller KL, Alfaro-Almagro F, Bangerter NK, et al. Multimodal population brain imaging in
the UK Biobank prospective epidemiological study. *Nat Neurosci* 2016; 19: 1523–1536.

42. Alfaro-Almagro F, Jenkinson M, Bangerter NK, et al. Image processing and Quality
Control for the first 10,000 brain imaging datasets from UK Biobank. *Neuroimage* 2018;
166: 400–424.

43. Littlejohns TJ, Holliday J, Gibson LM, et al. The UK Biobank imaging enhancement of
100,000 participants: rationale, data collection, management and future directions. *Nat*
Commun 2020; 11: 2624.

44. Klein A, Tourville J. 101 labeled brain images and a consistent human cortical labeling
protocol. *Front Neurosci*; 6. Epub ahead of print 2012. DOI: 10.3389/fnins.2012.00171.

45. Mori S, Wakana S, Nagae-Poetscher L, et al. *MRI atlas of human white matter*.
Amsterdam: Elsevier B.V., 2005.

46. Wakana S, Caprihan A, Panzenboeck MM, et al. Reproducibility of quantitative
tractography methods applied to cerebral white matter. *Neuroimage* 2007; 36: 630–644.

47. Griffanti L, Zamboni G, Khan A, et al. BIANCA (Brain Intensity AbNormality
Classification Algorithm): A new tool for automated segmentation of white matter
hyperintensities. *Neuroimage* 2016; 141: 191–205.

48. Fischl B. FreeSurfer. *Neuroimage* 2012; 62: 774–781.

49. Jenkinson M, Beckmann CF, Behrens TEJ, et al. FSL. *Neuroimage* 2012; 62: 782–790.

50. Patenaude B, Smith SM, Kennedy DN, et al. A Bayesian model of shape and
appearance for subcortical brain segmentation. *Neuroimage* 2011; 56: 907–922.

51. Bycroft C, Freeman C, Petkova D, et al. The UK Biobank resource with deep
phenotyping and genomic data. *Nature* 2018; 562: 203–209.

52. Ge T, Chen C-Y, Ni Y, et al. Polygenic prediction via Bayesian regression and
continuous shrinkage priors. *Nat Commun* 2019; 10: 1776.

53. Elliott P, Peakman TC, UK Biobank. The UK Biobank sample handling and storage
protocol for the collection, processing and archiving of human blood and urine. *Int J*
Epidemiol 2008; 37: 234–244.

54. Cullen B, Nicholl BI, Mackay DF, et al. Cognitive function and lifetime features of
depression and bipolar disorder in a large population sample: Cross-sectional study of
143,828 UK Biobank participants. *Eur Psychiatry* 2015; 30: 950–958.

55. Fawns-Ritchie C, Deary IJ. Reliability and validity of the UK Biobank cognitive tests.
medRxiv 2019; 19002204.

56. Alfaro-Almagro F, McCarthy P, Afyouni S, et al. Confound modelling in UK Biobank
brain imaging. *Neuroimage* 2021; 224: 117002.

57. O'Neal MA. Women and the risk of Alzheimer's disease. *Front Glob Womens Health*
2023; 4: 1324522.

58. Moutinho S. Women twice as likely to develop Alzheimer's disease as men - but
scientists do not know why. *Nat Med* 2025; 31: 704–707.

59. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. 1995; 57: 289–300.

60. Dadar M, Potvin O, Camicioli R, et al. Beware of white matter hyperintensities causing
systematic errors in FreeSurfer gray matter segmentations! *Hum Brain Mapp* 2021; 42:
2734–2745.

61. Nianogo RA, Rosenwohl-Mack A, Yaffe K, et al. Risk Factors Associated With
Alzheimer Disease and Related Dementias by Sex and Race and Ethnicity in the US.
JAMA Neurol 2022; 79: 584–591.

62. Pievani M, Galluzzi S, Thompson PM, et al. APOE4 is associated with greater atrophy
of the hippocampal formation in Alzheimer's disease. *Neuroimage* 2011; 55: 909–919.

63. O'Dwyer L, Lamberton F, Matura S, et al. Reduced hippocampal volume in healthy
young ApoE4 carriers: an MRI study. *PLoS One* 2012; 7: e48895.

64. Grasby KL, Jahanshad N, Painter JN, et al. The genetic architecture of the human
cerebral cortex. *Science*; 367. Epub ahead of print 20 March 2020. DOI:
10.1126/science.aay6690.

65. Wang H, Makowski C, Zhang Y, et al. Chromosomal inversion polymorphisms shape
human brain morphology. *Cell Rep* 2023; 42: 112896.

66. Hamer M, Batty GD. Association of body mass index and waist-to-hip ratio with brain
structure. *Neurology* 2019; 92: e594–e600.

67. García-García I, Michaud A, Dadar M, et al. Neuroanatomical differences in obesity:
meta-analytic findings and their validation in an independent dataset. *Int J Obes* 2019;
43: 943–951.

68. Zhang R, Beyer F, Lampe L, et al. White matter microstructural variability mediates the
relation between obesity and cognition in healthy adults. *Neuroimage* 2018; 172: 239–
249.

69.   Karlsson HK, Tuulari JJ, Hirvonen J, et al. Obesity is associated with white matter
atrophy: a combined diffusion tensor imaging and voxel-based morphometric study.
Obesity 2013; 21: 2530–2537.

70.   Daoust J, Schaffer J, Zeighami Y, et al. White matter integrity differences in obesity: A
meta-analysis of diffusion tensor imaging studies. *Neurosci Biobehav Rev* 2021; 129:
133–141.

71.   Dadar M, Zeighami Y, Yau Y, et al. White matter hyperintensities are linked to future
cognitive decline in de novo Parkinson's disease patients. *NeuroImage: Clinical* 2018;
20: 892–900.

72.   Dolatshahi M, Commean PK, Nguyen C, et al. Visceral abdominal adipose tissue and
Insulin Resistance respectively influence Alzheimer disease amyloid pathology and
neurodegeneration in midlife. *Alzheimers Dement* 2023; 19: e075841.

73.   Ungar L, Altmann A, Greicius MD. Apolipoprotein E, gender, and Alzheimer's disease:
an overlooked, but potent and promising interaction. *Brain Imaging Behav* 2014; 8: 262–
273.

74.   Mahley RW. Apolipoprotein E: cholesterol transport protein with expanding role in cell
biology. *Science* 1988; 240: 622–630.

75.   Sommer I, Teufer B, Szelag M, et al. The performance of anthropometric tools to
determine obesity: a systematic review and meta-analysis. *Sci Rep* 2020; 10: 12699.

76.   Nuttall FQ. Body mass index: Obesity, BMI, and health: A critical review. *Nutr Today*
2015; 50: 117–128.

77.   Ortega FB, Sui X, Lavie CJ, et al. Body mass index, the most widely used but also
widely criticized index: Would a criterion standard measure of total body fat be a better
predictor of cardiovascular disease mortality? *Mayo Clin Proc* 2016; 91: 443–455.

78.   Flegal KM, Shepherd JA, Looker AC, et al. Comparisons of percentage body fat, body
mass index, waist circumference, and waist-stature ratio in adults. *Am J Clin Nutr* 2009;
89: 500–508.

79.   Paniagua L, Lohsoonthorn V, Lertmaharit S, et al. Comparison of waist circumference,
body mass index, percent body fat and other measure of adiposity in identifying
cardiovascular disease risks among Thai adults. *Obes Res Clin Pract* 2008; 2: 215–223.

80.   Chin AL, Negash S, Hamilton R. Diversity and disparity in dementia: the impact of
ethnoracial differences in Alzheimer disease. *Alzheimer Dis Assoc Disord* 2011; 25:
187–195.

81.   Murrell JR, Price B, Lane KA, et al. Association of apolipoprotein E genotype and
Alzheimer disease in African Americans. *Arch Neurol* 2006; 63: 431–434.

82.   Fitipaldi H, Franks PW. Ethnic, gender and other sociodemographic biases in genome-
wide association studies for the most burdensome non-communicable diseases: 2005-
2022. *Hum Mol Genet* 2023; 32: 520–532.

83.   Peterson RE, Kuchenbaecker K, Walters RK, et al. Genome-wide association studies in
ancestrally diverse populations: Opportunities, methods, pitfalls, and recommendations.
Cell 2019; 179: 589–603.