Mild Behavioral Impairment as a Clinical Correlate of Early Neurodegeneration

*The submission is a preprint.

*The paper is under review at JAMA Network Open.

Yi Jin **Leow** (PGDip)[1], Seyed Ehsan **Saffari** (PhD)[1,4], Ashwati **Vipin** (PhD)[1], Pricilia **Tanoto** (MSc)[1], Rasyiqah **Binte Shaik Mohamed Salim** (BEng Hons)[1], Bocheng **Qiu** (MSc)[1], Zahinoor **Ismail** (MD, FRCPC)[5,6], Nagaendran **Kandiah**(FRCP)[1,2,3,4]

[1]*Dementia Research Centre (Singapore), Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore*

[2]*Neuroscience and Mental Health Programme, Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore*

[3]*National Healthcare Group, Singapore*

[4]*Duke-NUS Medical School, National University of Singapore, Singapore*

[5]*Hotchkiss Brain Institute, University of Calgary, Canada*

[6]*University of Exeter Faculty of Health and Life Sciences and NIHR Exeter Biomedical Research Centre, Exeter UK*

Corresponding Author: Associate Professor Nagaendran Kandiah

Address: Lee Kong Chian School of Medicine, Nanyang Technological University, 11 Mandalay Rd, Singapore 308232

Tel: 65 65922653

Email: nagaendran_kandiah@ntu.edu.sg

Word count: 3445 words

Importance:

Mild Behavioral Impairment(MBI), characterized by neuropsychiatric symptoms(NPS) that precede cognitive decline, may reflect early neurodegenerative changes. Identifying structural brain changes associated with MBI could enhance early risk stratification and intervention.

Objective:

To investigate the association between MBI symptoms and cortical thinning as a structural marker of neurodegeneration in a Southeast Asian cohort.

Design, Setting, and Participants:

This cross-sectional study analyzed baseline data from the Biomarkers and Cognition Study, Singapore(BIOCIS), a 5-year longitudinal study initiated in 2023 that recruits community participants across Singapore. Cross-sectional data from 1,145 participants(mean age: 62±13.5 years; 53% female) with normal cognition, subjective cognitive decline(SCD), or mild cognitive impairment(MCI) were included.

Main Outcomes and Measures:

Behavioral symptoms were assessed using the Mild Behavioral Impairment Checklist(MBI-C), which captures NPS across five domains: decreased motivation, affective dysregulation, impulse dyscontrol, social inappropriateness, and abnormal perception/false beliefs. Cortical thickness was measured using T1-weighted MRI, processed through FreeSurfer. The primary outcome was the relationship between MBI-C total and subdomain scores, and cortical thinning.

Results:

Higher MBI-C Belief scores were associated with cortical thinning in the right hemisphere(β=-0.0177;95% CI:-0.0342 to -0.0012;P=.035). Significant thinning was observed in specific regions - posterior banks of the superior temporal sulcus(β=-0.020;95% CI:-0.037 to -0.004;P=.013), fusiform gyrus(β=-0.021;95% CI:-0.037 to -0.005;P=.009), superior temporal gyrus(β=-0.020;95% CI:-0.037 to -0.002;P=.025), temporal pole(β=-0.018;95% CI:-0.034 to -0.003;P=.021), and transverse temporal gyrus(β=-0.020;95% CI:-0.036 to -0.004;P=.014). After false discovery rate(FDR) correction, the associations in the posterior banks of the superior temporal sulcus, fusiform gyrus, superior temporal gyrus, and transverse temporal gyrus remained significant(FDR P=.042–.045).

In addition to these temporal regions, higher MBI-C Belief scores were significantly associated with cortical thinning in the right postcentral gyrus(β=-0.020;P=.012) and right supramarginal gyrus(β=-0.021;P=.013), which remained significant post-FDR correction(FDR P=.039). The right insula(β=-0.0174;P=.034) showed significant thinning, confirmed by FDR correction(FDR P=.037).

Conclusions and Relevance:

Higher MBI-C scores are linked to cortical thinning in brain regions essential for memory, spatial

orientation, and emotional regulation. These findings support the use of MBI as an early marker of neurodegeneration, helping identify individuals at risk of cognitive decline. Incorporating MBI assessments into clinical practice could guide triaging of neuroimaging orders, optimizing healthcare resource use.

Mild Behavioral Impairment as a Clinical Correlate of Early Neurodegeneration

The brain and its behavior are fundamental to our humanity. Research in these areas deepens our knowledge of what it means to be human and aids in development of treatments for neurological and psychiatric conditions. Neuropsychiatric symptoms (NPS) and their subset, Mild Behavioral Impairment (MBI), and their relationship with morphological aspects of the brain such as cortical thickness could help in the understanding of pathobiological processes in Alzheimer's disease (AD) and other dementias. Recent studies emphasize the relationship between brain structure and behavioral features in predicting cognitive syndromes, underscoring the importance of multimodal approaches for early diagnosis and intervention [1,2].

NPS refers to a broad range of non-cognitive, behavioral, or psychiatric symptoms that frequently precede or accompany cognitive decline. MBI, a specific syndrome within NPS, highlights early behavioral changes that signal a high risk for developing cognitive impairment and dementia [3,4]. Characterized by the onset of sustained behavioral symptoms later in life, MBI has emerged as a reliable predictor of cognitive decline [5]. Notably, MBI is included in the National Institute on Aging and Alzheimer's Association (NIA-AA) AD research framework in Stage 2, as a potential preclinical manifestation of underlying neuropathology [6].

MBI and other NPS are well-established predictors of cognitive decline, even in cross-sectional analyses. Baseline NPS, such as depression, anxiety, and agitation, have been identified as significant predictors of incident mild cognitive impairment (MCI) in cognitively normal older adults [7]. More than 59% of individuals with MCI or AD present with NPS before diagnosis, with depression (24%) and irritability (21%) being the most common symptoms preceding cognitive impairment [8]. A review of 61 studies revealed that depression, present in 45.9% of cases, is a significant predictor of the transition from preclinical to prodromal stages of dementia [9]. Additionally, Ferreira et al. (2023) concluded that NPS could predict the etiology, severity, and progression of AD, reinforcing the role of NPS as critical early indicators of cognitive impairment.

In parallel, MBI has emerged as a robust, independent predictor of dementia, irrespective of baseline cognitive function [10,11]. The Mild Behavioral Impairment Checklist (MBI-C) developed by [12], categorizes MBI into five domains: decreased motivation, affective dysregulation, impulse dyscontrol, social inappropriateness, and abnormal perception or false beliefs. Specific MBI domains, such as impulse dyscontrol and affective dysregulation, are strong predictors of cognitive decline, underscoring the link between early behavioral changes and subsequent cognitive deterioration [13]. A four-year study showed that integrating the MBI-C into clinical assessments significantly improved the predictive accuracy for dementia [14]. Importantly, the MBI-C has been shown to capture behavioral

changes that are often the earliest detectable signs of cognitive decline [15,16], offering a comprehensive and refined tool for systematically assessing sustained behavioral shifts indicative of early neurodegeneration [3].

In addition to behavioral biomarkers, cortical thickness serves as a key structural marker of neurodegenerative processes by measuring the thickness of the cerebral cortex. Reductions in cortical thickness are consistently observed in individuals with cognitive impairment and dementia, especially in key regions. Early thinning in the entorhinal cortex, one of the first areas affected in AD, reliably predicts cognitive decline [17]. Likewise, significant thinning in the medial temporal lobe, crucial for memory, correlates with rapid memory deterioration [18]. Thinning in the frontal and temporal cortices is linked to impairments in executive function and language, while thinning in the parietal lobe, responsible for spatial orientation, correlates with visuospatial deficits [19].

Notably, cortical thinning in asymptomatic amyloid-positive individuals signals early preclinical AD changes, highlighting cortical thickness as a valuable early biomarker. Significant thinning in vulnerable regions of amyloid deposition, such as the precuneus and posterior cingulate cortex is seen even before cognitive symptoms appear, indicating early neurodegenerative processes [20]. These reductions in cortical thickness strongly predict future cognitive decline, identifying individuals at elevated risk for AD. Longitudinal studies demonstrate that thinning in brain regions such as the entorhinal cortex, medial temporal, lateral temporal, parietal, and frontal cortices robustly predicts cognitive decline, with more severe thinning linked to faster declines in memory and executive functions [21–23]. Monitoring cortical thickness is thus critical for early detection and tracking of neurodegenerative diseases.

Recent studies underscore the relationship between behavioral impairment and cortical thinning. Gill et al. [24] found that impulse dyscontrol in MBI is linked to thinning in the orbitofrontal cortex, anterior cingulate cortex (ACC) and insula, regions associated with poorer cognitive outcomes and dementia progression. Similarly, reduced cortical thickness and increased white matter hyperintensities have been observed in NPS across neurodegenerative and cerebrovascular diseases [25]. In the study by Donovan et al, cortical thinning in regions like the medial frontal cortex, ACC, and inferior parietal lobule predicts worsening apathy and hallucinations along the AD spectrum [26]. Apathy has also been associated with lower inferior temporal thickness in both MCI patients and cognitively normal elderly individuals, suggesting its potential as an early marker of degeneration.

Asian studies further support these findings. In Japan, emotional dysregulation in MBI was linked to thinning in the right supramarginal gyrus, indicating early neurodegenerative changes[27]. In Korea, higher MBI-C scores were associated with progression from amnestic MCI to AD, with thinning observed in the temporal, parietal, and frontal regions [28]. A Korean multi-center study found

that frontal behavioral impairments correlated with more severe thinning and faster cognitive decline, particularly in executive function and memory[29].

Many studies on cortical thickness have focused on clinical populations (sample sizes 100–500) using the Neuropsychiatric Inventory (NPI) or NPI-Q, which assess dementia-related behaviors. However, these tools, designed for behaviors occurring in later stages of dementia (e.g., biting, wandering), use short reference periods of 2–4 weeks, limiting their ability to detect prodromal neurodegenerative changes. In contrast, the MBI-C assesses symptoms over six months, allowing transient states (e.g., stress, medications) to subside and providing a more accurate measure of sustained impairments [12]. This extended timeframe improves early identification of NPS, enhancing diagnostic and prognostic accuracy for at-risk populations.

This study examines the association between MBI and cortical thinning, exploring how early NPS reflect neurodegenerative processes and guide clinical decision-making. Given growing evidence that behavioral symptoms are prodromal markers of cognitive decline, we hypothesize that early behavioral changes, measured by the MBI-C, may serve as biomarkers of neurodegeneration. Integrating the MBI-C into clinical workflows could facilitate timely interventions, reduce healthcare burdens, and optimize the use of advanced imaging tools. This study aims to improve the detection and prognosis of early-stage cognitive impairment, with a focus on a multi-ethnic Southeast Asian population.

Methods

Biomarkers and Cognition Study, Singapore

Participants were drawn from the Biomarkers and Cognition Study, Singapore (BIOCIS), a five-year longitudinal study assessing individuals aged 30 to 95 annually. BIOCIS recruits 2,500 participants through community outreach across Singapore. To explore pathology with or without minimal symptoms, both cognitively impaired and unimpaired individuals are included. Participants are classified as Cognitively Normal (CN), Subjective Cognitive Decline (SCD), or MCI. Assessments include neuroimaging, biomarker analysis, cognitive tests, behavioral profiling, retinal imaging, and microbiome analysis. Participant classification follows (Petersen, 2004) and the NIA-AA criteria. Further details are provided in the BIOCIS protocol (Leow et al., 2024).

Mild Behavioral Impairment-Checklist (MBI-C) and Cognitive Assessment
The MBI-C (Ismail et al., 2017), available at MBItest.org, consists of 34 items across five domains: decreased motivation (6 items), affective dysregulation (6 items), impulse dyscontrol (12 items), social inappropriateness (5 items), and abnormal perception (5 items). Each item is scored from 0

(absent) to 3 (severe), yielding a total score of 0–102. Participants self-reported symptoms lasting at least six months, reflecting changes from prior behavior.

Reduced cortical thickness may relate to both cognitive decline and behavioral changes, but whether these behavioral impairments are directly linked to brain structure or secondary to cognitive decline remains unclear. To isolate the role of cortical structure in behavior, global cognition was measured with the Montreal Cognitive Assessment (MoCA) (Nasreddine et al., 2005), and analyses were adjusted for MoCA scores.

Neuroimaging Acquisition and Processing

All participants underwent whole□brain MRI scans using a 3T Siemens Prisma Fit scanner (Siemens Healthineers, Erlangen, Germany). A T1-weighted MPRAGE sequence was used with the following parameters: repetition time=2000 ms, echo time=2.26 ms, inversion time=800 ms, flip angle=8°, matrix size=256×256, and voxel size=1.0×1.0×1.0 mm³. Scans were processed with FreeSurfer (v7.2.0) using the "recon-all" pipeline, which included motion correction, removal of non-brain tissue, automated Talairach transformation, intensity correction, volumetric segmentation, and cortical surface reconstruction and parcellation (Desikan et al., 2006; Fischl et al., 2002; Fischl, Salat, et al., 2004; Fischl, Van Der Kouwe, et al., 2004). The FreeSurfer cortical thickness estimations were obtained using FreeSurfer default automated settings. As per the Desikan- Killiany atlas [30] the cortical thickness from 34 brain regions were extracted (separately for the left and right hemispheres): Frontal Lobe (Caudal Middle Frontal, Lateral Orbitofrontal, Medial Orbitofrontal, Pars Opercularis, Pars Orbitalis, Pars Triangularis, Precentral, Rostral Middle Frontal, Superior Frontal, Frontal Pole); Parietal Lobe (Inferior Parietal, Paracentral, Postcentral, Precuneus, Superior Parietal, Supramarginal); Temporal Lobe (Posterior Banks of Superior Temporal Sulcus, Entorhinal, Fusiform, Inferior Temporal, Middle Temporal, Parahippocampal, Superior Temporal, Temporal Pole, Transverse Temporal); Occipital Lobe (Cuneus, Lateral Occipital, Lingual, Pericalcarine); Cingulate Cortex (Caudal Anterior Cingulate, Isthmus Cingulate, Posterior Cingulate, Rostral anterior Cingulate) and Other Regions (Insula). Mean cortical thickness for each of the regions mentioned were calculated separately for the right and left hemispheres.

Statistical Analyses

Continuous variables were summarized as mean and standard deviation (SD), while categorical variables were expressed as frequency and percentage. To address skewness in the behavioral data, logarithmic transformations were applied, with a constant of 1 added for variables including values below 1. Group comparisons of continuous variables between cognitive groups were performed using

independent sample t-tests or the Mann-Whitney U test, depending on the normality assumptions. Chi-square tests (or Fisher's exact test for low expected cell counts) were used to compare categorical variables. To examine the relationship between cortical thickness and MBI-C scores, multivariable linear regression models were employed, adjusting for potential confounders such as age, education, and global cognitive function as measured by the MoCA. Regression analyses reported standardized Beta coefficients (β) to facilitate comparability of effect sizes and enable nuanced interpretation of results. Additionally, 95% confidence intervals (CIs) and p-values were provided to quantify the precision and statistical significance of associations. The BIOCIS study was adequately powered for its primary endpoints, allowing for exploration of relationships between cortical thickness and MBI-C scores as secondary endpoints. To control for multiple comparisons and avoid inflation in type I error, False Discovery Rate (FDR) correction was applied, with adjusted p-values reported. All analyses were conducted using IBM SPSS Statistics for Windows, Version 29.0 (IBM Corp., Armonk, NY), with statistical significance set at $p < 0.05$ unless otherwise specified.

Results

Participant demographics

Data from 969 participants (mean age 61.99 years ± 10.19, 39.6% males, 87.2% Chinese) were analysed. 34.3% was classified as CN, 26.4% as SCD, and 39.3% as MCI. Table 1 presents descriptives of demographics, cognitive, and cortical thickness profiles of the BIOCIS cohort.

Table 1. Descriptives of Demographics, Cognitive, Behavioral and Cortical Thickness Profiles of the BIOCIS Cohort

	CN	SCD	MCI	Overall	Overall
	n=332	n=256	n=381	n=969	*p*
Age†	54.88±10.31	56.93±9.27	62.94±10.25	58.59±10.64	**<.001**
Sex (Male)	122 (37.1%)	101(39.45%)	161 (42.3%)	384 (39.8%)	0.37
Education†	15.74±3.2	15.07±3.03	13.93±3.88	14.85±3.53	**<.001**
MoCA†	27.03±2.18	26.24±2.41	25.01±2.81	26.03±2.65	**<.001**
MBI-C Total	1.38 ± 2.912	3.93 ± 7.088	2.59±4.763	2.53 ± 5.102	**<.001**
MBI-C Interest	0.29 ± 0.897	0.93 ± 1.985	0.47±1.193	0.53 ± 1.392	**<.001**
MBI-C Mood	0.41 ± 1.193	1.12 ± 2.011	0.75±1.523	0.73 ± 1.594	**<.001**
MBI-C Control	0.58 ± 1.261	1.44 ± 2.748	1.08±2.418	1.00 ± 2.223	**<.001**
MBI-C Social	0.08 ± 0.338	0.22 ± 0.885	0.17 ± 0.565	0.15 ± 0.612	**0.026**
MBI-C Beliefs	0.02 ± 0.189	0.22 ± 0.907	0.11 ± 0.540	0.11 ± 0.591	**<.001**
Cortical Thickness (Left hemisphere)					

Temporal†	2.3374±0.1	2.326±0.1	2.3023±0.11	2.3206±0.1	**<.001**
Cingulate Cortex†	2.4478±0.11	2.4274±0.1	2.4226±0.11	2.4325±0.11	0.005
Frontal†	2.4835±0.09	2.4794±0.08	2.4804±0.09	2.4812±0.09	0.835
Occipital†	1.8262±0.08	1.819±0.09	1.8081±0.09	1.8172±0.09	0.017
Parietal†	2.2962±0.1	2.2805±0.09	2.2583±0.11	2.2772±0.1	**<.001**
Other Regions - Insula†	2.9142±0.13	2.8846±0.12	2.8728±0.14	2.8901±0.13	**<.001**
Mean of Hemisphere†	2.4015±0.08	2.3894±0.07	2.3745±0.09	2.3877±0.08	**<.001**
Cortical Thickness (Right hemisphere)					
Temporal†	2.3691±0.1	2.3552±0.1	2.3298±0.11	2.35±0.11	**<.001**
Cingulate Cortex†	2.3971±0.12	2.3862±0.11	2.3781±0.12	2.3867±0.12	0.088
Frontal†	2.4673±0.09	2.4621±0.07	2.4734±0.09	2.4683±0.09	0.272
Occipital†	1.8556±0.08	1.8515±0.09	1.8383±0.09	1.8477±0.09	0.019
Parietal†	2.3071±0.1	2.2883±0.09	2.2661±0.12	2.286±0.11	**<.001**
Other Regions - Insula†	2.9217±0.14	2.9047±0.13	2.885±0.15	2.9028±0.14	0.002
Hemisphere†	2.3961±0.08	2.384±0.07	2.3702±0.09	2.3827±0.08	**<.001**

Note. P values significant at ≤ 0.05 are bolded, † = mean ± Standard Deviation, ☐ = Frequency (percentages), ☐ = non-adjusted. Cortical data is presented in millimetres.

Abbreviations: CN = Cognitively Normal, MCI = Mild Cognitive Impairment, SCD = Subjective Cognitive Decline, MoCA = The Montreal Cognitive Assessment, BIOCIS = Biomarkers and Cognition Study, Singapore, MBI-C = Mild Behavioral Impairment-Checklist

Cortical thickness and Mild Behavioral Impairment

Significant associations were found between cortical thinning and elevated MBI-C Belief scores, particularly in the right hemisphere. In the left hemisphere, none of the associations with MBI-C domains reached statistical significance. In the right hemisphere, cortical thinning was significantly associated with higher MBI-C Belief scores (β = -0.01768, 95% CI: -0.03416, -0.0012, p = 0.035). This indicates that a one standard deviation (SD) decrease in overall cortical thickness in the right hemisphere corresponds to a 0.01768-point increase in MBI-C Belief scores. Results are summarized in Table 2, with detailed findings on MBI-C domains and specific brain regions provided in Supplementary Table 1.

Table 2. Relationship Between Hemispheric Cortical Thickness and MBI in the BIOCIS Cohort

Left hemisphere means			**Right hemisphere means**		

	B	95% CI	p	B	95% CI	p
MBI-C Total	0.01016	-0.05208, 0.07232	0.749	0.0044	-0.05696, 0.06568	0.888
MBI-C Interest	-0.01472	-0.05024, 0.0208	0.416	-0.02144	-0.05648, 0.01352	0.229
MBI-C Mood	0.01392	-0.02528, 0.05304	0.486	0.01368	-0.02488, 0.05232	0.487
MBI-C Control	0.01328	-0.03088, 0.05736	0.556	0.00408	-0.03944, 0.04752	0.855
MBI-C Social	-0.00864	-0.02776, 0.0104	0.373	-0.00864	-0.02744, 0.01016	0.365
MBI-C Beliefs	-0.012	-0.02872, 0.00472	0.159	-0.01768	-0.03416, -0.0012	**0.035**

Note. Models were adjusted for age, education, and global cognition scores. Significant p-values appear bold.

Abbreviations: BIOCIS = Biomarkers and Cognition Study, Singapore, MBI-C = Mild Behavioral Impairment-Checklist

Further analysis of the significant MBI-C subdomain identified in Table 2 revealed additional associations between specific subregions and MBI-C-Belief scores. In the left hemisphere, thinning in the entorhinal cortex was associated with higher MBI-C Belief scores (β = -0.017, 95% CI: -0.032, -0.001, p = 0.034), although this did not remain significant post-FDR correction (FDR p = 0.153). Similarly, thinning in the fusiform gyrus (β = -0.020, 95% CI: -0.036, -0.004, p = 0.014) lost significance after correction (FDR p = 0.126).

In the right hemisphere, multiple regions exhibited significant associations with MBI-C Belief scores. These included the posterior banks of the superior temporal sulcus (β = -0.020, 95% CI: -0.037, -0.004, p = 0.013), fusiform gyrus (β = -0.021, 95% CI: -0.037, -0.005, p = 0.009), superior temporal gyrus (β = -0.020, 95% CI: -0.037, -0.002, p = 0.025), temporal pole (β = -0.018, 95% CI: -0.034, -0.003, p = 0.021), and transverse temporal gyrus (β = -0.020, 95% CI: -0.036, -0.004, p = 0.014). After FDR correction, associations for the posterior banks of the superior temporal sulcus, fusiform gyrus, superior temporal gyrus, and transverse temporal gyrus remained significant (FDR p = 0.042 - 0.045), highlighting the critical role these regions play in the behavioral manifestations of MBI. Moreover, thinning in the right postcentral gyrus (β = -0.020, 95% CI: -0.036, -0.004, p = 0.012) and right supramarginal gyrus (β = -0.021, 95% CI: -0.038, -0.005, p = 0.013) was significantly linked to higher MBI-C Belief scores, with both remaining significant after FDR correction (FDR p = 0.039). The right insula also exhibited a significant association (β = -0.01736, 95% CI: -0.03374, -0.00112, p = 0.034), retaining significance post-correction (FDR p = 0.037). Table 3 presents the results.

Table 3. Detailed Associations Between Cortical Thinning in Specific Brain Subregions and MBI-C Belief Scores in the BIOCIS Cohort

Brain Region	B	95% CI	p	Adj. p
Temporal				
L Entorhinal	-0.017	-0.032, -0.001	0.034	0.153
L Fusiform	-0.02	-0.036, -0.004	0.014	0.126
R Posterior Banks of Superior Temporal Sulcus	-0.02	-0.037, -0.004	0.013	**0.042**
R Fusiform	-0.021	-0.037, -0.005	0.009	**0.042**
R Superior Temporal	-0.02	-0.037, -0.002	0.025	**0.045**
R Temporal Pole	-0.018	-0.034, -0.003	0.021	**0.045**
R Transverse Temporal	-0.02	-0.036, -0.004	0.014	**0.042**
Parietal				
R Postcentral	-0.02	-0.036, -0.004	0.012	**0.039**
R Supramarginal	-0.021	-0.038, -0.005	0.013	**0.039**
Insula				
R Insula	-0.01736	-0.03374, -0.00112	0.034	**0.037**

Note. Models were adjusted for age, education, and global cognition scores. Significant p-values appear bold.

Abbreviations: BIOCIS = Biomarkers and Cognition Study, Singapore, Adj = Adjusted, L = left, R = right

Discussion

Our study reveals significant cortical thinning across cognitive stages (CN, SCD, and MCI), as well significant relationships between higher MBI-C Beliefs subdomain and cortical thinning in key brain regions. Cortical thinning in the temporal lobes, observed consistently across all groups, reflects its central role in early AD, as this region is critically involved in memory and other cognitive processes that are first affected in AD [31,32]. Thinning in the parietal lobes, crucial for spatial awareness and attention, further underscores the widespread nature of cortical atrophy associated with cognitive decline [18,20]. Notably, thinning in the left insula, a region essential for emotional and cognitive processing, reflects its role in the progression of neurodegenerative diseases [25]. Moreover, the overall thinning of the left and right hemispheres intensifies as the cognitive stage progresses, suggesting that mean cortical thinning is a robust marker of neurodegenerative progression. Collectively, these findings reinforce cortical thinning as a critical early biomarker for tracking disease evolution in AD and related dementias.

Our findings underscore a strong association between MBI-C Belief scores and cortical thinning in the temporal, parietal, and insula regions. This is consistent with previous research showing NPS, including MBI, as early indicators of neurodegeneration, particularly in Alzheimer's

disease [5,12]. Notably, thinning in both hemispheres of the temporal lobe—one of the earliest regions affected in AD—correlated with elevated MBI-C Belief scores, reinforcing the link between structural atrophy and cognitive-behavioral symptoms. Additionally, right parietal lobe thinning, which impacts spatial orientation[33] further underscore the role of early behavioral changes as manifestations of underlying neurodegenerative processes. Finally, thinning in the right insula, critical for interoception and emotional regulation, also showed significant associations with MBI-C Belief scores, highlighting its involvement in emotional dysregulation and paranoid delusions, common in MBI [34].

In the right hemisphere, cortical thinning in several regions showed significant associations with MBI-C Belief scores, highlighting the lateralized vulnerability of specific brain areas to neurodegenerative processes. The posterior banks of the superior temporal sulcus, responsible for integrating auditory and visual cues, plays a pivotal role in social cognition, and atrophy here is associated with deficits in social perception and delusional thinking [35]. Thinning of the temporal pole, a region crucial for emotional regulation and semantic memory, was also significantly associated with emotional misinterpretations and delusional behavior characteristic of MBI [36]. Additionally, thinning in the right postcentral gyrus, involved in somatosensory processing, and the right supramarginal gyrus, responsible for body perception, was associated with somatic delusions and spatial misperceptions [37]. Thinning in the right transverse temporal gyrus, which is involved in primary auditory processing, was significantly associated with MBI-C beliefs as well, reinforcing the role of auditory misperceptions in MBI-related abnormal perception and thoughts. These findings collectively emphasize the critical role of the right hemisphere in regulating self-perception, sensory integration, and emotional processing—functions that are often compromised in neurodegenerative diseases like dementia.

Collectively, these findings highlight the particular vulnerability of right hemisphere regions—key to emotional and social processing—to early neurodegeneration, consistent with prior research showing their involvement in behavioral symptoms such as delusions and paranoia [38]. The observed asymmetry in cortical thinning, with more regions in the right hemisphere linked to higher MBI-C Belief scores, is likely due to the right hemisphere's specialization in emotional regulation, visuospatial processing, and social cognition. Previous research suggests that neurodegenerative diseases, including AD, may preferentially affect the right hemisphere, particularly in regions responsible for behavior and emotion [39,40] This lateralization may also explain why neuropsychiatric symptoms—including apathy and delusions—are often more strongly associated with right hemisphere abnormalities [26].

Overall, our findings suggests that MBI is both an early behavioral manifestation of cognitive decline, as well as reflective of structural changes in the brain. The associations between cortical thinning in regions such as the right insula, right temporal pole and posterior banks of the superior

temporal sulcus and elevated MBI-C Belief scores provide strong evidence for the role of these regions in the development of abnormal beliefs and perceptions. This underscores the notion that structural brain changes underlie the neuropsychiatric symptoms observed in MBI, offering valuable insights for the early identification of individuals at high risk of cognitive decline.

Clinical and Public Health Implications

These findings reinforce the utility of the MBI-C as an early behavioral marker for dementia risk, emphasizing its clinical relevance. The right hemisphere's vulnerability to cortical thinning, particularly in the temporal and parietal regions, aligns with its role in emotional regulation, visuospatial perception, and social cognition. This suggests that behavioral symptoms observed in MBI—such as delusions or paranoia—reflect early structural brain changes rather than psychological responses to cognitive decline.

Incorporating the MBI-C into clinical workflows offers a cost-effective strategy for early identification and management of neurodegenerative diseases. By detecting behavioral changes that precede cognitive decline, the MBI-C can facilitate timely interventions, potentially slowing disease progression. When used alongside cognitive screening tools like MoCA, it enhances diagnostic accuracy, supporting more sensitive detection of conditions such as Alzheimer's disease. MBI-C scores can also guide triage, helping prioritize high-risk patients for neuroimaging or neuropsychological assessments, thus optimizing the use of costly healthcare resources.

From a public health perspective, routine use of the MBI-C in primary care and memory clinics can promote early detection and enable preventive interventions before significant cognitive decline occurs. Integrating the MBI-C into screening programs for older adults supports resource optimization, guiding the allocation of advanced diagnostic tools such as MRI to patients who would benefit most. Additionally, raising public awareness of behavioral shifts—such as delusions and emotional dysregulation—could reduce stigma, encouraging earlier engagement with healthcare services. Early management of neuropsychiatric symptoms not only improves patient outcomes by delaying cognitive and functional decline but also helps reduce long-term healthcare costs through timely interventions.

Limitations

While the inclusion of a Southeast Asian cohort adds valuable diversity to the literature, the generalizability of the findings may be limited. Future research should validate the MBI-C's predictive value by including larger and more ethnically diverse populations across different demographics. Additionally, the cross-sectional design limits causal interpretations, underscoring the need for longitudinal studies to establish temporal relationships between MBI symptoms and cortical thinning. The reliance on self-reported data via the MBI-C may introduce response bias; incorporating

supplementary methods, such as caregiver reports or interviews, could improve the accuracy of behavioral assessments. Moreover, excluding individuals with diagnosed psychiatric conditions from the BIOCIS cohort may have inadvertently omitted participants with subclinical MBI, potentially affecting the results.

Conclusion

The MBI-C emerges as a cost-effective and scalable tool for identifying individuals at risk of neurodegeneration, with strong correlations between MBI-C Belief scores and cortical thinning across key brain regions. These findings underscore the potential of the MBI-C to improve early detection, enhance triaging for neuroimaging, and optimize healthcare resource allocation. In particular, the observed asymmetry, with greater thinning in the right hemisphere, highlights the lateralized vulnerability of brain regions crucial for emotional regulation and social cognition in the context of neurodegeneration. The MBI-C offers a simple and scalable method to identify high-risk individuals for further evaluation, clinical trial enrichment, or preventive interventions.

Future research should broaden the applicability of these findings by studying more diverse populations and employing longitudinal designs to better understand the progression of MBI and its relationship with neurodegenerative biomarkers. Investigating the mechanisms linking behavioral symptoms, cortical thinning, and cognitive decline will provide further insights into the pathophysiological processes, paving the way for earlier diagnosis and more effective interventions in dementia care.

Article Information

Corresponding Author: Nagaendran Kandiah, Lee Kong Chian School of Medicine, Nanyang Technological University, 11 Mandalay Rd, Singapore 308232 (nagaendran_kandiah@ntu.edu.sg)

Author Contributions: Yi Jin Leow had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Yi Jin Leow, Nagaendran Kandiah

Acquisition, analysis, or interpretation of data: Yi Jin Leow, Ashwati Vipin, Pricilia Tanoto, Rasyiqah Binte Shaik Mohamed Salim, Bocheng Qiu, Zahinoor Ismail, Nagaendran Kandiah

Drafting of the manuscript: Yi Jin Leow, Seyed Ehsan Saffari, Nagaendran Kandiah

Statistical analysis: Yi Jin Leow, Seyed Ehsan Saffari

Supervision: Nagaendran Kandiah

Funding: This study received funding support from the Strategic Academic Initiative grant from the Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore, National Medical Research Council, Singapore under its Clinician Scientist Award (MOH-CSAINV18nov-0007), Ministry of Education Start-up Grant, Ministry of Education Academic Research Fund Tier 1 (RT02/21) and Ministry of Education Science of Learning Grant (MOESOL2022-0002).

Conflict of Interest Disclosures: All authors declare no competing interests.

Acknowledgements: The authors express their gratitude to all individuals who are currently or will be participating in research at the Dementia Research Centre (Singapore).

Availability of data and materials

Data may be made available upon reasonable request to the corresponding author.

Ethics approval and consent to participate

Informed consent was obtained from all participants according to the Declaration of Helsinki and local clinical research regulations, and procedures used in the study were in accordance with ethical guidelines. This study has been approved by the NTU Institutional Review Board (NTU-IRB-2021-1036).

References

1. Sui J, Jiang R, Bustillo J, Calhoun V. Neuroimaging-based individualized prediction of cognition and behavior for mental disorders and health: methods and promises. *Biol Psychiatry*. 2020;88(11):818-828.

2. Vipin A, Lee BTK, Kumar D, et al. The role of perfusion, grey matter volume and behavioural phenotypes in the data-driven classification of cognitive syndromes. *Alzheimers Res Ther*. 2024;16(1):40.

3. Ismail Z, Smith EE, Geda Y, et al. Neuropsychiatric symptoms as early manifestations of emergent dementia: provisional diagnostic criteria for mild behavioral impairment. *Alzheimers Dement*. 2016;12(2):195-202.

4. Ismail Z, McGirr A, Gill S, Hu S, Forkert ND, Smith EE. Mild behavioral impairment and subjective cognitive decline predict cognitive and functional decline. *J Alzheimers Dis*. 2021;80(1):459-469.

5. Creese B, Brooker H, Ismail Z, et al. Mild behavioral impairment as a marker of cognitive decline in cognitively normal older adults. *Am J Geriatr Psychiatry*. 2019;27(8):823-834.

6. Jack Jr CR, Bennett DA, Blennow K, et al. NIA-AA research framework: toward a biological definition of Alzheimer's disease. *Alzheimers Dement*. 2018;14(4):535-562.

7. Geda YE, Roberts RO, Mielke MM, et al. Baseline neuropsychiatric symptoms and the risk of incident mild cognitive impairment: a population-based study. *Am J Psychiatry*. 2014;171(5):572-581.

8. Wise EA, Rosenberg PB, Lyketsos CG, Leoutsakos JM. Time course of neuropsychiatric symptoms and cognitive diagnosis in National Alzheimer's Coordinating Centers volunteers. *Alzheimers Dement Diagn Assess Dis Monit*. 2019;11:333-339.

9. Ferreira DA, Macedo LBC, Foss MP. Neuropsychiatric symptoms as a prodromal factor in Alzheimer's type neurodegenerative disease: A scoping review. *Clin Neuropsychol*. 2024;38(5):1031-1059.

10. Kan CN, Cano J, Zhao X, Ismail Z, Chen CLH, Xu X. Prevalence, clinical correlates, cognitive trajectories, and dementia risk associated with mild behavioral impairment in Asians. *J Clin Psychiatry*. 2022;83(3):40123.

11. Sun Y, Xu W, Chen KL, et al. Mild behavioral impairment correlates of cognitive impairments in older adults without dementia: mediation by amyloid pathology. *Transl Psychiatry*. 2021;11(1):577.

12. Ismail Z, Agüera-Ortiz L, Brodaty H, et al. The Mild Behavioral Impairment Checklist (MBI-C): a rating scale for neuropsychiatric symptoms in pre-dementia populations. *J Alzheimers Dis*. 2017;56(3):929-938.

13. Banks SJ, Raman R, He F, et al. The Alzheimer's disease cooperative study prevention instrument project: longitudinal outcome of behavioral measures as predictors of cognitive decline. *Dement Geriatr Cogn Disord Extra*. 2014;4(3):509-516.

14. Ismail Z, Leon R, Creese B, Ballard C, Robert P, Smith EE. Optimizing detection of Alzheimer's disease in mild cognitive impairment: a 4-year biomarker study of mild behavioral impairment in ADNI and MEMENTO. *Mol Neurodegener*. 2023;18(1):50.

15. Leow YJ, Soo SA, Kumar D, et al. Mild Behavioral Impairment and Cerebrovascular Profiles Are Associated with Early Cognitive Impairment in a Community-Based Southeast Asian Cohort. *J Alzheimers Dis*. 2024;(Preprint):1-9.

16. Soo SA, Ng KP, Wong F, et al. The association between diabetes mellitus and mild behavioral impairment among mild cognitive impairment: Findings from Singapore. *J Alzheimers Dis*. 2021;82(1):411-420.

17. Velayudhan L, Proitsi P, Westman E, et al. Entorhinal cortex thickness predicts cognitive decline in Alzheimer's disease. *J Alzheimers Dis*. 2013;33(3):755-766.

18. Pacheco J, Goh JO, Kraut MA, Ferrucci L, Resnick SM. Greater cortical thinning in normal older adults predicts later cognitive impairment. *Neurobiol Aging*. 2015;36(2):903-908.

19. Fjell AM, Idland AV, Sala-Llonch R, et al. Neuroinflammation and tau interact with amyloid in predicting sleep problems in aging independently of atrophy. *Cereb Cortex*. 2018;28(8):2775-2785.

20. Dickerson BC, Bakkour A, Salat DH, et al. The cortical signature of Alzheimer's disease: regionally specific cortical thinning relates to symptom severity in very mild to mild AD dementia and is detectable in asymptomatic amyloid-positive individuals. *Cereb Cortex*. 2009;19(3):497-510.

21. Fjell AM, Westlye LT, Grydeland H, et al. Accelerating cortical thinning: unique to dementia or universal in aging? *Cereb Cortex*. 2014;24(4):919-934.

22. Márquez F, Yassa MA. Neuroimaging Biomarkers for Alzheimer's Disease. *Mol Neurodegener*. 2019;14(1):21. doi:10.1186/s13024-019-0325-5

23. Ossenkoppele R, Jansen WJ, Rabinovici GD, et al. Prevalence of amyloid PET positivity in dementia syndromes: a meta-analysis. *Jama*. 2015;313(19):1939-1950.

24. Gill S, Wang M, Mouches P, et al. Neural correlates of the impulse dyscontrol domain of mild behavioral impairment. *Int J Geriatr Psychiatry*. 2021;36(9):1398-1406.

25. Ozzoude M, Varriano B, Beaton D, et al. White matter hyperintensities and smaller cortical thickness are associated with neuropsychiatric symptoms in neurodegenerative and cerebrovascular diseases. *Alzheimers Res Ther*. 2023;15(1):114.

26. Donovan NJ, Wadsworth LP, Lorius N, et al. Regional cortical thinning predicts worsening apathy and hallucinations across the Alzheimer disease spectrum. *Am J Geriatr Psychiatry*. 2014;22(11):1168-1179.

27. Imai A, Matsuoka T, Narumoto J. Emotional dysregulation in mild behavioral impairment is associated with reduced cortical thickness in the right Supramarginal gyrus. *J Alzheimers Dis*. 2023;93(2):521-532.

28. Yoon EJ, Lee JY, Kwak S, Kim YK. Mild behavioral impairment linked to progression to Alzheimer's disease and cortical thinning in amnestic mild cognitive impairment. *Front Aging Neurosci*. 2023;14:1051621.

29. Kim SJ, Jung NY, Kim YJ, et al. Clinical effects of frontal behavioral impairment: Cortical thickness and cognitive decline in individuals with subjective cognitive decline and amnestic mild cognitive impairment. *J Alzheimers Dis*. 2019;69(1):213-225.

30. Desikan RS, Ségonne F, Fischl B, et al. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage*. 2006;31(3):968-980.

31. Du AT, Schuff N, Kramer JH, et al. Different regional patterns of cortical thinning in Alzheimer's disease and frontotemporal dementia. *Brain*. 2007;130(4):1159-1166.

32. Whitwell JL, Weigand SD, Shiung MM, et al. Focal atrophy in dementia with Lewy bodies on MRI: a distinct pattern from Alzheimer's disease. *Brain*. 2007;130(3):708-719.

33. Tumati S, Martens S, de Jong B, Aleman A. Lateral parietal cortex in the generation of behavior: Implications for apathy. *Prog Neurobiol*. 2019;175:20-34.

34. Cozza M, Boccardi V. A narrative review on mild behavioural impairment: an exploration into its scientific perspectives. *Aging Clin Exp Res*. 2023;35(9):1807-1821.

35. Pelphrey KA, Sasson NJ, Reznick JS, Paul G, Goldman BD, Piven J. Visual scanning of faces in autism. *J Autism Dev Disord*. 2002;32:249-261.

36. Olson IR, Plotzker A, Ezzyat Y. The enigmatic temporal pole: a review of findings on social and emotional processing. *Brain*. 2007;130(7):1718-1731.

37. Seghier ML. The angular gyrus: multiple functions and multiple subdivisions. *The Neuroscientist*. 2013;19(1):43-61.

38. Toga AW, Thompson PM. Mapping brain asymmetry. *Nat Rev Neurosci*. 2003;4(1):37-48.

39. Gainotti G. Emotional behavior and hemispheric side of the lesion. *Cortex*. 1972;8(1):41-55.

40. Thompson PM, Hayashi KM, De Zubicaray GI, et al. Mapping hippocampal and ventricular change in Alzheimer disease. *Neuroimage*. 2004;22(4):1754-1766.

.

Supplementary Table 1

Association of Cortical thickness and Mild Behavioral Impairment in the BIOCIS cohort

	Left hemisphere			Right hemisphere		
	B	95% CI	p	B	95% CI	p
MBI-C Total						
Temporal	-0.0046	-0.0649, 0.0557	0.881	0.02123	-0.04422, 0.08668	0.524
Cingulate Cortex	-0.03047	-0.0902, 0.02937	0.318	-0.0186	-0.0792, 0.042	0.547
Frontal	0.01422	-0.04509, 0.07353	0.639	0.00207	-0.05706, 0.06129	0.944
Occipital	0.00558	-0.05778, 0.06903	0.863	-0.02511	-0.08856, 0.03834	0.437
Parietal	-0.0009	-0.0638, 0.062	0.977	0	-0.05535, 0.05526	0.437
Other Regions - Insula	-0.03536	-0.09412, 0.02327	0.237	-0.01834	-0.07924, 0.0427	0.556
Mean of Hemisphere	0.01016	-0.05208, 0.07232	0.749	0.0044	-0.05696, 0.06568	0.888
MBI-C Interest						
Temporal	-0.0053	-0.0397, 0.0292	0.764	0.00649	-0.03091, 0.04389	0.735
Cingulate Cortex	-0.00803	-0.04224, 0.02607	0.643	-0.0096	-0.04428, 0.02496	0.584
Frontal	-0.00621	-0.04014, 0.02763	0.718	-0.02088	-0.05463, 0.01296	0.226
Occipital	-0.01161	-0.04788, 0.02457	0.528	-0.01791	-0.05409, 0.01836	0.333
Parietal	-0.0184	-0.0544, 0.0175	0.314	-0.02187	-0.05337, 0.00972	0.175
Other Regions - Insula	-0.02301	-0.05655, 0.01053	0.178	-0.01344	-0.0483, 0.02128	0.447
Mean of Hemisphere	-0.01472	-0.05024, 0.0208	0.416	-0.02144	-0.05648, 0.01352	0.229

		MBI-C Mood				
Temporal	-0.0017	-0.0397, 0.0363	0.93	0.0176	-0.02365, 0.05885	0.402
Cingulate Cortex	-0.00319	-0.04081, 0.03454	0.869	0.00108	-0.03708, 0.03924	0.954
Frontal	0.01422	-0.02313, 0.05148	0.456	0.0054	-0.03186, 0.04266	0.776
Occipital	0.00459	-0.03537, 0.04455	0.821	-0.01098	-0.05094, 0.02889	0.588
Parietal	0.0063	-0.0333, 0.0459	0.756	0.01062	-0.02421, 0.04545	0.55
Other Regions - Insula	-0.02236	-0.05928, 0.01456	0.236	-0.02198	-0.06034, 0.01652	0.263
Mean of Hemisphere	0.01392	-0.02528, 0.05304	0.486	0.01368	-0.02488, 0.05232	0.487
		MBI-C Control				
Temporal	0.0094	-0.0333, 0.0522	0.666	0.0077	-0.03872, 0.05401	0.746
Cingulate Cortex	-0.01518	-0.05753, 0.02728	0.484	-0.00348	-0.04644, 0.03948	0.875
Frontal	0.00333	-0.03879, 0.04536	0.878	-0.00468	-0.04671, 0.03726	0.826
Occipital	0.02115	-0.02385, 0.06606	0.356	0.00504	-0.03996, 0.04995	0.827
Parietal	0.0058	-0.0388, 0.0503	0.8	0.00423	-0.03501, 0.04347	0.832
Other Regions - Insula	-0.00091	-0.04251, 0.04069	0.966	-0.00224	-0.0455, 0.04102	0.918
Mean of Hemisphere	0.01328	-0.03088, 0.05736	0.556	0.00408	-0.03944, 0.04752	0.855
		MBI-C Social				
Temporal	-0.0109	-0.0294, 0.0076	0.248	-0.01496	-0.03498, 0.00517	0.144
Cingulate Cortex	0	-0.01826, 0.01837	0.996	0.0066	-0.012, 0.0252	0.486
Frontal	-0.00432	-0.0225, 0.01386	0.642	0.00216	-0.01602, 0.02034	0.817
Occipital	-0.00243	-0.02196, 0.01701	0.804	-0.01359	-0.03303, 0.00585	0.17
Parietal	-0.0123	-0.0316, 0.007	0.211	-0.01134	-0.02826, 0.00567	0.19
Other Regions - Insula	-0.01066	-0.0286, 0.00741	0.248	-0.00462	-0.02338, 0.014	0.624
Mean of Hemisphere	-0.00864	-0.02776, 0.0104	0.373	-0.00864	-0.02744, 0.01016	0.365
		MBI-C Beliefs				
Temporal	-0.0218	-0.0379, -0.0056	**0.008**	-0.02189	-0.03938, -0.00429	**0.015**
Cingulate Cortex	-0.0077	-0.02376, 0.00836	0.347	-0.00948	-0.0258, 0.00684	0.254
Frontal	-0.0054	-0.02133, 0.01053	0.504	-0.0045	-0.02043, 0.01143	0.577
Occipital	-0.00873	-0.02574, 0.00828	0.316	-0.01323	-0.03024, 0.00387	0.129

Parietal	-0.0121	-0.029, 0.0048	0.159	-0.0153	-0.03015, -0.00045	**0.043**
Other Regions - Insula	-0.0078	-0.02353, 0.00806	0.333	-0.01736	-0.03374, -0.00112	**0.037**
Mean of Hemisphere	-0.012	-0.02872, 0.00472	0.159	-0.01768	-0.03416, -0.0012	**0.035**

Note. Models were adjusted for age, education, and global cognition scores. Significant p-values appear bold.

Abbreviations: BIOCIS = Biomarkers and Cognition Study, Singapore, MBI-C = Mild Behavioral Impairment-Checklist