Older Employees' Perspectives on the wLiFE55+ Program: A Qualitative Evaluation of a

Workplace Lifestyle-integrated Exercise Intervention

Authors: Diana Pfister (University of Konstanz, Konstanz, Germany), Yvonne Ritter (University

of Konstanz, Konstanz, Germany), Greta M. Steckhan (Institute of Occupational Medicine, Charité

University Medicine Berlin, Berlin, Germany), Susanne Voelter-Mahlknecht (Institute of

Occupational Medicine, Charité University Medicine Berlin, Berlin, Germany and Preventive

Medicine, University Medical Center, Göttingen, Germany), Britta Weber (Institute for

Occupational Safety and Health of the German Social Accident Insurance (IFA), Sankt Augustin,

Germany), Rolf Ellegast (Institute for Occupational Safety and Health of the German Social

Accident Insurance (IFA), Sankt Augustin, Germany), Markus Gruber (University of Konstanz,

Konstanz, Germany), Michael Schwenk (University of Konstanz, Konstanz, Germany)

Corresponding author: Michael Schwenk (University of Konstanz, Konstanz, Germany)

(Michael.schwenk@uni-konstanz.de)

Clinical trial number: not applicable

Abstract

Background

The ageing workforce and increasing prevalence of sedentary behaviour highlight the need for workplace strategies that promote physical activity (PA) and prevent functional decline in older employees. Traditional workplace PA interventions (WPAI) often lack feasibility due to limited time, low motivation, and insufficient integration into daily routines. The wLiFE55+ program applies a lifestyle-integrated approach by embedding strength, balance, and PA activities into everyday work tasks. This study qualitatively explored participants' experiences to assess acceptability, perceived benefits, and priorities for refinement.

Methods

This qualitative sub-study was embedded in a four-week wLiFE55+ pilot intervention for employees aged 55 years and older. After completing the intervention, two focus groups (n = 8; mean age 58.5 years) were conducted. Discussions were audio-recorded, transcribed verbatim, and analysed using deductive qualitative content analysis informed by the Lifestyle-integrated Functional Exercise (LiFE) concept and the Theoretical Domains Framework. Analysis focused on three overarching categories: (1) overall program experiences, (2) perceptions of program content, and (3) behaviour-change processes.

Results

Participants perceived wLiFE55+ as meaningful, feasible, and aligned with naturally occurring work situations. Trainer support facilitated understanding, adaptation, and confidence in implementing activities. Reported challenges included insufficient preparatory information about the baseline assessment, rapid progression of activity difficulty, social discomfort in shared office environments, and the burden of paper-based documentation. Participants expressed a preference for clearer communication, more practice time between sessions, digital

53   documentation and feedback tools, and an expanded set of activities, including options for the

54   upper body.

Conclusion

Lifestyle-integrated approaches such as wLiFE55+ show promise for supporting functional

capacity and promoting physical activity among older employees. Embedding activities into routine

work tasks may help overcome common limitations of traditional workplace physical activity

interventions. Future refinements should include improved preparatory guidance, greater

environmental fit, and longer program durations to facilitate habit formation and sustainable

implementation.

Key Words

65   focus group, qualitative content analysis, Lifestyle-integrated Functional Exercise, older

66   employees, workplace physical activity intervention, sedentary lifestyle, health promotion

Background

The ageing of the global workforce represents one of the most significant demographic and

economic challenges of the coming decades [1]. As the proportion of employees aged 55 years

and older continues to rise, organisations increasingly rely on retaining older workers to respond

to labour shortages and maintain productivity [2-4]. Sustaining employability at older ages requires

sufficient functional capacity. However, age-related declines in neuromotor function, muscular

strength and aerobic capacity typically accelerate from about 55 years onwards [5-8]. These

changes increase the risk of mobility limitations, chronic disease and reduced work ability [9, 10].

Although age-related physical decline most strongly affects physically demanding occupations,

good physical functioning also supports work ability across all job types by reducing fatigue,

enhancing safety and facilitating routine movement tasks. Sedentary behaviour, which is highly

prevalent in office-based occupations, exacerbates this decline and may contribute to accelerated

functional deterioration among older employees [11-15].

Current World Health Organization (WHO) guidelines recommend multicomponent physical

activity (PA) for older adults, including strength and balance training, and emphasise that health

benefits can accrue from small bouts of activity accumulated throughout the day [16]. Yet

implementing these recommendations in real workplace settings remains challenging. Traditional

workplace physical activity interventions (WPAI) often struggle with limited time availability, low

relevance to employees' daily tasks, workflow disruptions and inconsistent managerial support,

resulting in modest adherence and limited behavioural change [17-23]. It is important to note that

most existing evidence originates from high-income countries in the Global North, whereas

findings from low- and middle-income countries highlight more diverse occupational PA patterns

among older adults. In many low- and middle-income settings, older employees engage in more

physically demanding work, whereas work in high-income countries is predominantly sedentary.

These structural differences underscore the need for context-sensitive and adaptable approaches

to workplace health promotion [24].

Lifestyle-integrated exercise approaches represent a promising alternative to traditional WPAI

because they integrate functional movements directly into everyday routines rather than requiring

additional training sessions. The Lifestyle-integrated Functional Exercise (LiFE) program

exemplifies this concept by embedding strength and balance activities into daily tasks through

personalised cues and contextual prompts [25]. Research across various LiFE adaptations—

individual, group-based and technology-supported—shows high acceptability, strong adherence

and improved functional outcomes among older adults [25, 26]. These findings suggest that

embedding meaningful movements into familiar routines may reduce cognitive and motivational

barriers, increase perceived relevance and enhance long-term adoption.

Building on this body of evidence, Ritter et al. developed wLiFE55+, a LiFE-based workplace

intervention tailored to office-based employees aged 55 years and older. The program aims to

integrate brief neuromotor, strength and PA tasks into commonly occurring work situations, while

being feasible to implement within the constraints of a standard workday. A pilot feasibility trial

demonstrated high adherence, improvements in neuromotor function and reductions in sedentary

time [27]. However, quantitative findings alone cannot capture how employees experience the

program in practice, how well activities fit into diverse workplace environments or which factors

facilitate or hinder sustained engagement.

Qualitative inquiry is therefore essential to provide a more nuanced understanding of employees'

perspectives and implementation processes. Insights into perceived relevance, feasibility,

behavioural mechanisms and contextual barriers can inform refinement of wLiFE55+ and support

future scale-up. The present study explored older employees' experiences with program content,

delivery and behavioural integration following a four-week workplace intervention. We aimed to

understand (1) overall program experiences and organisational context, (2) perceptions of specific

program components and materials, and (3) behavioural processes related to motivation, planning

and habit formation. These findings contribute to the optimisation of workplace lifestyle-integrated

PA interventions and the development of sustainable strategies to promote functional capacity and

active ageing in increasingly sedentary workforces.

Methods

Study design and setting

This qualitative study was embedded in the wLiFE55+ proof-of-concept pilot trial [27], which

evaluated the feasibility of a four-week workplace lifestyle-integrated exercise program for

employees aged 55 years and older. The four-week duration was intentionally chosen because

the pilot trial aimed to assess feasibility, acceptability and implementation processes rather than

to detect meaningful physiological changes, which would require a longer intervention period. The

qualitative component was designed to complement the quantitative pilot findings by providing an

in-depth understanding of participants' experiences, perceived facilitators and barriers, and

contextual factors influencing implementation.

A qualitative descriptive approach using focus group discussions was chosen to allow participants

to collectively reflect on shared and divergent experiences. Two focus groups with four participants

each were conducted after completion of the intervention and follow-up assessments. Small,

homogeneous focus groups are recommended in feasibility research because they promote

balanced participation and rich interaction [28, 29]. Sessions took place in seminar rooms at the

University of Konstanz, lasted approximately 60 minutes, and were moderated by an independent

researcher not involved in the intervention to minimise bias.

Participants

Participants were recruited from the wLiFE55+ pilot trial, which included 15 employees aged 55

years or older. Inclusion criteria were current employment, age ≥55, and no contraindications for

physical activity. A purposive sampling strategy ensured variation in working hours, physical

activity and functional capacity. Eight participants (four women, four men; mean age 58.5 years,

SD = 3.1) volunteered for the focus groups.

This sample size represents a typical proportion for embedded qualitative components in feasibility

studies. Thematic saturation was reached, as no new themes emerged in the second focus group

[30, 31]. All participants provided written informed consent.

Table 1 *Characteristics of focus group participants (n = 8)*

	Mean ± SD
Age (years)	58.5 ± 3.1
Gender (women/men)	4/4

BMI (kg/m^2)	26.0 ± 3.0
Working hours (h/week)	35.5 ± 8.4
IPAQ, Total physical activity (MET-minutes/week)	2835 ± 2239
IPAQ, Work activity (MET-minutes/week)	343 ± 809
CBM, score (0-96)	82.1 ±11.7
60 sec Chair Stand Test (repetitions)	44.3 ± 11.6

Note. BMI Body Mass Index, IPAQ The International Physical Activity Questionnaire, *CBM*

*C*ommunity Balance and Mobility scale

Program

wLiFE55+ integrates brief neuromotor, strength and physical activity tasks—such as one-leg

stands, squats and brisk walking—into naturally occurring work situations. Table 1 provides an

overview of activities; full program details are reported elsewhere [27].

Baseline and follow-up assessments were conducted either at the University of Konstanz or in the

workplace to evaluate real-world feasibility. In the pilot trial, wLiFE55+ showed high feasibility, with

100% adherence and no adverse events. Quantitative pre–post analyses indicated medium-to-

large improvements in neuromotor function (including significant CBM gains), reductions in

sedentary time and sedentary bouts, and positive trends in lower-body strength and daily activity.

As a feasibility study, the four-week duration was designed to assess acceptance and early

implementation rather than fully developed physiological change [27].

These quantitative findings demonstrate feasibility and potential benefit, but they do not capture

how participants experienced and implemented the program in daily work life—questions

addressed by the present qualitative study.

Participants attended four weekly individual sessions with a trained personal trainer after the

baseline assessment. Trainers introduced core activities, tailored difficulty levels, and supported

participants in linking activities to daily work routines using planning tools described in the program

manual. These sessions aimed to enhance understanding, personalize activities and support early

habit formation.

Administering the intervention

After the baseline assessment, participants received the wLiFE55+ manual and completed four

weekly individual sessions with a trained personal trainer at their workplace. Trainers introduced

core activities, assessed individual abilities and suitable difficulty levels, and supported

participants in linking strength, balance and physical activity tasks to everyday work situations

using tools such as the Daily Routine Chart and Activity Planner. Across sessions, trainers

reviewed previously introduced activities, adjusted difficulty, provided feedback and refined cue–

activity plans to support integration into daily routines. Documentation in the Activity Planner

supported action planning in line with the LiFE concept and the Health Action Process Approach

(HAPA). These sessions aimed to personalise activities, enhance capability and confidence, and

facilitate early habit formation [27].

Table 2 *Short overview of wLiFE55+ activities with situations in the working environment (detailed*

table in [27])

Domain	Activity	Situation in the working environment		
		On the way to work/at home	In the office	Break

Neuromotor function	*Tandem stand*	Do the *tandem stand* when you <u>wait for the bus</u>.	Do the *tandem stand* when you are <u>booting your computer.</u>	Do the *tandem stand* *when you* <u>wait for lunch in the mensa</u>.
	One-leg stand	Do the *one-leg stand* when you <u>tie your shoes.</u>	Do the *one-leg stand* at a height-adjustable desk when <u>writing mail.</u>	Do the *one-leg stand* when you <u>wait for the coffee machine</u>.
Strength	*Squat/ one-legged squatting*	Do the *squatting* when you <u>put down your working bag.</u>	Do the *squatting* when you do <u>phone calls with colleagues.</u>	Do the *squatting* when you <u>remove a cup</u>.
	lunging	Do the *lunging* when you <u>remove objects from your working bag.</u>	Do the *lunging* when you <u>wait at the copy machine.</u>	Do the *lunging* when you <u>throw something in the bin.</u>
Physical Activity	*Walk longer*	*Walk longer* to the <u>office.</u>	*Walk longer* when you are <u>going to the office of a colleague</u>.	*Walk longer* back to the <u>office after lunch</u>.
	Walk faster	*Walk fast* for 100 m when you are <u>on the way to the office.</u>	*Walk faster* when you <u>go to the restroom/ meeting.</u>	*Walk faster* when you are going to the <u>coffee machine.</u>

Data collection

An interdisciplinary team of researchers from sports science (DP, YR, MG, MS), occupational health (GS, SVM), occupational safety (BW, RE) and the occupational healthcare industry (CK, FB) developed a semi-structured interview guide (Table 3) based on previous qualitative evaluations of LiFE [3–5]. Two focus groups (n = 4 each) were conducted by an independent moderator (MS) who was not involved in the intervention to minimise bias. Participants were encouraged to discuss their experiences openly and to include both positive and critical

reflections. All discussions were audio-recorded and transcribed verbatim with a semantic focus

following Kuckartz's guidelines [32]. As the analysis targeted manifest content, non-verbal cues

were not coded. The moderator actively encouraged balanced participation and monitored for

dominance effects, which helped manage group dynamics in the small focus groups.

Table 3 *Semi-structured interview guide for wLiFE55+ focus groups*

Data analysis

Data were analysed using qualitative content analysis following Mayring [33], which provides a

systematic approach for examining implementation processes and health-related behaviours. A

primarily deductive category system was developed based on previous LiFE studies [26, 34, 35]

and the Theoretical Domains Framework (TDF) [36], which informed categories related to

motivation, planning, capability and habit formation.

Initial categories were derived from a targeted literature search on qualitative evaluations of LiFE-

based and workplace physical activity interventions (including aLiFE, gLiFE and iLiFE) [26, 34,

35] and were reviewed and refined by three researchers (DP, YR, MS), resulting in a structured

coding guide with definitions and anchor examples (Supplementary Material 1).

All transcripts were imported into MAXQDA© and coded line by line using the smallest meaningful

text segments as coding units, focusing exclusively on manifest content. During analysis,

categories were iteratively adjusted to ensure conceptual fit, leading to a final structure of three

overarching categories.

To ensure analytical rigour, two researchers initially coded the same material independently using

the shared coding guide to check for a common understanding of the categories. Differences in

coding were discussed and used to refine the coding framework. The revised framework was then

applied in a second coding round with the involvement of a third researcher. Remaining

differences were resolved through consensus discussions, a recognised quality-assurance

strategy in qualitative content analysis [37, 38]. As the final codes were established through

negotiated agreement rather than independent parallel coding, no intercoder reliability coefficient

was calculated.

After coding, the material was condensed within categories and summarised, and representative

quotations—including divergent perspectives—were selected to illustrate key themes in relation

to the research questions.

Researcher reflexivity and positionality

The research team consisted of sports science, occupational health and occupational safety

experts. Two of the researchers (DP and YR) were involved in developing and implementing the

wLiFE55+ pilot study, which could introduce interpretive bias. To mitigate this, an independent

researcher (MS) moderated the focus groups.

Results

Results focus groups

A total of eight participants contributed 313 codable statements across the two focus groups.

Three overarching themes emerged: (1) overall program experiences, (2) perceptions of program

content, and (3) behaviour-change processes. Speaking time was balanced across participants

(5–16%, M = 12.5%).

Table 4 *Overview number of statements*

Main category	Total number of statements *n*	Sub category	Number of statements *n*
Overall program	59	Overall program considerations	13
		Challenges with implementation	39

		Progress monitoring	7
Content of the program	137	Activities	26
		Intensity	38
		Personal trainer session	28
		Material	36
		Documentation	9
Changing behaviour	117	Motivation to participate in the program	20
		Planning actions	26
		Motivation and Barriers	51
		Habit formation and integration	20

1. Overall Program

Participants generally described wLiFE55+ as meaningful and practical in a workplace context.

The core idea of embedding strength and balance activities into everyday routines was

consistently valued. Several participants noted that the program helped them recognise

movement opportunities that "fit naturally into the day" (B3) and could be carried out without

preparation or equipment. Even those who considered themselves physically active at baseline

reported becoming more aware of intentionally integrating functional movements into their routine.

1.1 Overall program considerations

Participants suggested several ways to optimise organisational aspects of the program. A

recurring theme was the desire for more time between personal trainer sessions, as the weekly

interval felt too short to practise and consolidate new activities—especially when weekends

intervened (B2). One participant summarised that "more practice time and habituation time in

between" would have supported more stable learning (B3).

Another commonly mentioned issue was the need for a neutral, private room for trainer sessions.

Participants working in shared offices reported discomfort practising balance or strength activities

in front of colleagues. A separate space was described as more appropriate to avoid feeling

watched or self-conscious (B5, B7).

Participants also expressed interest in expanding the program to include more upper-body

activities, as exercises for the arms and trunk were perceived as underrepresented. They

requested clearer explanations of what types of upper-body tasks could be incorporated into daily

routines (B1, B4). Some further asked whether a follow-up study would be offered and expressed

willingness to participate again (B1, B2).

1.2 Challenges with implementation

Challenges formed one of the most prominent subcategories and included structural,

informational, and personal factors shaping early engagement. The baseline assessment was a

major source of uncertainty: several participants were surprised by the strong emphasis on

balance tests, especially the one-leg stand with closed eyes. Some described these tasks as

"overwhelming at first" (B5), while others said they would have appreciated clearer information

beforehand about what would be assessed and why (B1, B8). One participant even questioned

whether they belonged to the target group because of the difficulty of the assessments (B6).

Participants also questioned whether the program was optimally matched to their demographic.

Some felt that less active employees might benefit even more, noting that the group seemed

relatively fit (B5, B6).

Workplace structure was another recurring barrier. Participants in open-plan offices reported

feeling inhibited about practising visible exercises, such as tandem walking or stepping over

objects. Unpredictable workdays—characterised by spontaneous meetings, sudden tasks, or

periods of high workload—also disrupted activity plans. On stressful days, activities "became a

nuisance" and were deprioritised despite positive intentions (B4).

Several participants further noted that the number of new activities introduced during the four

weeks was high, leaving them feeling "overwhelmed" (B7) and unsure whether all tasks could

realistically be integrated into everyday routines.

1.3 Progress monitoring and feedback

Participants expressed a clear interest in more differentiated feedback throughout the program.

They wished to understand how their performance compared with age-matched reference values

(B7) and found it motivating when trainers explicitly acknowledged improvements (B1). Many

suggested the use of digital tools to enable automated feedback and easier progress tracking

(B4). Some expressed curiosity about how progress would look after six to eight weeks, indicating

a desire for a longer assessment horizon (B6).

2. Content of the Program

A total of 137 statements addressed participants' experiences with specific program components,

including activities, intensity, trainer support, materials, and documentation.

2.1 Activities

Participants expressed diverse views on the wLiFE55+ activities. Exercises that could be

incorporated seamlessly into daily routines—such as one-leg stands while waiting at the printer,

balance exercises during short pauses, or stair climbing between floors—were consistently

described as intuitive, feasible, and natural (B3, B4). By contrast, activities requiring props or

additional space were often viewed as impractical in an office setting (B2, B5). The tandem walk

was repeatedly described as uncomfortable or unsuitable (B3, B4). Feedback on jumping activities

was mixed: some found them energising (B2), whereas others felt they were unpleasant or too

demanding (B7). Neuromotor tasks were sometimes described as surprising yet engaging, as they

challenged participants in unexpected ways (B4).

Participants' physical conditions strongly shaped perceptions; those with knee or joint discomfort

adapted exercises accordingly (B5, B7).

2.2 Intensity

Intensity-related comments focused on both the baseline assessment and the weekly progression

of activities. The baseline assessment was generally considered manageable but tiring due to its

duration. The one-leg stand with closed eyes was repeatedly described as the most demanding

component (B5, B7), and some noted the cognitive strain of completing multiple tests in one

session.

During the intervention, perceptions of progression varied. Some participants appreciated the

systematic increase in difficulty, finding it motivating. Others felt the progression was "too fast"

(B3), particularly when they were still becoming familiar with earlier activities. Several mentioned

the cognitive overload of learning and remembering numerous exercises within a short time frame

(B7), which limited their ability to practise all tasks consistently.

2.3 Personal trainer sessions

Trainer sessions were consistently described as one of the strongest components of the program.

Participants highlighted trainers' empathy, responsiveness, and clarity of explanation. They valued

receiving alternative strategies when difficulties arose and appreciated the encouragement to

adapt exercises to personal ability and workplace constraints (B2, B3). Demonstrations were

described as especially helpful for understanding correct execution (B7), and the respectful

atmosphere reduced feelings of embarrassment or pressure.

A few participants noted that earlier, more explicit feedback in the first session would have reduced

uncertainty (B1). Others emphasised how trainers actively sought tailored solutions when

standard exercises were unsuitable (B2).

2.4 Materials and Documentation

Participants reported mixed experiences with the program materials. The wLiFE55+ manual was viewed as a helpful reference for recalling exercises once the trainer was absent, though participants recommended clearer illustrations and more detailed instructions.

The Activity Planner, however, was widely criticised. Participants described its structure as non-intuitive and difficult to integrate into daily routines. The need to navigate between multiple pages made it cumbersome (B4), and several admitted that they eventually stopped using it because documentation felt unnecessary for completing activities (B3, B7). Many strongly preferred a digital version that would reduce administrative burden and offer automated feedback (B4).

2.5 Documentation

Documentation requirements were often incompatible with participants' work routines. Activities were frequently performed spontaneously, without the planner at hand (B3). Some found documentation redundant, noting that whether or not they recorded an activity had little influence on performing it (B7). This further reinforced participants' preference for digital options.

3. Changing Behaviour

This category captures motivational dynamics, action planning, barriers, and habit formation.

3.1 Motivation to participate

Motivation at program start varied. Some participants described an immediate motivational increase after the baseline assessment, feeling encouraged to work on identified weaknesses (B2). Others maintained stable motivation throughout (B1). One participant expressed "cautious curiosity", stating, "Let's see how it turns out later" (B3).

A few participants indicated that because they already felt physically active, they perceived less

urgency to adopt new routines, explaining they were "already on the road quite a bit" (B6).

Nevertheless, many were curious about potential physical changes over time (B5).

3.2 Planning actions

Participants reported mixed experiences with the if–then planning strategy. Some found it a useful

tool for initiating behaviour change (B2). Others struggled to generate meaningful situational cues

during the trainer sessions and felt pressured to "come up with something spontaneously" (B2).

Several later discovered that their chosen cues did not align well with their actual work routines

(B1, B5). Over time, many participants independently adapted their activities to better match

naturally occurring opportunities (B1, B3). Some requested more time during sessions to reflect

on cues and test their applicability (B2, B4).

3.3 Motivation and barriers

This was the largest subcategory, illustrating the interplay between facilitators and barriers.

Early functional improvements—standing longer on one leg, increased stability on stairs, improved

navigation in narrow spaces—served as strong motivators. Participants described balance

improvements as occurring "very quickly, very well" (B3). Social experiences could also reinforce

motivation: one participant used colleagues as a positive cue, reporting, "I used colleagues… that

was really nice" (B4).

However, several barriers were identified. Negative experiences, such as stumbling, lowered

confidence: "I stumbled around… it demotivated me a bit." (B7). Rapid progression of difficulty

also discouraged engagement: "Some of the steps were too big… which demotivated me" (B3).

Social discomfort was a significant barrier. Participants avoided practising in visible areas due to

fear of being observed or receiving negative comments, as reflected in the remark, "My colleague

laughed in the background" (B7). Social dynamics were described as ambivalent—sometimes

supportive, sometimes uncomfortable (B2).

Unpredictable work demands frequently interfered with activity plans. As one participant noted,

"When tasks or stress suddenly get in the way… it felt more like a nuisance." (B4).

3.4 Habit formation and integration

Habit formation varied widely across participants. Activities tied to recurring natural situations—

particularly waiting times—were easiest to integrate and sometimes became semi-automatic. As

one participant stated, "Wherever I wait, I automatically do one of the balance exercises" (B3).

Participants emphasised that forced or artificial cue–activity combinations rarely worked (B5).

Many recognised early signs of routine development but felt that the four-week duration was

insufficient for stable habit formation. Several estimated that six to eight weeks would be

necessary (B6). Participants developed individualised strategies to flexibly integrate activities into

unpredictable workdays, adjusting exercise selection and timing according to personal routines

(B4, B6).

Discussion

This qualitative evaluation provides an in-depth understanding of how older employees

experienced wLiFE55+ and how lifestyle-integrated movements can be embedded into everyday

workplace routines. Participants assessed the program as feasible and meaningful, particularly

because the activities aligned with naturally occurring work tasks. These findings add to evidence

from LiFE-based interventions, which consistently show that anchoring functional exercises in

daily life enhances relevance, autonomy, and long-term adherence [25, 26, 34, 35, 39]. The

present study is the first to extend this concept to an occupational context among older employees,

providing new insights into how behaviour-change mechanisms manifest within organisational

environments.

Embedding lifestyle-integrated movement: alignment with work routines and behavioural theory

A central finding of this study was that perceived feasibility depended strongly on how well activities fitted into existing work routines. Activities that could be performed during naturally recurring situations—such as waiting times, short transitions, or routine movements—were described as intuitive and easy to adopt. In contrast, activities that required additional preparation, space, or visible performance were often perceived as awkward or impractical.

This finding directly reflects barriers consistently reported in workplace physical activity intervention research, including limited time, workflow disruptions, and misalignment with job demands [17, 40, 41]. Traditional workplace interventions are often perceived as add-ons rather than integral parts of the working day, which undermines sustained engagement. The present findings suggest that embedding movement into existing routines may overcome these structural barriers, supporting earlier claims that lifestyle-integrated approaches are particularly well suited for time-constrained populations [25, 39].

At the same time, participants' experiences highlight that contextual fit is not solely determined by task structure but also by physical and social environments. Open-plan offices, shared workspaces and visibility to colleagues frequently inhibited activity performance. These findings underscore the importance of considering environmental opportunity, as conceptualised in the Capability–Opportunity–Motivation–Behaviour (COM-B) model [42], and align with previous workplace studies showing that organisational context can either enable or constrain health-promoting behaviours [19, 25, 39, 43].

Behavioural mechanisms: autonomy, capability and early habit formation

Although behaviour change was not a primary aim of this four-week feasibility study, participants' accounts provide insight into mechanisms relevant to engagement and early integration. The

ability to adapt activities to personal routines and preferences was repeatedly emphasised and

aligns with Self-Determination Theory, which highlights autonomy and perceived competence as

key drivers of sustained behaviour [44].

Participants' descriptions further suggest that early indicators of habit formation emerged when

activities were linked to stable, frequently occurring cues. Waiting situations in particular appeared

to provide reliable triggers that required minimal cognitive effort. This observation is consistent

with habit formation theory, which posits that behavioural automatisation is facilitated by stable

contexts and repeated cue–behaviour pairing [45].However, most participants felt that four weeks

were insufficient for stable habit formation, estimating that at least six to eight weeks would be

required. This perception aligns with empirical evidence suggesting that habit formation typically

unfolds over longer timeframes [45, 46].

The if–then planning strategy was perceived as useful in principle but challenging to apply in

dynamic work environments. Many participants reported difficulty identifying meaningful cues

during training sessions, particularly because daily work routines were unpredictable. This finding

reflects known limitations of implementation intention approaches in highly variable contexts and

highlights the need for iterative refinement rather than one-time planning [47].

Role of personal trainer support in implementation

Personal trainer support emerged as a key facilitator of engagement and perceived feasibility.

Participants valued personalised instruction, adaptive problem-solving and empathetic

communication. Trainers supported participants in tailoring activities to individual capabilities and

workplace constraints, thereby enhancing perceived capability and confidence.

These findings are consistent with previous LiFE research, where guided personalisation was

shown to support autonomous self-regulation and adherence [26, 34]. From an implementation

perspective, trainer support appears particularly important in occupational settings, where

heterogeneity in work tasks and environments necessitates flexible adaptation. However,

participants also reported cognitive overload due to the rapid introduction of multiple activities

within a short timeframe. This suggests that future implementations should prioritise fewer

activities with slower progression to allow consolidation and reduce burden.

Perceived benefits and motivational dynamics

Participants reported early improvements in balance and stability, which served as strong

motivators for continued engagement. These mastery experiences enhanced self-efficacy and

reinforced motivation, consistent with behaviour change theories emphasising the role of feedback

and perceived competence [48, 49]. Conversely, negative experiences—such as stumbling or

rapid increases in difficulty—were described as demotivating, underscoring the importance of

carefully calibrated progression.

Social dynamics played an ambivalent role. While some participants experienced encouragement

from colleagues, others felt self-conscious or discouraged when being observed. Similar

ambivalence has been reported in prior workplace intervention studies, where social norms and

peer reactions strongly influenced participation [43, 50]. These findings highlight that social

context should be explicitly considered in workplace-integrated interventions.

Implications for the refinement of workplace lifestyle-integrated interventions

The present findings yield several implications for the optimisation of wLiFE55+ and similar

interventions. First, clearer preparatory communication is essential, particularly regarding

assessment procedures and the focus on neuromotor activities, which were unexpected for many

participants. Second, intervention duration and progression should be adjusted to better support

learning and habit formation. Third, activity planning should be treated as an ongoing, adaptive

process rather than a fixed component. Finally, environmental and social constraints within

workplaces must be acknowledged and addressed to enhance feasibility. Although some

participants expressed interest in digital documentation, digital tools should be considered optional

rather than central, as adoption among older employees may be limited by digital literacy,

perceived complexity, and additional cognitive burden.

These implications are consistent with implementation science frameworks emphasising flexibility,

contextual adaptation and participant-centred design in complex interventions [40, 51].

Conclusion

In summary, this qualitative evaluation demonstrates that workplace-integrated functional

activities can be meaningfully adopted by older employees, while also revealing structural and

organisational barriers that require refinement. Clearer preparatory communication, more gradual

progression and longer intervention periods may strengthen feasibility and support more stable

behavioural uptake. Future research should examine longer intervention durations and evaluate

implementation and scale-up in larger and more diverse organisational settings to support

sustainable workplace integration and active ageing in predominantly sedentary work

environments.

Strength and Limitation

A key strength of this study is the balanced participation within both focus groups, which ensured

that no single participant dominated the discussion and allowed for a broad range of perspectives

[52]. The open and reflective group atmosphere, including the willingness to discuss challenges,

suggests limited social desirability bias and supports the credibility of the findings. Methodological

rigour was further enhanced by the systematic application of qualitative content analysis following

Mayring's framework and by conducting two small, homogeneous focus groups, which are well

suited to generating rich, in-depth data [33]. The use of an independent facilitator to conduct the

focus groups reduced the risk of subjective bias and researcher influence [26].

Several limitations should be acknowledged. The four-week intervention period was appropriate

for feasibility testing but likely insufficient for full integration and stable habit formation, particularly

given the number of activities introduced. The qualitative sample was small and drawn from a

single pilot trial, which limits transferability to other workplace settings and occupational groups.

Methodologically, the primarily deductive coding approach may have constrained the emergence

of unanticipated themes, and no formal intercoder reliability coefficient was calculated, as coding

decisions were resolved through consensus. Finally, non-verbal communication was not analysed,

and social desirability effects—common in focus group research—cannot be fully excluded.

List of abbreviations

COM-B = Capability–Opportunity–Motivation–Behaviour model

PA = physical activity

WPAI = workplace physical activity intervention

wLAT55+ = work LiFE55+ Assessment Tool

wLiFE55+ = work Lifestyle-integrated Functional Exercise Program for employees over 55 years

Declarations

• Ethical approval and consent to participate: All participants provided written informed

consent prior to their involvement in the study. The study protocol was reviewed and

approved by the Ethics Committee of the University of Konstanz, under the ethics approval

number IRB23KN07-005w.

• Consent for publication: All the participants who provided their case reports provided

consent for publication.

• Availability of data and materials: All data generated or analyzed during this study are

included in this published article and its supplementary information files.

• Competing interests: The authors declare that they have no competing interests.

• Funding: This work was supported by the AFF Young Scholar Fund of the University of

Konstanz. The content of this paper is solely the responsibility of the authors.

• Authors' contributions:

YR developed the concept and design of the study and was responsible for study

management as well as the statistical analysis and interpretation of the quantitative data.

DP was responsible for writing the manuscript and was involved in study management,

qualitative data collection, and analysis, including the development of the interview guide,

transcription, and evaluation of the focus groups.

MS contributed to obtaining funding, the development of the study concept and design, the

implementation and moderation of the focus groups, and the interpretation of the data.

All authors contributed to the interpretation of the data, the writing of the article and the

final approval of the version to be published.

• Acknowledgments: The authors cordially thank the wLiFE55+ participants for making this

research possible by taking part in the study.

The authors further thank Alina Boden (AB) (University of Konstanz) for contribution in

supporting the coding of participant statements.

• Authors' details (optional): Not applicable

560　Literature

1.　Vasconcelos AF. Older workers: some critical societal and organizational challenges.

Journal of Management Development. 2015;34(3):352-72.

2.　Harasty C, Ostermeier M. Population ageing: Alternative measures of dependency and

implications for the future of work: ILO Working Paper; 2020.

3.　Sippli K, Schmalzried P, Rieger MA, Voelter-Mahlknecht S. Challenges arising for older

workers from participating in a workplace intervention addressing work ability: a qualitative study

from Germany. International archives of occupational and environmental health. 2021;94:919-33.

4.   Westermeier C, Wolf M. Betriebe wollten 2018 deutlich mehr rentenberechtigte Mitarbeiter

halten als 2015. IAB-Kurzbericht; 2020.

5.   de Fátima Ribeiro Silva C, Ohara DG, Matos AP, Pinto ACPN, Pegorari MS. Short physical

performance battery as a measure of physical performance and mortality predictor in older adults:

a comprehensive literature review. International journal of environmental research and public

health. 2021;18(20):10612.

6.   Kenny GP, Yardley JE, Martineau L, Jay O. Physical work capacity in older adults:

implications for the aging worker. American journal of industrial medicine. 2008;51(8):610-25.

7.   Marées Hd, Heck H. Sportphysiologie.[Unveränd. Nachdr. d.] korr. Nachdr d. 2006;9.

8.   Suorsa K, Mattila V-M, Leskinen T, Heinonen OJ, Pentti J, Vahtera J, et al. Work ability and

physical fitness among aging workers: the Finnish Retirement and Aging Study. European journal

of ageing. 2022;19(4):1301-10.

9.   Bauman A, Merom D, Bull FC, Buchner DM, Fiatarone Singh MA. Updating the evidence

for physical activity: summative reviews of the epidemiological evidence, prevalence, and

interventions to promote "active aging". The gerontologist. 2016;56(Suppl_2):S268-S80.

10.   Claudino JG, Afonso J, Sarvestan J, Lanza MB, Pennone J, Filho CAC, et al. Strength

training to prevent falls in older adults: a systematic review with meta-analysis of randomized

controlled trials. Journal of clinical medicine. 2021;10(14):3184.

11.   Ferrucci L, Cooper R, Shardell M, Simonsick EM, Schrack JA, Kuh D. Age-related change

in mobility: perspectives from life course epidemiology and geroscience. Journals of gerontology

series a: biomedical sciences and medical sciences. 2016;71(9):1184-94.

12.   Shariat A, Cleland JA, Danaee M, Kargarfard M, Sangelaji B, Tamrin SBM. Effects of

stretching exercise training and ergonomic modifications on musculoskeletal discomforts of office

workers: a randomized controlled trial. Brazilian journal of physical therapy. 2018;22(2):144-53.

13.   Smith L, Hamer M, Ucci M, Marmot A, Gardner B, Sawyer A, et al. Weekday and weekend

patterns of objectively measured sitting, standing, and stepping in a sample of office-based

workers: the active buildings study. BMC public health. 2015;15:1-9.

14.   Thorp AA, Healy GN, Winkler E, Clark BK, Gardiner PA, Owen N, et al. Prolonged

sedentary time and physical activity in workplace and non-work contexts: a cross-sectional study

of office, customer service and call centre employees. International journal of behavioral nutrition

and physical activity. 2012;9:1-9.

15.   Wackerhage H, Sitzberger C, Kreuzpointner F, Oberhoffer-Fritz R. WHO-Leitlinien zu

körperlicher Aktivität und sitzendem Verhalten. Bayerisches Ärzteblatt. 2021(3):91-3.

16.   WHO O. WHO guidelines on physical activity and sedentary behaviour. Geneva: World

Health Organization. 2020:1-582.

17.   Garne-Dalgaard A, Mann S, Bredahl TVG, Stochkendahl MJ. Implementation strategies,

and barriers and facilitators for implementation of physical activity at work: a scoping review.

Chiropractic & manual therapies. 2019;27:1-13.

18.   Holtermann A, Straker L, Lee I-M, Van Der Beek AJ, Stamatakis E. Long overdue

remarriage for better physical activity advice for all: bringing together the public health and

occupational health agendas. BMJ Publishing Group Ltd and British Association of Sport and

Exercise Medicine; 2020. p. 1377-8.

19.   Justesen JB, Eskerod P, Christensen JR, Sjøgaard G. Implementing workplace health

promotion–role of middle managers. International Journal of Workplace Health Management.

2017;10(2):164-78.

20.   Kinnafick F-E, Thøgersen-Ntoumani C, Shepherd SO, Wilson OJ, Wagenmakers AJ, Shaw

CS. In it together: a qualitative evaluation of participant experiences of a 10-week, group-based,

workplace HIIT program for insufficiently active adults. Journal of Sport and Exercise Psychology.

2018;40(1):10-9.

21.   Neupane S, Kyrönlahti S, Oakman J, Siukola A, Riekhoff A-J, Kuivalainen S, et al. Can

workplace intervention prolong work life of older workers? A quasi-experimental study.

International Archives of Occupational and Environmental Health. 2023;96(2):237-46.

22.   Taylor WC, Horan A, Pinion Jr C, Liehr P. Evaluation of booster breaks in the workplace.

Journal of occupational and environmental medicine. 2014;56(5):529-34.

23.   Tudor-Locke C, Hendrick CA, Duet MT, Swift DL, Schuna Jr JM, Martin CK, et al.

Implementation and adherence issues in a workplace treadmill desk intervention. Applied

physiology, nutrition, and metabolism. 2014;39(10):1104-11.

24.   Asiamah N, Awuviry-Newton K, Vieira ER, Bateman A, Khan HT, Mensah HK, et al. Older

adults' vigorous occupational physical activity levels in six countries are explained by country and

'having multiple jobs'. International Journal of Environmental Research and Public Health.

2022;19(21):14065.

25.   Schwenk M, Bergquist R, Boulton E, Van Ancum JM, Nerz C, Weber M, et al. The adapted

lifestyle-integrated functional exercise program for preventing functional decline in young seniors:

development and initial evaluation. Gerontology. 2019;65(4):362-74.

26.   Reicherzer L, Kramer-Gmeiner F, Labudek S, Jansen C-P, Nerz C, Nystrand MJ, et al.

Group or individual lifestyle-integrated functional exercise (LiFE)? A qualitative analysis of

acceptability. BMC geriatrics. 2021;21:1-12.

27.   Ritter Y, Pfister D, Steckhan GM, Voelter-Mahlknecht S, Weber B, Ellegast R, et al. The

work Lifestyle-integrated Functional Exercise program for preventing functional decline in

employees aged 55 years and older: development and initial evaluation. European Review of

Aging and Physical Activity. 2024;21(1):21.

28.   Krueger RA, Casey MA. Focus group interviewing. Handbook of practical program

evaluation. 2015:506-34.

29.   Morgan DL. Focus groups as qualitative research: Sage; 1997.

30.   Guest G, Bunce A, Johnson L. How many interviews are enough? An experiment with data
saturation and variability. Field methods. 2006;18(1):59-82.

31.   Hennink MM, Kaiser BN, Marconi VC. Code saturation versus meaning saturation: how
many interviews are enough? Qualitative health research. 2017;27(4):591-608.

32.   Kuckartz U. Qualitative inhaltsanalyse: methoden, praxis, computerunterstützung: Beltz
Juventa; 2012.

33.   Mayring P. Qualitative Inhaltsanalyse. Grundlagen und Techniken (Beltz Pädagogik, 12.,
überarb. Aufl.). Weinheim: Beltz; 2015.

34.   Boulton E, Hawley-Hague H, French DP, Mellone S, Zacchi A, Clemson L, et al.
Implementing behaviour change theory and techniques to increase physical activity and prevent
functional decline among adults aged 61–70: The PreventIT project. Progress in cardiovascular
diseases. 2019;62(2):147-56.

35.   Kramer F, Labudek S, Jansen C-P, Nerz C, Fleig L, Clemson L, et al. Development of a
conceptual framework for a group-based format of the Lifestyle-integrated Functional Exercise
(gLiFE) programme and its initial feasibility testing. Pilot and feasibility studies. 2020;6:1-18.

36.   Atkins L, Francis J, Islam R, O'Connor D, Patey A, Ivers N, et al. A guide to using the
Theoretical Domains Framework of behaviour change to investigate implementation problems.
Implementation science. 2017;12:1-18.

37.   O'Connor C, Joffe H. Intercoder reliability in qualitative research: Debates and practical
guidelines. International journal of qualitative methods. 2020;19:1609406919899220.

38.   Schreier M. Qualitative content analysis in practice. 2012.

39.   Clemson L, Singh MAF, Bundy A, Cumming RG, Manollaras K, O'Loughlin P, et al.
Integration of balance and strength training into daily life activity to reduce rate of falls in older
people (the LiFE study): randomised parallel trial. Bmj. 2012;345.

40.   Mackenzie K, Such E, Norman P, Goyder E. The development, implementation and

evaluation of interventions to reduce workplace sitting: a qualitative systematic review and

evidence-based operational framework. BMC Public Health. 2018;18:1-18.

41.   Pieper C, Schröer S, Eilerts A-L. Evidence of workplace interventions—a systematic review

of systematic reviews. International journal of environmental research and public health.

2019;16(19):3553.

42.   Michie S, Van Stralen MM, West R. The behaviour change wheel: a new method for

characterising and designing behaviour change interventions. Implementation science.

2011;6(1):42.

43.   Bredahl TVG, Særvoll CA, Kirkelund L, Sjøgaard G, Andersen LL. When intervention

meets organisation, a qualitative study of motivation and barriers to physical exercise at the

workplace. The Scientific World Journal. 2015;2015(1):518561.

44.   Ryan RM, Deci EL. Self-determination theory and the facilitation of intrinsic motivation,

social development, and well-being. American psychologist. 2000;55(1):68.

45.   Lally P, Van Jaarsveld CH, Potts HW, Wardle J. How are habits formed: Modelling habit

formation in the real world. European journal of social psychology. 2010;40(6):998-1009.

46.   Van der Weiden A, Benjamins J, Gillebaart M, Ybema JF, De Ridder D. How to form good

habits? A longitudinal field study on the role of self-control in habit formation. Frontiers in

Psychology. 2020;11:494700.

47.   Gardner B, Lally P, Wardle J. Making health habitual: the psychology of 'habit-

formation'and general practice. The British Journal of General Practice. 2012;62(605):664.

48.   Bandura A. Self-efficacy: The exercise of control: Freeman; 1997.

49.   Schwarzer R. Self-efficacy in the adoption and maintenance of health behaviors:

theoretical approaches and a new model. Self-efficacy: thought control of action. Washington, DC,

US: Hemisphere Publishing Corp. 1992:217-43.

50.   Burton E, Lewin G, Clemson L, Boldy D. Effectiveness of a lifestyle exercise program for

older people receiving a restorative home care service: a pragmatic randomized controlled trial.

Clinical interventions in aging. 2013:1591-601.

51.   Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M. Developing and

evaluating complex interventions an introduction to the new Medical Research Council guidance.

Evidence–Based Public Health: Effectiveness and Efficiency. 2010;1:185-203.

52.   Smithson J. Using and analysing focus groups: limitations and possibilities. International

journal of social research methodology. 2000;3(2):103-19.

Please insert the following table on page 9 under the title:

Table 3 *Semi-structured interview guide for* wLiFE55+ *focus groups*

Topic	Initial question	Prompts
Baseline assessment	Can you tell us what you thought of the baseline assessment at the beginning of the study?	How easy or difficult did you find the baseline assessment? How did you feel about the duration of the assessment? How motivated were you to continue participating in the study after you had taken part in the baseline assessment?
Personal trainer sessions	Can you tell us how you found the session with the personal trainer?	How challenging or undemanding did you find the personal trainer session? How high was your motivation during the execution of the activities?

		What did you think of the duration of the personal trainer sessions? What did you like? Were there any difficulties? What worked well? What did you not like?
Activities	How did you find the activities?	How fitting or unfitting did you find the activities? If unfitting: Can you provide a reason for this response? Are there activities that you would suggest as alternatives? How did you find the difficulty levels of the activities?
Experiences between the weekly personal trainer sessions	How did you perceive the time between the personal trainer sessions, during which you could integrate the activities into your daily life on your own?	How easy or difficult was it for you to integrate the activities into your daily life? Which activities could be integrated into daily life particularly well or less well? In which everyday situation (e.g., at work or home) was the implementation of the program particularly easy or difficult?
Improvement suggestions	Do you have any ideas on how the wLiFE55+ program could be further improved?	Do you have ideas on how the personal trainer sessions can be improved? Do you have ideas on how the materials can be improved? How helpful did you find the activity planner? How helpful did you find the participant's manual? Are there any additional pieces of information that

you think are important and might be missing?