Toward Digital Twins in the Intensive Care Unit: A Medication Management Case Study

Behnaz Eslami, MS[1], Majid Afshar, M.D., MSCR[2], M. Samie Tootooni, Ph.D.[1], Timothy Miller, Ph.D.[3], Matthew Churpek, M.D., MPH, PH.D.[2], Yanjun Gao, Ph.D.[4], Dmitriy Dligach, Ph.D.[1]

[1] Loyola University Chicago, [2] University of Wisconsin School of Medicine and Public Health, [3] Boston Children's Hospital and Harvard Medical School, [4] University of Colorado

ABSTRACT

Digital twins, computational representations of individuals or systems, offer promising applications in the intensive care unit (ICU) by enhancing decision-making and reducing cognitive load. We developed digital twins using a large language model (LLM), LLaMA-3, fine-tuned with Low-Rank Adapters (LoRA) on physician notes from different ICU specialties in the MIMIC-III dataset. This study hypothesizes that specialty-specific training improves treatment recommendation accuracy compared to training on other ICU specialties. Additionally, we evaluated a zero-shot baseline model, which relied solely on contextual instructions without training. Discharge summaries were analyzed, and medications were masked to create datasets for model training and testing. The medical ICU dataset (1,000 notes) was used for evaluation, and performance was measured using BERTScore and ROUGE-L. LLMs trained on medical ICU notes achieved the highest BERTScore (0.842), outperforming models trained on other specialties or mixed datasets, while untrained zero-shot models showed the lowest performance. These results underscore the value of context-specific training for digital twins, offering foundational insights into LLMs for personalized clinical decision support.

INTRODUCTION

A digital twin is a virtual model, continuously synchronized with real-time data, that replicates a physical object, system, or process. By integrating diverse data sources and leveraging predictive analytics, it enables simulation, monitoring, and analysis of its physical counterpart, fostering improved decision-making and optimization.[1] A digital twin in the medical context is a dynamic, data-driven representation of real-time clinical scenarios that evolves in response to patient data and advancing medical knowledge.[2] Functioning as a bridge between healthcare providers and artificial intelligence systems, such as large language models (LLMs), a digital twin continuously adapts to the types of patients seen by a provider as electronic health records (EHR) expand with new data and disease insights. By integrating its internal medical knowledge with EHR data, the digital twin can assist in diagnosis and treatment reasoning, helping to alleviate the cognitive burden and information overload often encountered in the care of complex critical care patients.[3,4] Digital twins can be tailored for varying levels of granularity, ranging from a single provider to an entire medical specialty or healthcare organization, offering scalable and personalized support across diverse clinical settings.

In critical care medicine, a digital twin can have a particularly high impact given the dynamic and time-sensitive nature of intensive care units (ICU), drawing on massive amounts of data that are often beyond the processing abilities of an individual physician. The typical ICU is not a monolith, but rather a consortium of multi-disciplinary specialists with different training paths and experience. For example, a patient with a major cerebrovascular accident would be treated in the Neuro ICU, a severe trauma in the surgical ICU, and septic shock in the Medical ICU. Our choice of the level of granularity at which we perform our modeling is driven by the availability of training data; we resort to the Medical Information Mart for Intensive Care III (MIMIC-III) corpus[5] of over two million critical care notes that can be further stratified into eleven ICU specialties (**Figure 1**), facilitating the creation of ICU specialty-specific digital twins.

We begin by introducing a new and challenging modeling task: ICU medication prediction. The total number of *unique* medications mentioned in MIMIC Medical ICU discharge summaries exceeds 14,000, making the task challenging for both humans and machines. The task was designed as a sequence generation task where the input to an LLM are sections from the discharge summaries with *all* medication mentions masked using a special token. The generated output from the LLM was a prediction of the masked medications. A synthetic example that illustrates this task is shown in **Figure 2** (our data use agreement with PhysioNet does not allow us to include actual MIMIC notes). Best practices for common ICU medications like vasopressors or sedation often vary between ICU specialties. We trained digital twin classifiers on notes from different ICU specialties and assessed their performance on specialty-specific notes, comparing zero-shot learners with LLMs adapted to distinct ICU specialties to measure shifts in treatment preferences. Open-source LLMs were used in zero-shot experiments to represent a physician without any knowledge of preferences that may exist within an ICU specialty. The digital twin was trained by adapting an LLM, LLaMA3[6], to reflect treatment preferences specific to an ICU specialty using Low-Rank Adapters (LoRA)[7]. For example, we used notes written by physicians working in a cardiothoracic ICU to train a cardiothoracic digital twin. We evaluated our digital twins by measuring their performance on the notes from the most common ICU type, Medicine. Our hypothesis was that there are measurable differences across the treatment courses that are used across different ICU specialties and an LLM adapter can be trained to capture those preferences, effectively creating a digital twin. The datasets are available to the community to facilitate future work on this task [GitHub repo available upon acceptance].

Fig. 1: Frequency Distribution of Service Types in MIMIC-III Note Events

RESULTS

Training data consisted of five datasets of ICU notes, as detailed in **Table 1**. For evaluation, the development and test sets included only Medical ICU notes, with 1,500 notes allocated to the development set and 1,000 notes to the test set. The performance of various models was assessed using accuracy, with the results on the test set of Medical ICU notes presented in **Table 1**. The flow diagram and final set of experiment architectures are shown in **Figure 3**. Models trained on the Medical ICU specialty demonstrated the best performance with a Bidirectional Encoder Representations from Transformers Score (BERTScore)[8] of 0.842, achieving markedly better results than those trained on the notes from other ICU specialties. Even the training scenario that used all ICU specialties had a lower BERTScore at 0.783. The Cardiothoracic ICU digital twin had the worst performance, revealing a distinct gap between physician preferences for medication use in Cardiothoracic versus medical ICU specialty. The zero-shot models performed substantially worse, particularly in terms of accuracy, which was much lower than other models.

Fig. 3: LLMs Medication Prediction Flow Diagram

We attributed this performance gap to challenges in designing prompts that effectively extracted medication names, as demonstrated by instances where the model returned medication categories (e.g., "analgesics" instead of "dilaudid," or "antibiotic" instead of "ceftriaxone"). This was supported by a BERTScore of 0.713 for the zero-shot model that was closer to the performance of the supervised models but lower than the other models. Customizing the zero-shot prompt to specific ICU specialties, as shown in the bottom three rows of **Table 1**, did not improve the performance compared to the default (ICU-agnostic prompt), highlighting the need of supervised learning for creating digital twins.

While the medical ICU digital twin achieved relatively high performance with the BERTScore, it gave partial credit to approximate matches such as "ibuprofen" and "NSAID," and in some scenarios such credit may not be warranted. While BERTScore effectively measures semantic similarity between two sequences, its inherent variability with longer sequences — likely influenced by the high variance in sequence lengths across notes — underscores the importance of utilizing n-gram overlap metrics like the Recall-Oriented Understudy for Gisting Evaluation - Longest Common Subsequence (ROUGE-L).[9] ROUGE-L provides a more stable assessment of token-level overlap and alignment, ensuring a

comprehensive evaluation of model performance despite differences in note length. Our results remained consistent when examining ROUGE-L scores and accuracy, with the medical ICU digital twin model trained on all medicine notes performing the best **(Table 1).**

To assess variability around the point estimate scores, 95% confidence intervals were calculated using 1000-iteration bootstrapping. The BERTScore showed high variance as demonstrated by the wide confidence intervals across all scenarios. To further examine the variance, the individual confidence intervals across different medication sequence lengths generated by the LLM are shown in **Figure 4**. Shorter medication sequences were associated with greater BERTScore variability, reflected in wider confidence intervals, indicating higher uncertainty in the score estimation. Conversely, longer sequences resulted in narrower confidence intervals, suggesting more robust and consistent BERTScore measurements. The variability in confidence intervals for sequence length in generated medications is an inherent characteristic of BERTScore. This metric relies on pairwise semantic comparisons, and as the number of available medication entities increases, the likelihood of identifying closely matching pairs rises. Consequently, longer sequences enable more comparisons, contributing to statistical variability in the BERTScore. In at least 20% of the test set notes, the models generated medication sequences did not match the reference medication sequence lengths **(Table 2).** The number of mismatched sequences were as high as 55% for the model trained on cardiothoracic ICU notes.

Method	Prompt	Training Corpus	Accuracy	ROUGE-L	BERTScore
Supervised Fine Tuning (SFT)	Default	Medical ICU (4,118)	0.219 (95% CI 0.211-0.227)	0.593 (95% CI 0.588-0.598)	0.820 (95% CI 0.544- 0.923)
		Cardiothoracic ICU (4,118)	0.116 (95% CI 0.111-0.121)	0.533 (95% CI 0.528-0.538)	0.783 (95% CI 0.524- 0.913)
		Surgical ICU (4,118)	0.163 (95% CI 0.156-0.170)	0.560 (95% CI 0.556-0.565)	0.797 (95% CI 0.568- 0.938)
		All ICU Random Sample (4,118)	0.187 (95% CI 0.180-0.194)	0.577 (95% CI 0.572-0.582)	0.813 (95% CI 0.575- 0.931)
		All Medical ICU (16,330)	0.313 (95% CI 0.304-0.322)	0.636 (95% CI 0.631-0.642)	0.842 (95% CI 0.601-0.951)
Zero-Shot	Default	N/A	0.010 (95% CI 0.008-0.012)	0.345 (95% CI 0.337-0.352)	0.715 (95% CI 0.428- 0.882)
	Medicine		0.008 (95% CI 0.006-0.010)	0.328 (95% CI 0.320-0.336)	0.713 (95% CI 0.397-0.883)
	Surgery		0.012 (95% CI 0.010-0.014)	0.356 (95% CI 0.349-0.363)	0.721 (95% CI 0.397-0.883)
	Cardiothoracic		0.009	0.359	0.723

	(95% CI 0.007-0.011)	(95% CI 0.352-0.366)	(95% CI 0.397-0.885)

Table 1. Medication prediction performance on Medicine notes for supervised fine-tuning (SFT) and Zero-shot learning experiments. For SFT, LoRA adapters were trained to reflect medication preferences within a single ICU specialty. For zero-shot learning, prompts were designed for an ICU specialty-agnostic prompt ("Default" prompt above) and ICU-specific prompts (e.g. a prompt that begin with "You are a Cardiothoracic ICU physician …"). The accuracy was measured as the proportion of exact matches between predicted and ground truth medications, and the 95% confidence intervals were calculated using bootstrap resampling with 10,000 iterations.

Fig. 4: BERTScore with confidence intervals for different ICU specialties across medication sequences of different lengths binned into 10 intervals. The number below each interval is the sample size for that range. Overall, shorter sequences are associated with wider confidence intervals. Longer sequences, on the other hand, have tighter confidence intervals.

ICU Specialty	Notes with mismatched length, %
Medical ICU (4,118)	22.2
Cardiothoracic ICU (4,118)	55.1
Surgical ICU (4,118)	21.4
All ICU Random Sample (4,118)	20.4
All Medical ICU (16,330)	7.9

Table 2. Percentage of test notes for which the number of generated medication sequences does not match the number of reference medications.

DISCUSSION

Our study demonstrates the feasibility of developing ICU-specific digital twins using LLMs enhanced with LoRA adapters, focusing on medication prediction as a proof-of-concept task. Our findings underscore the superior performance of specialty-specific digital twins, such as the Medical ICU model, compared to both multi-disciplinary and other specialty models when predicting medications. This highlights the utility of tailored digital twins in augmenting decision-support systems and enhancing medication management in critical care. By leveraging the internal medical knowledge of LLMs and the

real-world practice data embedded in EHRs, digital twins have the potential to alleviate cognitive burden and streamline complex decision-making for ICU providers.

Digital twins in healthcare traditionally represent virtual counterparts of patients, aiding in simulating outcomes, forecasting treatment responses, and advancing precision medicine.[10] Vallée et al.[3] demonstrates the potential of digital twins for real-time monitoring and predictive analytics in chronic disease management, highlighting their role in providing proactive and individualized care. Our approach extends this concept to provider-specific digital twins that adapt to ICU specialties, incorporating both clinical expertise and local practice behaviors captured in the EHR. LoRA adapters proved instrumental in this task, offering efficient fine-tuning with fewer parameters and enabling models to reflect provider-specific practices. This adaptability is critical, as ICU specialties vary significantly in their treatment protocols and clinical priorities. Additionally, the flexibility of LoRA allows for dynamic updates, ensuring that digital twins remain current with evolving EHR data and guidelines.

LLMs like GPT-4[11] and LLaMA[12] have shown great promise in medical document processing[13], becoming increasingly important in medicine and medical informatics by enhancing diagnostic precision and treatment decisions through data-driven insights[14]. Our experiments indicated that LLMs can be trained to reflect the differences in treatment methods across different ICU specialties, similar to work by others. Liu et al.[15] proposed a framework where LLMs, fine-tuned with a prompt template, train a smaller student model via knowledge distillation to recommend medications by adjusting output probabilities. Moreover, Dou et al.[16] introduced ShennongGPT, an advanced LLM designed for medication guidance and adverse drug reaction prediction, using a two-phase training approach with drug databases and real-world patient data. Additionally, Ahmed et al.[17] leveraged the latest advancements in LLMs for domain-specific factual knowledge, presenting MED-Prompt — a novel prompt engineering framework designed to achieve curate and precise medicine predictions. The growing evidence supports using LLMs as a foundation for creating tailored digital twins in critical care settings.

The implications of digital twins extend beyond medication prediction. These systems can support dynamic, real-time simulations for monitoring and treatment planning, optimizing care in areas such as glycemic control, cardiovascular support, and mechanical ventilation.[18,19] Digital twins additionally support the automation of care workflows, improving operational efficiency while addressing human factors to ensure effective usability in the dynamic ICU setting. Successful implementation relies on achieving robust interoperability among medical devices and ongoing research aimed at refining model accuracy and adaptability to the complexities of critical care environments[18]. While it is conceivable that a digital twin can be created via crafting a prompt that explicitly states the purpose and the specifications of the digital twin, we find that supervision is critical to the creation of digital twins. The reasoning process of physicians incorporates (1) knowledge representation in medicine, typical of medical board exam questions and textbooks; (2) practice experience represented by years of clinical training at the bedside, and (3) familiarity with best practices within an individual ICU and health system. Incorporating the EHR via adapters brings in many years of practice experience recorded in the documentation of EHR care notes within an ICU specialty and adds unique knowledge to the LLM. The local practice behaviors are also represented in the EHR as health systems have their own unique formulary of medicines available and guidelines such as antibiograms to follow. Our use of LoRA across multiple medical specialties allows the flexible interchange of LLMs to represent multiple digital twins and allow easy updating of the model as more EHR data arrives.

Finally, our work highlights the limitations of current metrics for evaluating generative models. Balancing the need to account for semantic variability in language with the risk of being overly permissive necessitates using multiple metrics, as no single metric provides a comprehensive assessment of

generative performance. Exact-match metrics like accuracy and ROUGE-L are likely to underestimate performance, while soft-match metrics like BERTScore may overestimate it, particularly for long sequences. At the expense of a more stringent, straightforward evaluation such as single medication prediction, our task was more clinically relevant to use document-level input from the EHR and generate all relevant medications. Our iterative approach refined the evaluation framework, settling on clinically relevant tasks that better reflect real-world applications. This approach revealed sensitivities in model performance, such as the impact of repeated medication mentions and long sequences on prediction accuracy, emphasizing the need for robust and context-sensitive evaluation methods.

Conclusion: This study introduces a novel approach to building ICU-specific digital twins with LLMs, providing a foundation for advancing decision-support systems in critical care. By focusing on medication prediction, we demonstrated the importance of context-specific training and the value of supervised learning techniques like LoRA adapters in enhancing model efficiency and adaptability. These findings pave the way for extending digital twin capabilities to include multi-tasking functionalities, such as predicting procedures, lab orders, and other ICU events. Ultimately, our work represents an early but significant step toward developing interactive digital twins that integrate seamlessly with LLMs to support personalized, real-time decision-making in critical care settings. Further research is needed to expand these capabilities and optimize interactions between digital twins and healthcare providers.

METHODS

Data Corpus and Study Setting

MIMIC-III database is an open-access resource that provides de-identified health data from over 40,000 patients treated in intensive care units, including free-text admission notes that detail patient encounters.[20] For this study, we obtained the counts of discharge summaries in MIMIC-III across different ICU specialties (**Figure 1**) and selected the three most frequent ones: (1) Medical ICU; (2) Cardiothoracic ICU; and (3) Surgical ICU. Using a simple rule-based approach, we identified the five sections in the discharge summaries that are most relevant to clinical care during the hospital course: (1) Chief Complaint (CC); (2) Brief Hospital Course (BHC); (3) History of Present Illness (HPI); (4) Allergies; and (5) Discharge Diagnosis (DD). These sections are ubiquitous to discharge summaries at many health systems and contain the most detailed content of the hospital and ICU course, including the medications prescribed.

SparkNLP[21] was applied to automatically identify all medications mentioned in the discharge summaries. A critical care physician and clinical informaticist (MA) reviewed a random sampling of notes and confirmed high medication detection accuracy of the SparkNLP tool. For each note, the tagged medication mentions were masked with the special token [MEDICATION] (see **Figure 2** for a sample note). In addition to medication mentions, SparkNLP tagged other medication-related entities: DOSAGE, STRENGTH, ROUTE, FORM, FREQUENCY, and DURATION. To avoid revealing the identity of the masked medication, the additional entities were masked with a special token [MEDICATION_INFO]. The total number of unique medications for each ICU specialty, along with the top 10 medications and their frequencies, are shown in **Table 3**. The masked medication mentions were retained and used as the reference labels for the prediction targets. This preprocessing resulted in a dataset of 18,830 Medical ICU notes. To create our training and evaluation data, each note was paired (with all medication mentions masked) with a comma-separated list of medication mentions from that note that served as the prediction targets. The dataset was split into training (16,330), development (1500), and test (1000) sets. An additional four training sets were curated, each containing 4,118 notes, for the most common ICU specialties: Surgery, Medicine, and Cardiothoracic, and a random sample from all 3 ICU specialties.

Training set	# of Unique Medications	Top 10 Medications
Medical ICU (4K)	6,029	lasix (2673), vancomycin (2431), coumadin (2140), heparin (1940), metoprolol (1709), antibiotics (1680), insulin (1672), aspirin (1345), levofloxacin (1221), asa (1123)
Cardiothoracic ICU	2,138	coumadin (1707), amiodarone (1185), heparin (1180), antibiotics (657), lasix (627), beta blocker (617), propofol (611), vancomycin (606), diuretics (515), plavix (514)
Surgical ICU	3,078	antibiotics (1087), coumadin (912), heparin (886), lasix (820), vancomycin (785), zosyn (623), flagyl (591), dilaudid (552), penicillins (529), pressors (496)
All ICU Random Sample	4,331	coumadin (1594), lasix (1390), heparin (1348), vancomycin (1283), antibiotics (1133), insulin (773), metoprolol (772), amiodarone (704), aspirin (689), flagyl (591)
All Medical ICU (16,330)	14,811	lasix (10521), vancomycin (9613), coumadin (8492), heparin (7662), metoprolol (6993), insulin (6576), antibiotics (6419), aspirin (5324), levofloxacin (4653), ivf (4605)

Table 3: Unique Medication Counts and Frequencies of Top 10 Medications. The high number of unique medications mentions in discharge summaries makes our medication prediction task very challenging.

Instruction	Please list the medications in the text below that are masked with the [MEDICATION] token.
Input	**Chief Complaint:** 65 yo M with a history of severe COPD admitted for respiratory failure due to exacerbation. The patient was taking **[MEDICATION]**, **[MEDICATION]**, and **[MEDICATION]** prior to admission.
	History of Present Illness: This 65-year-old man has a long-standing history of severe COPD, having experienced multiple hospitalizations over the past decade for exacerbations requiring intensive care. In early August 2023, he experienced progressively worsening dyspnea and increased sputum production despite regular use of home nebulizer treatments and oral corticosteroids. Given his failure to respond to outpatient therapy and his deteriorating oxygenation (SaO2 dropping to 85% on room air), he was admitted to the ICU on 08/26/2023. Initial management included high-flow nasal oxygen and escalation to non-invasive ventilation, but his condition necessitated intubation. His hospital course involved repeated bronchoscopies revealing significant secretions and mucous plugging, necessitating a percutaneous tracheostomy on 09/02/2023. His medications on admission included **[MEDICATION]**, **[MEDICATION]**, and **[MEDICATION]**.
	Brief Hospital Course: 65 y/o male admitted on 08/26/2023 due to severe COPD exacerbation leading to respiratory failure. Initial management involved high-flow nasal oxygen and non-invasive ventilation but required intubation on 08/27/2023. Bronchoscopies on 08/28 and 08/30 showed copious secretions necessitating a percutaneous tracheostomy on 09/02. Post-procedure complications included transient hypotension managed with **[MEDICATION_INFO]** **[MEDICATION]** and **[MEDICATION]**. Developed a secondary infection with MRSA treated with **[MEDICATION_INFO]** **[MEDICATION]**. Pain management involved **[MEDICATION_INFO]** **[MEDICATION]**, later managed with **[MEDICATION]** due to effective tracheostomy secretion management. Physical therapy initiated early, leading to improved mobility. Cardiovascular management included controlled hypertension through **[MEDICATION_INFO]** **[MEDICATION]** and a stable heart rate with no ischemic changes

	noted. Addressed hyperlipidemia and hyperglycemia through **[MEDICATION_INFO] [MEDICATION]** and **[MEDICATION_INFO] [MEDICATION_INFO]**. Nutritional support delivered via parenteral nutrition transitioning to enteral feeding after tracheostomy optimization. **Allergies: [MEDICATION], [MEDICATION]** **Discharge Diagnosis:** - COPD exacerbation - Respiratory failure requiring tracheostomy - Hypertension - Type 2 Diabetes Mellitus - Coronary Artery Disease - MRSA pneumonia - Hyperlipidemia - Peptic ulcer disease
Output	Albuterol, Ipratropium, Prednisone, Ipratropium, Prednisone, fluids, vasopressors, Vancomycin, Dilaudid PCA, Acetaminophen, Hydralazine, Statin therapy, Insulin, Penicillin, Morphine

Fig. 2: Sample synthetic note with masked medication mentions like the notes used for supervised fine-tuning. We train LoRA adapters using thousands of similar notes to predict the identity of the masked medication mentions.

Experiments

Supervised fine-tuning (SFT): Llama3 LLM (llama3-8B-Instruct)[22] was fined tuned using a LoRA adapter to accept notes from a single type of ICU and predict all medication mentions that were masked. This method leveraged low-rank factorization, reducing the number of trainable parameters and mitigating overfitting risks. We performed an extensive hyperparameter search to optimize the training and LoRA adapter parameters to ensure stable and efficient model training. After completing the model selection step using our validation set, a final evaluation was conducted using the test set, which also consisted of Medical ICU notes, and reported the results in **Table 1**. The average training runtime across the five corpora was approximately 5 hours, utilizing 8 NVIDIA RTX A6000 GPUs, each with 48 GB of RAM. The longest runtime was observed for the "All Medical ICU [16,330 notes]" dataset, which took approximately 13 hours.

Zero-Shot Learning: To understand better how well the supervised models captured the preferences of ICU physicians from a given specialty of ICU, the fine-tuned models were compared to a model that did not have access to ICU-specific training data using zero-shot learning. Using the same LLM as in SFT, the prompt for zero-shot was refined using the development set. The final prompt is shown in **Figure 5**. Sequential numbering was applied to the masked medication entities within each clinical note to assist the model in predicting the correct number of medications. Additionally, we tested whether zero-shot learning can induce ICU-specific behavior by prepending to the prompt the sentence "You are a [ICU_specialty] ICU physician.", where [ICU_specialty] is Medicine, Cardiothoracic, or Surgery. Following the generation of results, we conducted a post-processing step to remove any extraneous elements, thereby refining the final output for more accurate evaluation.

Prompt	Based on the clinical note provided, predict the medication names for all placeholders labeled as [MEDICATION_X], where X is a sequential number.
Input	**Brief Hospital Course:** 65 y/o male admitted on 08/26/2023 due to severe COPD exacerbation leading to respiratory failure. Initial management involved high-flow nasal oxygen and non-invasive ventilation but required intubation on 08/27/2023. Bronchoscopies on 08/28 and 08/30 showed copious secretions necessitating a

percutaneous tracheostomy on 09/02. Post-procedure complications included transient hypotension managed with [MEDICATION_INFO] **[MEDICATION_7]** and **[MEDICATION_8]**. Developed a secondary infection with MRSA treated with [MEDICATION_INFO] **[MEDICATION_9]**. Pain management involved [MEDICATION_INFO] **[MEDICATION_10]**, later managed with **[MEDICATION_11]** due to effective tracheostomy secretion management. Physical therapy initiated early, leading to improved mobility. Cardiovascular management included controlled hypertension through [MEDICATION_INFO] **[MEDICATION_12]** and a stable heart rate with no ischemic changes noted. Addressed hyperlipidemia and hyperglycemia through [MEDICATION_INFO] **[MEDICATION_13]** and [MEDICATION_INFO] **[MEDICATION_14]** [MEDICATION_INFO]. Nutritional support delivered via parenteral nutrition transitioning to enteral feeding after tracheostomy optimization.

Fig. 5: A sample synthetic note, similar to the notes used for zero-shot learning. An LLM is given a prompt that asks to identify the masked medication mentions.

Evaluation metrics: Evaluating the performance of the models presented several challenges. First, some medication names were synonyms and were used interchangeably (e.g., "ibuprofen" and "advil"). Next, medication classes were frequently substituted for specific medication names (e.g., "NSAID" instead of "aspirin"). Finally, there was no guarantee that the length of the generated sequence of medication mentions would match the length of the reference sequence. To address these complexities, we employed multiple evaluation metrics. First, the exact match accuracy was calculated by assuming a one-to-one alignment between the generated and reference sequences. The assumption of one-to-one match was violated in at least 20% of the cases and thus the exact match accuracy likely underestimated the model's performance. The accuracy was calculated as the proportion of exact matches between the predicted medication sequences and the ground truth sequences. To get a better estimate of the model performance, we opted for two metrics that are routinely used for evaluating generative models: BERTScore and ROUGE-L. BERTScore is a soft match metric, with SapBERT[23] as the base encoder. We chose SapBERT because it was trained on the Unified Medical Language System (UMLS)[24] and had a strong correlation with human judgments.[25] In our experiments, we found that BERTScore exhibited a high variance between longer and shorter sequences.[26] This limitation highlighted the need for complementary metrics; therefore, we also reported ROUGE-L, which provided a more balanced assessment by focusing on the longest common subsequence, thereby mitigating the problem of aligning sequences. Our primary evaluation metric was the exact match accuracy score, and we used it for model selection. We compute 95% confidence intervals using bootstrap resampling with 10,000 iterations.

DATA AVAILABILTY

It is available upon request.

CODE AVAILABILITY

It is available upon request.

REFERENCE:

1.   Grieves, M. & Vickers, J. Digital twin: Mitigating unpredictable, undesirable emergent behavior in complex systems. *Transdisciplinary perspectives on complex systems: New findings and approaches* 85–113 (2017).

2. Pellegrino, G., Gervasi, M., Angelelli, M. & Corallo, A. A Conceptual Framework for Digital Twin in Healthcare: Evidence from a Systematic Meta-Review. *Information Systems Frontiers* 1–26 (2024).

3. Vallée, A. Digital twin for healthcare systems. *Front Digit Health* **5**, 1253050 (2023).

4. Schwartz, S. M., Wildenhaus, K., Bucher, A. & Byrd, B. Digital twins and the emerging science of self: implications for digital health experience design and "small" data. *Front Comput Sci* **2**, 31 (2020).

5. Johnson, A. E. W. *et al.* MIMIC-III, a freely accessible critical care database. *Sci Data* **3**, 1–9 (2016).

6. Meta, A. I. Introducing meta llama 3: The most capable openly available llm to date, 2024. *URL https://ai. meta. com/blog/meta-llama-3/. Accessed on April* **26**, (2024).

7. Hu, E. J. *et al.* Lora: Low-rank adaptation of large language models. *arXiv preprint arXiv:2106.09685* (2021).

8. Zhang, T., Kishore, V., Wu, F., Weinberger, K. Q. & Artzi, Y. Bertscore: Evaluating text generation with bert. *arXiv preprint arXiv:1904.09675* (2019).

9. Lin, C.-Y. Rouge: A package for automatic evaluation of summaries. in *Text summarization branches out* 74–81 (2004).

10. Venkatesh, K. P., Raza, M. M. & Kvedar, J. C. Health digital twins as tools for precision medicine: Considerations for computation, implementation, and regulation. *NPJ Digit Med* **5**, 150 (2022).

11. Achiam, J. *et al.* Gpt-4 technical report. *arXiv preprint arXiv:2303.08774* (2023).

12. Touvron, H. *et al.* Llama: Open and efficient foundation language models. *arXiv preprint arXiv:2302.13971* (2023).

13. Singhal, K. *et al.* Large language models encode clinical knowledge. *Nature* **620**, 172–180 (2023).

14. Nazi, Z. Al & Peng, W. Large language models in healthcare and medical domain: A review. in *Informatics* vol. 11 57 (2024).

15. Liu, Q. *et al.* Large language model distilling medication recommendation model. *arXiv preprint arXiv:2402.02803* (2024).

16. Dou, Y. *et al.* ShennongGPT: A Tuning Chinese LLM for Medication Guidance. in *2023 IEEE International Conference on Medical Artificial Intelligence (MedAI)* 67–72 (2023).

17. Ahmed, A., Zeng, X., Xi, R., Hou, M. & Shah, S. A. MED-Prompt: A novel prompt engineering framework for medicine prediction on free-text clinical notes. *Journal of King Saud University-Computer and Information Sciences* **36**, 101933 (2024).

18. Geoffrey Chase, J. *et al.* Digital twins and automation of care in the intensive care unit. *Cyber–Physical–Human Systems: Fundamentals and Applications* 457–489 (2023).

19. Thangaraj, P. M., Benson, S. H., Oikonomou, E. K., Asselbergs, F. W. & Khera, R. Cardiovascular care with digital twin technology in the era of generative artificial intelligence. *Eur Heart J* ehae619 (2024).

20. Aden, I., Child, C. H. T. & Reyes-Aldasoro, C. C. International Classification of Diseases Prediction from MIMIIC-III Clinical Text Using Pre-Trained ClinicalBERT and NLP Deep Learning Models Achieving State of the Art. *Big Data and Cognitive Computing* **8**, 47 (2024).

21. Kocaman, V. & Talby, D. Improving clinical document understanding on COVID-19 research with spark NLP. *arXiv preprint arXiv:2012.04005* (2020).

22. Dubey, A. *et al.* The llama 3 herd of models. *arXiv preprint arXiv:2407.21783* (2024).

23. Liu, F., Shareghi, E., Meng, Z., Basaldella, M. & Collier, N. Self-alignment pretraining for biomedical entity representations. *arXiv preprint arXiv:2010.11784* (2020).

24. Bodenreider, O. The unified medical language system (UMLS): integrating biomedical terminology. *Nucleic Acids Res* **32**, D267–D270 (2004).

25. Croxford, E. *et al.* Development of a Human Evaluation Framework and Correlation with Automated Metrics for Natural Language Generation of Medical Diagnoses. *medRxiv* (2024).

26. Guo, X. & Vosoughi, S. Length does matter: Summary length can bias summarization metrics. in *Proceedings of the 2023 Conference on Empirical Methods in Natural Language Processing* 15869–15879 (2023).

APPENDIX

Hyperparameter	Value
Cut-Off	2048, 3072, **4096**
Top-P	0.95
Top-K	20
LoRA Alpha	16, **32**
LoRA Rank	8, **16**
Learning Rate	5×10^{-5}
Epochs	2,4,**8**, 16
Temperature	0.0001
Quantization Bit	**4**, 8
Length Penalty	1.2

Table 4: Evaluated hyperparameter values with the optimal values highlighted. The optimal values, determined through extensive evaluation, are bolded to indicate the best configuration for model training.