The yield on re-interpretation of genetic variants in pediatric cardiomyopathy

Takanori Suzuki, MD, PhD[1,] Robert Lesurf, PhD[1], Raj Akilen, PhD[1], Xiaoqiao Xu, MHSc[2],

Rebekah Jobling, MD[3-5], Laura Zahavich, MSc[6], Seema Mital, MD[1,3,7*]

[1]Genetics and Genome Biology Program, The Hospital for Sick Children, Toronto, Ontario,

Canada

[2]The Centre for Computational Medicine, The Hospital for Sick Children, Toronto,

Ontario, Canada

[3]Ted Rogers Centre for Heart Research, The Hospital for Sick Children, Toronto, ON,

Canada

[4]Clinical and Metabolic Genetics, Department of Pediatrics, The Hospital for Sick

Children, Toronto, ON, Canada

[5]Genome Diagnostics, Department of Pediatric Laboratory Medicine, The Hospital for Sick

Children, Toronto, ON, Canada

[6]Department of Genetic Counselling, Hospital for Sick Children, University of Toronto,

Toronto, Ontario, Canada

[7]Division of Cardiology, Department of Pediatrics, The Hospital for Sick Children,

University of Toronto, Toronto, Ontario, Canada

Short title: Variant reevalution in pediatric cardiomyopathy

Corresponding Author:

Seema Mital, MD

Hospital for Sick Children,

555 University Avenue,

Toronto, Ontario M5G 1X8, Canada.

Tel: 416-813-7418; Fax: 416-813-5857

*Email: seema.mital@sickkids.ca

Twitter: @seema_mital

Linkedin: https://www.linkedin.com/in/seema-mital-970989b/

Word count: 3477

ABSTRACT

Background: Regular re-interpretation of disease-causing genetic variants is recommended to

determine if variant pathogenicity changes as new information becomes available. Our aim was to

determine the yield on systematic re-evaluation of genetic variants in pediatric patients with

cardiomyopathy.

Methods: The study cohort included 227 unrelated pediatric patients with cardiomyopathy enrolled in

the Heart Centre Biobank who harbored a pathogenic/likely pathogenic (P/LP) variant and/or a variant

of uncertain significance (VUS) on clinical genetic testing between 2005-2022. Variant pathogenicity

was re-evaluated using the American College of Medical Genetics and Genomics (ACMG) guidelines.

The re-evaluation included updates to ClinGen gene curation, ClinVar variant interpretation, population

allele frequency, in silico tools for variant deleteriousness, segregation with disease or de novo status,

and genotype-phenotype evaluation.

Results: Re-evaluation was performed in 382 variants (110 P/LP, 272 VUS) reported in the study cohort

of 227 patients. 12 of 110 P/LP (10.9%) variants were downgraded to VUS in 14 patients. Leading

criteria were high population allele frequency and variant not located in mutational hotspot or critical

functional gene domain. 37 of 272 VUS (13.6%) were upgraded to P/LP in 35 patients. Leading criteria

were variant in mutational hotspot for gene, variant interpreted as deleterious using in silico prediction

and novel missense change at an amino acid residue where a different missense change was previously

determined to be pathogenic. Overall, the reclassification of 49 variants affected 49 of 227 (21.6%)

patients. Of note, only 8 reclassified variants had been reported back by the clinical genetic testing

laboratory.

Conclusions: One in five patients with cardiomyopathy had a clinically relevant change from the

original variant classification on systematic re-evaluation. This has implications for clinical screening

and cascade genetic testing of family members of affected patients. These findings underscore the

importance of regular variant re-interpretation on follow-up in cardiomyopathy patients.

Keywords: Cardiomyopathy, genetic testing, pathogenic variants, variants of uncertain significance,

variant reinterpretation

INTRODUCTION

Cardiomyopathy is primarily a genetic disorder and a common cause of childhood heart failure.

Cardiomyopathy phenotypes include arrhythmogenic cardiomyopathy (ACM), dilated cardiomyopathy

(DCM), hypertrophic cardiomyopathy (HCM), restrictive cardiomyopathy (RCM), and left ventricular

non-compaction cardiomyopathy (LVNC). Dozens of genes are implicated in causing the disease, with

most cases being autosomal dominant and caused by a rare DNA variant [1], with considerable overlap

in the genes associated with different cardiomyopathy subtypes [1, 2].

The genotype of a patient informs cascade clinical and genetic screening of first-degree relatives. When

a patient is genotype-positive i.e., harbors a pathogenic/likely pathogenic (P/LP) variant in a

cardiomyopathy gene, family members are offered cascade testing for the variant, and only genotype-

positive family members require ongoing echocardiographic screening for cardiomyopathy. When a

patient is genotype-negative i.e., does not harbor a P/LP variant, then all family members are

recommended serial echocardiographic screening. A previous study in HCM reported that, among 1,361

probands with non-benign variants, 917 were P/LP (67.3%) and 444 (32.6%) were variants of uncertain

significance (VUS), the latter reflecting incomplete or conflicting evidence for pathogenicity [3].

However, variant interpretation can change over time as new knowledge, additional functional

validation data, and more accurate computational prediction tools emerge. A change in variant

interpretation can change family screening recommendations. The 2024 American Heart Association &

American College of Cardiology guidelines for HCM emphasize the importance of periodic variant re-

evaluation [4]. To improve the efficiency of variant re-evaluation, clinicians are advised to document the

date when the genetic test results were initially assessed and to consider any new medical and scientific

information that has become available since that time [5]. Small pediatric cardiomyopathy cohort studies

have reported variant reclassification rates ranging from 8.9%-10.3% **[6,7]**. Here we performed a

systematic reevaluation of variants in a larger cohort of 227 unrelated pediatric cardiomyopathy cases to

identify the frequency of and criteria for clinically 'actionable' variant reclassification on follow-up

using contemporary criteria.

METHODS

Study cohort

The study cohort included 227 unrelated patients with primary cardiomyopathy <18 years old, enrolled

through the Heart Centre Biobank Registry at the Hospital for Sick Children (Toronto, Ontario, Canada)

(**Table 1**). Patients who harbored a P/LP variant or a VUS on clinical genetic testing were included.

Patients with secondary cardiomyopathy caused by chromosomal malformations, neuromuscular

disorders, mitochondrial or metabolic disorders, congenital heart defects, or reversible causes were

excluded. Patient demographics, diagnosis, type of clinical testing (panel, exome), year of clinical

genetic testing, the original (and updated) variant classification by the genetic testing laboratory were

captured. The study was approved by the Institutional Research Ethics Board of the Hospital for Sick

Children. Written informed consent was obtained from all patients and/or their parents/legal guardians

and the study protocol adhered to the Declaration of Helsinki.

Re-evaluation of variant pathogenicity

Variants identified as P/LP or VUS on clinical genetic testing were reinterpreted for pathogenicity using

the American College of Medical Genetics and Genomics/Association for Molecular Pathology

(ACMG/AMP) guidelines for interpretation of variant pathogenicity. Using these criteria, a variant can

be interpreted as pathogenic, likely pathogenic, a VUS, likely benign (LB), or benign (B) **[8, 9]**. For

variants with multiple isoforms, the Matched Annotation from NCBI and EMBL-EBI transcript i.e.,

MANE Select and MANE Plus Clinical) was used to annotate and evaluate variant effects in

standardized and clinically relevant gene transcripts. A clinically actionable variant reclassification was

defined as either the upgrading of a VUS to P/LP, or the downgrading of a P/LP variant to a non-

pathogenic variant i.e., VUS or B/LB variant.

Several criteria were applied for variant re-interpretation including confirmation of patient phenotype,

the current association between gene and cardiomyopathy phenotype on the most recent ClinGen

version 11 curation **[10]**, updated ClinVar (2024-01-29 Web Release) records, literature evidence,

control population allele frequencies, and prediction scores derived from computational tools. When

available, the results of clinical genetic tests from parents or siblings were utilized to ascertain *de novo*

status and to perform segregation analysis for a familial variant. P/LP variants downgraded to

VUS/LB/B, or VUS upgraded to P/LP were deemed clinically actionable. Reclassifications were

verified by Return of Research Results committee comprising pediatric cardiologists, clinical

geneticists, genetic counselors, and bioinformaticians **[11].**

A detailed approach to variant reinterpretation is described below. Of note, all eligible variants reported

on clinical testing were single nucleotide variants and small insertions-deletions.

1) PVS1 criterion was applied to loss-of-function variants in haploinsufficiency-intolerant genes

using gnomAD 4.1.0, defined as a probability of loss-of-function intolerant (pLI) score $\geq$0.9.

PVS1 criterion was not applied to *MYH7* variants as per ClinGen guidelines because the

contribution of loss-of-function variants in this gene to inherited cardiomyopathy remains

incompletely understood **[12]**. *TTN* variants were only considered pathogenic if they were

protein-truncating variants (frameshift, nonsense, canonical splice site) predominantly in the A-

band and I-band regions and isoforms **[13, 14].**

2) The PS2 criterion was applied when the variant was confirmed through genetic testing to be *de*

novo with parentage confirmed and no family history of cardiomyopathy.

3) The PS4 criterion of higher variant burden in cases compared to controls could not be applied

since there was insufficient in-house data for burden analyses.

4) The PM1 criterion was applied if the variant was located in a mutational hotspot or critical

functional domain for the gene. This was determined using DiscoVari, the Cardiomyopathy

Variant Curation Expert Panel from ClinGen, and/or literature evidence **[15, 16]**. For *MYH7*, the

relevant transcripts analyzed were ENST00000355349 and NM_000257.4, covering codons 167

to 931. In the case of *MYBPC3*, the analysis focused on transcripts ENST00000545968 and

NM_000256.3, specifically targeting codons 485 to 502 and 1248 to 1266. For *TNNI3*, the key

transcripts used were ENST00000344887 and NM_000363.5, covering codons 141 to 209. For

TNNT2, the transcripts ENST00000367318 (codons 79 to 179), and

ENST00000656932.1/NM_001276345.2 (codons 89 to 189) were included in the analysis.

5) PM2 criterion was applied if population variant allele frequency in gnomAD 4.1.0 was <0.0001

for autosomal dominant and <0.01 for recessive inheritance, while accounting for Grpmax

filtering allele frequencies (95% confidence) as ancestry-specific allele frequencies **[17]**.

6) The PM5 criterion was applied when the variant incorporated a novel missense change at an

amino acid residue where a different missense change is determined to be pathogenic in ClinVar

or Leiden Open Variation Database v.3.0 supported by literature **[9, 18]**.

7) The PP1 criterion was applied when the variant segregated with disease amongst family

members. According to the ClinGen guidelines, in cardiomyopathies with reduced penetrance,

PP1 is applied depending on the number of affected individuals demonstrating co-segregation

with the variant (not including the proband). Specifically, PP1 supporting is applied when co-

segregation is observed in a single affected family member, PP1 moderate for 2-3 affected

members, and PP1 strong for ≥4 affected members **[19]**.

8) The PP2 criterion was defined as a missense variant in a gene with a low rate of benign missense

variation, where missense variants are a known mechanism of disease. Based on ClinGen

recommendations, PP2 was applied only to variants in *TPM1* among the sarcomere genes listed

in the Criteria Specification Registry for cardiomyopathy **[16]**.

9) The PP3 criterion for missense variants which requires multiple lines of computational evidence

supporting a deleterious effect on the gene was applied when all three pathogenicity scores

(CADD v1.7, REVEL, and AlphaMissense.v2023.hg38) reached pre-defined thresholds [**20-22**]

These thresholds were adopted from the ClinGen recommendations for CADD (>25.3) and

REVEL (>0.644) **[23]**. As the threshold for AlphaMissense was not specified in this

recommendation, it was derived from the original publication as a score >0.564 **[21]**.

Statistical Analysis

Frequency of variant reclassification was compared using Fisher exact test by year of clinical testing

(before vs. after 2015 i.e., publication of ACMG guidelines), cardiomyopathy phenotype (HCM vs.

DCM), ancestry (European vs. non-European), and sex (male vs. female). A nominal p-value <0.05 was

considered significant.

RESULTS

Study cohort characteristics

The baseline characteristics of the study cohort (n=227) are summarized in **Table 1**. Among the

participants, 133 (59%) were male; 64% were of European descent, 25% were of Asian descent, 10%

were of African descent, and 1% were of mixed descent or other ancestries. Of 227 participants, 203

underwent one or more gene panel testing, 4 underwent whole exome sequencing, and 20 underwent

other types of genetic tests. The 227 patients harbored a total of 382 variants (110 P/LP, 272 VUS)

(**Table S1**). 105 participants (46.3%) harbored at least one P/LP variant, 122 (53.7%) harbored only

VUS, and 34 (15%) harbored a combination of P/LP variants and VUS. Three variants that were found

in two families each - *MYBPC3* (c.442G>A, p.Gly148Arg), *MYH7* (c.1759G>A, p.Asp587Asn) and

LMNA (c.868G>A, p.Glu290Lys) were reported differently for each family by the clinical laboratory

i.e., P/LP for one family and VUS in another family.

Reinterpretation of variant pathogenicity

Reclassification of P/LP variants: After reinterpretation, 12 of 110 P/LP (10.9%) variants were

downgraded to VUS in 14 of 227 patients (6.2%) **(Table 2)**. Genes affected by the downgrade included

MYH7 (5 variants), *MYBPC3* (4 variants), and one variant each in *LMNA*, *DSG2*, and *RRAGD*. Notably,

the *MYBPC3* c.3628-41_3628-17del (Δ25bp) variant was identified in six South Asian cases. The

criteria primarily contributing to downgrading of P/LP variants were an allele frequency higher than

expected in gnomAD (n=7), the variant not being in a mutational hotspot or critical functional gene

domain (n=5), the variant not being predicted as deleterious by *in silico* tools (n=4), the variant recently

reported as VUS in ClinVar (n=3), and failure to meet *MYH7*-specific ACMG/AMP criteria (n=1)

(**Figure 1a**).

Reclassification of VUS: 37 of 272 VUS (13.6%) were upgraded to P/LP in 35 of 227 patients (15.4%)

(**Table 3**). Genes affected by the upgrade included *ACTC1* (1 variant), *ACTN2* (2 variants), *LMNA* (1

variant), *MYBPC3* (2 variants), *MYH7* (13 variants), *NEXN* (3 variants), *PLN* (1 variant), *TNNC1* (1

variant), *TNNI3* (2 variants), *TNNT2* (4 variants), *TPM1* (3 variants), *TTN* (1 variant), and *VCL* (2

variants). The primary criteria contributing to upgrading of VUS to P/LP were the variant location in a

mutational hotspot for the gene (n=23), variant interpretation as deleterious by modern *in silico*

prediction tools (n=19), a novel missense change occurring at an amino acid residue where a different

pathogenic missense change had been observed before (n=11), a change in the gene association with

cardiomyopathy according to ClinGen (n=8), published experimental findings suggesting that the variant

affects gene function (n=7), co-segregation with disease in multiple affected family members (n=6), the

variant being *de novo* (n=4), the variant being recently reported as pathogenic in ClinVar (n=2), and the

finding of a different variant that results in the same amino acid change being recently reported as

pathogenic in ClinVar (n=1) (**Figure 1b**). 47 VUS were downgraded to B/LB variants, but these were

not further evaluated since they do not reflect a clinically actionable change. An overview of variant

reclassification frequency is provided in **Figures 2a**.

Patient level reclassification: The variant reclassifications described above affected 49 of 227 patients

(21.6%) (**Figure 2b**). 15.5% of patients with only P/LP variants had one or more variants downgraded to

a VUS. 23.8% of patients with only VUS had one or more variants upgraded to P/LP. 26.5% of patients

with a combination of P/LP variants and VUS had either a downgrade of their P/LP variants and/or

upgrade of their VUS resulting in a change in affected gene.

Of note, only 8 reclassified variants (*MYBPC3* c.3628-41_3628-17del, *MYH7* c.1759G>A,

p.Asp587Asn, *MYH7*　c.3667G>A p.Glu1223Lys, *MYH7* c.4276G>A p.Glu1426Lys, *MYH7*

c.2683C>G p.Gln895Glu, *NEXN* c.1473+1G>T, *ACTN2* c.784-2A>G and *MYBPC3* c.442G>A,

p.Gly148Arg) in 12 patients had been previously reported back by the clinical genetic testing laboratory

at the time of our study. Among these variants, *MYH7* c.3667G>A p.Glu1223Lys reclassification was

initiated by the clinical laboratory, and *ACTN2* c.784-2A>G reclassification was prompted by the

patient's clinical team. For the remaining variants, the reasons triggering reclassification were not

available.

Frequency of variant reclassification by subgroups

The year of clinical testing was known for 358 variants. In this group, frequency of variant

reclassification did not vary by year of clinical testing i.e. before or after 2015 (**Figure 3a and 3b).**

Additionally, there was no difference in frequency of variant reclassification by cardiomyopathy

phenotype (HCM vs DCM), ancestry (European vs non-European descent), or by sex (male vs female)

(p > 0.05) (**Figure 4**).

DISCUSSION

We conducted a systematic reinterpretation of variants identified on clinical genetic testing in a cohort

of 227 pediatric cardiomyopathy patients. The frequency of clinically actionable variant reclassification

was 12.8% and affected 49 (21.6%) patients. This included downgrading of 10.9% P/LP variants to

VUS and upgrading of 13.6% VUS to P/LP variants. Previous pediatric studies have reported a rate of

variant reclassification ranging from 8.9%-10.3% in 71-128 patients **[6, 7]**. Our study with a larger

cohort size reveals a somewhat higher reclassification rate and reinforces the importance of routine

reinterpretation of variants on follow-up.

A detailed description of evaluation of variant pathogenicity is important to guide confidence in

pathogenicity predictions **[24]**. In our study, we examined the criteria predominantly affecting the

change in pathogenicity of the variants based on the ACMG evaluation methodology. The most common

criterion for reclassification of a variant from P/LP to VUS was a higher-than-expected population allele

frequency. The increase in genome sequencing including those of previously underrepresented

populations has enabled more accurate assessment of variant frequencies in the general population **[25,**

26]. Further, population-specific databases are increasingly being utilized **[27]**, which continues to

improve the accuracy of variant frequency assessments across different ancestries and will continue to

enhance the precision of pathogenicity evaluation with time. The most common criterion for

reclassification of a variant from VUS to P/LP was location of the variant in a mutational hotspot for the

gene. Often the evaluation of a hotspot involves determining whether a variant is located within a

constitutive domain of the protein or assessing the role of the domain in which the variant is located,

based on literature and experimental data. In this study, reclassification was carried out using the

hotspots defined by DiscoVari, the Cardiomyopathy Variant Curation Expert Panel from ClinGen, and

published literature where hotspots had been identified [**15, 16**]. The number of genes with defined

hotspots is still limited, and future efforts may enable the accurate assessment of variant hotspots in

additional genes. Segregation analysis of the variant in family members and reevaluation of inheritance

contributed to a change in pathogenicity in 10 out of 81 variants (12.3%). This consitutes a substantial

proportion, and as recommended in the ACMG statement **[5]**, reclassification should include a

reassessment of familial phenotype and variant inheritance on follow-up.

In our study, the *MYBPC3* c.3628-41_3628-17del (Δ25bp) intronic deletion was reclassified from

pathogenic to a VUS. This 25-base pair deletion is located within intron 32 of the *MYBPC3* gene and is

observed in 4% to 8% of individuals of South Asian ancestry [28, 29]. Given its allele frequency which

exceeds the incidence of pediatric cardiomyopathy, this variant was reclassified to a VUS (BS1

criterion). It is now known that this variant alone without *MYBPC3* c.1224-52G>A as part of a

haplotype is not sufficient to produce a phenotype [30], although in the adult population, it has been

associated with myocardial diastolic dysfunction [31].

We identified several recurrent VUS i.e., the same VUS affecting multiple individuals, that have the

potential to be reclassified in the future with new evidence. For example, *ACTC1* c.372C>G,

p.Ile124Met and *MYH7* c.1597A>G, p.Ile533Val VUS would be classified as P/LP if variant burden in

cases exceeds that in controls (PS4 criterion) which may transpire in the future as more patients are

sequenced. Therefore, it is crucial to flag variants with the potential for reclassification and to document

the criteria that have the potential to change over time. It is equally important to revisit not just variant

pathogenicity but also the availability of updated gene testing panels that may capture disease-associated

genes that were not captured at the time of previous genetic testing. The ClinGen curation of the

literature and new evidence serve an important role in informing the size and nature of gene panels.

There were no significant differences in the proportion of patients reclassified before and after 2015,

when the ACMG/AMP guidelines were first published. Of note, 2 of 11 variants (18%) identified on

clinical testing in 2022, the most recent year included, showed a change in pathogenicity within 2 years

of clinical reporting. The downgrading of *RRAGD* c.289_303del p.Thr97_Cys101del from LP to VUS

was based on the application of in silico tools (PP3), which indicated that the variant was not

deleterious. Conversely, the upgrading from VUS to LP of *MYH7* c.1423C>A p.Gln475Lys was because

of evidence of a mutational hotspot at the codon (PM1) and a differential missense change at the same

codon (*MYH7* c.1425G>T p.Gln475His), which had been previously reported as pathogenic (PM5) in

2022, further supported by in silico analysis using AlphaMissense issued in 2023 (PP3).   Hence, it is

important to reevaluate variants every few years due to rapid advances in genomic characterization of

cases and controls. In this regard, there remains a gap in regular reinterpretation of genetic variants by

testing labs. Only 8 of the 49 variants reclassified by us were reported back to the clinical team by the

original testing laboratory. The ACMG and the other reports recommend that clinical laboratories

reevaluate variants at least every three to five years and the costs of recontacting physicians can be built

into original testing costs **[5, 32, 33]**. The Canadian College of Medical Geneticists recommends that

laboratories reanalyze variants when a healthcare provider initiates the request [**34**]. This is important

because laboratories do not always have access to updated family phenotype information. This

emphasizes that periodic reinterpretation should, where feasible, be built into the ongoing clinical care

of patients by their healthcare providers. The ACMG in its most recent policy statement emphasizes that

"re-contact is a shared responsibility" **[35]**, indicating that both clinicians and clinical laboratories

should take on an increasing role in assessing and communicating variant reinterpretations to patients. In

addition, patients and families should be advised to notify their physician if family genotype or

phenotype information changes on follow-up.

It is important to acknowledge as a limitation that we were not privy to the exact rationale and criteria

for the original variant classification used by the clinical laboratory (especially for ACMG criteria that

were only published in 2015) or their threshold for recontacting physicians with updated

reclassifications.

Clinical significance

The findings of clinically actionable change in variant pathogenicity in 21.6% patients is significant. The

14 patients in whom P/LP variants were downgraded will require all previously genotype-negative

family members to return for ongoing clinical screening. For the 35 patients in whom VUS were

upgraded to P/LP, family members can now be offered cascade genetic testing to determine who is at

risk and needs ongoing clinical surveillance and who is genotype-negative and can be discharged from

follow-up. Providing timely feedback to patients regarding changes in variant classification can lead to

changes in follow-up protocols and may improve prognosis for other family members. A reevaluation

system for a broad range of variants, including those that emerge as children transition into adulthood is

essential given the long-term prognosis of pediatric patients.

Conclusions

One in 5 patients with childhood onset cardiomyopathy had a clinically actionable change in variant

pathogenicity on follow-up that impacts clinical and genetic family screening recommendations. This

highlights the importance of serial reevaluation of variant pathogenicity by clinical providers as well as

by clinical genetic testing laboratories.

ACKNOWLEDGEMENTS

We acknowledge the patients and families participating in the Labatt Family Heart Centre Biobank at the Hospital for Sick Children for access to study data.

SOURCES OF FUNDING

This project received support from the Canadian Institutes of Health Research's Canadian Heart Function Alliance Network Grant (HFN 181992) (SM), the Ted Rogers Centre for Heart Research (SM, RJ), and the Heart & Stroke Foundation of Canada / Robert M Freedom Chair of Cardiovascular Science (SM). TS was funded by the Japan Heart Foundation Research Grant, Canadian Heart Function Alliance, and Philip Witchel Research Fellowship in Heart Failure at the Hospital for Sick Children.

DISCLOSURES

SM is on the Scientific Advisory Board of Bristol Myers Squibb, Rocket Pharmaceuticals, and Tenaya Therapeutics. The remaining authors declare that they have no competing interests.

ETHICS APPROVAL

The study was approved by the Institutional Research Ethics Board of The Hospital for Sick Children. Written informed consent to participate was obtained from all patients and/or their parents/legal guardians. The study protocol adhered to the Declaration of Helsinki.

DATA AVAILABILITY

De-identified data analyzed in this study are available in the main and supplemental tables, and additional data are available from the corresponding author on reasonable request.

AUTHOR CONTRIBUTIONS

TS, RL, and SM conceptualized and planned the study, performed data analysis, drafted the initial

manuscript, and made significant revisions. RA collected the clinical data. PX, LZ and RJ assisted with

bioinformatics and clinical data analysis. SM secured funding for the project. All authors approved the

final manuscript as submitted and agree to be accountable for all aspects of the work.

REFERENCES

1. Watkins H, Ashrafian H, Redwood C. Inherited cardiomyopathies. N Engl J Med. 2011;364:1643–1656.

2. Morimoto S. Sarcomeric proteins and inherited cardiomyopathies. Cardiovasc Res. 2008;77:659–666.

3. Alfares AA, Kelly MA, McDermott G, Funke BH, Lebo MS, Baxter SB, Shen J, McLaughlin HM, Clark EH, Babb LJ, et al. Results of clinical genetic testing of 2,912 probands with hypertrophic cardiomyopathy: expanded panels offer limited additional sensitivity. Genet Med. 2015;17:880–888.

4. Ommen SR, Ho CY, Asif IM, Balaji S, Burke MA, Day SM, Dearani JA, Epps KC, Evanovich L, Ferrari VA, et al. 2024 AHA/ACC/AMSSM/HRS/PACES/SCMR Guideline for the Management of Hypertrophic Cardiomyopathy: A Report of the American Heart Association/American College of Cardiology Joint Committee on Clinical Practice Guidelines. Circulation. 2024;149:e1239–e1311.

5. Deignan JL, Chung WK, Kearney HM, Monaghan KG, Rehder CW, Chao EC, ACMG Laboratory Quality Assurance Committee. Points to consider in the reevaluation and reanalysis of genomic test results: a statement of the American College of Medical Genetics and Genomics (ACMG). Genet Med. 2019;21:1267–1270.

6. Westphal DS, Pollmann K, Marschall C, Wacker-Gussmann A, Oberhoffer-Fritz R, Laugwitz K-L, Ewert P, Wolf CM. It Is Not Carved in Stone-The Need for a Genetic Reevaluation of Variants in Pediatric Cardiomyopathies. J Cardiovasc Dev Dis. 2022;9:41.

7. Cherny S, Olson R, Chiodo K, Balmert LC, Webster G. Changes in genetic variant results over time in pediatric cardiomyopathy and electrophysiology. J Genet Couns. 2021;30:229–236.

8. Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, Grody WW, Hegde M, Lyon E, Spector E, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus

recommendation of the American College of Medical Genetics and Genomics and the Association for

Molecular Pathology. Genet Med. 2015;17:405–424.

9. Arbustini E, Behr ER, Carrier L, van Duijn C, Evans P, Favalli V, van der Harst P, Haugaa KH,

Jondeau G, Kääb S, et al. Interpretation and actionability of genetic variants in cardiomyopathies: a

position statement from the European Society of Cardiology Council on cardiovascular genomics. Eur

Heart J. 2022;43:1901–1916.

10. Rehm HL, Berg JS, Brooks LD, Bustamante CD, Evans JP, Landrum MJ, Ledbetter DH, Maglott

DR, Martin CL, Nussbaum RL, et al. ClinGen--the Clinical Genome Resource. N Engl J Med.

2015;372:2235–2242.

11. Papaz T, Liston E, Zahavich L, Stavropoulos DJ, Jobling RK, Kim RH, Reuter M, Miron A,

Oechslin E, Mondal T, et al. Return of genetic and genomic research findings: experience of a pediatric

biorepository. BMC Med Genomics. 2019;12:173.

12. Kelly MA, Caleshu C, Morales A, Buchan J, Wolf Z, Harrison SM, Cook S, Dillon MW, Garcia J,

Haverfield E, et al. Adaptation and validation of the ACMG/AMP variant classification framework for

MYH7-associated inherited cardiomyopathies: recommendations by ClinGen's Inherited

Cardiomyopathy Expert Panel. Genet Med. 2018;20:351–359.

13. Inoue S, Ko T, Shindo A, Nomura S, Yamada T, Jimba T, Dai Z, Nakao H, Suzuki A, Kashimura T,

et al. Association Between Clonal Hematopoiesis and Left Ventricular Reverse Remodeling in

Nonischemic Dilated Cardiomyopathy. JACC: Basic to Translational Science.

2024;S2452302X24001839.

14. Jolfayi AG, Kohansal E, Ghasemi S, Naderi N, Hesami M, MozafaryBazargany M, Moghadam

MH, Fazelifar AF, Maleki M, Kalayinia S. Exploring TTN variants as genetic insights into

cardiomyopathy pathogenesis and potential emerging clues to molecular mechanisms in

cardiomyopathies. Sci Rep. 2024;14:5313.

15. Kurzlechner LM, Kishnani S, Chowdhury S, Atkins SL, Moya-Mendez ME, Parker LE, Rosamilia

MB, Tadros HJ, Pace LA, Patel V, et al. DiscoVari: A Web-Based Precision Medicine Tool for

Predicting Variant Pathogenicity in Cardiomyopathy- and Channelopathy-Associated Genes. Circ

Genom Precis Med. 2023;16:317–327.

16. Walsh R, Mazzarotto F, Whiffin N, Buchan R, Midwinter W, Wilk A, Li N, Felkin L, Ingold N,

Govind R, et al. Quantitative approaches to variant classification increase the yield and precision of

genetic testing in Mendelian diseases: the case of hypertrophic cardiomyopathy. Genome Med.

2019;11:5.

17. Walsh R, Thomson KL, Ware JS, Funke BH, Woodley J, McGuire KJ, Mazzarotto F, Blair E, Seller

A, Taylor JC, et al. Reassessment of Mendelian gene pathogenicity using 7,855 cardiomyopathy cases

and 60,706 reference samples. Genet Med. 2017;19:192–203.

18. Fokkema IFAC, Kroon M, López Hernández JA, Asscheman D, Lugtenburg I, Hoogenboom J, den

Dunnen JT. The LOVD3 platform: efficient genome-wide sharing of genetic variants. Eur J Hum Genet.

2021;29:1796–1803.

19. Biesecker LG, Byrne AB, Harrison SM, Pesaran T, Schäffer AA, Shirts BH, Tavtigian SV, Rehm

HL, ClinGen Sequence Variant Interpretation Working Group. ClinGen guidance for use of the

PP1/BS4 co-segregation and PP4 phenotype specificity criteria for sequence variant pathogenicity

classification. Am J Hum Genet. 2024;111:24–38.

20. Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. A general framework for

estimating the relative pathogenicity of human genetic variants. Nat Genet. 2014;46:310–315.

21. Cheng J, Novati G, Pan J, Bycroft C, Žemgulytė A, Applebaum T, Pritzel A, Wong LH, Zielinski

M, Sargeant T, et al. Accurate proteome-wide missense variant effect prediction with AlphaMissense.

Science. 2023;381:eadg7492.

22. Ioannidis NM, Rothstein JH, Pejaver V, Middha S, McDonnell SK, Baheti S, Musolf A, Li Q,

Holzinger E, Karyadi D, et al. REVEL: An Ensemble Method for Predicting the Pathogenicity of Rare

Missense Variants. Am J Hum Genet. 2016;99:877–885.

23. Pejaver V, Byrne AB, Feng B-J, Pagel KA, Mooney SD, Karchin R, O'Donnell-Luria A, Harrison

SM, Tavtigian SV, Greenblatt MS, et al. Calibration of computational tools for missense variant

pathogenicity classification and ClinGen recommendations for PP3/BP4 criteria. Am J Hum Genet.

2022;109:2163–2177.

24. Harrison SM, Rehm HL. Is "likely pathogenic" really 90% likely? Reclassification data in ClinVar.

Genome Med. 2019;11:72.

25. 1000 Genomes Project Consortium, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM,

Korbel JO, Marchini JL, McCarthy S, McVean GA, et al. A global reference for human genetic

variation. Nature. 2015;526:68–74.

26. Liu Y, Wang H, Wang X, Liu J, Li J, Wang X, Zhang Y, Bai Z, Zhou Q, Wu Y, et al. Prevalence

and reclassification of BRCA1 and BRCA2 variants in a large, unselected Chinese Han breast cancer

cohort. J Hematol Oncol. 2021;14:18.

27. Tadaka S, Kawashima J, Hishinuma E, Saito S, Okamura Y, Otsuki A, Kojima K, Komaki S, Aoki

Y, Kanno T, et al. jMorp: Japanese Multi-Omics Reference Panel update report 2023. Nucleic Acids

Res. 2024;52:D622–D632.

28.   Dhandapany PS, Sadayappan S, Xue Y, Powell GT, Rani DS, Nallari P, Rai TS, Khullar M, Soares

P, Bahl A, et al. A common MYBPC3 (cardiac myosin binding protein C) variant associated with

cardiomyopathies in South Asia. Nat Genet. 2009;41:187–191.

29.   Viswanathan SK, Puckelwartz MJ, Mehta A, Ramachandra CJA, Jagadeesan A, Fritsche-Danielson

R, Bhat RV, Wong P, Kandoi S, Schwanekamp JA, et al. Association of Cardiomyopathy With

MYBPC3 D389V and MYBPC3Δ25bpIntronic Deletion in South Asian Descendants. JAMA Cardiol.

2018;3:481–488.

30.   Harper AR, Bowman M, Hayesmoore JBG, Sage H, Salatino S, Blair E, Campbell C, Currie B,

Goel A, McGuire K, et al. Reevaluation of the South Asian MYBPC3Δ25bp Intronic Deletion in

Hypertrophic Cardiomyopathy. Circ Genom Precis Med. 2020;13:e002783.

31.   Bazrafshan S, Sibilia R, Girgla S, Viswanathan SK, Puckelwartz MJ, Sangha KS, Singh RR,

Kakroo M, Jandarov R, Harris DM, et al. South Asian-Specific MYBPC3 Δ25bp Deletion Carriers

Display Hypercontraction and Impaired Diastolic Function Under Exercise Stress. Front Cardiovasc

Med. 2021;8:766339.

32.   Richards CS, Bale S, Bellissimo DB, Das S, Grody WW, Hegde MR, Lyon E, Ward BE, Molecular

Subcommittee of the ACMG Laboratory Quality Assurance Committee. ACMG recommendations for

standards for interpretation and reporting of sequence variations: Revisions 2007. Genet Med.

2008;10:294–300.

33. Stranneheim H, Lagerstedt-Robinson K, Magnusson M, Kvarnung M, Nilsson D, Lesko N, Engvall

M, Anderlid B-M, Arnell H, Johansson CB, et al. Integration of whole genome sequencing into a

healthcare setting: high diagnostic rates across multiple clinical entities in 3219 rare disease patients.

Genome Med. 2021;13:40.

34. Hume S, Nelson TN, Speevak M, McCready E, Agatep R, Feilotter H, Parboosingh J, Stavropoulos

DJ, Taylor S, Stockley TL, et al. CCMG practice guideline: laboratory guidelines for next-generation

sequencing. J Med Genet. 2019;56:792–800.

35. David KL, Best RG, Brenman LM, Bush L, Deignan JL, Flannery D, Hoffman JD, Holm I, Miller

DT, O'Leary J, et al. Patient re-contact after revision of genomic test results: points to consider-a

statement of the American College of Medical Genetics and Genomics (ACMG). Genet Med.

2019;21:769–771.

Table 1. Baseline study cohort characteristics

Characteristics	N=227 (%)
Male	133 (58.6%)
Ancestry	
European	140 (61.7%)
Asian/South Asian	56 (24.7%)
African	22 (9.7%)
Mixed	5 (2.2%)
Unknown	4 (1.8%)
Cardiomyopathy phenotype	
DCM	104 (45.8%)
HCM	76 (33.5%)
LVNC	20 (8.8%)
RCM	10 (4.4%)
ACM	9 (4.0%)
Mixed type	8 (3.5%)
Type of test	
DCM panel	25 (11.0%)
HCM panel	54 (23.8%)
Multiple panels	124 (54.6%)
Whole exome sequencing	4 (1.8%)
Others	3 (1.3%)
Unknown	17 (7.5%)
Variants identified	
P/LP	110 (28.8%)
VUS	272 (71.2%)
Patient genotype	
Harbored P/LP variant	71 (31.2%)
Harbored VUS	122 (53.7%)
Harbored P/LP variant and VUS	34 (15.0%)

ACM; arrhythmogenic cardiomyopathy, DCM; dilated cardiomyopathy, HCM; hypertrophic cardiomyopathy, LVNC; left ventricular non-compaction, RCM; restrictive cardiomyopathy, VUS; variant of uncertain significance.

Table 2. Pathogenic or likely pathogenic variants reclassified to VUS

Participant ID	Cardiomyopathy phenotype	Year of clinical test	Affected Gene	CDS Change	Amino Acid Change	Current ACMG criteria	Final variant classification
95	RCM	2016	*DSG2*	c.3039C>A	p.Tyr1013*	PVS1, BS1	VUS
216	DCM	2009	*LMNA*	c.1912G>A	p.Gly638Arg	PM1, PM2	VUS
73	DCM	2013	*MYBPC3*	c.1654G>T	p.Ala552Ser	PM2, PP1 (supporting)	VUS
26/76/101/116/145/175	DCM (3), RCM/HCM, HCM (2)	2010/2010/2013/2017/2018/2020	*MYBPC3*	c.3628-41_3628-17del	NA	PP1 (supporting), PP5, BS1	VUS*
59	HCM	2009	*MYBPC3*	c.3548T>G	p.Phe1183Cys	PM2, PP1 (supporting), PP3 (moderate)	VUS
211	HCM	2009	*MYBPC3*	c.2170C>T	p.Arg724Trp	PM1, PM2	VUS
62	DCM	2010	*MYH7*	c.1322C>G	p.Thr441Arg	PM1, PM2	VUS
67	DCM	2009	*MYH7*	c.5507C>T	p.Ser1836Leu	PM2	VUS
168	HCM	2009	*MYH7*	c.1003G>T	p.Ala335Ser	PM1, PM2	VUS
16	LVNC	Unknown	*MYH7*	c.732+1G>A	NA	PM2, PP5	VUS
209	LVNC/DCM	2021	*MYH7*	c.5395G>A	p.Glu1799Lys	PM2, PP3 (moderate)	VUS
12	DCM/RCM	2022	*RRAGD*	c.289_303del	p.Thr97_Cys101del	PM2, PM4	VUS

* One variant was reclassified by the clinical laboratory
ACM; arrhythmogenic cardiomyopathy, ACMG; American College of Medical Genetics and Genomics, CDS; coding sequence, CMP; cardiomyopathy, DCM; dilated cardiomyopathy, HCM; hypertrophic cardiomyopathy, LVNC; left ventricular non-compaction, RCM; restrictive cardiomyopathy, VUS; variant of uncertain significance.
454

Table 3. VUS reclassified to pathogenic or likely pathogenic

Participant ID	Cardiomyopathy phenotype	Year of clinical test	Affected Gene	CDS Change	Amino Acid Change	Current ACMG criteria	Final variant classification
207	LVNC/DCM	2014	ACTC1	c.809G>A	p.Gly270Asp	PS2, PM1, PM2, PP3 (moderate)	Pathogenic
150	DCM	2018	ACTN2	c.784-2A>G	NA	PVS1, PM2	Likely pathogenic*
167	DCM	Unknown	ACTN2	c.36C>G	p.Tyr12*	PVS1, PM2	Likely pathogenic
87	DCM	2012	LMNA	c.868G>A	p.Glu290Lys	PS2, PM2, PP3 (moderate)	Likely pathogenic†
213	HCM	2007	MYBPC3	c.442G>A	p.Gly148Arg	PS3, PP1 (moderate), PP5	Likely pathogenic*†
217	HCM	2010	MYBPC3	c.1828G>A	p.Asp610Asn	PS3, PM1, PM2, PM5, PP3 (supporting)	Pathogenic
31	DCM	2011	MYH7	c.2171T>A	p.Ile724Asn	PS1, PM1, PM2, PP3 (moderate), PP5	Pathogenic
57	DCM	2016	MYH7	c.5158A>G	p.Asn1720Asp	PM1, PM2, PP3 (moderate)	Likely pathogenic
115	DCM	2019	MYH7	c.3667G>A	p.Glu1223Lys	PM1, PM2, PM5, PP3 (supporting)	Likely pathogenic*
126	DCM	2019	MYH7	c.4276G>A	p.Glu1426Lys	PS3, PM2, PP3 (moderate), PP5	Likely pathogenic*
197	DCM	2022	MYH7	c.1423C>A	p.Gln475Lys	PM1, PM2, PM5, PP3 (supporting)	Likely pathogenic
222	DCM	2009	MYH7	c.2683C>G	p.Gln895Glu	PM1, PM2, PP1 (strong)	Likely pathogenic*

ID	Phenotype	Year	Gene	cDNA	Protein	Criteria	Classification
84	HCM	2017	*MYH7*	c.4136C>A	p.Ala1379Asp	PM1, PM2, PP1 (supporting), PP3 (supporting)	Likely pathogenic
151	HCM	Unknown	*MYH7*	c.530C>T	p.Thr177ile	PM1, PM2, PP3 (moderate)	Likely pathogenic
205	HCM	2006	*MYH7*	c.1759G>A	p.Asp587Asn	PM1, PM2, PM5	Likely pathogenic*†
221	HCM	2010	*MYH7*	c.1220G>T	p.Gly407Val	PM1, PM2, PM5, PP3 (supporting), PP5	Likely pathogenic
179	LVNC	2021	*MYH7*	c.1118C>A	p.Ala373Glu	PM1, PM2, PM5, PP3 (supporting)	Likely pathogenic
202	LVNC	2015	*MYH7*	c.532G>A	p.Gly178Arg	PM1, PM2, PP1 (moderate), PP3 (moderate)	Likely pathogenic
79	LVNC/DCM	2015	*MYH7*	c.1180G>A	p.Asp394Asn	PM1, PM2, PM5	Likely pathogenic
126	DCM	2019	*NEXN*	c.1473+1G>T	NA	PVS1, PM2	Likely pathogenic*
210	DCM	2015	*NEXN*	c.373C>T	p.Arg125*	PVS1, PM2	Likely pathogenic
209	LVNC/DCM	2021	*NEXN*	c.201G>A	p.Trp67*	PVS1, PM2	Likely pathogenic
54	HCM	2015	*PLN*	c.2T>C	p.Met1	PVS1, PM2	Likely pathogenic
104	DCM	2018	*TNNC1*	c.376G>A	p.Glu126Lys	PS2, PM2	Likely pathogenic
5	RCM	2011	*TNNC1*	c.23C>T	p.Ala8Val	PS3, PM2, PP3 (supporting), PP5	Pathogenic
117	DCM	2008	*TNNI3*	c.550G>A	p.Glu184Lys	PS2, PM1, PM2	Likely pathogenic
76	RCM/HCM	2013	*TNNI3*	c.484C>T	p.Arg162Trp	PM1, PM2, PM5, PP5	Likely pathogenic
14	DCM	Unknown	*TNNT2*	c.481C>T	p.Arg161Cys	PM1, PM2, PP3 (moderate)	Likely pathogenic
89	DCM	2016	*TNNT2*	c.552G>T	p.Lys184Asn	PM1, PM2, PP1 (supporting), PP3 (supporting)	Likely pathogenic

	Disease	Year	Gene	CDS	Protein	ACMG criteria	Classification
137	DCM	2020	TNNT2	c.316G>A	p.Glu106Lys	PS3, PM1, PM2, PP3 (moderate), PP5	Pathogenic
215	LVNC/DCM	2014	TNNT2	c.862C>T	p.Arg288Cys	PS3, PM1, PM5, PP5	Likely pathogenic
28	DCM	2009	TPM1	c.45G>T	p.Lys15Asn	PS3, PM1, PM2, PP2, PP3 (moderate)	Likely pathogenic
151	HCM	Unknown	TPM1	c.413A>G	p.Glu138Gly	PM2, PM5, PP2, PP3 (strong)	Likely pathogenic
161	HCM	2020	TPM1	c.560A>T	p.Glu187Val	PM1, PM2, PM5, PP2, PP3 (moderate)	Likely pathogenic
106	DCM	2012	TTN	c.16353C>G	p.Tyr5451*	PVS1, PM2, PP1 (supporting)	Likely pathogenic
101	DCM	2010	VCL	c.2949del	p.Gln971Profs*	PVS1, PM2	Likely pathogenic
122	DCM	2018	VCL	c.313C>T	p.Arg105*	PVS1, PM2	Likely pathogenic

Seven variants were reclassified by the clinical laboratory

†3 variants were reported as P/LP in one family and as VUS in another family by the clinical testing laboratories.

ACM; arrhythmogenic cardiomyopathy, ACMG; American College of Medical Genetics and Genomics, CDS; coding sequence, CMP; cardiomyopathy, DCM; dilated cardiomyopathy, HCM; hypertrophic cardiomyopathy, LVNC; left ventricular non-compaction, RCM; restrictive cardiomyopathy, VUS; variant of uncertain significance.

  FIGURE LEGENDS

Figure 1. Criteria for variant reclassification. (a) Downgrading P/LP variants to VUS, **(b)** Upgrading

VUS to P/LP variants.

ACMG, American College of Medical Genetics and Genomics; P/LP, Pathogenic/Likely pathogenic;

VUS, variants of uncertain significance.

Figure 2. Variant and patient level reclassification: (**a**) *Variant level reclassification*: Of 110 P/LP

variants, 12 were downgraded to VUS (10.9%). Of 272 VUS, 37 were upgraded to P/LP (13.6%). In

addition, 47 VUS were downgraded to B/LB. (**b**) *Patient level reclassification*: Of 71 patients harboring

only P/LP variants, 11 patients were downgraded to having only VUS. Of 122 patients harboring only

VUS, 16 patients were upgraded to having only P/LP variants and 13 patients were upgraded to

P/LP+VUS. Of 34 patients harboring both P/LP+VUS, 2 patients were downgraded to only VUS, 3

patients were upgraded to only P/LP variants and *4 patients had a change in the gene harboring a P/LP

variant.

B/LB, Benign/Likely benign; P/LP, Pathogenic/Likely pathogenic; VUS, variant of unceratin

significance.

Figure 3. Variant reclassification by year of clinical testing. The graphs show the **(a)** frequency of

variants reclassified by year of clinical testing (2005-2022); **(b)** percentage of variants reclassified by

year of clinical testing. Year of testing was known for 358 variants. The average annual reclassification

rate was 15%.

Figure 4. Proportion of variants reclassified by year of clinical testing, cardiomyopathy phenotype,

ancestry, and sex. Proportion of variants reclassified did not differ by year of testing (before and after

2015), cardiomyopathy phenotype (HCM vs DCM), ancestry (European vs non-European descent), and

sex (males vs females) ($p > 0.05$ for all).

DCM, dilated cardiomyopathy; HCM, hypertrophic cardiomyopathy

SUPPLEMENTAL MATERIAL

Table S1 All Cardiomyopathy variants reevaluated (n=382)

Figure 1.
a.

492
493 **b.**
494

495
496

497 **Figure 2.**
498 **a.**
499

500
501 **b.**
502

504   **Figure 3.**
505   **a.**

506
507   **b.**

508
509
510
511

512   **Figure 4.**
513

514
515
516
517