Umbilical cord blood pTau217 and BD-tau are associated with markers of neonatal hypoxia: a prospective cohort study

Emma Payne,[1] Fernando Gonzalez-Ortiz,[2,3] Kaitlin Kramer,[4,5] Thomas Payne,[4,5] Shreeya Marathe,[5] Neha Mahajan,[5] Ashly Liu,[5] Jessica Barry,[5] Andrew Duckworth,[5] Mitchell Brookes,[5] Bradley de Vries,[6,7] Benjamin Moran,[8,9] Helen Manning,[10] Adrienne Gordon,[4,6] Kaj Blennow,[2,3] Henrik Zetterberg,[11,12] David Zalcberg,[4,5] Robert D. Sanders[4,5]

Author affiliations:

1. St George Hospital, South Eastern Sydney Local Health District, New South Wales, Australia

2. Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, the Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden

3. Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden

4. Central Clinical School, Faculty of Medicine and Health, The University of Sydney, New South Wales, Australia

5. Department of Anaesthetics, Royal Prince Alfred Hospital, Sydney Local Health District, New South Wales, Australia

6. Sydney Institute for Women, Children and their Babies, Sydney Local Health District, New South Wales, Australia

7. NHMRC Clinical Trials Centre, The University of Sydney, New South Wales, Australia

8. Critical Care Program, The George Institute of Global Health, Sydney, Australia.

9. Department of Intensive Care, Gosford Hospital, Gosford, Australia.

10. Dept of Obstetrics and Gynaecology, Central coast local health district, NSW

11. Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, the Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden

12. Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden

* Corresponding authors: Emma Payne, David Zalcberg

email: emmapayne4114@gmail.com

address: St George Hospital, Gray St, Kogarah, NSW, 2217

phone: +61449554114

Word Count: 2903

Abstract

Objective: Current methods for early detection of hypoxic–ischemic encephalopathy (HIE)

are limited by lack of specificity, cost, and time constraints. Blood tau protein concentrations

reflect neuropathology in adults. This study examines tau as a potential HIE biomarker in

neonates by relating cord blood levels to short-term fetomaternal outcomes. We aimed to

examine 1) association of BD-tau with non-reassuring fetal status; 2) correlations between

cord blood tau and other hypoxia biomarkers; 3) associations between tau levels and risk

factors for fetomaternal morbidity; 4) associations between tau levels and short-term

fetomaternal outcome.

Methods: 107 maternal participants were prospectively recruited at Royal Prince Alfred

Hospital—a large Australian tertiary referral centre. Simoa analysis detected umbilical cord

blood pTau217 and brain-derived (BD)-tau levels.

Results: Of 509 deliveries, cord blood was analysed in 107/110 recruited maternal

participants. BD-tau correlated with non-reassuring fetal status (OR=3.0;95%CI=1.6–

5.7;p=0.001), though not when adjusting for mode of delivery and gestational age. BD-tau

was higher in vaginal deliveries, and positively associated with pTau217, NfL, and lactate

(p<0.001), and negatively associated with pH and base excess. pTau217 was higher in

preterm neonates and was associated with neurofilament light chain (Spearman's

rho=0.44,p<0.001). BD-tau and pTau217 were associated with maternal hypertension and

placental abnormalities.

Conclusions: Cord blood BD-tau correlates with surrogate markers of fetal hypoxia, whilst

pTau217 may represent a marker of neurodevelopment. Further studies could explore

whether these findings translate to clinical use of tau as an HIE biomarker.

Funding: US National Institutes of Health (grant:R01AG063849-01).

Keywords: tau; asphyxia; hypoxic–ischemic encephalopathy; HIE.

Introduction

Hypoxic–ischemic encephalopathy (HIE)—a subtype of neonatal encephalopathy—is a syndrome of central nervous system (CNS) dysfunction caused by abnormalities in cerebral blood flow and impaired gas exchange perinatally, resulting in multiple organ failure.[1] Approximately 40–60% of affected infants face mortality or severe neurodevelopmental disability by two years of age.[2]

Risk factors for HIE can broadly be categorised into antepartum (including fetal and maternal characteristics) and intrapartum factors. Understanding antepartum risk factors can complement intrapartum monitoring techniques used to detect 'non-reassuring fetal status' as defined by pathological fetal heart rate traces and fetal scalp blood sampling. Together, this information can identify fetuses at risk of hypoxic injury and guide decision-making regarding the urgency of delivery. However, intrapartum monitoring techniques are limited by a lack of specificity for pathological neurological damage, measurement difficulties, and poor inter-user reliability.[3,4]

The only proven neuroprotective treatment for HIE is therapeutic hypothermia (TH), which has only been validated in term or late preterm neonates.[2,5] Severity of HIE can be classified as mild, moderate, or severe based on clinical signs and grading systems such as the modified Sarnat score.[6] However, these scoring criteria are subjective, and grading may fail to identify neonates at risk of long-term adverse neurological sequelae; research has demonstrated abnormal neurological outcome in over 22% of cohorts of mild HIE.[5] At this

stage, TH is validated only in moderate to severe HIE, and there is insufficient evidence to

recommend TH for mild HIE cohorts.[5] Cord blood biomarkers may be helpful in identifying

subgroups of mild HIE who would benefit from TH. Further aids to establish aetiology and

severity of HIE include electroencephalogram (EEG) monitoring, and neuroimaging studies

including MRI and cranial ultrasound.[3,7] These tools are limited by time and cost constraints,

require expert interpretation, and may be difficult to access in emergency situations.[7] This

indicates an ongoing need for objective, cost-effective, and rapid measures of identifying HIE

in order to appropriately identify subgroups of HIE patients that could benefit from early

intervention.

Real-time physiological biomarkers could address this gap in diagnosis and prognostication

by providing objective point-of-care testing. Current cord or whole blood markers such as

pH, lactate, and base excess remain non-specific for neurological injury.[2] Tau is a promising

biomarker of poor neurological outcome in adult cohorts with acute and chronic

neurological disease.[8,9] Studies of tau in neonates has, to date, focused on correlation of

total blood tau and cerebral injury-related outcome measures. However, compared to total

blood tau, specific tau subtypes—in particular pTau217 and 'brain-derived tau' (BD-tau)—

have demonstrated greater specificity for adult neurological injury.[10,11] This is the first study

to examine tau subtypes in a neonatal cohort.

Given the paucity of evidence addressing tau subtypes and fetomaternal outcome, we

conducted a prospective cohort study with the following aims:

1. Address feasibility of neonatal cord blood tau level measurement in an

114   Australian cohort.

115   2. Assess the correlation between BD-tau and non-reassuring fetal status.

116   3. Assess correlations between cord blood tau—BD-tau, pTau217—and other

117   cord blood biomarkers, maternal and neonatal risk factors for

118   encephalopathy, and adverse outcome.

Materials and Methods

Study Population and Design

We performed a prospective cohort study of maternal and neonatal patients at the Royal

Prince Alfred Hospital (RPAH), a tertiary referral centre in Sydney, Australia. Maternal

participants of any age who were planned to birth at RPAH were eligible for study

recruitment. Recruitment occurred prospectively from October 2022 at any time from 4

weeks prior to estimated date of delivery to the immediate post-partum period (with the

option of retrospective consent). Trained recruiters obtained written, informed consent in

English for all maternal participants in antenatal visits or on the birthing unit. Participants

were excluded if they were non-English speaking, or had a history of psychological illness or

other conditions that may interfere with capacity to provide informed consent after

appropriate counselling in English.

Recruitment was undertaken as part of the BABBies (Benefits of Analysing Brain Biomarkers

in perinatal care) Study—a prospective observational cohort study exploring the biomarkers

neurofilament light chain (NfL), BD-tau, and pTau217. Data were stored in Sydney Local

Health District (SLHD) REDCap electronic database. Ethics approval was obtained from SLHD

Research and Ethics Committee (RPAH zone) (approval number: 2022/ETH01100). Our study

was performed according to the National Statement on Ethical Conduct in Human Research

(2007) and the CPMP/ICH Note for Guidance on Good Clinical Practice. Data reporting

adhered to Strengthening the Reporting of Observational Studies in Epidemiology (STROBE)

guidelines.

Cord Blood Collection and Analysis

Umbilical venous cord blood was collected by the attending midwife in the immediate

postpartum period. 1–3mL samples were collected and centrifuged in the RPAH Department

of Anaesthetics laboratory, and plasma samples were then stored in deidentified cryovials.

Tau was measured on the Simoa HD-X platform with two-fold dilution factor in plasma.

Plasma pTau217 and BD-tau were measured with previously validated assays.[12,13] Signal

variations within and between analytical runs were assessed using three internal quality

control samples at the beginning and the end of each run.

Outcome Data Collection

Demographic and outcome data were collected from RPAH electronic and paper medical

records. Mode of delivery was classified categorically: vaginal delivery, emergency caesarean

section, and elective caesarean section. Placental abnormalities were defined as any

abnormality detected on placental ultrasound, on examination in the birth unit or in the

operating theatre, or histopathology. The composite term 'histopathological placental

abnormality' refers to only abnormalities detected on anatomical or histopathological

examination per the 2014 Amsterdam Working Group nosology for classification of placental

clinicopathological disorders.[14] 'Non-reassuring fetal status' was defined as fetal blood

sampling showing a pH ≤7.20 or lactate >4.7,[15,16] or intrapartum cardiotocography (CTG)

changes identified as 'red zone' criteria according to NSW Health electronic fetal monitoring

guidelines (Table S1).[17] 'Non-reassuring fetal status' did not include participants incidentally

found to have abnormal umbilical cord blood values without meeting other criteria for non-

reassuring fetal status.

Outcomes

Our primary outcome was the association of BD-tau with non-reassuring fetal status. We

adjusted this analysis for gestational age and mode of delivery, given these variables have

previously been associated with both non-reassuring fetal status and cord blood

concentrations of other biomarkers of acidosis and anaerobic metabolism.[18] Secondary

outcomes included the association of BD-tau and pTau217 with antenatal factors

(gestational age, head circumference, birthweight), other cord blood biomarkers, maternal

factors, delivery factors, and perinatal outcome. Resuscitation at birth was defined as

cardiorespiratory support required following drying, warming, and mechanical stimulation.[19]

Power Analysis

Our power analysis was based on the association of NfL with non-reassuring fetal status,

which is reported separately.[20] We used the previously reported baseline incidence of non-

reassuring fetal status of 16%.[15] Based on a two-sided t-test with α=0.025, a sample size of

110 participants provides 80% power to detect a Cohen's d of 0.8 (i.e., a large effect size).

The power analysis also applies to the primary outcome of this study, with a preserved type

1 error rate of α=0.05, as this value was halved a priori for the power calculation to account

for additional analysis.

Statistical Methods

Primary outcome

Both BD-tau and pTau217 concentrations showed a strong positive skew, hence were $\log_{10}$-

transformed for all analyses. To determine the association of BD-tau with non-reassuring

fetal status after adjusting for other variables, we used logistic regression (binomial family

with logit link) with maximum likelihood estimation of the model parameters. The

parametric G-formula was used to compute the mean risk difference from logistic regression

models.[21] We included a BD-tau*mode of delivery interaction in our model, as the

relationship between neuronal biomarkers and fetal outcome would likely vary by birth

route. To provide interpretable effect measures, we calculated the risk difference for the

outcome per increase in BD-tau by its interquartile range (Q3 – Q1). The relative association

of cord biomarkers with non-reassuring fetal status was quantified using the area under the

receiver operator curve (AUROC). The 95% confidence interval for the AUROC was calculated

using 2000 stratified bootstrap replicates.

Secondary outcomes

Bivariate biomarker correlations were assessed using rank-based nonparametric methods.

The relative strength of biomarker correlations was tested using the method described by

Meng, Rosenthal, and Rubin.[22] For continuous secondary outcomes, we used linear

regression with the ordinary least squares estimator for the model parameters. Associations

between biomarker concentrations and placental abnormalities were adjusted for low birth

weight (defined as birth weight <2.5kg, per World Health Organisation definition).[23] Binary

secondary outcomes were tested for linearity, and analysed in the same fashion as our

primary outcome.

We used a p-value <0.05 to denote statistical significance. No adjustments for multiple

comparisons were made in this preliminary study. All analyses were conducted in R using

RStudio (Version 2024.04.0; R Foundation for Statistical Computing, Vienna, Austria). The

'stats' package was used for linear and generalised linear models. The 'cocor' package was

used to compare the strength of bivariate biomarker concentrations.[24] The 'plotROC' and

'pROC' packages were used for calculation and plotting of the AUROC.[25]

Results

Over the study period 24[th] October 2022 to 9[th] December 2022, 110 maternal study

participants were recruited from a total of 509 deliveries at RPAH. Cord blood was collected

for a total of 108 participants, of whom 107 had either BD-tau or pTau217 levels available;

105 had BD-tau values and 106 had pTau217 values (STROBE diagram: Figure S1). Cohort

demographic information is summarised in Table S2.

Primary outcome

Increased BD-tau was associated with a higher risk of non-reassuring fetal status; each IQR

increase in BD-tau was associated with an increased odds of fetal distress (OR=3.0; 95% CI:

1.6, 5.7; p=0.001). However, this relationship was no longer observed when adjusting for

mode of delivery and gestational age (OR=1.1; 95%CI=0.6, 1.2; p=0.667) (Table S3). BD-tau

was not associated with non-reassuring fetal status when stratifying within each individual

mode of delivery (Table S3). Comparative performances of BD-tau pH, lactate, and base

excess in predicting non-reassuring fetal status are displayed in Figure 1. There was no

association demonstrated between pTau217 and non-reassuring fetal status (Wilcoxon

p=0.62).

Secondary outcomes

Antenatal Factors

Relationship between tau and fetal factors

pTau217 was negatively associated with gestational age (Spearman's rho=−0.25, p=0.0096).

pTau217 was also negatively associated with birthweight (Spearman's rho=−0.23, p=0.019)

and head circumference (Spearman's rho=−0.24, p=0.016), however, these associations were

not significant when controlling for gestational age (Table S4). BD-tau did not demonstrate

evidence of association with fetal parameters (Figure 2).

Relationship between tau and maternal factors

A total of 8 participants were classified as having maternal hypertension—either chronic

hypertension, gestational hypertension, or pre-eclampsia. Both BD-tau and pTau217 were

positively associated with maternal hypertension (median BD-tau: 81.0 vs. 54.9pg/mL,

Wilcoxon p=0.026; median pTau217: 10.3 vs. 7.7pg/mL, Wilcoxon p=0.029), but were not

associated with other maternal health conditions (Figures S2 and S3).

Relationship between tau and intrapartum factors

BD-tau was associated with mode of delivery (Kruskal–Wallis p<0.001) (Figure 3), with BD-

tau levels being higher with vaginal delivery compared to CS. On pairwise comparisons, no

statistically significant difference was detected in BD-tau level between elective and

emergency CS (median 48.4 vs. 53.6pg/mL, Wilcoxon p=0.22). pTau217 did not demonstrate

a significant association with mode of delivery (Kruskal–Wallis p=.420). pTau217 was

negatively associated with the duration of the second stage of labour (Spearman rho −0.4,

p=0.042), whilst BD-tau and other biomarkers were not (Figure S4).

Correlation between tau and other biomarkers

Correlations between cord blood tau and other biomarkers are demonstrated in Figure 4.

BD-tau demonstrated a positive association with pTau217 (Spearman's rho=0.66, p<0.001),

NfL (Spearman's rho=0.58, p<0.001), and lactate (Spearman's rho=0.34, p<0.001), and a

negative association with cord pH (Spearman's rho=−0.29, p=0.003) and base excess

(Spearman's rho=−0.35, p<0.001). pTau217 demonstrated a positive correlation with NfL

(Spearman's rho=0.44, p<0.001), but was not associated with other biomarkers. We did not

observe evidence of a difference between pTau217 and BD-tau in the strength of the

positive correlation with NfL (z=1.48, p=0.140) (Table S5).

Relationship between tau and fetoplacental outcome

A total of 11 participants had reported placental abnormalities, 6 of which were defined as

histopathological placental abnormalities (Appendix 1). Adjusting for low birth weight, the

presence of histopathological placental abnormalities was positively associated with BD-tau

and pTau217 (Table S6).

pTau217 levels were associated with preterm birth (median 7.7 in term participants vs.

12.2pg/mL in preterm participants; Wilcoxon p=0.010). In addition, pTau217 levels were

positively associated with respiratory complications (median 7.6 vs. 10.4pg/mL; Wilcoxon

p=0.012) and resuscitation requirement at birth (median pTau217: 7.6 vs. 10.2pg/mL;

Wilcoxon p=0.027), however these associations were not observed when adjusting for

preterm birth (Table S7).

Discussion

Main findings

Our study adds credence to the use of cord blood tau as a biomarker of neurodevelopment,

perinatal hypoxia, and CNS injury. It is the first to examine associations between

fetomaternal outcomes and cord blood subtypes of tau—BD-tau and pTau217. We observed

a correlation between BD-tau and surrogate markers of hypoxia, including serum biomarkers

of anaerobic metabolism and non-reassuring fetal status. Additionally, we found associations

between tau and maternal and fetoplacental factors that could contribute to increased

baseline risk of encephalopathy, including preterm birth, placental abnormalities, and

maternal hypertension. Understanding the relationship between the pathogenesis of

hypoxic brain injury, neuron-specific biomarkers, and clinically observed outcome could aid

diagnosis, prognosis, and treatment of patients suffering from HIE.

Clinical implications and future research

By addressing surrogate markers of fetal and neonatal hypoxia, our study of predominantly

'healthy' participants aims to link blood tau levels with risk factors for HIE, and provide a

foundation for future research into the clinical use of cord blood tau in HIE diagnosis. Our

study demonstrated a strong correlation between tau levels and non-specific markers of

acidosis and anaerobic metabolism, such as neonatal pH and lactate levels, extending

previous research.[1,18,26–28] Importantly, our study is the first to examine BD-tau specifically,

which offers an advantage over these traditional biomarkers due to its greater specificity for

central nervous system (CNS) injury.[18,26–28] Previous studies have mainly involved case-

control analyses of HIE cohorts, demonstrating a positive correlation between early serum

tau levels, HIE diagnosis based on EEG and clinical criteria, as well as HIE severity.[26,28]

The proposed role of tau as a surrogate marker of fetal hypoxia is further evidenced by the

correlation between early cord blood BD-tau and non-reassuring fetal status observed in this

study. This finding was not statistically significant when adjusting for mode of delivery and

gestational age. However, operative delivery is commonly performed for fetal distress, and

adjusting for mode of delivery could therefore cause the association between BD-tau and

fetal distress to be underestimated. Fetal distress, indicated by pathological fetal heart rate

abnormalities, may signal early hypoxia compensation and correlate with neonatal acidosis,

low APGAR scores, NICU admission, and resuscitation.[13,29] However, CTG interpretation is

subjective, operator-dependent, and has low predictive value for long-term outcomes.[29]

Biomarkers like BD-tau could improve diagnostic specificity and guide interventions such as

therapeutic hypothermia. BD-tau may offer a more specific marker of neurological damage

compared current blood biomarkers.[18,26–28]

Our study also explored potential correlations between neonatal tau levels and antenatal

risk factors for adverse neurological outcomes. Macroscopic (e.g., cord knots, marginal cord

insertion) and microscopic (e.g., avascular villi) abnormalities in the 'placenta–brain axis'

can contribute to intrauterine growth restriction and increased susceptibility of the fetal

brain to injury during acute stress.[30,31] We observed increased pTau217 levels in participants

with histopathological placental abnormalities. Our findings should prompt further

investigation into whether excessive tau phosphorylation may be related to aberrant

placental function, indicating potential increased risk of neurological injury.

Microscopic placental changes may also occur as a consequence of pre-eclampsia (PET).[32]

Non-PET hypertensive disorders may show some similar microvascular changes to a lesser

unknown degree, manifesting in increased neonatal stroke risk.[32] Associations between

increased tau and hypertension should be further explored to identify patients with

microvascular placental changes who may be more vulnerable to physiologic stress and

sentinel intrapartum events.

Phosphorylated tau (pTau217) showed a positive correlation with gestational age but not

hypoxic markers. While tau phosphorylation is essential for neuronal function, excessive

phosphorylation may contribute to neuronal dysfunction.[33,34] In adults, pTau217 elevation

reflects blood-brain barrier disruption and protein aggregation rather than direct neuronal

injury.[33,34] It remains unclear whether high neonatal pTau217 signals abnormal neuronal

development or increased neuropathology risk. Further research is needed to clarify these

mechanisms and define threshold levels for clinical significance.

Strengths and limitations

This study is the first to distinguish between specific tau subtypes—rather than relying solely

on total tau levels—paving the way for more precise diagnosis and prognosis. Correlations

between BD-tau and surrogate markers of hypoxia form a strong foundation for future

research into the use of cord blood tau in birth asphyxia screening. The successful sample

collection and analysis, using validated assays, highlights the potential for tau cord blood

measurement to feasibly be incorporated into neonatal care in an Australian context.

Limitations of this study included the relatively small sample size, from a single centre. In

addition, the largely 'healthy' maternal cohort and short-term follow-up limited adverse

outcome data. Also, our study only included static biomarker measurement at time of

delivery, missing serial neonatal samples. Late serum tau rises have been linked to

neurodevelopmental issues at 1–2 years,[35] highlighting the need for further research on

serial tau measurements.

Conclusions

This study establishes BD-tau and pTau217 as potential biomarkers for fetal hypoxia and

neurological vulnerability. BD-tau correlated with fetal distress, cord lactate, and pH, while

phosphorylated tau levels linked to maternal hypertension and placental abnormalities.

Larger, longitudinal studies are needed to validate these findings and explore the integration

of tau biomarkers into neonatal screening tools.

Acknowledgements

HZ is a Wallenberg Scholar and a Distinguished Professor at the Swedish Research Council

supported by grants from the Swedish Research Council (#2023-00356; #2022-01018 and

#2019-02397), the European Union's Horizon Europe research and innovation programme

under grant agreement No 101053962, Swedish State Support for Clinical Research

(#ALFGBG-71320), the Alzheimer Drug Discovery Foundation (ADDF), USA (#201809-

2016862), the AD Strategic Fund and the Alzheimer's Association (#ADSF-21-831376-C,

#ADSF-21-831381-C, #ADSF-21-831377-C, and #ADSF-24-1284328-C), the European

Partnership on Metrology, co-financed from the European Union's Horizon Europe Research

and Innovation Programme and by the Participating States (NEuroBioStand, #22HLT07), the

Bluefield Project, Cure Alzheimer's Fund, the Olav Thon Foundation, the Erling-Persson

Family Foundation, Familjen Rönströms Stiftelse, Stiftelsen för Gamla Tjänarinnor,

Hjärnfonden, Sweden (#FO2022-0270), the European Union's Horizon 2020 research and

innovation programme under the Marie Skłodowska-Curie grant agreement No 860197

(MIRIADE), the European Union Joint Programme – Neurodegenerative Disease Research

Conflicts of interest

HZ has served at scientific advisory boards and/or as a consultant for Abbvie, Acumen,

Alector, Alzinova, ALZPath, Amylyx, Annexon, Apellis, Artery Therapeutics, AZTherapies,

Cognito Therapeutics, CogRx, Denali, Eisai, LabCorp, Merry Life, Nervgen, Novo Nordisk,

Optoceutics, Passage Bio, Pinteon Therapeutics, Prothena, Red Abbey Labs, reMYND, Roche,

Samumed, Siemens Healthineers, Triplet Therapeutics, and Wave, has given lectures

sponsored by Alzecure, BioArctic, Biogen, Cellectricon, Fujirebio, Lilly, Novo Nordisk, Roche,

and WebMD, and is a co-founder of Brain Biomarker Solutions in Gothenburg AB (BBS),

which is a part of the GU Ventures Incubator Program (outside submitted work).

Contribution of Authorship

RDS designed the study in consultation with BdV, KK, BM, FG, and HM. FG, HZ and KB

supplied the assays and managed biofluid analysis. KK consulted hospital records for

relevant cohort data. EP and TP conducted the statistical analysis. EP, FG, and TP drafted the

manuscript. All authors provided critical feedback on the manuscript.

Ethics approval

Ethics approval was granted by the Sydney Local Health District Human Research and Ethics

Committee (approval reference: 2022/ETH01100). Data were reported according to the

Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines.

Funding

The work is supported by US National Institutes of Health (NIH) grant R01 AG063849-01 (RDS). NIH R01 grants are subject to external peer review for scientific quality, and priority score is based advisory council assessment. KB is supported by the Swedish Research Council (#2017-00915 and #2022-00732), the Swedish Alzheimer Foundation (#AF-930351, #AF-939721, #AF-968270, and #AF-994551), Hjärnfonden, Sweden (#FO2017-0243 and #ALZ2022-0006), the Swedish state under the agreement between the Swedish government and the County Councils, the ALF-agreement (#ALFGBG-715986 and #ALFGBG-965240), the European Union Joint Program for Neurodegenerative Disorders (JPND2019-466-236), the Alzheimer's Association 2021 Zenith Award (ZEN-21-848495), the Alzheimer's Association 2022-2025 Grant (SG-23-1038904 QC), La Fondation Recherche Alzheimer (FRA), Paris, France, and the Kirsten and Freddy Johansen Foundation, Copenhagen, Denmark. HZ is a Wallenberg Scholar and a Distinguished Professor at the Swedish Research Council supported by grants from the Swedish Research Council (#2023-00356; #2022-01018 and #2019-02397), the European Union's Horizon Europe research and innovation programme under grant agreement No 101053962, Swedish State Support for Clinical Research (#ALFGBG-71320), the Alzheimer Drug Discovery Foundation (ADDF), USA (#201809-2016862), the AD Strategic Fund and the Alzheimer's Association (#ADSF-21-831376-C, #ADSF-21-831381-C, #ADSF-21-831377-C, and #ADSF-24-1284328-C), the Bluefield Project, Cure Alzheimer's Fund, the Olav Thon Foundation, the Erling-Persson Family Foundation, Stiftelsen för Gamla Tjänarinnor, Hjärnfonden, Sweden (#FO2022-0270), the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 860197 (MIRIADE), the European Union Joint Programme –

References

1. Gaulee P, Yang Z, Sura L, et al. Concentration of Serum Biomarkers of Brain Injury in Neonates With a Low Cord pH With or Without Mild Hypoxic-Ischemic Encephalopathy. *Front Neurol*. 2022;13:934755. doi:10.3389/fneur.2022.934755

2. Jacobs SE, Berg M, Hunt R, Tarnow-Mordi WO, Inder TE, Davis PG. Cooling for newborns with hypoxic ischaemic encephalopathy. *Cochrane Database Syst Rev*. 2013;2013(1):CD003311. doi:10.1002/14651858.CD003311.pub3

3. Leijser LM, Vein AA, Liauw L, Strauss T, Veen S, Wezel-Meijler G van. Prediction of short-term neurological outcome in full-term neonates with hypoxic-ischaemic encephalopathy based on combined use of electroencephalogram and neuro-imaging. *Neuropediatrics*. 2007;38(5):219-227. doi:10.1055/s-2007-992815

4. Sarnat HB, Sarnat MS. Neonatal encephalopathy following fetal distress. A clinical and electroencephalographic study. *Arch Neurol*. 1976;33(10):696-705. doi:10.1001/archneur.1976.00500100030012

5. Finder M, Boylan GB, Twomey D, Ahearne C, Murray DM, Hallberg B. Two-Year Neurodevelopmental Outcomes After Mild Hypoxic Ischemic Encephalopathy in the Era of Therapeutic Hypothermia. *JAMA Pediatr*. 2020;174(1):48. doi:10.1001/jamapediatrics.2019.4011

6. Sarnat HB, Flores-Sarnat L, Fajardo C, Leijser LM, Wusthoff C, Mohammad K. Sarnat Grading Scale for Neonatal Encephalopathy after 45 Years: An Update Proposal. *Pediatr Neurol*. 2020;113:75-79. doi:10.1016/j.pediatrneurol.2020.08.014

7. Fiala M, Baumert M, Surmiak P, Walencka Z, Sodowska P. Umbilical markers of perinatal

hypoxia. *Ginekol Pol*. 2016;87(3):200-204. doi:10.17772/gp/60552

8. Gonzalez-Ortiz F, Dulewicz M, Ashton NJ, et al. Association of Serum Brain-Derived Tau

With Clinical Outcome and Longitudinal Change in Patients With Severe Traumatic Brain

Injury. *JAMA Netw Open*. 2023;6(7):e2321554.

doi:10.1001/jamanetworkopen.2023.21554

9. Gundersen JK, Gonzalez-Ortiz F, Karikari T, et al. Clinical value of plasma brain-derived tau

and p-tau217 in acute ischemic stroke. *Cereb Circ - Cogn Behav*. 2024;6:100291.

doi:10.1016/j.cccb.2024.100291

10. Ashton NJ, Brum WS, Di Molfetta G, et al. Diagnostic Accuracy of a Plasma

Phosphorylated Tau 217 Immunoassay for Alzheimer Disease Pathology. *JAMA Neurol*.

Published online January 22, 2024:e235319. doi:10.1001/jamaneurol.2023.5319

11. Rubenstein R, Chang B, Yue JK, et al. Comparing Plasma Phospho Tau, Total Tau, and

Phospho Tau–Total Tau Ratio as Acute and Chronic Traumatic Brain Injury Biomarkers.

JAMA Neurol. 2017;74(9):1063. doi:10.1001/jamaneurol.2017.0655

12. Gonzalez-Ortiz F, Turton M, Kac PR, et al. Brain-derived tau: a novel blood-based

biomarker for Alzheimer's disease-type neurodegeneration. *Brain*. 2023;146(3):1152-

1165. doi:10.1093/brain/awac407

13. Gonzalez-Ortiz F, Ferreira PCL, González-Escalante A, et al. A novel ultrasensitive assay

for plasma p-tau217: Performance in individuals with subjective cognitive decline and

early Alzheimer's disease. *Alzheimers Dement*. 2024;20(2):1239-1249.

doi:10.1002/alz.13525

14. Khong TY, Mooney EE, Ariel I, et al. Sampling and Definitions of Placental Lesions:

Amsterdam Placental Workshop Group Consensus Statement. *Arch Pathol Lab Med*.

2016;140(7):698-713. doi:10.5858/arpa.2015-0225-CC

15. East CE, Leader LR, Sheehan P, Henshall NE, Colditz PB, Lau R. Intrapartum fetal scalp

lactate sampling for fetal assessment in the presence of a non-reassuring fetal heart rate

trace. *Cochrane Database Syst Rev*. 2015;2015(5):CD006174.

doi:10.1002/14651858.CD006174.pub3

16. Fetal blood sampling in intrapartum care: SESLHD clinical procedure. Published online

2019.

17. New South Wales Ministry of Health. Heart rate monitoring guide. Published online

2018. https://www1.health.nsw.gov.au/pds/Pages/doc.aspx?dn=GL2018_025

18. Toorell H, Zetterberg H, Blennow K, Sävman K, Hagberg H. Increase of neuronal injury

markers Tau and neurofilament light proteins in umbilical blood after intrapartum

asphyxia. *J Matern Fetal Neonatal Med*. 2018;31(18):2468-2472.

doi:10.1080/14767058.2017.1344964

19. ANZCOR (Australian and New Zealand Committee on Resuscitation). Guideline 13.1 –

Introduction to Resuscitation of the Newborn. Published online 2024.

https://www.anzcor.org/home/neonatal-resuscitation/guideline-13-1-introduction-to-

resuscitation-of-the-newborn/

20. Zalcberg D, Kramer K, Payne E, et al. The association of umbilical cord blood

neurofilament light with non-reassuring fetal status: a prospective observational study.

Published online January 25, 2025. doi:10.1101/2025.01.23.25320706

21. Hernan M, Robins J. *Causal Inference: What If*. Boca Raton: Chapman & Hall/CRC; 2020.

https://www.hsph.harvard.edu/miguel-hernan/wp-

content/uploads/sites/1268/2024/01/hernanrobins_WhatIf_2jan24.pdf

22. Meng X li, Rosenthal R, Rubin DB. Comparing correlated correlation coefficients. *Psychol*

Bull. 1992;111(1):172-175. doi:10.1037/0033-2909.111.1.172

23. World Health Organization. Low birth weight. Published online 2024. Accessed

September 19, 2024. https://www.who.int/data/nutrition/nlis/info/low-birth-

weight#:~:text=Low%20birth%20weight%20has%20been,2500%20grams%20(5.5%20po

unds).

24. Diedenhofen B, Musch J. cocor: a comprehensive solution for the statistical comparison

of correlations. *PloS One*. 2015;10(3):e0121945. doi:10.1371/journal.pone.0121945

25. Sachs M. plotROC: A Tool for Plotting ROC Curves. *J Stat Softw Code Snippets*.

2017;79(2):1-19. doi:doi:10.18637/jss.v079.c02

26. Wu H, Li Z, Yang X, Liu J, Wang W, Liu G. SBDPs and Tau proteins for diagnosis and

hypothermia therapy in neonatal hypoxic ischemic encephalopathy. *Exp Ther Med*.

2017;13(1):225-229. doi:10.3892/etm.2016.3911

27. Chavez-Valdez R, Miller S, Spahic H, et al. Therapeutic Hypothermia Modulates the

Relationships Between Indicators of Severity of Neonatal Hypoxic Ischemic

Encephalopathy and Serum Biomarkers. *Front Neurol*. 2021;12:748150.

doi:10.3389/fneur.2021.748150

28. Toorell H, Carlsson Y, Hallberg B, et al. Neuro-Specific and Immuno-Inflammatory

Biomarkers in Umbilical Cord Blood in Neonatal Hypoxic-Ischemic Encephalopathy.

Neonatology. Published online September 29, 2023:1-9. doi:10.1159/000533473

29. Jonsson M, Ågren J, Nordén-Lindeberg S, Ohlin A, Hanson U. Neonatal encephalopathy

and the association to asphyxia in labor. *Am J Obstet Gynecol*. 2014;211(6):667.e1-8.

doi:10.1016/j.ajog.2014.06.027

30. Penn AA, Wintermark P, Chalak LF, et al. Placental contribution to neonatal

encephalopathy. *Semin Fetal Neonatal Med*. 2021;26(4):101276.

doi:10.1016/j.siny.2021.101276

31. Fox A, Doyle E, Geary M, Hayes B. Placental pathology and neonatal encephalopathy. *Int

J Gynaecol Obstet Off Organ Int Fed Gynaecol Obstet*. 2023;160(1):22-27.

doi:10.1002/ijgo.14301

32. Mann JR, McDermott S, Pan C, Hardin JW. Maternal hypertension and intrapartum fever

are associated with increased risk of ischemic stroke during infancy. *Dev Med Child*

Neurol. 2013;55(1):58-64. doi:10.1111/j.1469-8749.2012.04409.x

33. Flavin WP, Hosseini H, Ruberti JW, Kavehpour HP, Giza CC, Prins ML. Traumatic brain

injury and the pathways to cerebral tau accumulation. *Front Neurol*. 2023;14:1239653.

doi:10.3389/fneur.2023.1239653

34. Palmqvist S, Janelidze S, Quiroz YT, et al. Discriminative Accuracy of Plasma Phospho-

tau217 for Alzheimer Disease vs Other Neurodegenerative Disorders. *JAMA*.

2020;324(8):772-781. doi:10.1001/jama.2020.12134

35. Juul SE, Voldal E, Comstock BA, et al. Association of High-Dose Erythropoietin With

Circulating Biomarkers and Neurodevelopmental Outcomes Among Neonates With

Hypoxic Ischemic Encephalopathy: A Secondary Analysis of the HEAL Randomized

Clinical Trial. *JAMA Netw Open*. 2023;6(7):e2322131.

doi:10.1001/jamanetworkopen.2023.22131

Figure Captions

Figure 1:

Receiver–operator curve (ROC) analysis of BD-tau, pH, base excess, and lactate in predicting binary non-reassuring fetal status. (A) ROC for the generalised linear model predicting non-reassuring fetal status and including the regressors: biomarker, mode of delivery, and gestational age. (B) ROC for the generalised linear model predicting non-reassuring fetal status and including the regressors: biomarker and mode of delivery. ROC for the biomarkers only, including (C) all births, (D) excluding preterm births, (E) excluding elective caesarean section, and (F) excluding elective CS and preterm birth.

Figure 2:

Correlation between BD-tau and pTau217 and fetal factors—birthweight, head circumference, and gestational age. Spearman correlation coefficients are shown. Red dot: elective Caesarean section (CS); blue dot: emergency CS; green dot: vaginal delivery.

Figure 3:

Correlation between tau and mode of delivery. A: Correlation between Log_{10} BD-tau and mode of delivery. Wilcoxon p-values are shown for comparisons between individual modes of delivery. B: Correlation between Log_{10} pTau217 and mode of delivery. Red dot: elective caesarean section (CS); blue dot: emergency CS; green dot: vaginal delivery.

588   Figure 4:

589   Correlation between cord blood biomarkers. Association of BD-tau with Neurofilament light

590   chain (NfL) (n=105), pH (n=104), base excess (n=94), and lactate (n=103); association of

591   pTau217 and NfL (n=106), pH (n=105), base excess (n=95), and lactate (n=104). Red dot:

592   elective caesarean section (CS); blue dot: emergency CS; green dot: vaginal delivery.