Linkage between HLA-B8 and HLA-DQ2.5 Contributes to Ancestry-Dependent Genetic Risk for Celiac Disease

Authors: Xin Long[1], Hemanth Karnati[1], Wenjing Ying[1], Mary-Joe Touma[1], Ioana Smith[1], Suzanne Lewis[2], Chao Xing[4], Ezra Burstein[1], Peter HR Green[2], Michele J. Alkalay[3], Alexandre Bolze[4], Xiao-Fei Kong[1,4]

Institutions:
1. Division of Digestive and Liver Diseases, Dept of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, TX, 75390
2. Celiac Disease Center, Columbia University, New York, NY, 10032
3. Division of Pediatric Gastroenterology, Hepatology, and Nutrition, Department of Pediatrics, University of Texas Southwestern Medical Center, Dallas, Texas 75235
4. McDermott Center for Human Growth and Development, University of Texas Southwestern Medical Center, Dallas, TX, 75390

Correspondence:
Xiao-Fei Kong, MD, Ph.D
Division of Digestive and Liver Diseases, Dept of Internal Medicine
McDermott Center for Human Growth and Development
UT Southwestern Medical Center, 5323 Harry Hines Blvd, Suite J5.150
Dallas, TX, 75390-9151; E-mail: Xiao-Fei.Kong@utsouthwestern.edu

Keywords: Celiac disease, Precision medicine, Genetic risk stratification

Author Contributions:

All authors provided important insights in the analysis and interpretation of the data and were involved in editing the manuscript. X.F.K. wrote the manuscript.

Disclosures: The authors have no relevant financial or non-financial interests to disclose.

Statement: The data and code used in this study are available as a shared workspace to registered researchers of the *All of Us* Researcher Workbench. For information about access, please visit https://www.researchallofus.org/. Computational codes are available from: https://github.com/Kong-Lab-UTSW/

Funding

This work was supported by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health (K08DK128631 and R03DK144282 to X.-F.K.) and by the Disease-Oriented Clinical Scholars Program at the University of Texas Southwestern Medical Center.

Acknowledgments

The *All of Us* research program is supported by the National Institutes of Health, Office of the Director: Regional Medical Centers: 1 OT2 OD026549; 1 OT2 OD026554; 1 OT2 OD026557; 1OT2 OD026556; 1 OT2 OD026550; 1 OT2 OD 026552; 1 OT2 OD026553; 1 OT2 OD026548; 1OT2 OD026551; 1 OT2 OD026555; IAA #: AOD 16037; Federally Qualified Health Centers: HHSN 263201600085U; Data and Research Center: 5 U2C OD023196; Biobank: 1 U24OD023121; The Participant Center: U24 OD023176; Participant Technology Systems Center: 1U24 OD023163; Communications and Engagement: 3 OT2 OD023205; 3 OT2 OD023206; and Community Partners: 1 OT2 OD025277; 3 OT2 OD025315; 1 OT2 OD025337; 1 OT2OD025276. The *All of Us* Research Program would not be possible without the partnership established with its participants.

Summary

Using data from the *All of Us Research Program*, we found that HLA-DQ2.5, a key genetic risk factor for celiac disease, is present in European (11.6%), Admixed American (8.1%), and African (8.1%) populations, but its disease penetrance differs significantly by ancestry.

Abstract

Background: Most genetic studies on celiac disease (CeD) have focused on individuals of European descent. Limited data are available for the Hispanic and black populations.

Methods: We analyzed whole-genome sequencing data, electronic health records (EHR), and laboratory results from the *All of Us* Research Program. We identified 3,481 individuals with CeD through EHR, self-reporting, or both. Of these, 2,899 carried one of the four well-established risk haplotypes, including 262 of admixed American (89% Hispanic) and 108 of African (70% black) ancestry. Five sex-, age-, and ancestry-matched controls per case were selected for the assessment of genetic and clinical risk factors.

Results: An enrichment in the *DQB1*02:01* allele was observed in CeD patients across all ancestries, with the strongest association in Europeans (32.3% vs. 11.6%), followed by Americans (18.5% vs. 8.1%) and Africans (15.7% vs. 8.1%). Among individuals carrying the DQ2.5 (*DQA1*05:01-DQB1*02:01* haplotype), HLA-B8 was present in 72.3% of Europeans, 42.3% of Admixed Americans, and lower in Africans. This linkage disequilibrium was higher in CeD patients than in controls across all three ancestries. A polygenic risk score distinguished seropositive CeD from controls with 86% accuracy. Incorporating clinical risk factors, including family history, hypothyroidism, diarrhea, vitamin D deficiency, and anemia, increased predictive accuracy to 92%. The model identified 93% of CeD patients with tTG-IgA levels greater than 10 IU/mL.

Conclusion: Linkage between HLA-B8 and DQ2.5 differs significantly among individuals of European, admixed American, and African ancestry, contributing to ancestry-dependent genetic risk for CeD.

Introduction

Celiac disease (CeD) has a worldwide prevalence of about 1%, ranging from 0.7 to 2%, depending on the tools used for screening and the country(1,2). However, most CeD cases remain undiagnosed(3). Clinically, the recognition of CeD early in life would be beneficial to patients because undiagnosed CeD can lead to various complications, including osteoporosis, iron deficiency anemia, poor quality of life, and even cancer(4,5). The only treatment available for CeD is adherence to a gluten-free diet (GFD), which is not always effective, as only 67% of individuals achieve mucosal healing after five years(6). Recent advances have led to the development of additional nomenclatures for describing the various phenotypes related to CeD, such as seronegative CeD, potential CeD, and refractory CeD, further complicating the diagnosis of this condition(7–9). The genetic basis of this clinical variability remains unclear, and the optimal clinical management for these phenotypes remains to be determined(10).

Human genetics plays a key role in the development of CeD. Twin studies have shown that genetic factors account for 70% of the overall risk of CeD(11,12), with CeD occurring in 10% of the first-degree relatives of index cases(13). *HLA-DQA1* and *HLA-DQB1* are two adjacent genes that encode the alpha and beta chains, respectively, to form HLA-DQ heterodimers. The four CeD-compatible HLA risk heterodimers encoded by four haplotypes include DQ2.5 (*DQA1*05:01-DQB1*02:01*), DQ2.2 (*DQA1*02:01-DQB1*02:02*), DQ8.1 (*DQA1*03:01-DQB1*03:02*), and DQ7.5 (*DQA1*05:05-DQB1*03:01*)(14), for which testing is recommended by current guidelines in various clinical scenarios (15,16). Throughout the manuscript, we use standard clinical shorthand

(e.g., 'DQ2.5') to denote HLA-DQ heterodimers. These are not gene symbols but heterodimers designations referring to specific *DQA1*/*DQB1* allele combinations, as defined by the WHO HLA nomenclature. HLA-DQ2.5 has made a major contribution to our understanding of the pathogenesis of CeD in recent decades, as 95% of European CeD patients carry HLA-DQ2.5 in a *cis* or *trans* configuration (17,18). However, most of these studies were performed in Europe and included participants from the United Kingdom, Italy, and the Netherlands, with very few individuals of other ancestries represented(18,19). This lack of ethnic diversity in genetic studies has limited our understanding of disease risk across different genetic backgrounds. *All of Us,* a large population-based genetic study, has enrolled more than 700,000 individuals from the United States, 46% of whom belong to underrepresented racial and minority ethnic groups (20). This NIH-supported project provides an unprecedented opportunity for advancing disease prevention and treatment and enhancing diversity in medical studies (20,21). We therefore made used of the *All of Us* datasets to investigate ancestry-specific genetic risks for CeD and to evaluate a risk score that integrates genetic variants, including HLA genotypes, along with clinical risk factors for the prediction of CeD risk across diverse populations.

Methods

Ethics Statement

The Ethics Committee/Institutional Review Board (IRB) of the *All of Us* Research Program (AoURP) gave its approval for the collection and analysis of data from human participants (AoU IRB Protocol Number: 2021-02-TN-001). This study was approved by the *All of Us* Research Program Science Committee. The results are reported in

accordance with the *All of Us* Data and Statistics Dissemination Policy and are displayed only for groups of at least 20 individuals.

Data sources

The AoURP aims to build one of the largest and most diverse health databases by enrolling one million adults and children from all backgrounds who have consented to participate and share their electronic health records (EHRs). These records include inpatient and outpatient visits, international classification of diseases (ICD) diagnostic codes, physician notes, laboratory results, and more[20,22–24]. Interested participants complete a basic survey. Numerous institutions across the United States are collaborating with *All of Us* to facilitate in-person visits, serving as enrollment centers where individuals can provide physical measurements and biological samples.

We analyzed controlled-tier datasets from the Curated Data Repository (CDR, C2024Q3R4 version), which includes survey responses, EHRs, measurements, and genomic data from participants enrolled between May 31, 2017, and Oct 01, 2023. Whole-genome sequencing (WGS) data were available for 414,830 participants (Fig. 1A), including compressed reference-oriented alignment map (CRAM) files and variant data store (VDS) files. Variant analysis was performed with the Hail framework. The control group consisted of participants without CeD. Individuals with autoimmune thyroid disorders, type 1 diabetes (T1D), selective IgA deficiency, inflammatory bowel disease (IBD), HIV infection, common variable immunodeficiency, cancer, Down syndrome, or DiGeorge syndrome were excluded from the control group. Genetic ancestry was inferred from SNP data by principal component analysis. Propensity score matching based on age,

sex and genetic ancestry was performed to identify appropriate controls. Tissue transglutaminase 2 (tTG) and deamidated gliadin peptide (DGP) antibody levels (tTG-IgA, DGP-IgA, and DGP-IgG) were analyzed to identify participants with seropositive CeD. Because those measurements can vary across laboratories and reagents, the cutoffs differ accordingly. Although the specific tTG-IgA assay kits used for these participants were not available, individuals with a reported positive result or with tTG-IgA, DGP-IgA, or DGP-IgG levels ≥20 IU/mL were classified as seropositive.

Ancestry-specific accuracy of HLA imputation

Participants with potential celiac disease, seronegative celiac disease, gluten intolerance, or non-celiac gluten sensitivity may have been labeled as CeD. To better characterize the cohort, we performed HLA typing using genomic data (20). We tested three approaches on 794 samples to obtain a fast accurate method for HLA typing: HLA genotype imputation with attribute bagging (HIBAG)(25), HLA*LA(26) and the tagSNPs(27) approach (S.Fig1).

We first investigated ancestry-specific HLA imputation accuracy by comparing tagSNP-based inference with HIBAG. One commonly used tagSNP, rs2187668 (27), is widely applied to infer HLA-DQ2.5 and showed excellent concordance in participants of European and African ancestry (98.5% and 97.3%, respectively), but substantially lower concordance in those of admixed American ancestry (62.7%, S.Table2). Consistent with the prior report (27), tagSNPs perform well for DQ7.5 and DQ8.1 but are suboptimal for DQ2.2 in individuals of European ancestry. Concordance between tagSNP and HIBAG was lower in participants of non-European ancestry (S.Table2).

HIBAG is a machine learning-based imputation tool that uses ancestry-specific, pre-trained classifiers to predict HLA genotypes from SNP data (28), whereas HLA-LA is a graph-based method that uses all the HLA reference sequences from the IMGT/HLA database to enable accurate high-resolution typing across diverse ancestries (26). For HIBAG, genetic data from subjects of European (n=2668), Asian (n=720), Hispanic (n=439) and African (n=173) ancestries were used in a machine-learning approach to develop accurate HLA prediction. With WGS data from 797 participants, concordance between HIBAG and HLA-LA exceeded 90% at the *HLA-A*, *B*, *C*, *DQB1*, and *DRB1* loci (S. Fig. 1). The largest discrepancy was observed for *HLA-DQA1*, largely reflecting reduced performance in distinguishing *DQA1*05:01 and DQA1*05:05*. We then leveraged known linkage disequilibrium (LD) among *HLA-DQA1*, *HLA-DQB1*, and *HLA-DRB1* to evaluate internal consistency. For the DQ2.5, *DQA1*05:01–DQB1*01:01–DRB1*03:01* linkage was strong (>99%) across individuals of European and admixed American ancestry (Fig 2A). In contrast, individuals with African ancestry showed lower linkage: among 119 *DQB1*02:01* alleles, 103(86.6%) were linked to *DQA1*05:01* (Fig 2A). This pattern suggests greater allelic diversity at the *HLA-DQA1* locus in African populations, which may also contribute to reduced *DQA1* concordance. We checked the internal consistency of the HLA-DQ haplotype by applying ancestry-specific linkage disequilibrium to *HLA-DQB1/DRB1*(29). The most common three-locus haplotypes are provided in the Supplementary Table3. Because HIBAG is cost-effective and supports ancestry-specific classifiers, whereas HLA-LA provides superior accuracy at a higher computational cost, we used HIBAG for HLA typing in the full cohort.

Statistical analysis

We evaluated genetic differences between the CeD and non-CeD participants, regardless of genetic ancestry and HLA-DQ risk genotypes and taking these factors into account. Chi-squared tests were performed for categorical variables. ANOVA was performed to compare differences in numerical variables across more than three groups, followed by Tukey's honestly significant difference post-hoc test to identify group means and their corresponding 95% confidence intervals. False discovery rate (FDR) correction was applied to adjust for multiple comparisons. Logistic regression analyses were performed to identify independent genetic risks and comorbid conditions, with age, sex, and the principal components of genetic ancestry as covariates. The prediction model was developed with a logistic regression framework combined with machine learning techniques to identify the optimal features. All analyses were conducted within the *All of Us* Researcher Workbench, a cloud-based platform providing data access(20,22). The analysis was performed with Python and the pandas, statsmodels, numpy, scikit-learn, matplotlib, and seaborn packages.

Results

Characteristics of participants with celiac disease in All of Us Research Program

The AoURP collaborates with hospitals, commercial laboratory networks, and community organizations to enroll one million U.S. residents across diverse racial and ethnic backgrounds. While the program is designed to be inclusive and now provides WGS data for 414,830 participants, not all participants completed the surveys. Because electronic health record (EHRs) systems vary widely across the nation, not all participants have EHRs or laboratory data. We identified 3,481 of these participants as

having CeD (Fig. 1A, S. Table 1) based on the diagnosis in their EHRs or self-report in surveys. As an exploratory cohort, 78% were female and 86% were white, both proportions being significantly higher than in the rest of the cohort (60% female, 57% white; S. Table1). The rate of CeD varied by ethnicity in the AoURP: 1.2% in white individuals, 0.4% in Hispanic individuals, and 0.2% in black individuals (S. Table 1). A clinical diagnosis of CeD typically requires serological and histological evidence(7,30). Serological data were unavailable for the majority of participants, 991 CeD participants have at least one available measurement (tTG-IgA, DGP-IgA, or DGP-IgG). Histological data concerning villous atrophy or crypt hyperplasia were unavailable.

CeD patients with compatible HLA-DQ genotypes as primary analytic cohort

Among 3,481 individuals with documented or self-reported CeD, the HLA-DQ2.5 haplotype was enriched in participants with EHR-confirmed or self-reported CeD compared to non-CeD controls (both 43% vs. 23%, S. Fig. 2). No significant differences were observed in sex at birth, genetic ancestry, or HLA-DQ genotype distribution between the two CeD groups. Therefore, we included all 2,899 (83.3%) participants with one of the four well-established HLA-DQ risk haplotypes, and 14,495 propensity-matched non-CeD controls were included for further analysis (Fig. 1A-B, S.Fig.3). In this CeD cohort, 1904 had an ICD-coded CeD diagnosis in their EHRs, 1,666 self-reported CeD in surveys, and 771 had CeD documented in both sources (Fig. 1C). 333 individuals have positive serology. In seropositive and HLA-compatible cases as the primary analytic cohort, we found that 2,495 of the individuals with CeD were of European ancestry, 262 were of admixed American ancestry (89% self-identified as Hispanic), 108 were of

African ancestry (71% self-identified as black), and 34 belonged to other ancestries (Fig. 1B).

The prevalent AH8.1 ancestry haplotype in CeD

Only 13 of HLA alleles identified had a frequency that was more than 1% higher in CeD cases than in non-CeD controls (Fig. 2B, S.Table 4). Notably, an enrichment in *DQA1*05:01* and *DQB1*02:01*, which form the DQ2.5 heterodimer, was observed in CeD patients across all ancestries (Fig. 2B). The allele frequencies of *DQB1*02:01* varied by ancestry among individuals with CeD and non-CeD controls: 32.2% vs 11.6% in Europeans (OR=3.6, FDR p= 3.76E-257), 18.5% vs 8.1% in admixed Americans (OR=2.6, FDR p= 1.84E-10, Fig.2C). Among participants of African ancestry, the *DQB1*02:01* allele had a frequency of 15.7% (34/216) in CeD cases versus 8.1% (85/1,054) in non-CeD controls (OR = 2.1; FDR p = 0.005).

In non-CeD controls of European ancestry, the *DQA1*05:01-DQB1*02:01 (*DQ2.5*)* haplotype was in strong LD with A*01:01(A1), B*08:01(B8), C07:01(C7), and DRB1*03:01(DR3). B8 was linked to DQ2.5 in 74.2% of controls. This extended haplotype A1-B8-DR3-DQ2, known as ancestral haplotype 8.1 (AH8.1), has been implicated in multiple autoimmune diseases (14,31). This linkage was less pronounced in non-CeD controls of admixed American ancestry (42.1%, S.Table5). B8 was rare among individuals of African ancestry, with only 27 B8 alleles observed in the entire cohort (S.Table5). All observed B8 alleles were linked to *DQB1*02:01*, with an estimated B8–DQ2.5 linkage frequency of 9.7%–27.1% in non-CeD controls.

Next, we found that the frequency of the B8–DQ2.5 linkage was higher in individuals with CeD than in non-CeD controls across all ancestry groups: 78.2% vs 74.2% in Europeans (OR = 1.24, 95% CI 1.07–1.43, FDR p = 0.01), 54.6% vs 42.1% in admixed Americans (OR = 1.66, 95% CI 1.02–2.69, FDR p = 0.039; Table S2). For participants of African ancestry, B8–DQ2.5 linkage was more frequent in CeD cases (OR = 2.88, 95% CI 1.16–7.13, FDR p = 0.03, S.Table 6); however, the number of African CeD participants carrying DQ2.5 was small (n = 32, S.Table 7). Consequently, this estimate is based on fewer than 20 B8–DQ2.5–positive CeD cases, yields wide confidence intervals, and should be interpreted with caution.

Although DQ2.5 was observed across ancestries (11.6% in Europeans, 8.1% in admixed Americans, and 7.2% in Africans), the B8–DQ2.5 haplotype was much less frequent: 8.6% in Europeans, 3.4% in admixed Americans, and <1.9% in Africans (S. Table 5). Given the corresponding CeD prevalence estimates (1.2% in Europeans, 0.4% in admixed Americans, and 0.2% in Africans), these ancestry-specific LD patterns may contribute to differences in CeD prevalence across ancestries.

Different clinical presentations associated with HLA-DQ risk genotypes

Our analysis of HLA-DQ genotypes showed that the highest risk of CeD was conferred by homozygosity for DQ2.5, followed by the DQ2.5/DQ2.2 genotype (Fig. 3A, S.Table 8). The combination of DQ2.5 and DQ2.2 was associated with a higher risk than DQ2.5/DQ7.5, consistent with a stronger additive effect of the DQB1*02 allele (32). The frequency of DQ2.5 carriers, either in *cis* (DQ2.5) or in *trans* (DQ2.2/7.5), was notably higher in CeD patients of European ancestry (62.5% vs 24.5%, Fig 3B), but this

difference was less pronounced in those of American (38.9% vs 20.1%) or African ancestry (36.1% vs 16.7%, Fig. 3B). Genotypes such as homozygosity or heterozygosity for DQ2.2 or DQ7.5 were more frequent in controls than in CeD patients, consistent with their risks as permissive genotypes (17,18,32–35). These results again suggest while individuals of different ancestry having the DQ2.5 heterodimer, their risk for celiac disease is different, consistent with the observation that ancestry-specific B8-DQ2.5 LD patterns may contribute to differences in CeD prevalence. We then analyzed laboratory test results and comorbid conditions for the individuals for whom data were available. Based on the distribution of HLA-DQ genotypes in CeD cases and controls, we categorized participants into three risk groups: high-, moderate-, and low- risk (Fig. 3A). Serum autoantibody levels can decline significantly after the introduction of a GFD (36). We therefore used the highest recorded value for each antibody (tTG-IgA, DGP-IgA, and DGP-IgG) when multiple measurements were available for a participant. Individuals with high- or moderate-risk HLA-DQ genotypes had significantly higher mean levels of these CeD-specific antibodies than those with low-risk genotypes, highlighting a strong correlation between HLA-DQ genotype and serological markers of CeD (Fig. 4A). Interestingly, individuals with low-risk genotypes had fewer CeD-related visits documented in their EHRs (Fig. 4A). We compared clinical characteristics between CeD patients in three risk groups. High-risk CeD patients were more likely to have type 1 diabetes (T1D) and a family history of CeD, whereas those in the low-risk group more frequently presented with symptoms such as irritable bowel syndrome (IBS), diarrhea, or migraines, features more consistent with functional gastrointestinal disorders (Fig. 4B, S.Table 9). These findings suggest that, in some cases, the CeD diagnosis code may have

been added to the EHR as part of diagnostic investigations rather than as a confirmed diagnosis. In summary, the high- and moderate- HLA-DQ risk genotypes are strongly associated with the presence of CeD-specific autoantibodies. Individuals with low-risk genotypes are more likely to display symptoms of functional gastrointestinal disorders, such as IBS, rather than true CeD.

Performance of a polygenic risk score (PRS) for predicting celiac disease across ancestries

A PRS derived from published GWAS and UK Biobank data, combining HLA-DQ genotypes with 38 non-HLA SNPs, effectively identifies individuals with CeD of European ancestry (17,18,32–35). We evaluated this log-additive PRS model in our cohort and confirmed that it could discriminate between 333 seropositive CeD cases and 676 non-CeD controls, achieving an area under the curve (AUC) of 0.86 (Fig. 5A- D). However, the model's performance declined when applied to the full cohort or to individual ancestry groups. The median PRS in seropositive CeD cases was approximately 5.07, consistent with previous findings (32). PRS scores were significantly higher in CeD patients of European ancestry, intermediate in those of American ancestry, and discriminated poorly between cases and controls of African ancestry (Fig. 5B). Individuals of African ancestry had the lowest mean PRS scores, consistent with the observed lower risk of developing CeD in this population(B-D). Using the threshold that optimized prediction accuracy, we found that 33% of European CeD patients and 15% of seropositive CeD cases did not exceed the PRS cutoff. In addition, 26% of non-CeD European controls had PRS scores above the threshold, which may be partly explained by

the presence of undiagnosed CeD among some controls. These findings suggest that, although the current PRS performs well in those with seropositive CeD (Fig 3D), the genetic risk score varies significantly in different populations, further work is needed for understanding the genetic risks for individuals of non-European ancestry.

CeD prediction model combining clinical and genetic risk factors

We then developed a predictive model incorporating HLA-DQ genotypes and a total of 50 features, including CeD-associated comorbid conditions. We evaluated machine-learning approaches, including decision trees and logistic regression, using seropositive CeD patients as positive cases (S. Methods). Logistic regression identified seven independent predictors: high- or moderate-risk HLA-DQ genotype, family history of CeD, diarrhea, vitamin D deficiency, anemia, and hypothyroidism. This model achieved an AUC of 0.87 for CeD prediction (Fig. 6A). As a means of improving prediction further, we developed a composite log-additive AoU-CeD score that integrates family history, four clinical diagnoses, and PRS. This combined model outperformed the PRS alone. Using a threshold score of 4.17 to achieve 90% sensitivity for the detection of seropositive CeD, we assessed the performance of our model in patients lacking strong serological biomarkers. The threshold was reached for 77% of CeD patients of European ancestry, with lower proportions attained in the American and African ancestry groups (Fig. 6C).

We then validated the model by investigating its relationship to tTG-IgA levels. We found that 186 CeD patients had tTG-IgA levels >20 IU/mL, and 45 had levels between 10–20 IU/mL. Among those with tTG-IgA levels >10 IU/mL, 93% surpassed the

predictive threshold, versus 77% with levels <5 IU/mL. These results support the utility of this model for stratifying seropositive CeD cases.

Discussion

A major advantage of the *All of Us* dataset is its enormous volume of EHRs, genomic and survey data, including the most comprehensive set of information available for populations historically underrepresented in medical research. This enabled us to study CeD patients from diverse ancestries in this first study to investigate extensive genetic data from patients from African and admixed American ancestry.

The *All of Us* data confirm that CeD diagnosis is positively associated with sex (more frequent in women) and socioeconomic status but negatively associated with black or Hispanic ethnicity (2,28,37). Clinical epidemiological studies, based principally on the serological marker tTG-IgA, have shown that the prevalence of seropositive CeD varies across countries and racial/ethnic group. The prevalence of seropositivity ranges from 0.3% in Germany, to 2.4% in Finland (38), and 2% in Norway (HUNT study) (38), with an overall estimated prevalence of 1% in Europe, 0.7% in the United States, 0.5% in South America, 0.34% to 1.1% in Africa, and 0.36% in China (2). In the All of Us cohort, CeD prevalence varied by self-reported race/ethnicity—1.2% in White individuals, 0.4% in Hispanic individuals, and 0.2% in Black individuals—which is consistent with findings from the National Health and Nutrition Examination Survey (NHANES) 2009–2012 (39). In NHANES, based on tTG and endomysial (EMA) IgA antibody testing in 14,701 participants, CeD prevalence was highest among White individuals (1.08%) and

substantially lower among Mexican Americans (0.23%), other Hispanics (0.38%), and Black individuals (0.22%) (39).

This raises the question as to what drives ancestry-specific variation in seropositivity rates. HLA-DQ2.5 frequency is only modestly lower across ancestries—11.6% in Europeans, 8.1% in individuals of admixed American ancestry, and 7.2% in Africans—yet CeD prevalence in Africans and admixed Americans appears disproportionately lower than would be expected based on DQ2.5 carriage alone. One possible explanation lies in the LD patterns of HLA alleles. The high-risk haplotype (B8: C7: DR3: DQ2.5) is present in 72.3% of DQ2.5-positive individuals of European ancestry, but in less than 20% of those of African ancestry. Our findings suggest a potential explanation: individuals of African ancestry may carry the canonical DQ2.5 risk heterodimer, yet reduced linkage to B8—or to B8-associated genetic factors—could influence the likelihood or magnitude of a tTG-IgA production. For example, the *MICA*5*.1 allele is in strong linkage disequilibrium with B8 and may modify DQ2.5-mediated antigen presentation (40). An additional possibility is that patients with CeD of non-European ancestry may not mount tTG-IgA responses comparable to those of European ancestry. In NHANES, a discordant pattern between tTG-IgA and EMA-IgA was observed among participants of African ancestry(39), and in an Alabama-based study (in which 26% of residents were of African ancestry), tTG-IgA was more frequently reported as negative (41). Undermeasurement and seronegative disease have therefore been proposed as alternative explanations for the apparently lower frequency of CeD among individuals of African descent (42). Finally, tTG-IgA quantification is not fully standardized: upper limits of normal vary

substantially across laboratories, commercial assays, and serum dilution protocols, complicating the definition of reliable cutoff values(36,43,44). This inter-assay variability may contribute to false-negative serologic results and may disproportionately hinder accurate CeD ascertainment in individuals of non-European ancestry.

HLA-DQ testing is included in both the American College of Gastroenterology (ACG) and American Gastroenterological Association (AGA) guidelines for the diagnostic evaluation of CeD (15,16). According to these guidelines, genetic testing for CeD-compatible HLA haplotypes is not necessary in all cases but may be useful in specific situations, such as cases in which there is a discrepancy between serology and histology results, or in individuals who have already switched to a GFD before evaluation. A negative result of HLA-DQ genotypes effectively rules out CeD. The presence of a CeD-compatible haplotype may support gluten challenge in appropriate clinical contexts. We found that 17% of individuals with a CeD diagnosis did not carry any of the four historically defined CeD-compatible HLA-DQ haplotypes. Conversely, about 70% of non-CeD controls carried these haplotypes. Our data suggest that CeD is frequently misdiagnosed in the United States; in particular, many patients lack serologic evidence of CeD and may carry low-risk HLA-DQ genotypes.

Can predictive models improve CeD risk stratification? Prediction models for CeD have been developed based on symptoms, comorbid conditions (45,46), or polygenic risk scores (32). Our preliminary prediction model supports a case-finding approach, with effective risk stratification and targeted identification of high-risk individuals. A cost-

effectiveness analysis showed that testing children with a pre-test probability ≥10% by both HLA typing and tTG-IgA was the most effective approach (47). HLA testing before tTG-IgA testing was the most cost-effective approach for predicting CeD (48). As genome sequencing becomes more affordable, it will become possible to use HLA risk stratification to optimize population-level CeD screening further (44). Further studies are required to assess whether AoU-CeD score combining clinical factors and PRS can aid in clinical management in independent cohorts. For example, can the score identify patients with an ambiguous diagnosis who are most likely to benefit from a gluten challenge, and how does its predictive performance compare with established endpoints such as seropositivity and histologic response? In addition, is the score concordant with tTG-IgA levels when screening first-degree relatives?

However, there are important limitations in our study. Key clinical details are missing for many participants, such as pathology reports for endoscopy examinations and dietary information, including whether patients are already following a gluten-free diet or how long they have been on it. Future studies incorporating dietary factors would be highly valuable. Other limitations include incomplete survey participation and limited availability of EHRs for review; moreover, only a small subset of patients had serologic measurements of tTG and DGP antibodies. Another concern is the potential inaccuracy of EHR-derived phenotypes: diagnostic codes may be assigned primarily for billing purposes rather than to reflect a confirmed diagnosis. Some patients labeled with CeD may have been diagnosed with seronegative or potential CeD, rather than seropositive and histologically confirmed disease. We observed that chronic conditions associated with functional gastrointestinal disorders were prevalent, affecting 35.8% of individuals

with low-risk *HLA-DQ* genotypes. This suggests that a subset of patients labeled with CeD may actually have other conditions, such as non-celiac gluten sensitivity(49) or functional gastrointestinal disorders(48). These findings highlight the need for more stringent criteria for identifying CeD patients in future genetic studies, particularly among patients with non-European ancestry.

In conclusion, genetic risks can differ significantly across individuals of diverse ancestry, and integrating polygenic risk scores with family history, symptoms, and comorbid conditions provides a promising approach for improving the screening and diagnosis of celiac disease.

References

1. Singh P, Arora A, Strand TA, Leffler DA, Catassi C, Green PH, et al. Global Prevalence of Celiac Disease: Systematic Review and Meta-analysis. Clin Gastroenterol Hepatol [Internet]. 2018 Jun 1 [cited 2024 Apr 1];16(6):823-836.e2. Available from: http://www.cghjournal.org/article/S1542356517307838/fulltext

2. Gatti S, Rubio-Tapia A, Makharia G, Catassi C. Patient and Community Health Global Burden in a World With More Celiac Disease. Gastroenterology [Internet]. 2024 Jun 1 [cited 2025 Mar 30];167(1):23–33. Available from: https://pubmed.ncbi.nlm.nih.gov/38309629/

3. Choung RS, Larson SA, Khaleghi S, Rubio-Tapia A, Ovsyannikova IG, King KS, et al. Prevalence and Morbidity of Undiagnosed Celiac Disease From a Community-Based Study. Gastroenterology. 2017 Mar 1;152(4):830-839.e5.

4. Lebwohl B, Sanders DS, Green PHR. Coeliac disease. www.thelancet.com [Internet]. 2018 [cited 2020 Oct 9];391. Available from: https://www.coeliac.org.uk/home

5. Caio G, Volta U, Sapone A, Leffler DA, De Giorgio R, Catassi C, et al. Celiac disease: a comprehensive current review. BMC Med. 2019 Jul 23;17(1):142.

6. Rubio-Tapia A, Rahim MW, See JA, Lahr BD, Wu TT, Murray JA. Mucosal

Recovery and Mortality in Adults with Celiac Disease after Treatment with a Gluten-Free Diet. Am J Gastroenterol [Internet]. 2010 Jun [cited 2024 Apr 1];105(6):1412. Available from: /pmc/articles/PMC2881171/

7. Elli L, Leffler D, Cellier C, Lebwohl B, Ciacci C, Schumann M, et al. Guidelines for best practices in monitoring established coeliac disease in adult patients. Nat Rev Gastroenterol Hepatol 2023 213 [Internet]. 2023 Dec 18 [cited 2024 Apr 1];21(3):198–215. Available from: https://www.nature.com/articles/s41575-023-00872-2

8. Schiepatti A, Sanders DS, Baiardi P, Caio G, Ciacci C, Kaukinen K, et al. Nomenclature and diagnosis of seronegative coeliac disease and chronic non-coeliac enteropathies in adults: the Paris consensus. Gut [Internet]. 2022 Nov 1 [cited 2024 Apr 1];71(11):2218–25. Available from: https://gut.bmj.com/content/71/11/2218

9. Husby S, Murray JA, Katzka DA. AGA Clinical Practice Update on Diagnosis and Monitoring of Celiac Disease—Changing Utility of Serology and Histologic Measures: Expert Review. Gastroenterology [Internet]. 2019 Mar 1 [cited 2024 Apr 1];156(4):885–9. Available from: http://www.gastrojournal.org/article/S0016508518354088/fulltext

10. Lundin KEA, Green PHR. Seronegative coeliac disease and non-coeliac enteropathies: precision medicine, precision medicine, where are you? Gut [Internet]. 2022 Nov 1 [cited 2024 Apr 1];71(11):2148–9. Available from: https://gut.bmj.com/content/71/11/2148

11. Greco L, Romino R, Coto I, Di Cosmo N, Percopo S, Maglio M, et al. The first large population based twin study of coeliac disease. Gut. 2002;50(5):624–8.

12. Kuja-Halkola R, Lebwohl B, Halfvarson J, Wijmenga C, Magnusson PKE, Ludvigsson JF. Heritability of non-HLA genetics in coeliac disease: A population-based study in 107 000 twins. Gut. 2016 Nov 1;65(11):1793–8.

13. Rubio-Tapia A, Van Dyke CT, Lahr BD, Zinsmeister AR, El-Youssef M, Moore SB, et al. Predictors of Family Risk for Celiac Disease: A Population-Based Study. Clin Gastroenterol Hepatol [Internet]. 2008 Sep [cited 2020 Jan 12];6(9):983–7. Available from: www.cghjournal.org

14.   Sollid LM. The roles of MHC class II genes and post-translational modification in celiac disease. Immunogenetics [Internet]. 2017 Aug 1 [cited 2023 Jun 2];69(8–9):605–16. Available from: https://pubmed.ncbi.nlm.nih.gov/28695286/

15.   Rubio-Tapia A, Hill ID, Kelly CP, Calderwood AH, Murray JA. ACG clinical guidelines: Diagnosis and management of celiac disease. Am J Gastroenterol [Internet]. 2013 May [cited 2025 Mar 30];108(5):656–76. Available from: https://journals.lww.com/ajg/fulltext/2013/05000/acg_clinical_guidelines__diagnosis_and_management.7.aspx

16.   Leonard MM, Lebwohl B, Rubio-Tapia A, Biagi F. AGA Clinical Practice Update on the Evaluation and Management of Seronegative Enteropathies: Expert Review. Gastroenterology [Internet]. 2021 Jan 1 [cited 2025 Mar 30];160(1):437–44. Available from: https://www.gastrojournal.org/action/showFullText?pii=S0016508520352203

17.   Trynka G, Hunt KA, Bockett NA, Romanos J, Mistry V, Szperl A, et al. Dense genotyping identifies and localizes multiple common and rare variant association signals in celiac disease. Nat Genet [Internet]. 2011 Dec [cited 2023 Jun 2];43(12):1193–201. Available from: https://pubmed.ncbi.nlm.nih.gov/22057235/

18.   Withoff S, Li Y, Jonkers I, Wijmenga C. Understanding Celiac Disease by Genomics. Trends Genet [Internet]. 2016;32(5):295–308. Available from: http://dx.doi.org/10.1016/j.tig.2016.02.003

19.   Sollis E, Mosaku A, Abid A, Buniello A, Cerezo M, Gil L, et al. The NHGRI-EBI GWAS Catalog: knowledgebase and deposition resource. Nucleic Acids Res [Internet]. 2023 Jan 6 [cited 2024 Jun 2];51(D1):D977–85. Available from: https://pubmed.ncbi.nlm.nih.gov/36350656/

20.   Genomic data in the All of Us Research Program. Nature [Internet]. 2024 Mar 1 [cited 2024 Jun 2];627(8003):340–6. Available from: https://pubmed.ncbi.nlm.nih.gov/38374255/

21.   Bianchi DW, Brennan PF, Chiang MF, Criswell LA, D'Souza RN, Gibbons GH, et al. The All of Us Research Program is an opportunity to enhance the diversity of US biomedical research. Nat Med. 2024 Feb;30(2):330–3.

22.   Park J, Karnati H, Rustgi SD, Hur C, Kong XF, Kastrinos F. Impact of population

screening for Lynch syndrome insights from the All of Us data. Nat Commun [Internet]. 2025 Jan 9 [cited 2025 Feb 16];16(1):523. Available from: https://pubmed.ncbi.nlm.nih.gov/39788943/

23. Denny JC, Rutter JL, Goldstein DB, Philippakis A, Smoller JW, Jenkins G, et al. The "All of Us" Research Program. N Engl J Med [Internet]. 2019 Aug 15 [cited 2020 Apr 26];381(7):668–76. Available from: http://www.nejm.org/doi/10.1056/NEJMsr1809937

24. Ramirez AH, Sulieman L, Schlueter DJ, Halvorson A, Qian J, Ratsimbazafy F, et al. The All of Us Research Program: Data quality, utility, and diversity. Patterns (New York, NY) [Internet]. 2022 Aug 12 [cited 2024 Jun 2];3(8). Available from: https://pubmed.ncbi.nlm.nih.gov/36033590/

25. Zheng X, Shen J, Cox C, Wakefield JC, Ehm MG, Nelson MR, et al. HIBAG--HLA genotype imputation with attribute bagging. Pharmacogenomics J [Internet]. 2014 [cited 2024 May 16];14(2):192–200. Available from: https://pubmed.ncbi.nlm.nih.gov/23712092/

26. Dilthey AT, Mentzer AJ, Carapito R, Cutland C, Cereb N, Madhi SA, et al. HLA*LA-HLA typing from linearly projected graph alignments. Bioinformatics [Internet]. 2019 Nov 1 [cited 2023 Jun 2];35(21):4394–6. Available from: https://pubmed.ncbi.nlm.nih.gov/30942877/

27. Monsuur AJ, de Bakker PIW, Zhernakova A, Pinto D, Verduijn W, Romanos J, et al. Effective Detection of Human Leukocyte Antigen Risk Alleles in Celiac Disease Using Tag Single Nucleotide Polymorphisms. PLoS One [Internet]. 2008 May 28 [cited 2024 Jun 2];3(5). Available from: /pmc/articles/PMC2386975/

28. Zylberberg HM, Miller EBP, Ratner A, Hammill BG, Mehta P, Alesci S, et al. Correlations Between Relative Prevalence of Celiac Disease and Sociodemographic Variables in the United States. Am J Gastroenterol. 2024 Feb 1;119(2):374–7.

29. Gonzalez-Galarza FF, McCabe A, Santos EJM Dos, Jones J, Takeshita L, Ortega-Rivera ND, et al. Allele frequency net database (AFND) 2020 update: gold-standard data classification, open access genotype data and new query tools. Nucleic Acids Res [Internet]. 2020 Jan 1 [cited 2025 Mar 30];48(D1):D783–8.

Available from: https://pubmed.ncbi.nlm.nih.gov/31722398/

30. Rubio-Tapia A, Hill ID, Semrad C, Kelly CP, Greer KB, Limketkai BN, et al. American College of Gastroenterology Guidelines Update: Diagnosis and Management of Celiac Disease. Am J Gastroenterol. 2023 Jan 1;118(1):59–76.

31. Price P, Witt C, Allcock R, Sayer D, Garlepp M, Kok CC, et al. The genetic basis for the association of the 8.1 ancestral haplotype (A1, B8, DR3) with multiple immunopathological diseases. Immunol Rev [Internet]. 1999 Feb 1 [cited 2024 Sep 27];167(1):257–74. Available from: https://onlinelibrary.wiley.com/doi/full/10.1111/j.1600-065X.1999.tb01398.x

32. Sharp SA, Jones SE, Kimmitt RA, Weedon MN, Halpin AM, Wood AR, et al. A single nucleotide polymorphism genetic risk score to aid diagnosis of coeliac disease: a pilot study in clinical care. Aliment Pharmacol Ther [Internet]. 2020 Oct 1 [cited 2024 Apr 2];52(7):1165–73. Available from: https://pubmed.ncbi.nlm.nih.gov/32790217/

33. Dubois PCA, Trynka G, Franke L, Hunt KA, Romanos J, Curtotti A, et al. Multiple common variants for celiac disease influencing immune gene expression. Nat Genet. 2010 Apr;42(4):295–302.

34. Gutierrez-Achury J, Zhernakova A, Pulit SL, Trynka G, Hunt KA, Romanos J, et al. Fine mapping in the MHC region accounts for 18% additional genetic risk for celiac disease. Nat Genet [Internet]. 2015 May 27 [cited 2024 Nov 25];47(6):577–8. Available from: https://pubmed.ncbi.nlm.nih.gov/25894500/

35. van Heel DA, Franke L, Hunt KA, Gwilliam R, Zhernakova A, Inouye M, et al. A genome-wide association study for celiac disease identifies risk variants in the region harboring IL2 and IL21. Nat Genet. 2007 Jul;39(7):827–9.

36. Gong C, Saborit C, Long X, Wang A, Zheng B, Chung H, et al. Serological Investigation of Persistent Villous Atrophy in Celiac Disease. Clin Transl Gastroenterol [Internet]. 2023 Sep 27 [cited 2023 Oct 3]; Available from: https://pubmed.ncbi.nlm.nih.gov/37753949/

37. Roy A, Mehra S, Kelly CP, Tariq S, Pallav K, Dennis M, et al. The association between socioeconomic status and the symptoms at diagnosis of celiac disease: a retrospective cohort study. Therap Adv Gastroenterol [Internet]. 2016 Jul 1 [cited

2024 Jun 2];9(4):495–502. Available from:
https://pubmed.ncbi.nlm.nih.gov/27366218/

38.   Mustalahti K, Catassi C, Reunanen A, Fabiani E, Heier M, McMillan S, et al. The
       prevalence of celiac disease in Europe: Results of a centralized, international mass
       screening project. Ann Med [Internet]. 2010 Dec [cited 2025 Mar 30];42(8):587–
       95. Available from:
       https://www.tandfonline.com/doi/abs/10.3109/07853890.2010.505931

39.   Mardini HE, Westgate P, Grigorian AY. Racial Differences in the Prevalence of
       Celiac Disease in the US Population: National Health and Nutrition Examination
       Survey (NHANES) 2009–2012. Dig Dis Sci 2015 606 [Internet]. 2015 Jan 11
       [cited 2025 Nov 2];60(6):1738–42. Available from:
       https://link.springer.com/article/10.1007/s10620-014-3514-7

40.   Hüe S, Mention JJ, Monteiro RC, Zhang SL, Cellier C, Schmitz J, et al. A direct
       role for NKG2D/MICA interaction in villous atrophy during celiac disease.
       Immunity [Internet]. 2004 [cited 2026 Jan 24];21(3):367–77. Available from:
       https://pubmed.ncbi.nlm.nih.gov/15357948/

41.   Cartee AK, Beasley TM, Estes D, Herman TT, Morgan DR. Black Patients Are
       More Likely to Have Negative Serologies With Celiac Consistent Histology in a
       Southeast US Tertiary Center. Gastro Hep Adv. 2024;3(1):4–6.

42.   Lebwohl B. Celiac Disease and the Forgotten 10%: The "Silent Minority." Dig Dis
       Sci [Internet]. 2015 Jun 5 [cited 2024 Nov 27];60(6):1517. Available from:
       https://pmc.ncbi.nlm.nih.gov/articles/PMC4456229/

43.   Silvester JA, Kurada S, Szwajcer A, Kelly CP, Leffler DA, Duerksen DR. Tests
       for Serum Transglutaminase and Endomysial Antibodies Do Not Detect Most
       Patients With Celiac Disease and Persistent Villous Atrophy on Gluten-free Diets:
       a Meta-analysis. Gastroenterology [Internet]. 2017 Sep 1 [cited 2023 Jun
       2];153(3):689-701.e1. Available from: https://pubmed.ncbi.nlm.nih.gov/28545781/

44.   Chou R, Bougatsos C, Blazina I, Mackey K, Grusing S, Selph S. Screening for
       Celiac Disease: Evidence Report and Systematic Review for the US Preventive
       Services Task Force. JAMA [Internet]. 2017 Mar 28 [cited 2024 Apr
       1];317(12):1258–68. Available from: https://pubmed.ncbi.nlm.nih.gov/28350935/

45. Ludvigsson JF, Yao J, Lebwohl B, Green PHR, Yuan S, Leffler DA. Coeliac disease: complications and comorbidities. Nat Rev Gastroenterol Hepatol 2025 [Internet]. 2025 Jan 28 [cited 2025 Feb 16];1–13. Available from: https://www.nature.com/articles/s41575-024-01032-w

46. Elwenspoek MMC, O'Donnell R, Jackson J, Everitt H, Gillett P, Hay AD, et al. Development and external validation of a clinical prediction model to aid coeliac disease diagnosis in primary care: An observational study. EClinicalMedicine [Internet]. 2022 Apr 1 [cited 2024 Nov 26];46. Available from: https://pubmed.ncbi.nlm.nih.gov/35434586/

47. Elwenspoek MMC, Thom H, Sheppard AL, Keeney E, O'Donnell R, Jackson J, et al. Defining the optimum strategy for identifying adults and children with coeliac disease: systematic review and economic modelling. Health Technol Assess [Internet]. 2022 [cited 2024 Nov 26];26(44):vii–164. Available from: https://pubmed.ncbi.nlm.nih.gov/36321689/

48. Black CJ, Drossman DA, Talley NJ, Ruddy J, Ford AC. Functional gastrointestinal disorders: advances in understanding and management. Lancet [Internet]. 2020 Nov 21 [cited 2024 Nov 27];396(10263):1664–74. Available from: http://www.thelancet.com/article/S0140673620321152/fulltext

49. Catassi C, Elli L, Bonaz B, Bouma G, Carroccio A, Castillejo G, et al. Diagnosis of Non-Celiac Gluten Sensitivity (NCGS): The Salerno Experts' Criteria. Nutrients. 2015 Jun 18;7(6):4966–77.

Figures and legends

Figure 1. CONSORT flow diagram and cohort demographics for CeD analysis

(A) CONSORT-style flow diagram illustrating the steps of data processing and cohort selection. Propensity score matching was performed based on age, sex, and genetic ancestry. Confounding was reduced by excluding individuals with comorbidities

commonly associated with CeD, including T1D, autoimmune thyroid disorders, selective IgA deficiency, IBD, CVID, cancer, and HIV infection, from the non-CeD control group.

(B) Demographic characteristics of 5:1 propensity matched Non-CeD controls and CeD patients from *All of Us*. GED, General Educational Development.

(C) Venn diagram showing the route of CeD diagnosis for the included participants. "Seropositive CeD" refers to individuals with tTG-IgA or DGP IgA or IgG levels >20 IU/mL documented in their EHRs.

(D) Age, sex, and genetic ancestry of the 1:5 propensity score–matched CeD cases and controls.

A

DQA1*05:01-DQB1*02:01-DRB1*03:01 haplotype

Linkage	European	Admixed American	African
Total *DQB1*02:01* (n)	4504	311	119
With *DRB1*03:01* (n, %)	4501 (99.9%)	310 (99.7%)	105 (88.2%)
With *DQA1*05:01* (n, %)	4476 (99.4%)	309 (99.4%)	103 (86.6%)
With both *DRB1*03:01* and *DQA1*05:01* (n, %)	4475 (99.4%)	309 (99.4%)	103 (86.6%)

B

HLA Allele	Non-CeD (n=14,495)	CeD (n=2,899)	OR(95%CI)
DQA1*05:01	30.67%	11.16%	3.52(3.30-3.77)
DQB1*02:01	30.25%	11.11%	3.47(3.24-3.71)
DRB1*03:01	30.25%	11.15%	3.46(3.23-3.70)
B*08:01	24.39%	10.02%	2.90(2.70-3.11)
C*07:01	27.73%	15.27%	2.13(1.99-2.27)
DPB1*01:01	11.28%	5.69%	2.11(1.92-2.32)
DQA1*03:01	17.28%	10.34%	1.81(1.68-1.96)
A*01:01	23.04%	14.56%	1.76(1.64-1.88)
DQB1*02:02	14.13%	10.29%	1.43(1.32-1.56)
DQB1*03:02	13.82%	10.56%	1.36(1.25-1.48)

C

Figure 2. HLA alleles and linkage disequilibrium patterns with DQ2.5 in CeD patients and matched controls across ancestries. (A) Cross-ancestry LD between HLA-DQA1, HLA-DQB1, and HLA-DRB1 in DQ2.5 carriers.

(B) Table showing HLA alleles from the HLA-*A, -B, -C, -DPB1, -DQA1, -DQB1*, and -*DRB1* loci overrepresented in individuals with CeD, with a frequency at least 1% higher in such individuals than in matched controls. (C) Allele frequencies for HLA-DQ2.5 and other alleles in linkage disequilibrium with HLA-DQ2.5 (A1, B8, C7, DRB3 and DPB1) across different genetic ancestries and CeD statuses.

A

	DQ	HLA-DQ haplotypes	CeD (n=2,899)	Non-CeD (n=14,495)	Odds ratio [95% CI]
High	2.5/2.5	*DQA1*05:01-DQB1*02:01/DQA1*05:01-DQB1*02:01*	184 (6.4%)	160 (1.1%)	8.1 [6.3–10.3]
	2.5/2.2	*DQA1*05:01-DQB1*02:01/DQA1*02:01-DQB1*02:02*	264 (9.1%)	337 (2.3%)	5.5 [4.5–6.7]
Moderate	2.5/X	*DQA1*05:01-DQB1*02:01/X*	795 (27.4%)	1784 (12.3%)	3.1 [2.7–3.6]
	2.2/7.5	*DQA1*02:01-DQB1*02:02/DQA1*05:05-DQB1*03:01*	149 (5.1%)	389 (2.7%)	2.7 [2.2–3.4]
	2.5/7.5	*DQA1*05:01-DQB1*02:01/DQA1*05:05-DQB1*03:01*	156 (5.4%)	456 (3.2%)	2.4 [1.9–3.0]
	2.5/8.1	*DQA1*05:01-DQB1*02:01/DQA1*03:01-DQB1*03:02*	166 (5.7%)	316 (2.2%)	3.7 [3.0–4.6]
	8.1/8.1	*DQA1*03:01-DQB1*03:02/DQA1*03:01-DQB1*03:02*	69 (2.4%)	173 (1.2%)	2.8 [2.1–3.8]
	2.2/8.1	*DQA1*02:01-DQB1*02:02/DQA1*03:01-DQB1*03:02*	93 (3.2%)	303 (2.1%)	2.2 [1.7–2.8]
Low	8.1/X	*DQA1*03:01-DQB1*03:02/X*	344 (11.9%)	1662 (11.5%)	1.5 [1.2–1.7]
	7.5/8.1	*DQA1*05:05-DQB1*03:01/DQA1*03:01-DQB1*03:02*	57 (2.0%)	407 (2.8%)	1.0 [0.7–1.3]
	2.2/X	*DQA1*02:01-DQB1*02:02/X*	249 (8.6%)	1547 (10.7%)	1.1 [0.9–1.4]
	7.5/X	*DQA1*05:05-DQB1*03:01/X*	313 (10.8%)	2202 (15.2%)	Reference
	2.2/2.2	*DQA1*02:01-DQB1*02:02/DQA1*02:01-DQB1*02:02*	22 (0.8%)	161 (1.1%)	1.0 [0.6–1.5]
	7.5/7.5	*DQA1*05:05-DQB1*03:01/DQA1*05:05-DQB1*03:01*	38 (1.3%)	290 (2.0%)	0.9 [0.6–1.3]
None	X/X	*X/X*	0 (0.0%)	4308 (29.7%)	

B

Figure 3. HLA-DQ genotype distribution and ancestry-specific effects in CeD. (A) Distribution of HLA-DQ genotypes based on the four known CeD-compatible HLA-DQ haplotypes among individuals with and without CeD. Genotypes were further categorized by risk level: high, moderate, low, or no risk (none). HLA-DQ 7.5/X was used as the reference, and p-values were FDR-adjusted. (B) HLA-DQ genotype distribution among seropositive CeD patients and across genetic ancestry groups (European, Admixed American, and African) for both the CeD and control cohorts. The frequency of individuals carrying HLA-DQ2.5, either in *cis* or *in trans*, is presented for each group.

Figure 4. The frequencies of serological markers and comorbid clinical conditions differ between HLA-DQ genotypes in individuals with CeD

(A) ANOVA was performed to compare differences in serological markers (tTG-IgA, DGP-IgA, DGP-IgG) and the number of clinical events classified as CeD recorded in the EHRs across HLA-DQ genotype categories. Tukey's honestly significant difference post-hoc test was then performed to obtain mean values (black dots) and 95% confidence intervals (lines) for each group. Blue dotted lines indicate an arbitrary threshold value of 20 IU/mL. Visit counts represent unique encounters documented in the EHR on distinct dates. (D) Frequency of comorbid conditions across HLA-DQ risk categories in CeD patients, grouped into 'high-risk', 'moderate-risk' and 'low risk'. A bubble plot displays the odds ratios and FDR adjusted p-values, using the high-risk group as the reference.

Figure 5. Evaluation of a reported PRS prediction model in participants from the

***All of Us* project with CeD participants**

(A) ROC curves showing the performance of the PRS for distinguishing CeD cases from

controls. The green line represents the ROC curve obtained with seropositive CeD cases

considered positive and their matched non-CeD controls considered negative. PRS

performance is evaluated across different ancestry subgroups: European, admixed

American, and the combined cohort (regardless of ancestry).

(B) Violin plots comparing PRS distributions between CeD patients and matched controls.

PRS scores from seropositive CeD patients or those with European (Eur), admixed

American (Amr), or African (Afr) ancestry were compared to their respective matched non-CeD controls. Mean PRS values are indicated by white dots. The dotted line represents the optimal PRS threshold with the highest accuracy for distinguishing seropositive cases from non-CeD controls. Percentages reflect the proportion of individuals in each group with PRS values above this threshold. The black boxes represent the interquartile range (25th to 75th percentile) of PRS values.

(C) Platt post hoc probability calibration was applied by fitting a logistic regression that maps the model's raw score (logit) to calibrated event probabilities. Calibration performance is summarized by the Brier score (mean squared error between predicted probabilities and observed outcomes), with lower values indicating better probabilistic accuracy. Bins centered at higher mean predicted risk indicate that the model assigns higher average risk in that subgroup (e.g., mean predicted risk >0.8 in the seropositive group).

(D) Performance of the PRS for predicting CeD across subgroups defined by genetic ancestry and serologic status.

Figure 6. The AoU-CeD score combines clinical risk factors and PRS to improve CeD prediction over and above that achieved with the PRS alone

(A) Logistic regression and machine-learning approaches were used to identify the variables that best predicted CeD. The input variables included sex, age, genetic ancestry, HLA-DQ genotype, and 42 clinical features, including symptoms, comorbid conditions and complications related to malabsorption. Seropositive CeD participants were used as positive cases. A ROC curve was plotted and the model coefficients for the top predictors are reported.

(B) We evaluated the extent to which the incorporation of clinical data improved prediction over that achieved with the PRS alone, by testing models including family history (FH) and seven clinical diagnoses (Dx). ROC curves comparing three scoring methods: PRS alone, PRS + FH, and PRS + FH + Dx. Scores were calculated with a log-additive model, combining log-transformed values weighted by logistic regression coefficients. **(C)** Violin plots showing AoU-CeD score distributions in seropositive CeD

cases versus matched non-CeD controls, using a threshold capturing 90% of seropositive CeD patients. The AoU-CeD score was then applied to CeD patients without high levels of serological markers (>20 IU/mL), stratified by ancestry. **(D)** Validation of the AoU-CeD score in CeD patients from *All of Us* with available but variable tTG-IgA values recorded in their EHRs.

A

	DQ	HLA-DQ haplotypes	CeD (n=2,899)	Non-CeD (n=14,495)	Odds ratio [95% CI]
High	2.5/2.5	*DQA1*05:01-DQB1*02:01/DQA1*05:01-DQB1*02:01*	184 (6.4%)	160 (1.1%)	8.1 [6.3–10.3]
	2.5/2.2	*DQA1*05:01-DQB1*02:01/DQA1*02:01-DQB1*02:02*	264 (9.1%)	337 (2.3%)	5.5 [4.5–6.7]
Moderate	2.5/X	*DQA1*05:01-DQB1*02:01/X*	795 (27.4%)	1784 (12.3%)	3.1 [2.7–3.6]
	2.2/7.5	*DQA1*02:01-DQB1*02:02/DQA1*05:05-DQB1*03:01*	149 (5.1%)	389 (2.7%)	2.7 [2.2–3.4]
	2.5/7.5	*DQA1*05:01-DQB1*02:01/DQA1*05:05-DQB1*03:01*	156 (5.4%)	456 (3.2%)	2.4 [1.9–3.0]
	2.5/8.1	*DQA1*05:01-DQB1*02:01/DQA1*03:01-DQB1*03:02*	166 (5.7%)	316 (2.2%)	3.7 [3.0–4.6]
	8.1/8.1	*DQA1*03:01-DQB1*03:02/DQA1*03:01-DQB1*03:02*	69 (2.4%)	173 (1.2%)	2.8 [2.1–3.8]
	2.2/8.1	*DQA1*02:01-DQB1*02:02/DQA1*03:01-DQB1*03:02*	93 (3.2%)	303 (2.1%)	2.2 [1.7–2.8]
Low	8.1/X	*DQA1*03:01-DQB1*03:02/X*	344 (11.9%)	1662 (11.5%)	1.5 [1.2–1.7]
	7.5/8.1	*DQA1*05:05-DQB1*03:01/DQA1*03:01-DQB1*03:02*	57 (2.0%)	407 (2.8%)	1.0 [0.7–1.3]
	2.2/X	*DQA1*02:01-DQB1*02:02/X*	249 (8.6%)	1547 (10.7%)	1.1 [0.9–1.4]
	7.5/X	*DQA1*05:05-DQB1*03:01/X*	313 (10.8%)	2202 (15.2%)	Reference
	2.2/2.2	*DQA1*02:01-DQB1*02:02/DQA1*02:01-DQB1*02:02*	22 (0.8%)	161 (1.1%)	1.0 [0.6–1.5]
	7.5/7.5	*DQA1*05:05-DQB1*03:01/DQA1*05:05-DQB1*03:01*	38 (1.3%)	290 (2.0%)	0.9 [0.6–1.3]
None	X/X	*X/X*	0 (0.0%)	4308 (29.7%)	

B

A

HLA-DQ risk

tTG-IgA

High n=148
Moderate n=471
Low n=278

IU/mL — 10 20 30 40 50 60

F= 20.7
p= 1.6E-09

DGP-IgG

High n=60
Moderate n=206
Low n=124

IU/mL — 5 10 15 20 25 30 35 40

F= 3.7
p= 2.6E-02

DGP-IgA

High n=73
Moderate n=247
Low n=167

IU/mL — 10 20 30 40 50 60 70

F= 14.9
p= 5.04E-07

Counts of CeD visits in EHR

High n=326
Moderate n=958
Low n=620

6 8 10 12 14

F= 17.7
p= 2.33E-08

B

Conditions	High-risk n=448	Moderate-risk n=1248	Low-risk n=1023
Positive family history	25.8%	21.3%	18.3%
Type 1 Diabetes	5.5%	4.7%	<2.0%
Stomach ulcers	5.5%	5.9%	7.6%
Anxiety reaction/panic disorder	31.2%	32.8%	38.2%
Vitamin D deficiency	47.2%	52.2%	52.4%
Allergies	54.3%	60.7%	64.5%
Acid reflux	37.0%	46.0%	49.6%
Reactions to Anesthesia	6.1%	5.0%	7.3%
Memory impairment	4.6%	4.8%	7.8%
Migraine	29.3%	36.4%	42.4%
Diarrhea	21.5%	23.2%	27.1%
Yeast infections	9.0%	11.0%	13.8%
Asthma	23.1%	29.2%	33.9%
Chronic sinus infections	15.9%	23.5%	26.8%
IBS	20.6%	27.7%	35.8%

vs high: moderate, low

FDR p-value
○ >0.05
○ ≤0.05
○ ≤0.01
○ ≤0.001

Odds ratio
2.0
1.5
0.6

A

All of Us v8 CDR
↓
WGS available
414830 participants
↓
3481 CeD
EHR-only: 1411
Self-Report-only: 1228
Both: 842
↓
Exclude individuals without CeD HLA risk
↓
2899 CeD
333sero+

413349 Non-CeD
↓ *Exclude individuals with selected Dx*
311572 Non-CeD
↓ *Propensity Match 5:1*
14495 Non-CeD

↓
Test prediction models
Clinical and genetic risks
↓
AoU-CeD model
↓
tTG-IgA level
↓
Validation

B

	CeD n(%) n=2899	Non-CeD n(%) n=14495
Gender		
Woman	2298 (79.3%)	11509 (79.4%)
Man	566 (19.5%)	2785 (19.2%)
Other	35 (1.2%)	201 (1.4%)
Race		
White	2371 (81.8%)	12119 (83.6%)
More than one population	155 (5.3%)	356 (2.5%)
Black	77 (2.7%)	432 (3.0%)
other	296 (10.1%)	1588 (11.0%)
Ethinicity		
Hispanic	272 (9.4%)	1326 (9.1%)
Education		
Advanced Degree	951 (32.8%)	3975 (27.6%)
College Graduate	829 (28.6%)	4076 (28.3%)
College One to Three	738 (25.5%)	3721 (25.8%)
Twelve Or GED	284 (9.8%)	1849 (12.8%)
Less than High School	67 (2.3%)	580 (4.0%)
No Answer	30 (1.0%)	213 (1.5)
Genetic Sex		
Female	2317 (80.1%)	11626 (80.3%)
Genetic Ancestry		
European	2495 (86.1%)	12460 (86.0%)
Admixed American	262 (9.0%)	1326 (9.1%)
African	108 (3.7%)	527 (3.6%)
Other	34(1.2%)	182(1.3%)

C

All of Us CeD cohort n=2899

EHR   EHR+ Self-Report   Self-Report

1133   771   995

Sero+ (n=333)

A

DQA1*05:01-DQB1*02:01-DRB1*03:01 haplotype

Linkage	European	Admixed American	African
Total *DQB1*02:01* (n)	4504	311	119
With *DRB1*03:01* (n, %)	4501 (99.9%)	310 (99.7%)	105 (88.2%)
With *DQA1*05:01* (n, %)	4476 (99.4%)	309 (99.4%)	103 (86.6%)
With both *DRB1*03:01* and *DQA1*05:01* (n, %)	4475 (99.4%)	309 (99.4%)	103 (86.6%)

B

HLA Allele	CeD (n=2,899)	Non-CeD (n=14,495)	OR(95%CI)
DQA1*05:01	30.67%	11.16%	3.52(3.30-3.77)
DQB1*02:01	30.25%	11.11%	3.47(3.24-3.71)
DRB1*03:01	30.25%	11.15%	3.46(3.23-3.70)
B*08:01	24.39%	10.02%	2.90(2.70-3.11)
C*07:01	27.73%	15.27%	2.13(1.99-2.27)
DPB1*01:01	11.28%	5.69%	2.11(1.92-2.32)
DQA1*03:01	17.28%	10.34%	1.81(1.68-1.96)
A*01:01	23.04%	14.56%	1.76(1.64-1.88)
DQB1*02:02	14.13%	10.29%	1.43(1.32-1.56)
DQB1*03:02	13.82%	10.56%	1.36(1.25-1.48)

C