7   Differential fMRI neural synchrony associated with migraine during naturalistic stimuli with
8   negative emotional valence
9
10

11   Keva Klamer [a]., Joshua Craig [a]., Christina Haines [a]., KiAnna Sullivan [a]., Peter Seres[b], & Chelsea
12   Ekstrand* [a].
13
14
15
16
17   [a]Ekstrand Neuroimaging Lab, Department of Neuroscience, University of Lethbridge, 4401

18   University Dr W, Lethbridge, AB, Canada, T1K 6T5

19   [b]Peter S. Allen MRI Research Centre, University of Alberta, 116 St & 85 Ave, Edmonton, AB,

20   Canada, T6G 2R3

22   *Correspondence to: Chelsea Ekstrand, chelsea.ekstrand@uleth.ca, 4401 University Dr W,

23   Lethbridge, AB, Canada, T1K 3M4

26   Key words: migraine, neural synchrony, fMRI, multisensory integration, emotional processing
27
List of abbreviations:
pSTG: posterior superior temporal gyrus
SPL: superior parietal lobule
IPS: intraparietal sulcus
MTG: middle temporal gyrus
ISC: intersubject correlation analysis
fMRI: functional magnetic resonance imaging
EPI: echo planar imaging
T1w: T1 weighted
ANTs: advanced normalization tools
LME: linear mixed effects
GLM: general linear model
LOC: lateral occipital cortex
IFG: inferior frontal gyrus
FDR: false discovery rate
SMA: supplementary motor area
44

Declarations

Ethics approval and consent to participate: Ethics approval was obtained from the University of

Alberta Research Ethics Board (Pro00120063). Written informed consent was obtained from all

participants prior to their participation in the study.

Consent for publication: Not applicable.

Availability of data and materials: The datasets used and/or analysed during the current study are

available from the corresponding author on reasonable request.

Competing interests: The authors declare that they have no competing interests.

Funding: This work was supported by a NSERC Discovery Grant and NSERC Discovery

Launch Supplement to C.E. (RGPIN-2021-03568 and DGECR-2021-00297) and an NSERC

PGS-D Scholarship to K.K. (Application No: PGS D – 589897 – 2024). The funding source had

no involvement in the study design; in the collection, analysis and interpretation of data; in the

writing of the report; and in the decision to submit the article for publication.

Author contributions: KK wrote and prepared the original draft of the manuscript,

conceptualized the study, completed formal analysis, and was involved in data acquisition. JC

completed review and editing of the manuscript, was involved in analysis of data and data

acquisition. CH was involved in methodology planning and data acquisition. KS was involved in

methodology planning and data acquisition. PS was involved in data acquisition. CE was

69   involved in conceptualization, data curation, funding acquisition, project administration, review

70   and editing, resources, and supervision. All authors read and approved the final manuscript.

72   Acknowledgements: Not applicable.

Abstract

Migraine is a common neurological disorder that impacts approximately 12% of the general population and is characterized by moderate to severe headaches, nausea, mood changes, and fatigue. It impacts lower-level visual and auditory processing, causing hypersensitivities that lead to heightened audiovisual multisensory integration. However, the impact of migraine on the processing of complex, audiovisual stimuli is still unclear. Additionally, migraine may induce hypersensitivities to emotional arousal and valence, though the relative significance of these factors remains unknown. The current study seeks to identify how migraine impacts synchronous neural processing of complex, audiovisual stimuli, and how this differs based on the emotional arousal and valence of the stimulus. To do so, we collected functional magnetic resonance imaging data from 22 migraineurs and 21 healthy controls during the passive viewing of three audiovisual films of differing emotional arousal and valence. We identified that, in response to a negative valence, high arousal emotional stimulus, the migraine group showed greater neural synchrony in regions associated with multisensory integration, including the bilateral posterior superior temporal gyrus (pSTG), superior parietal lobule (SPL), and left middle temporal gyrus (MTG). There were no significant differences in neural synchrony between the migraine and control groups in response to positive valence, high arousal and neutral valence, low arousal stimuli. These findings suggest that migraine involves hypersensitivity to audiovisual movies as a function of negative emotional valence, where negative/aversive emotional states may drive greater synchrony in multisensory integration. Overall, this research highlights distinct pathways through which emotion and arousal impact neural processing in migraine.

97   Differential fMRI neural synchrony associated with migraine during naturalistic stimuli with

98   negative emotional valence

Migraine is a common neurological disorder that impacts approximately 12% of the

general population and is more common in females than males (Szabó et al., 2019). It is

characterized by unilateral moderate to severe headaches, nausea, mood changes, and fatigue

(Silberstein, 1995). Migraineurs experience enhanced perception and altered cerebral processing

of sensory stimuli, commonly referred to as generalized hyperexcitable cortex. Hyperexcitable

cortex refers to a reduction of intracortical inhibition, which leads to the excitability of neurons

(Ciuffini et al., 2020) and is present even during the interictal period (i.e., in between migraine

attacks). As a result, migraineurs experience hypersensitivities to lower-level stimuli (e.g.,

visual/auditory stimuli) and higher-level stimuli (e.g., painful/emotional stimuli; Wang et al.,

2017). This suggests that migraineurs are producing a different neurological reaction than non-

migraineurs when perceiving the same visual and auditory stimuli.

There are several different subtypes of migraine, as highlighted by The International

Classification of Headache Disorders (3[rd] edition; ICHD-3; Headache Classification Committee

of the International Headache Society, 2013), which has been shown to display excellent

specificity (Göbel et al., 2020). The ICHD criteria includes several subtypes of migraine,

including migraine without aura, probable migraine without aura, migraine with aura, and

probable migraine with aura. Migraine without aura is defined as a headache lasting 4-72 hours

that may involve unilateral location, pulsating quality, moderate or severe pain intensity,

aggravation by physical activity, and the headache may be accompanied by nausea and/or

vomiting and photophobia and phonophobia. Probable migraine without aura is chararacterized

by migraine-like attacks missing one of the features required to fill all criteria for a type or

subtype of migraine, indicating that individuals with probable migraine without aura are on the

spectrum of migraine disorders. Probable migraine without aura has symptom and epidemiologic

profiles that overlap with migraine, and is therefore considered a prevalent form of migraine

(Patel et al., 2004). Migraine with aura is defined as recurrent attacks of unilateral fully-

reversible visual sensory or other symptoms that usually develop gradually and are followed by

headache and associated migraine symptoms (Headache Classification Committee of the

International Headache Society, 2013). Similarly, probable migraine with aura refers to aura-like

episodes that are missing one criterion required for a full diagnosis of migraine with aura, but

still share key clinical features (Headache Classification Committee of the International

Headache Society, 2013). Individuals with probable migraine with aura are considered part of the

broader spectrum of migraine with aura disorders. All of these conditions involve recurrent

headache attacks with overlapping symptomatology and underlying pathophysiological

mechanisms.

133   Migraine is associated with atypical unimodal processing of lower level stimuli during

the interictal period, specifically auditory and visual hypersensitivities. In the domain of visual

hypersensitivities, migraineurs have an increased sensitivity to normal light, flickering lights

(e.g., computer screens), bright light (e.g., sunlight), and patterns (e.g., plaid, achromatic

gratings; Friedman & De Ver Dye, 2009; Chong et al., 2016). Within the scope of auditory

hypersensitivities, approximately 75% of migraineurs report sensitivities to sound during the

interictal period (Vingen et al., 1998), and studies have shown that migraineurs are more

sensitive to sound compared to healthy controls (Ashkenazi et al., 2009). Further, migraineurs

exhibit lower light and auditory discomfort thresholds compared to controls (Meylakh &

Henderson, 2022). These findings highlight that migraineurs experience atypical processing of

both visual and auditory stimuli, leading to heightened sensitivity and lower discomfort

thresholds during the interictal period.

Atypical unimodal processing of lower level stimuli during the interictal period in

migraine has been increasingly linked to altered multisensory integration. Multisensory

integration refers to the production of a single perception of the environment through the co-

modulation of different modes of incoming stimuli (Schwedt, 2013). In other words,

multisensory integration allows for the formation of a coherent perception of everyday life by

combining inputs from different sensory modalities. Multisensory integration is a crucial

component of daily lived experience, and several pieces of evidence suggest that migraineurs

show altered multisensory integration compared to healthy controls (Schwedt, 2013). First, the

presence of one migraine symptom (e.g., visual hypersensitivity) positively correlates with the

intensity of other migraine symptoms (e.g., auditory hypersensitivity, pain severity; Kelman &

Tanis, 2006). Second, exposure to visual, auditory, and olfactory stimuli can trigger migraine

attacks (Harle et al., 2006). Third, exposure to one mode of sensory stimulation alters the

sensitivity to concurrent sensory stimuli of other modalities (Ashkenazi et al., 2010). Fourth,

studies have shown that audiovisual integration operates over a longer time window in

migraineurs compared to controls (O'Hare, 2017) and that controls are better than migraineurs at

motion discrimination, which involves integration of a stimulus over time (O'Hare, 2018). More

specifically, research has shown that audiovisual integration is significantly elevated, and

audiovisual suppression is weaker, in migraineurs compared to controls (Yang et al., 2014). This

may be the result of visual and auditory hypersensitivities, which reflect an increased excitability

state toward headache attacks (i.e., cortical hyperexcitability; Boran et al., 2021).

Several brain regions are implicated in the integration and coordination of visual and

auditory inputs, and their involvement may vary depending on the complexity of the stimulus

(Gao et al., 2023). These regions include the bilateral posterior superior temporal gyrus (pSTG),

bilateral superior parietal lobule/intraparietal sulcus (SPL/IPS), right lingual gyrus, right

thalamus, left middle temporal gyrus (MTG), and bilateral frontal gyri. The pSTG is considered

the hub of multisensory integration, where it integrates auditory and visual speech (Ozker et al.,

2017) and is activated during multisensory integration of simple and complex speech, and

complex nonspeech stimuli (Gao et al., 2023). The SPL/IPS is more generally involved in

multisensory integration as it processes both auditory and visual stimuli, whereby auditory inputs

arrive before visual inputs (Molholm et al., 2006). In addition, the SPL shows greater activation

in response to congruent stimuli across different senses (e.g., a visual cue that matches an

auditory cue) compared to congruent stimuli within a single sense (e.g., two visual cues) during

fMRI (Saito et al., 2005). The right frontal gyri are activated in response to simple stimuli, while

the left MTG and frontal gyri respond to complex non-speech stimuli. In contrast, complex

speech stimuli activate the right lingual gyrus, right thalamus, and left inferior occipital gyrus

(Gao et al., 2023). These findings demonstrate the complex interplay of various brain regions in

the integration of visual and auditory stimuli, where the pSTG consistently plays a central role,

regardless of stimulus complexity.

Emotional features of stimuli can also influence multisensory processing. Emotional

arousal, which refers to the intensity of an emotional response (Citron et al., 2014) has a

relationship with multisensory processing, where multisensory stimulation has been shown to

amplify the emotional arousal experienced when compared to unimodal stimulation (Schreuder

et al., 2016). Further, high levels of emotional arousal may generally alter perception, where they

are linked to increased attentional engagement (Zsidó, 2024). For example, research suggests

that multisensory integration may be enhanced by positive emotional valence stimuli

(Takeshima, 2020). In addition, brain regions involved in multisensory integration, such as the

pSTG, show altered activation levels in response to emotionally charged stimuli, suggesting that

emotional content can enhance or modify multisensory processing in these areas (Klasen et al.,

2014). Further, neural network models designed for multisensory integration also show improved

performance when emotional features are incorporated. However, negative emotional valence

stimuli may slow or impair audiovisual multisensory integration (Taffou et al., 2017), suggesting

a complex relationship between emotional valence, arousal, and multisensory integration.

197   In line with this, migraineurs show increased neural activation in the visual cortex and

cerebellum to emotional static picture stimuli (e.g., emotional faces), and this heightened neural

response becomes especially pronounced when the valence of the incoming stimulus is negative

(Szabó et al., 2019; Wilcox et al., 2016). Therefore, it has been theorized that migraineurs may

have a general hypersensitivity to aversive/negative stimuli, regardless of whether the stimulus is

low-level (e.g., visual, auditory) or high-level (e.g., emotions; Wilcox et al., 2016). However, this

theory contrasts another prominent theory that suggests that migraineurs have a heightened

sensitivity to emotionally arousing stimuli, regardless of if the emotional valence is positive or

negative (Steppacher et al., 2016). Several studies have also supported the theory of enhanced

responses to higher arousal stimuli in migraine, including an electroencephalography study

investigating the response to emotional pictures (Steppacher et al., 2016), and studies

investigating visual (Sezai et al., 2023) and auditory (Masson et al., 2020) hypersensitivities to

low-level arousing stimuli. This enhanced response to emotional stimuli could potentially

contribute to the psychiatric comorbidities often observed in migraineurs, such as anxiety and

depression (Breslau & Davis, 1993). However, it remains uncertain which of these theories (i.e.,

a general hypersensitivity to negative stimuli or a heightened sensitivity to emotionally arousing

stimuli) may underlie this phenomenon and its effect on multisensory integration in migraineurs.

While the aforementioned research has provided crucial insights into the relationship

between migraine, audiovisual multisensory integration, and emotional valence/arousal, the

precise mechanisms underlying these interactions remain largely unexplored. Naturalistic

stimuli, such as audiovisual films, podcasts, and virtual reality, show promise for exploring how

heightened sensitivity to emotional stimuli in migraine may impact multisensory integration.

Naturalistic stimuli are rich, multimodal, and dynamic, and better represent our daily lived

experience (Sonkusare et al., 2019). Consequently, naturalistic stimuli can evoke brain responses

that are highly reproducible within and across subjects (Hasson et al., 2004). Importantly,

naturalistic paradigms engage a broader set of brain regions and more diverse modes of network

interactions than task or resting-state paradigms (Zhang et al., 2021). Of particular note, Finn et

al. (2018) used naturalistic stimuli to characterize differences in brain synchrony between

participants who varied based on personality traits, in this case low and high trait paranoia. Their

results showed characteristic patterns of neural synchrony unique to the high paranoia group,

suggesting that underlying traits can influence the processing of naturalistic stimuli in the brain.

Further, naturalistic stimuli allow for the modulation of emotional valence and arousal, providing

a means to explore differences in neural responses related to the higher-level emotional content

of the stimulus. Therefore, using naturalistic stimuli will offer insight into neurological

processing that is more akin to real-life experience, allowing for a more comprehensive

understanding of the interplay between the emotional content of external stimuli, migraine, and

multisensory integration.

One common approach for analyzing neural responses to naturalistic stimuli is

intersubject correlation analysis (ISC). ISC examines the correlation of hemodynamic responses

to an audiovisual stimulus over time across multiple subjects. This approach helps identify neural

synchrony, which is the shared neural activity among subjects (Hasson et al., 2009). ISC is

particularly effective for naturalistic paradigms as it detects shared neural responses with

minimal assumptions, as opposed to GLM-based approaches (Finn et al., 2020). ISC isolates the

impact of the stimulus, making it easier to link observed neural activity directly to the stimulus.

Therefore, ISC is a highly suitable technique for examining how migraine impacts the

synchronous neurological responses to complex audiovisual stimuli.

The current study seeks to investigate differences in neural synchrony between

migraineurs and controls in response to complex audiovisual stimuli. Additionally, it aims to

explore how emotional arousal and valence influence these differences. In doing so, we will

evaluate two competing theories that propose migraineurs may be predominantly sensitive either

to emotional arousal or emotional valence. To do so, we acquired fMRI data from 22 individuals

with migraine and probable migraine and 21 healthy controls during the passive viewing of a

negative valence, high arousal film, a positive valence, high arousal film, and a neutral valence,

low arousal film. We hypothesized that if migraineurs are hypersensitive to high arousal stimuli,

that they will exhibit heightened neural synchrony in regions associated with multisensory

integration (right pSTG, bilateral SPL/IPS, left MTG, and left frontal cortices) during both high

arousal negative and positive film conditions. However, if migraineurs are hypersensitive to

negative valence stimuli, we hypothesize that they will show heightened neural synchrony in

these same multisensory integration regions only during the negative valence, high arousal film.

This research will determine how migraine impacts neural synchrony in response to complex

audiovisual stimuli and examine whether arousal or valence more profoundly influences

neurological processing.

<div align="center">**Methods**</div>

Participants and Stimuli

We recruited 46 participants to watch three short films during fMRI. We had to restart at

least one of the films for 2 participants who were subsequently removed from the study. One

participant dropped out during MRI scanning and was removed from the study, resulting in 43

total participants (28 females/15 males, mean age = 27.15, standard deviation = 8.32). All

participants had no history of neurological illnesses, no hearing impairments and normal or

corrected-to-normal vision. To assess migraine status, participants completed the Mickleborough

migraine questionnaire (Mickleborough et al., 2016). This questionnaire includes questions that

align with the ICHD-3 (Headache Classification Committee of the International Headache

Society, 2013) criteria for migraine without aura, migraine with aura, probable migraine without

aura, probable migraine with aura, and chronic migraine. 9 participants met the ICHD-3 criteria

for migraine without aura (8 females/1 male, mean age = 27.7), 4 participants met the ICHD-3

criteria for migraine with aura (4 females/0 males, mean age = 27.75), 9 participants met the

ICHD-3 criteria for probable migraine without aura (5 females/4 males, mean age = 30.33). As

probable migraine without aura has symptom and epidemiologic profiles that overlap with

migraine and is considered a prevalent form of migraine (Patel et al., 2004), we included

participants who met the ICHD-3 classification for migraine without aura, migraine with aura,

and probable migraine without aura in the migraine group, resulting in 22 participants in the

migraine group (17 females/5 males, mean age = 28.77). The migraine and control groups were

approximately matched on age ($t(41) = -1.287$, $p = 0.205$) and level of education ($t(41) = 1.147$,

$p = 0.258$). In ISC analysis, a sample size of 22 migraineurs is sufficient and can be robust due to

the focus on shared neural responses across subjects (Nastase et al., 2019) and with 20 subjects

per group, the ISC statistic converges close to a large sample ISC statistic with 130 subjects

(Pajula & Tohka, 2016). The migraine group averaged 42.14 (standard deviation = 47.66)

headaches a year, with the average headache lasting 16.52 hours (standard deviation = 23.26). 21

participants who did not meet the ICHD-3 criteria served as the control group (11 females/10

males, mean age = 25.52). The control group averaged 18.45 (standard deviation = 26.45)

headaches a year, with the average headache lasting 1.69 hours (standard deviation = 1.36). All

participants in the control group reported headaches of only mild-moderate pain intensity, and 18

participants in the migraine group reported moderate-severe pain intensity. No participants

reported having a headache within 48 hours prior to participation. This study was approved by

the University of Alberta Research Ethics Board (Pro00120063).

Participants watched three short films of differing emotional arousal and valence during

fMRI. The first short film, titled "Wicken" (Hashmic House Films, 2019; a short horror film),

was negative emotional valence and high arousal. The second short film, titled "One-Minute

Time Machine" (Avery, 2023; a short comedy film), was positive emotional valence and high

arousal. The third short film, titled "How to Create an Effective Training Video" (iSpring, 2020;

a short tutorial), was neutral emotional valence and low arousal. These films were

counterbalanced across participants to ensure any possible order effects were minimized and we

verbally checked in with participants before and after each film to confirm they wanted to

continue. Following each film, participants completed the Modified Differential Emotions Scale

(Fredrickson, 2001) to assess emotional responses, resulting in a 2-3 minute break from watching

movies in between films. Participants were presented with 16 emotional categories and asked to

make a verbal response on a scale of 1 to 7 of the intensity they felt that emotion over the time

course of the film. The verbal response was recorded as a keypress response from the

experimenter before moving to the next item. All films were edited using VEED.IO (Veed.io,

2024) to add subtitles, remove opening credits/title scenes and remove closing credits. One

additional scene was removed from the negative valence, high arousal film to avoid potential

motion artifacts because it contained a jump scare.

Data Acquisition

Neuroimaging data was acquired on a 3-Tesla Siemens Skyra Magnetom scanner at the

University 3T MRI Centre on the University of Lethbridge campus. After an initial localizer

scan, a high-resolution three-dimensional anatomical image was collected using a T1-weighted

magnetization-prepared rapid gradient echo sequence (192 continuous sagittal slices, slice

thickness = 1.0mm, matrix size = 256 x 256, field of view = 256mm, voxel size = 1.0mm x

1.0mm x 1.0mm, TR = 2300ms, TE = 2.26ms, flip angle = 8 degrees). Functional images were

acquired using a multiband T2*-sensitive gradient-recalled single-shot echo-planar imaging

(EPI) sequence (TR = 1500ms, TE = 34.0ms, voxel size = 2.4mm isotropic, flip angle = 90

degrees, matrix size = 85 x 85, field of view = 204mm, multiband factor = 4).

We acquired three different functional scans, one throughout the duration of each of the

films watched by participants (i.e., negative valence/high arousal, positive valence/high arousal,

neutral valence/low arousal). Stimuli were presented on a 40" MRI compatible LCD screen and

the audio portion of the stimulus was delivered through in-ear MRI compatible headphones. A

mirror was fixed to the head coil to allow participants to have a direct view of the screen. The

position of the LCD screen was marked on the ground with tape to ensure that the positioning of

the screen remained consistent across participants. Prior to the functional scans, we tested the

alignment of the mirror and LCD screen to ensure that participants had a direct view of the entire

screen.

Data Preprocessing

Initial preprocessing was performed using fMRIPrep 21.0.1 (Esteban et al., 2019). Full

preprocessing details can be found in Supplementary Materials A. Briefly, the T1-weighted

images were corrected for intensity non-uniformity, skull-stripped, and spatially normalized to

MNI152NLin2009cAsym space using advanced normalization tools (ANTs; Avants et al., 2008).

For the functional data, we used fMRIPrep 21.0.11 (Esteban et al., 2019) to apply slice-timing

correction, head-motion correction with six degrees of freedom. The BOLD data were co-

registered to T1-weighted images, and nuisance regressors (e.g., FD, DVARS, CompCor) were

computed. Frames exceeding 0.5mm FD or 1.5 standardized DVARS were flagged as motion

outliers. Finally, the BOLD time-series were spatially normalized to standard space using ANTs,

with final outputs including both volumetric and surface-based resampling. Additional

preprocessing was performed using python code that included spatial smoothing with a 6mm

FWHM Gaussian kernel, detection and removal of spikes, denoising through regression using

motion parameters, cerebrospinal fluid signal, and polynomial trends.

Behavioral Questionnaires

After passively viewing each film during the fMRI session, participants completed the

Modified Differential Emotions Scale (Fredrickson, 2001) to quantify their emotional

experience. This 16-question scale comprehensively measures positive, negative, and neutral

emotions, as well as subjective arousal, providing a detailed assessment of participants emotional

responses to each film. To ensure that each film elicited the intended arousal level, we compared

responses to the 'interested, concentrated, alert' item between all films using paired-samples *t*-

tests in SPSS (version 27; IBM Corp., 2020).

To ensure that the negative valence film elicited the intended emotional response, we

used paired-samples *t*-tests in SPSS (version 27; IBM Corp., 2020) to test differences within

participants on the average of their answers to the Modified Differential Emotions Scale items

'fearful, scared, afraid' and 'anxious, tense, nervous' between all films. To ensure that the

positive valence film elicited the intended emotional response, we used paired-samples *t*-tests in

SPSS (version 27; IBM Corp., 2020) to test differences within participants on the average of

their answers to the Modified Differential Emotions Scale items 'joyful, happy, amused' and

'warmhearted, gleeful, elated' between all films. We also assessed whether migraineurs and

controls differed in their emotional responses to the films. To do so, we conducted independent

samples *t*-tests comparing migraine and control groups on arousal, negative valence and positive

valence ratings for each film using the Modified Differential Emotions Scale items previously

mentioned.

Following each MRI scanning session, participants also completed the Beck Depression

Inventory-II (BDI; Beck et al., 2011), the Perceived Stress Scale (PSS; Cohen et al., 2014), and

the PANAS-GEN questionnaire (Watson et al., 1988). To assess group differences in depression,

perceived stress, and positive and negative affect, we conducted independent samples *t*-tests on

scores using SPSS (version 27; IBM Corp., 2020).

Intersubject Correlation Analyses

ISC was computed for all unique pairs of participants using 3dTcorrelate by AFNI (Cox

& Hyde, 1997) which produced 903 ($n*(n$-1)/2, where n = 43) unique ISC maps overall. Of

these, 231 maps originated from the migraine group ($n*(n$-1)/2, where n = 22), 210 originated

from the control group ($n*(n-1)/2$, where n = 21), and 462 ISC maps were obtained from

between-group pairings. ISC is a model-free approach used to analyze complex fMRI data

acquired in naturalistic, audiovisual stimulus environments (Kauppi, 2010). It allows us to

measure shared content across experimental conditions by filtering out subject-specific signals

and revealing voxels with a consistent, stimulus-evoked response time series across subjects

(Nastase et al., 2019a). It does this by calculating pairwise correlation coefficients between all

pairs of participants for each voxel throughout the brain. Ultimately, it provides us with a map of

neural synchrony, indicating regions with synchronous ISC across participants throughout the

entire time course of the film.

Linear Mixed Effects Modelling

381   After running ISC for all pairs of participants, we used the Linear Mixed Effects

Modelling (LME) implemented via the *3dISC* module in AFNI (Cox & Hyde, 1997; described by

Chen et al., 2020) to identify significant ISC associated with our contrasts of interest. LME is a

parametric method based on the general linear model (GLM) and is used to model the

relationship between the fMRI time series data and the experimental conditions at each voxel,

accounting for the complex covariance structure of ISC data. As LME accounts for variability in

the data due to random effects, it allows for more precise estimation of fixed effects (Koerner &

Zhang, 2017). The random effects specified in the LME model indicate random intercepts for

each subject. These random effects account for individual differences in how participants

respond to the same movie stimulus. LME was run separately for each condition (i.e., the

audiovisual films) resulting in the following contrasts: comparison of ISC between migraineurs

and controls during the negative valence, high arousal film; comparison of ISC between

migraineurs and controls during the positive valence, high arousal film; and comparison of ISC

between migraineurs and controls during the neutral valence, low arousal film. To address the

imbalance in sex distribution between the migraine and control groups, we included sex as a

quantitative explanatory variable in our LME model. We used sex as a main effect to account for

its potential influence on the results. Random effects for subjects were also included to account

for individual variability, allowing us to isolate the effects of interest more accurately. Significant

ISCs were defined using a voxelwise false discovery rate (FDR) of $q < 0.05$. Significant results

were transformed into surface space for visualization purposes only.

<div align="center">

Results

</div>

Behavioral Results

After passively viewing each film during the fMRI session, participants completed the Modified

Differential Emotions Scale (Fredrickson, 2001) to quantify their emotional experience. To

ensure that each film elicited the intended arousal level, we used paired-samples t-tests in SPSS

(version 27; IBM Corp., 2020) to test differences on the Modified Differential Emotions Scale

item 'interested, concentrated, alert' between all films. There were no significant differences

found between negative (mean = 4.907, standard deviation = 1.428) and positive (mean = 5.116,

standard deviation = 1.418; $t(42) = 0.802$, $p = 0.427$) films. There was a significant difference

found between negative and neutral (mean = 3.279, standard deviation = 1.723; $t(42) = 5.185$, p

< 0.001), and positive and neutral ($t(42) = 5.468$, $p < 0.001$) films, suggesting that the positive

and negative high arousal films elicited significantly higher arousal levels than the neutral, low

arousal film.

To ensure that the negative valence film elicited the intended negative emotional

response, we averaged the responses from the Modified Differential Emotions Scale items

'fearful, scared, afraid' and 'anxious, tense, nervous' and used paired-samples t-tests in SPSS to

test for differences between the three films. There was a significant difference found between

negative (mean = 4.778, standard deviation = 1.329) and positive (mean = 1.256, standard

deviation = 0.581; $t(42) = 16.795$, $p < 0.001$), and negative and neutral (mean = 1.081, standard

deviation = 1.329; $t(42) = 17.453$, $p < 0.001$) films. There was no significant difference found

between positive and neutral films ($t(42) = 1.801$, $p = 0.079$). To ensure that the positive valence

film elicited the intended emotional response, we used paired-samples t-tests in SPSS to test

differences within participants on the average of their answers to the Modified Differential

Emotions Scale items 'joyful, happy, amused' and 'warmhearted, gleeful, elated' between all

films. There was a significant difference found between positive (mean = 4.884, standard

deviation = 1.262) and negative (mean = 1.361, standard deviation = 0.601; $t(42) = 15.528$, $p <$

0.001), positive and neutral (mean = 1.942, standard deviation = 1.221; $t(42) = 10.658$, $p <$

0.001) and neutral and negative ($t(42) = -3.019$, $p = 0.004$), suggesting that the positive film

elicited significantly higher positive emotions.

Next, we assessed whether migraineurs and controls differed in their emotional responses

to the films. To do so, we conducted independent samples t-tests comparing migraine and control

groups on the arousal and valence Modified Differential Emotions Scale response previously

mentioned. Migraineurs did not differ from controls on arousal during the negative film ($t(41) = -$

0.737, $p = 0.466$), the positive film ($t(41) = -0.221$, $p = 0.826$) or the neutral film ($t(41) = 0.375$,

$p = 0.710$). See Figure 1 for the means and standard deviations of the arousal modified

differential scores across groups.

Figure 1

Arousal Modified Differential Scores

Note. Means and standard errors for the perceived arousal of each movie type in migraine and control groups. Negative- Migraine: migraine group's response to the negative valence, high arousal film. Negative- Control: control group's response to the negative valence, high arousal film. Positive- Migraine: migraine group's response to the positive valence, high arousal film. Positive- Control: control group's response to the positive valence, high arousal film. Neutral- Migraine: migraine group's response to the neutral valence, low arousal film. Neutral- Control: control group's response to the neutral valence, low arousal film.

For negative valence ratings, migraineurs differed from controls during the negative

valence film ($t(41) = -2.117$, $p = 0.040$), but did not differ during the positive ($t(41) = -0.193$, $p =$

0.848) or neutral films ($t(41) = 1.214$, $p = 0.232$). For positive valence ratings, migraineurs did

not differ from controls during the negative ($t(41) = -0.035$, $p = 0.972$), positive ($t(41) = -0.980$,

$p = 0.33$), or neutral films ($t(41) = -0.943$, $p = 0.351$). Thus, migraineurs and controls did not

differ in their emotional responses in terms of arousal or positive valence for any of the films, but

they did differ in their negative emotional response during the negative film. Refer to Figure 2

for the means and standard deviations of the valence modified differential scores across groups.

Figure 2

Valence Modified Differential Scores

Note. Means and standard errors for the perceived negative and positive valence of each movie type in migraine and control groups. Negative- Migraine: migraine group's response to the negative valence, high arousal film. Negative- Control: control group's response to the negative valence, high arousal film. Positive- Migraine: migraine group's response to the positive valence, high arousal film. Positive- Control: control group's response to the positive valence, high arousal film. Neutral- Migraine: migraine group's response to the neutral valence, low arousal film. Neutral- Control: control group's response to the neutral valence, low arousal film.

Following each MRI scanning session, participants completed the Beck Depression Inventory-II (BDI; Beck et al., 2011), the Perceived Stress Scale (PSS; Cohen et al., 2014), and the PANAS-GEN questionnaire (Watson et al., 1988). To assess group differences in depression, perceived stress, and positive and negative affect, we conducted independent samples t-tests on participants' scores using SPSS (version 27; IBM Corp., 2020). There were no significant differences in BDI scores ($t(41) = -1.899$, $p = 0.065$), PSS scores ($t(41) = -1.798$, $p = 0.079$), or

PANAS-GEN scores ($t(41) = -0.643$, $p = 0.521$).

Results from LME

Results from the LME examining group differences in processing the negative valence,

high arousal film are shown in Figure 3 and Tables S1 and S2 (Supplementary Materials B).

Migraineurs showed significantly greater ISCs than healthy controls in the bilateral pSTG,

bilateral SPL, and left MFG. Healthy controls showed significantly greater ISCs than

migraineurs in the right precuneus and right occipital pole. Results from the LME examining

group differences in processing the positive valence, high arousal film and neutral valence, low

arousal film found no significant differences.

484
Figure 3. Voxels showing significant ISC within participants with migraine and probable
migraine compared to healthy controls across the time course of the negative valence, high

arousal audiovisual stimulus. Warm colors refer to regions that are more synchronous in the
migraine than the control group. Cool colors refer to regions that are more synchronous in the
control than the migraine group. Results are displayed as a voxelwise false discovery (FDR)
threshold of $q < 0.05$.
491

Discussion

This study sought to examine differences in ISCs between migraineurs and healthy

controls in response to films of differing emotional valence and arousal (i.e., negative

valence/high arousal, positive valence/high arousal, neutral valence/low arousal). To do so, we

used ISC analysis and LME modelling to determine how neural synchrony varies between the

migraine and control groups across the three film conditions that varied based on emotional

valence and arousal. Results from this study show that migraineurs exhibit more similar

processing in regions associated with audiovisual multisensory integration, including the

bilateral pSTG and SPL, while processing negative valence, high arousal audiovisual stimuli,

which may reflect an increased sensitivity to negative emotional information in migraine.

Behaviourally, results from the Modified Differential Emotions Scale confirmed that the

high arousal films were found to be more arousing than the neutral film and did not differ in

arousal from each other. Further, each film used was found to elicit their intended emotional

responses in terms of negative, positive, and neutral valence. When comparing the emotional

response between groups, there were no significant group differences for arousal during any of

the films. There were significant group differences for negative valence emotions, with the

migraine group scoring higher during the negative film. These results validate the fMRI findings

which suggest that migraineurs may be hypersensitive to negative valence emotional stimuli.

There were no significant group differences for positive valence emotions during any of the

films. Additionally, there were no significant group differences for BDI, PSS, or PANAS-GEN

scores.

In response to the negative valence, high arousal film, migraineurs showed heightened

neural synchrony in the bilateral pSTG and SPL compared to controls, which are key

multisensory integration areas. The pSTG is considered a hub of multisensory integration,

combining visual and auditory inputs (Ozker et al., 2017), while the SPL is known to integrate

auditory and visual speech (Molholm et al., 2006). Additionally, migraineurs showed heightened

neural synchrony in the left MTG, a region implicated in multisensory integration and, more

specifically, in audiovisual integration during complex non-speech stimuli (Gao et al., 2023),

which is consistent with the type of stimuli used in this study. It is well known that migraineurs

experience visual and auditory sensitivities, resulting in altered multisensory integration (Yang et

al., 2014). As migraineurs may experience an enhanced response to negative valence stimuli

(Steppacher et al., 2016), it is possible that multisensory integration is enhanced during

negative/aversive emotional states. Importantly, no differences in neural synchrony were

observed in the bilateral pSTG or SPL during the neutral valence, low arousal or positive

valence, high arousal films. Together, these findings support the theory that migraineurs have a

heightened sensitivity to negative valence emotional stimuli (Wilcox et al., 2016) during the

interictal period. Therefore, heightened synchrony in the bilateral pSTG, SPL and left MTG

during only the negative valence, high arousal film may reflect an increased effort to integrate

the increased sensory input from both auditory and visual modalities. This suggests that in

migraineurs, amplified sensory processing associated with hyperexcitable cortex during negative

emotional states may also increase the demand on multisensory integration processes, however,

more research is needed to determine the exact mechanisms underlying this heightened

synchrony.

In response to the negative valence, high arousal film, controls had heightened neural

synchrony in the left planum temporale and right occipital pole compared to migraineurs. The

left planum temporale is involved in early auditory processing (Binder et al., 1996), where it

plays a pivotal role in processing auditory and visual linguistic stimuli (Wani, 2024). The right

occipital lobe is involved in early visual processing and is considered the primary distributor of

visual information that reaches other cortical areas (Tong, 2003). Given the heightened

sensitivity to auditory and visual stimuli in migraine, the increased synchrony observed in these

regions in the control group during negative valence audiovisual processing may reflect more

stable and uniform early-stage processing, however more research is needed to fully understand

the implications of heightened neural synchrony in these regions.

This study sought to examine differences in how migraineurs and controls process

naturalistic audiovisual stimuli of differing emotional arousal and valence, however, there are

several limitations to be discussed. The films used in this study were chosen to elicit specific

valence and arousal responses but were not matched in terms of visual and auditory

characteristics. For example, the negative valence, high arousal film had dark lighting, a

suspenseful musical score, and less dialogue compared to the positive valence film. These

differences may have influenced neural synchrony independently of emotional content, making it

difficult to isolate the effects of valence and arousal. Therefore, we did not directly compare

across films, as we cannot control for these differences between stimuli. Cross-film comparisons

would be challenging to interpret accurately because differences in neural synchrony may result

from varying visual and auditory elements rather than emotional valence or arousal alone. Future

research should use stimuli where the visual and auditory components are more closely matched,

such as using films with similar lighting, color grading, sound design, and dialogue length, which

would ensure that only emotional valence differs between conditions and enable direct

comparisons across films.

Additionally, based on our sample size, we were unable to examine differences in neural

synchrony associated with different subtypes of migraine (e.g., migraine with and without aura).

Future research should extend this paradigm to larger samples of each subtype to understand how

different migraine pathologies impact the processing of naturalistic stimuli. We also did not

assess anxiety levels. Future studies should collect data on trait anxiety to account for its

potential influence on ISC results. We also did not collect information about which stage in the

migraine cycle each participant was in at the time of the study or medication use, which may be

relevant as different phases are associated with sensory processing changes, particularly in the

preictal phase (Mykland et al., 2019). Future studies should account for these factors.

In conclusion, this study demonstrates that migraineurs exhibit greater neutral synchrony

when processing negative emotional stimuli, particularly in the bilateral pSTG and SPL,

suggesting migraineurs have altered multisensory integration in response to negative emotional

content. In other words, the results suggest that negative emotional valence, but not high

emotional arousal (regardless of valence), alter processing in migraine as migraineurs exhibited

heightened neural synchrony in regions linked to audiovisual multisensory integration when

responding to negative valence audiovisual stimuli. Overall, this study highlights how migraine

uniquely shapes the brain's response to negative emotional valence, revealing distinct but

interconnected differences in neural processing. These findings underscore the broader

implications of migraine-related sensory hypersensitivity, particularly in the context of

emotionally salient stimuli, and provide valuable insight into how heightened valence sensitivity

may drive differences in multisensory integration and visual information processing.

References

Ashkenazi, A., Mushtaq, A., Yang, I., & Oshinsky, M. (2009). Ictal and Interictal Phonophobia in

Migraine—A Quantitative Controlled Study. *Cephalalgia*, *29*(10), 1042–1048.

https://doi.org/10.1111/j.1468-2982.2008.01834.x

Ashkenazi, A., Yang, I., Mushtaq, A., & Oshinsky, M. L. (2010). Is phonophobia associated with

cutaneous allodynia in migraine? *Journal of Neurology, Neurosurgery & Psychiatry*,

81(11), 1256–1260. https://doi.org/10.1136/jnnp.2009.198481

Beck, A. T., Steer, R. A., & Brown, G. (2011). *Beck Depression Inventory–II* [Dataset].

https://doi.org/10.1037/t00742-000

Binder, J. R., Frost, J. A., Hammeke, T. A., Rao, S. M., & Cox, R. W. (1996). Function of the left

planum temporale in auditory and linguistic processing. *Brain: A Journal of Neurology*,

119 (Pt 4), 1239–1247. https://doi.org/10.1093/brain/119.4.1239

Boran, H. E., Bolay, H., Gantenbein, H. A. R., & Pohl, H. (2021). Sensory Processing and

Sensorimotor Integration in Migraine. In G. Coppola & W.-T. Chen (Eds.),

Neurophysiology of the Migraine Brain (pp. 113–131). Springer International Publishing.

https://doi.org/10.1007/978-3-030-56538-1_9

Boulloche, N., Denuelle, M., Payoux, P., Fabre, N., Trotter, Y., & Géraud, G. (2010).

Photophobia in migraine: An interictal PET study of cortical hyperexcitability and its

modulation by pain. *Journal of Neurology, Neurosurgery & Psychiatry*, *81*(9), 978–984.

https://doi.org/10.1136/jnnp.2009.190223

Breslau, N., & Davis, G. C. (1993). Migraine, physical health and psychiatric disorder: A

prospective epidemiologic study in young adults. *Journal of Psychiatric Research*, *27*(2),

211–221. https://doi.org/10.1016/0022-3956(93)90009-Q

Chen, G., Taylor, P. A., Qu, X., Molfese, P. J., Bandettini, P. A., Cox, R. W., & Finn, E. S. (2020).

Untangling the relatedness among correlations, part III: Inter-subject correlation analysis

through Bayesian multilevel modeling for naturalistic scanning. *NeuroImage*, *216*,

116474. https://doi.org/10.1016/j.neuroimage.2019.116474

Citron, F. M. M., Gray, M. A., Critchley, H. D., Weekes, B. S., & Ferstl, E. C. (2014). Emotional

valence and arousal affect reading in an interactive way: Neuroimaging evidence for an

approach-withdrawal framework. *Neuropsychologia*, *56*(100), 79–89.

https://doi.org/10.1016/j.neuropsychologia.2014.01.002

Ciuffini, R., Stratta, P., Pacitti, F., Bologna, M., & Marrelli, A. (2020). Emotional dysregulation,

affective status and personality traits in a sample of migraine patients. *Journal of*

Psychopathology, 141–146. https://doi.org/10.36148/2284-0249-384

Cohen, S., Kamarck, T., & Mermelstein, R. (2014). *Perceived Stress Scale* [Dataset].

https://doi.org/10.1037/t02889-000

Cox, R. W. (1996). AFNI: Software for Analysis and Visualization of Functional Magnetic

Resonance Neuroimages. *Computers and Biomedical Research*, *29*(3), 162–173.

https://doi.org/10.1006/cbmr.1996.0014

Cox, R. W., & Hyde, J. S. (1997). Software tools for analysis and visualization of fMRI data.

NMR in Biomedicine, *10*(4–5), 171–178. https://doi.org/10.1002/(SICI)1099-

1492(199706/08)10:4/5<171::AID-NBM453>3.0.CO;2-L

Dell'Acqua, C., Moretta, T., Dal Bò, E., Messerotti Benvenuti, S., & Palomba, D. (2022).

Emotional processing prospectively modulates the impact of anxiety on COVID-19

pandemic-related post-traumatic stress symptoms: An ERP study. *Journal of Affective*

Disorders, *303*, 245–254. https://doi.org/10.1016/j.jad.2022.02.027

Esteban, O., Markiewicz, C. J., Blair, R. W., Moodie, C. A., Isik, A. I., Erramuzpe, A., Kent, J. D., Goncalves, M., DuPre, E., Snyder, M., Oya, H., Ghosh, S. S., Wright, J., Durnez, J., Poldrack, R. A., & Gorgolewski, K. J. (2019). fMRIPrep: A robust preprocessing pipeline for functional MRI. *Nature Methods*, *16*(1), 111–116. https://doi.org/10.1038/s41592-018-0235-4

Finn, E. S., Glerean, E., Khojandi, A. Y., Nielson, D., Molfese, P. J., Handwerker, D. A., & Bandettini, P. A. (2020). Idiosynchrony: From shared responses to individual differences during naturalistic neuroimaging. *NeuroImage*, *215*, 116828. https://doi.org/10.1016/j.neuroimage.2020.116828

Fredrickson, B. L. (2001). The role of positive emotions in positive psychology: The broaden-and-build theory of positive emotions. *American Psychologist*, *56*(3), 218–226. https://doi.org/10.1037/0003-066X.56.3.218

Friedman, D. I., & De Ver Dye, T. (2009). Migraine and the Environment. *Headache: The Journal of Head and Face Pain*, *49*(6), 941–952. https://doi.org/10.1111/j.1526-4610.2009.01443.x

Gao, C., Green, J. J., Yang, X., Oh, S., Kim, J., & Shinkareva, S. V. (2023). Audiovisual integration in the human brain: A coordinate-based meta-analysis. *Cerebral Cortex*, *33*(9), 5574–5584. https://doi.org/10.1093/cercor/bhac443

Göbel, C. H., Karstedt, S. C., Münte, T. F., Göbel, H., Wolfrum, S., Lebedeva, E. R., Olesen, J., & Royl, G. (2020). ICHD-3 is significantly more specific than ICHD-3 beta for diagnosis of migraine with aura and with typical aura. *The Journal of Headache and Pain*, *21*(1), 2. https://doi.org/10.1186/s10194-019-1072-2

Harle, D. E., Shepherd, A. J., & Evans, B. J. W. (2006). Visual stimuli are common triggers of

migraine and are associated with pattern glare. *Headache*, *46*(9), 1431–1440.

https://doi.org/10.1111/j.1526-4610.2006.00585.x

Hasson, U., Avidan, G., Gelbard, H., Vallines, I., Harel, M., Minshew, N., & Behrmann, M.

(2009). Shared and idiosyncratic cortical activation patterns in autism revealed under

continuous real-life viewing conditions. *Autism Research*, *2*(4), 220–231.

https://doi.org/10.1002/aur.89

Hasson, U., Nir, Y., Levy, I., Fuhrmann, G., & Malach, R. (2004). Intersubject Synchronization

of Cortical Activity During Natural Vision. *Science*, *303*(5664), 1634–1640.

https://doi.org/10.1126/science.1089506

Headache Classification Committee of the International Headache Society (IHS). (2013). The

International Classification of Headache Disorders, 3rd edition (beta version).

Cephalalgia, *33*(9), 629–808. https://doi.org/10.1177/0333102413485658

Kauppi. (2010). Inter-subject correlation of brain hemodynamic responses during watching a

movie: Localization in space and frequency. *Frontiers in Neuroinformatics*.

https://doi.org/10.3389/fninf.2010.00005

Kelman, L., & Tanis, D. (2006). The Relationship between Migraine Pain and Other Associated

Symptoms. *Cephalalgia*, *26*(5), 548–553. https://doi.org/10.1111/j.1468-

2982.2006.01075.x

Klasen, M., Kreifelts, B., Chen, Y.-H., Seubert, J., & Mathiak, K. (2014). Neural processing of

emotion in multimodal settings. *Frontiers in Human Neuroscience*, *8*.

https://doi.org/10.3389/fnhum.2014.00822

Koerner, T., & Zhang, Y. (2017). Application of Linear Mixed-Effects Models in Human

Neuroscience Research: A Comparison with Pearson Correlation in Two Auditory

Electrophysiology Studies. *Brain Sciences*, *7*(12), 26.

https://doi.org/10.3390/brainsci7030026

Kumar, R., Asif, S., Bali, A., Dang, A. K., & Gonzalez, D. A. (n.d.). The Development and

Impact of Anxiety With Migraines: A Narrative Review. *Cureus*, *14*(6), e26419.

https://doi.org/10.7759/cureus.26419

Letts, E., Basharat, A., & Barnett-Cowan, M. (2022). *Evaluating the Effect of Semantic*

Congruency and Valence on Multisensory Integration. https://doi.org/10.1163/22134808-

bja10073

Masson, R., Lévêque, Y., Demarquay, G., ElShafei, H., Fornoni, L., Lecaignard, F., Morlet, D.,

Bidet-Caulet, A., & Caclin, A. (2020). Auditory attention alterations in migraine: A

behavioral and MEG/EEG study. *Clinical Neurophysiology*, *131*(8), 1933–1946.

https://doi.org/10.1016/j.clinph.2020.05.024

Meylakh, N., & Henderson, L. A. (2022). Exploring alterations in sensory pathways in migraine.

The Journal of Headache and Pain, *23*(1), 5. https://doi.org/10.1186/s10194-021-01371-

y

Mickleborough, M. J. S., Ekstrand, C., Gould, L., Lorentz, E. J., Ellchuk, T., Babyn, P., &

Borowsky, R. (2016). Attentional Network Differences Between Migraineurs and Non-

migraine Controls: fMRI Evidence. *Brain Topography*, *29*(3), 419–428.

https://doi.org/10.1007/s10548-015-0459-x

Molholm, S., Sehatpour, P., Mehta, A., Shpaner, M., Gomez-Ramirez, M., Ortigue, S., Dyke, J.,

Schwartz, T., & Foxe, J. J. (2006). Audio-visual multisensory integration in superior

parietal lobule revealed by human intracranial recordings. *Journal of Neurophysiology*.

https://www.semanticscholar.org/paper/Audio-visual-multisensory-integration-in-

superior-Molholm-Sehatpour/955a30e7bbbde0e36cae9c3e6546517f063ce59b

Mykland, M. S., Bjørk, M. H., Stjern, M., Omland, P. M., Uglem, M., & Sand, T. (2019).

Fluctuations of sensorimotor processing in migraine: A controlled longitudinal study of

beta event related desynchronization. *The Journal of Headache and Pain*, *20*(1), 77.

https://doi.org/10.1186/s10194-019-1026-8

Nastase, S. A., Gazzola, V., Hasson, U., & Keysers, C. (2019a). Measuring shared responses

across subjects using intersubject correlation. *Social Cognitive and Affective*

Neuroscience, *14*(6), 667–685. https://doi.org/10.1093/scan/nsz037

Nastase, S. A., Gazzola, V., Hasson, U., & Keysers, C. (2019b). Measuring shared responses

across subjects using intersubject correlation. *Social Cognitive and Affective*

Neuroscience, nsz037. https://doi.org/10.1093/scan/nsz037

O'Hare, L. (2017). *Multisensory Integration in Migraine: Recent Developments*.

https://doi.org/10.1163/22134808-00002570

O'Hare, L. (2018). Temporal Integration of Motion Streaks in Migraine. *Vision*, *2*(3), Article 3.

https://doi.org/10.3390/vision2030027

Ozker, M., Schepers, I. M., Magnotti, J., Yoshor, D., & Beauchamp, M. (2017). A Double

Dissociation between Anterior and Posterior Superior Temporal Gyrus for Processing

Audiovisual Speech Demonstrated by Electrocorticography. *Journal of Cognitive*

Neuroscience. https://www.semanticscholar.org/paper/A-Double-Dissociation-between-

Anterior-and-Superior-Ozker-Schepers/2f132c6a403792c50aa4d821d007e992d1faf52a

Pajula, J., & Tohka, J. (2016). How Many Is Enough? Effect of Sample Size in Inter-Subject

Correlation Analysis of fMRI. *Computational Intelligence and Neuroscience*, *2016*, 1–10.

https://doi.org/10.1155/2016/2094601

Patel, N. V., Bigal, M. E., Kolodner, K. B., Leotta, C., Lafata, J. E., & Lipton, R. B. (2004).

Prevalence and impact of migraine and probable migraine in a health plan. *Neurology*,

63(8), 1432–1438. https://doi.org/10.1212/01.WNL.0000142044.22226.54

Saito, D. N., Yoshimura, K., Kochiyama, T., Okada, T., Honda, M., & Sadato, N. (2005). Cross-

modal Binding and Activated Attentional Networks during Audio-visual Speech

Integration: A Functional MRI Study. *Cerebral Cortex*, *15*(11), 1750–1760.

https://doi.org/10.1093/cercor/bhi052

Schreuder, E., van Erp, J., Toet, A., & Kallen, V. L. (2016). Emotional Responses to

Multisensory Environmental Stimuli: A Conceptual Framework and Literature Review.

Sage Open, *6*(1), 2158244016630591. https://doi.org/10.1177/2158244016630591

Schwedt, T. J. (2013). Multisensory Integration in Migraine. *Current Opinion in Neurology*,

26(3), 248–253. https://doi.org/10.1097/WCO.0b013e328360edb1

Sezai, T., Murphy, M. J., Riddell, N., Nguyen, V., & Crewther, S. G. (2023). Visual Processing

During the Interictal Period Between Migraines: A Meta-Analysis. *Neuropsychology*

Review, *33*(4), 765–782. https://doi.org/10.1007/s11065-022-09562-3

Silberstein, S. D. (1995). Migraine Symptoms: Results of a Survey of Self☐Reported

Migraineurs. *Headache: The Journal of Head and Face Pain*, *35*(7), 387–396.

https://doi.org/10.1111/j.1526-4610.1995.hed3507387.x

Sonkusare, S., Breakspear, M., & Guo, C. (2019). Naturalistic Stimuli in Neuroscience:

Critically Acclaimed. *Trends in Cognitive Sciences*, *23*(8), 699–714.

https://doi.org/10.1016/j.tics.2019.05.004

Steppacher, I., Schindler, S., & Kissler, J. (2016). Higher, faster, worse? An event-related

potentials study of affective picture processing in migraine. *Cephalalgia*, *36*(3), 249–257.

https://doi.org/10.1177/0333102415587705

Szabó, E., Galambos, A., Kocsel, N., Édes, A. E., Pap, D., Zsombók, T., Kozák, L. R., Bagdy, G.,

Kökönyei, G., & Juhász, G. (2019). Association between migraine frequency and neural

response to emotional faces: An fMRI study. *NeuroImage: Clinical*, *22*, 101790.

https://doi.org/10.1016/j.nicl.2019.101790

Taffou, M., Ondřej, J., O'Sullivan, C., Warusfel, O., Dubal, S., & Viaud-Delmon, I. (2017).

Multisensory aversive stimuli differentially modulate negative feelings in near and far

space. *Psychological Research*, *81*(4), 764–776. https://doi.org/10.1007/s00426-016-

0774-1

Takeshima, Y. (2020). Emotional information affects fission illusion induced by audio-visual

interactions. *Scientific Reports*, *10*(1), 998. https://doi.org/10.1038/s41598-020-57719-y

Tong, F. (2003). Primary visual cortex and visual awareness. *Nature Reviews Neuroscience*, *4*(3),

219–229. https://doi.org/10.1038/nrn1055

Vingen, J. V., Pareja, J., Støren, O., White, L., & Stovner, L. (1998). Phonophobia in Migraine.

Cephalalgia, *18*(5), 243–249. https://doi.org/10.1111/j.1468-2982.1998.1805243.x

Wang, M., Su, J., Zhang, J., Zhao, Y., Yao, Q., Zhang, Q., Zhang, H., Wang, S., Li, G.-F., Liu, J.-

R., & Du, X. (2017). Visual cortex and cerebellum hyperactivation during negative

emotion picture stimuli in migraine patients. *Scientific Reports*, *7*(1), 41919.

https://doi.org/10.1038/srep41919

Wani, P. D. (n.d.). From Sound to Meaning: Navigating Wernicke's Area in Language

Processing. *Cureus*, *16*(9), e69833. https://doi.org/10.7759/cureus.69833

Watson, D., Clark, L. A., & Tellegen, A. (1988). Development and validation of brief measures

of positive and negative affect: The PANAS scales. *Journal of Personality and Social*

Psychology, *54*(6), 1063–1070. https://doi.org/10.1037//0022-3514.54.6.1063

Wilcox, S. L., Veggeberg, R., Lemme, J., Hodkinson, D. J., Scrivani, S., Burstein, R., Becerra,

L., & Borsook, D. (2016). Increased Functional Activation of Limbic Brain Regions

during Negative Emotional Processing in Migraine. *Frontiers in Human Neuroscience*,

10. https://doi.org/10.3389/fnhum.2016.00366

Yang, W., Chu, B., Yang, J., Yu, Y., Wu, J., & Yu, S. (2014). Elevated audiovisual temporal

interaction in patients with migraine without aura. *The Journal of Headache and Pain*,

15(1), 44. https://doi.org/10.1186/1129-2377-15-44

Zhang, Y., Kim, J.-H., Brang, D., & Liu, Z. (2021). Naturalistic stimuli: A paradigm for

multiscale functional characterization of the human brain. *Current Opinion in Biomedical*

Engineering, *19*, 100298. https://doi.org/10.1016/j.cobme.2021.100298

Zsidó, A. N. (2024). The effect of emotional arousal on visual attentional performance: A

systematic review. *Psychological Research*, *88*(1), 1–24. https://doi.org/10.1007/s00426-

023-01852-6

779