A Nutrient Ratio–Based, Web-Enabled Food Quality Score Performs Favorably Compared with Leading Nutrient Profiling Systems for Weight and Blood Pressure

Christopher J. Damman[1,2], Cara L. Frankenfeld[3]
1 University of Washington, Department of Medicine, Division of Gastroenterology, Seattle, WA;
2 GutBites MD, Seattle, WA; 3 MaineHealth Institute for Research, Scarborough, ME

Contact Information:
Cara L. Frankenfeld
Center for Interdisciplinary Population & Health Research
MaineHealth Institute for Research
81 Research Drive
Scarborough, ME 04074
Cara.Frankenfeld@mainehealth.org

Corresponding Author:
Christopher J. Damman
Division of Gastroenterology
UW Medicine
1959 NE Pacific St
Box 356424
Seattle, WA 98195-6524
cdamman@medicine.washington.edu

Keywords:
Nutrient ratios; Obesity; Hypertension; Nutrient profiling systems; Food quality score

Cara Frankenfeld is an Associate Editor for the Journal of Nutrition and played no role in the Journal's evaluation of the manuscript.

Abstract

Background: Rising prevalence of chronic cardiometabolic conditions may be partly driven by shifts in dietary patterns. Nutrient profiling systems (NPSs) aim to guide healthier food choices through labeling and consumer-facing technologies but vary in accessibility and how well they distinguish food healthfulness.

Objectives: The primary objective was to compare a ratio-based, web-enabled NPS, Nutrient Consume Score (NCS), and its underlying nutrient ratios, with four leading NPSs—Nutri-Score (NS), Health Star Rating (HS), NOVA Classification (NC), and Food Compass 2.0 (FC)—by examining associations with obesity and blood pressure. Secondary objectives included assessing associations with cardiometabolic biomarkers and identifying food categories contributing most to each score.

Methods: NHANES 2015–2016 data for adults aged ≥20 years were analyzed. Dietary intake was assessed via 24-hour recalls, and NPS scores were calculated. Multivariable regression models adjusted for sociodemographic and health factors examined associations with obesity, blood pressure, and cardiometabolic biomarkers. Compositional analysis evaluated food categories driving the scores.

Results: The ratio-based NCS performed favorably compared with others. All NPSs significantly associated with healthier weight, while blood pressure associations were more variable. Including alcohol and adjusting for bioactives strengthened the ratio-based NCS's association with weight outcomes. Across NPSs, food categories contributing to high and low scores were broadly consistent, generally reflecting minimally processed foods for high scores and more processed foods for low scores.

27 **Conclusions:** The Nutrient Consume Score (NCS) showed favorable associations with weight

28 and blood pressure relative to other NPSs, highlighting its potential for improving diet-related

29 risk factors for chronic conditions. As a web-enabled NPS, NCS may be well suited for

30 consumer-facing technologies, such as smartphones, and could facilitate interventional trials to

31 assess the causal impact of diet quality on cardiometabolic outcomes.

Introduction

The rising incidence and prevalence of obesity (1), metabolic syndrome (2), and other

noncommunicable disease (3) in the United States and globally (4) has been linked to shifts in

the types of foods available to consumers including increased ultra-processed (5) and

hyperpalatable foods (6) with resulting changes in dietary habits (7). Complex, confusing (8),

and conflicting (9) nutrition guidance make it difficult for individuals to make informed, healthy

choices on food purchases, preparation, and consumption (10).

Nutrient profiling systems (NPSs) aim to simplify nutrition complexity into easier-to-follow ratings

that assess the health quality of foods and beverages. These systems can offer real-time

guidance through package labeling (11), online websites (12), and smartphone technologies

(13,14), helping individuals make healthier food choices. Additionally, they can support public

health recommendations (15) and guide food companies in creating healthier products (16,17).

While there are over 100 reported NPSs (15,18), a small subset has been rigorously evaluated

for their association with health outcomes. The most studied NPSs with strongest links to

improved dietary choices and better health are Nutri-Score (NS) (19), Health Star (HS) (20),

NOVA Classification (NC) (21), and Food Compass (FC) (22,23).

Nutri-Score (24), popular in several European countries, and Health Star (25), widely used in

Australia and New Zealand, rank foods based on nutrient profiles and food categories. The

NOVA classification categorizes foods by their degree of processing, popularizing the term ultra-

processed food (26). Food Compass evaluates foods across nine health domains, including

categories like nutrient ratios, fiber, and phytochemicals (23,27). These systems are variably

effective in assessing the relative quality of different ultra-processed foods (28) and not

available on front of product packaging in many countries including the United States(23).

Despite the availability of multiple NPSs, a gap remains for a rigorously studied, web-enabled

algorithm that is independent of country-specific labeling practices and that can scale, based on

relative rather than absolute nutrient measures. The Nutrient Consume Score (NCS) (12) is a

recently developed web-based algorithm that focuses on nutrient ratios – carbohydrate-to fiber

(29), saturated fat-to-unsaturated fat (30), sodium-to-potassium (31), and calorie-to-weight (32)

– as proxies for the degree of food processing and previously identified as strong predictors of

food quality (33). It also incorporates alcohol and adjusts for food categories high in bioactive

components that support a healthy gut microbiome, including polyphenols, bioactive fats,

fermentable fibers, and fermentation products (34).

The primary aim of this study was to compare NCS, a nutrient ratio-based, web-enabled food

quality score with other leading NPSs in relation to weight and blood pressure, key prognostic

factors in cardiometabolic disease and mortality.

Methods

Data Sources and Analytic Population

The National Health and Nutrition Examination Survey (NHANES) is a repeated, cross-sectional

analysis that has been conducted continuously since 1999 and data are released in two-year

cycles, with exceptions for recent cycles due to COVID-19 disruptions in survey data

collection(35). The survey includes interview and physical measurement components and is

conducted in a nationally representative sample of approximately 5000 persons each year from

persons located in 15 counties that are randomly selected each year.

For the primary analysis to compare NCS and other nutrient profiling scores, data from the

continuous survey data releases 2015-2016 was used for this analysis because complete data

was available for all evaluated NPS scores for this NHANES data cycle (36). In the 2015-2016

data cycle, 9971 individuals participated. The following exclusions were applied in this order to

obtain the analytic subsample used (n excluded): age <20 years (n=4252), missing education

(n=5), missing poverty-to-income ratio (n=0), missing day 1 dietary intake data (n=699),

physiologically unsustainable low dietary intake of <500 kcal (n=42), physiologically

unsustainable high dietary intake >5000 kcal (n=57), missing smoking information (n=20),

missing waist circumference (n=203), missing body mass index (n=9), missing blood pressure

(n=51), or missing physical activity (n=35). NHANES does not have missing data for age,

gender, or race/ethnicity. The analytic sample for primary analyses used included 4598 adult

individuals. A secondary analysis was undertaken to evaluate associations of the NCS score

with other cardiometabolic biomarkers in NHANES cycles 2005-2018. Additional exclusions to

the 4598 individuals in the analytic sample were missing LDL, HDL, or total cholesterol, plasma

glucose, or blood triglycerides (n=2647). The analytic sample for secondary analyses of

cardiometabolic biomarkers included 1951 adult individuals.

Dietary Intake Assessment

Dietary intake is assessed for NHANES using two multiple pass 24-hour recalls. The first 24-

hour recall is conducted in person during the visit to the Mobile Examination Center (MEC). The

second 24-hour recall is conducted via phone 3 to 10 days after the MEC visit. All NHANES

participants are eligible for the 24-hour recall interviews. The interviews are conducted using a

midnight-to-midnight time frame for the 24-hour period prior to the interview. The USDA Food

Survey Research Group conducts the dietary data collection methodology, maintenance of

databases used to code and process data, and data review and processing. The dietary data is

released in two files: individual foods and total nutrients. The individual foods file lists each food

reported by the participant, along with details of the consumption such as a USDA FNDDS code

(food code), eating occasion, and amount of food/beverage consumed in grams. The total

nutrient intake file is a daily aggregate of nutrients from the reported foods consumed for that

24-hour recall.

Nutrient Profiling Systems and Nutrient Ratios

This study evaluated five nutrient profiling systems, each rescaled to a 100-point scale for

comparison. Nutri-Score (A–E = 100–20), Health Star Rating (0.5–5 = 10–100), NOVA

classification (1–4 = 80–20), and Food Compass 2 (1–100 = 1–100) scores were converted

from previously reported values (23,27). Alcoholic beverages were assigned null values in

systems that did not evaluate them: NS, HS, NC, and FC. Nutrient Consume Score values

were retrieved using the NCS algorithm available online at Gutbites.org (12). Additionally,

nutrient ratios—carbohydrate-to-fiber, saturated fat-to-unsaturated fat, sodium-to-potassium,

and calorie-to-weight—were mapped to a quantized scale of 1 to 3 in 0.125 increments before

analysis.

Nutri-Score is a scoring algorithm that balances positively weighted nutrients (e.g., fruits,

vegetables, fiber) against negatively weighted ones (e.g., saturated fat, sugars, sodium),

resulting in a color-coded rating from A to E (24). Health Star Rating separates foods into three

groups – dairy products, non-dairy beverages, and all other foods – and assigns points based

on levels of protein, fats, saturated fats, energy, carbohydrates, sugar, and sodium, yielding in a

5-star scale with half-star increments (25). The NOVA Classification ranks foods by processing

level, distinguishing between minimally or unprocessed foods, processed ingredients,

processed foods, and ultra-processed foods, resulting in a 4-level scale (26). Food Compass

assesses foods across nine domains – nutrient ratios, vitamins, minerals, food-based

ingredients, additives, processing, specific lipids, fiber & protein, phytochemicals – producing a

100-point score (23,27).

Nutrient Consume Score (NCS) is an online nutrient profiling system based on nutrient ratios –

Carbohydrate-to-Fiber, Saturated Fat-to-Unsaturated Fat, Sodium-to-Potassium, and Calorie-to-

Weight – and other nutrients (protein, alcohol, vitamin D, iron, calcium) available on U.S.

Nutrition Facts labels (12). Positive adjustments are made for food categories like fruits,

vegetables, nuts, seeds, and whole grains to account for microbiome-supportive and bioactive

factors not listed on food labels: polyphenols, prebiotic fibers, bioactive fats, and fermentation

products. In addition, negative adjustments are made for alcohol, soft drinks, processed meats,

and processed potatoes to reflect additives and food components linked to negative health

outcomes. Three versions of NCS were evaluated in the analysis: NCS proper, NCS without

alcohol included in calculations to compare to other algorithms that don't include alcohol, and

NCS without bioactive adjustments.

All scores and nutrient ratios were evaluated in three ways: weighted by calorie (kcal), weighted

by weight (grams), and unweighted. Unweighted scores were calculated for each individual as

the average score of each food consumed in the 24-hour period divided the total number of food

items. Scores weighted by energy and weight were calculated as the sum of the individual food

items' scores multiplied by the kcal or grams provided by the food item and then divided by the

total kcal or grams for the 24-hour period. Scores were analyzed as continuous variables on a

10-unit scale.

Food Categories and Contributions to Scores

To evaluate which foods were contributing to most of the variation in the scores, a compositional

approach was applied. Individual foods reported were assigned to food group classifications

based on the What We Eat in America (WWEIA) food categories (37). Total intake of food

categories was calculated for each person based on percentage energy (kcals) provided by all

the foods in that category for the 24-hour dietary intake recall period. Center-log transformed

ratios of percentage energy contributed by food categories to overall energy were calculated

(38) using the R *compositions* package (39) to include in regression models.

Outcomes

Anthropometry and blood pressure were evaluated as continuous outcomes and as binary

outcomes based on established cut-offs. Anthropometric characteristics are measured during

the in-person visit by trained technicians. Blood pressure was measured one to four times by

trained technicians. Details about measurement and analysis are available from NHANES

established procedures. The average of the available measurements for each person was used

for analysis.

Obesity was defined as BMI 30 kg/m^2 (40) and abdominal obesity was defined as waist

circumference >88 cm for women and >102 cm for men (41). High blood pressure was defined

as the presence of systolic blood pressure 130 mm/Hg or diastolic blood pressure 85 mm/Hg

(41,42).

Other cardiometabolic risk factors (comparison with NCS only) evaluated included low-density

serum lipoprotein (LDL), high-density lipoprotein (HDL), total cholesterol, and triglycerides, and

plasma glucose. These risk factors were evaluated as continuous measures and in relation to

established cut-offs for high or low values (41,42). Additionally, metabolic syndrome was

assessed and was classified as the presence of three or more of abdominal obesity, high blood

pressure, high LDL, low HDL, or high serum glucose (41).

Covariates

Personal characteristics were included to describe the analytic group and as adjustment

variables in multivariable analyses. Characteristics included were: age (in categories for

description and in years for multivariable analyses), gender (female and male), race or ethnicity

(non-Hispanic White, non-Hispanic Black, non-Hispanic other, and Hispanic). Education (< high

school graduate; high school graduate or equivalent; some college or associate degree; and,

four-year college graduate or more), poverty-to-income ratio (PIR, <1, 1 to <2, 2, to <3, and 3+),

smoking status (less than 100 cigarettes in lifetime; former smoker; current <20 cigarettes per

day; and, current 20+ cigarettes per day), and physical activity (<15, 15 to <75, 75 to <165,

165+ metabolic equivalent score (METS) per week).

Statistical Analysis

For visualization of overall differences in relation to increasing NCS score, descriptive statistics

for covariates were calculated by tertiles of the energy-weighted NCS score. Means and

standard errors were calculated for continuous variables and frequencies and percentages were

calculated for categories variables. To evaluate associations of NPSs and ratio scores with

anthropometry, blood pressure, and cardiometabolic biomarkers, multivariable linear and logistic

regression analyses were performed for continuous and binary outcomes, respectively, and

adjusted for personal characteristics (age, gender, race/ethnicity, education, PIR, smoking, and

exercise. For food category contributions to scores, linear regression analyses were used to

evaluate the contribution of center-log ratio transformed food categories of the energy-weighted

scores and subscores. Except for the compositional analysis of food category contributions to

NPSs and ratios scores, analyses were conducted using population weights supplied by

NHANES and incorporating the complex survey design. A subpopulation command to maintain

appropriate population weights for the study population with complete data for this analysis.

Statistical significance was considered p<0.05 for analyses with cardiometabolic and mortality

outcomes. Statistical analysis was conducted in Stata (StataCorp, TX, USA, version 18).

Results

Study Population Characteristics

For visualization, study population characteristics by tertiles of energy-weighted NCS scores are

presented (Table 1). Categories more represented in the highest score tertiles included age

greater than 65, female gender, Hispanic, non-Hispanic other, four-year college graduates,

poverty-to-income ratio of three or higher, and never smokers. Categories more represented in

the lowest score tertiles included younger age, non-Hispanic White, non-Hispanic Black, lower

education level, current smoker, and highest physical activity level (165+ METS).

Weight and Blood Pressure Associations with Scores and Ratios

All energy-weighted NPS scores (NCS, FC, HS, NC and NS) were significantly associated with

continuous measures of anthropometry and blood pressure (Figure 1). Similar associations

were observed with grams-weighted and unweighted scores (Supplementary Table 1). NCS and

FC showed the largest beta-coefficients for BMI and waist circumference. NCS without alcohol

and NS showed the largest beta-coefficients for systolic and diastolic blood pressure. Removing

bioactives from NCS scoring reduced beta-coefficients for association with anthropometry and

blood pressure. Removing alcohol from NCS scoring reduced beta-coefficients for association

with anthropometric measures, but had the opposite effect on blood pressure.

Higher scores for all energy-weighted NPS scores (NCS, FC, HS, NC, and NS) were associated

with lower odds of obesity, abdominal obesity, and high blood pressure (Figure 2), with similar

associations observed with grams-weighted and unweighted scores (Supplementary Table 1).

NCS and FC showed the lowest odds for obesity, while NCS and NS showed the lowest odds

for abdominal obesity. NCS without alcohol and NS showed the lowest odds for high blood

pressure. Removing bioactives or from NCS scoring reduced odds of obesity, abdominal

obesity, and high blood pressure. Removing alcohol from NCS scoring reduced odds of obesity,

but had the opposite effect on odds of high blood pressure.

Nutrient ratio scores (energy, salt, fat, fiber; Figure 3 and Figure 4) accounted for much of the

signal observed across the full scoring systems, with outcome-specific effects. Ratios were

scaled such that higher values reflected less favorable nutrient profiles. Energy, salt and fat

ratios were favorably associated with anthropometric measures; while associations with blood

pressure varied: salt and fiber ratios were favorable, while energy ratios were unfavorable.

Other Cardiometabolic Risk Factor Associations with Scores and Ratios

Among cardiometabolic risk factors (Supplemental Figures 2A, 2B, 3A, and 3B), there was an

association between NCS, FC and higher total cholesterol. Fiber ratio was favorably associated

with higher HDL and salt ratio was favorably associated with lower glucose.

Food Category Contributors to Scores

The top food categories contributing to NPS variation were largely consistent across scores,

(Figure 5, Supplemental Figure 4). Among negatively contributing categories, NCS provided

relatively higher weight than most other NPS's to "pizza", "soft drinks", "frankfurters" and

"egg/breakfast sandwiches". Among positively contributing categories, NCS provided relatively

higher weight than most other NPS's to "Rice", "Beans, peas, and legumes", "nuts and seeds".

A summary of NCS food scores within different categories can be found in Supplemental Figure

1.

Discussion

Nutrient profiling systems assess the quality of food and beverages, and aim to help consumers, industry, and government select, create, and promote healthier options to improve public and individual health (18). This study shows that leading NPSs are associated with weight and blood pressure, key prognostic factors in cardiometabolic disease, neurological disorders, cancer, and mortality. A ratio-based web-available score performed favorably relative to leading NPSs and also associated with a higher cholesterol likely driven by high favorable HDL. Notably, the food categories contributing greatest to scores were largely consistent across systems. Those foods contributing most to positive scores included minimally processed foods and those that contributed most to negative scores included ultra-processed foods.

A key finding is that much of the power of NPSs may be captured in the following nutrient ratios: carbohydrate-to fiber (29), saturated fat-to-unsaturated fat (30), sodium-to-potassium (31), and calorie-to-weight (32), which all associated with weight and variably associated with blood pressure. Nutrient ratios reflect nutritional balance in whole foods and the degree to which certain nutrients are concentrated or depleted in processed foods. Ratios are explicitly incorporated in systems like NCS and FC and implicitly applied in systems like NS and HS, which assign positive and negative weights to under- and overrepresented nutrients. Nutrient ratios may complement the NOVA classification, which has been critiqued for grouping all ultra-processed foods together, by offering an empirical approach to assessing the relative healthfulness of different ultra-processed foods (43). The unfavorable association of calorie-to-weight with blood pressure could be related to beverages (including juices and sodas) having lower nutrient density and future work may benefit from treating nutrient density of beverages and foods differently.

Another key finding is that including alcohol in a ratio-based score may enhance its health-

discriminatory power for weight, but not blood pressure, perhaps also owing to complexities

around beverages. In NPSs that excluded alcohol from their calculations, alcoholic beverages

contributed to the scores, likely due to co-consumption of alcohol with other foods. This result

also highlights an important point about the role of alcohol in overall dietary intake for individuals

that choose to consume alcohol. Alcohol contributes at least 16% of the average US adult's

daily caloric intake (44), a figure that is likely higher since the impact of COVID-19 on

consumption (45). However, many are unaware of alcohol's impact (46). Given the prevalence

of alcohol consumption and its association with obesity (47) and other adverse health

outcomes(48), there is a need for consumer education and NPSs that account for alcohol's

caloric contribution to encourage healthier dietary choices.

A third key takeaway is that removing adjustments for food groups serving as proxies for

bioactive factors diminished the ability of NCS to discriminate associations with weight and

blood pressure measures. These food group adjustments included both those linked to

positively associated microbiome-active and bioactive factors (e.g., phytochemicals,

fermentable fibers, and bioactive fats) and those linked to negatively associated bioactives (e.g.,

alcohol, fructose, nitrites, acrylamide, advanced glycosylation end products, and trans fats). A

better understanding and measurement of these positive and negative health-associated

bioactive factors, along with their explicit inclusion in food labels and NPS algorithms, are key

focuses of current nutrition research (34).

NPSs have been shown to positively influence consumer buying habits (49) and prospective

cohort studies have shown they can positively impact health measures like obesity and mortality

(19,50). They can play a key role in public health messaging through front-of-package labels

(11). In countries where front of package labels don't yet exist, web-enabled portals (12), and

smartphone technologies that scan product UPCs to determine food quality could be useful

(13,14). A ratio-based score in particular could intuitively encourage individuals to incorporate

more fiber-, potassium-, unsaturated fat-rich foods, potentially countering the health impact of

foods high in simple carbohydrates, sodium, and saturated fat. A ratio-based NPS could also

guide the food industry in developing healthier products by providing empirical guide rails to

rebalance simple carbohydrates with fiber (51,52) and sodium with potassium (53).

Nutrient profiling systems (NPSs) hold promise for improving health outcomes, but they have

limitations. Some systems generate outlier results for certain foods that conflict with

epidemiological data (54). Most exclude alcohol, despite its strong association with adverse

health outcomes (19,27,55), while others are critiqued for being overly simple or too complex

(22). Additionally, most NPSs rely on nutritional databases that lack quantification of the

bioactive components of foods, such as microbiome-active components (i.e. polyphenols,

fermentable fibers, and fermentation products), and are missing data on food additives that may

harm the microbiome and individual (56). Most NPSs fail to account for a food's matrix or ultra-

structure, which may be important for understanding the health benefits of unprocessed foods

(57). NPS algorithms don't currently provide personalized advice which may be important to

account for individual variations in nutrient requirements, including those influenced by the gut

microbiome (58,59). Tailoring algorithms in low- and middle-income country cohorts, where

baseline macro- and micronutrient intakes differ significantly, may also be necessary.

This study design also has limitations including selection (60), recall (61), and reporting (62)

biases. While NHANES attempts to survey respondents to obtain representativeness to the US

population, there is still self-selection of participants who are invited to participate in NHANES,

and individuals may misreport information in surveys and interviews due to misremembering or

aligning responses with social desirability. Additionally, the single assessment of dietary intake

may not be reflective of longer term or usual intake. However, the large sample size of NHANES

does mitigate some of these biases. Temporality of associations cannot be determined due to

the cross-sectional design of NHANES, and weight measures in particular can be subject to

reverse causation bias (63). Results regarding weight should be interpreted in this context, and

it is possible that individuals of different body sizes eat differently rather than the scores predict

weight. Cohort studies demonstrating prospective association between Nutri-score and

abdominal obesity suggest that reverse causation may not entirely explain the associations

observed (19).

In conclusion, this study highlights the potential of nutrient profiling systems (NPSs) including a

nutrient ratio-based web-enabled score to support personal and public health efforts in curbing

weight gain and improving blood pressure. Ratio-based systems, in particular, may guide

individuals, food companies, and governments in rebalancing nutrients in our diet. As a ratio-

based system, NCS may be particularly useful at measuring the spectrum of health of different

processed and ultra-processed foods (Supplementary Figure 1). A ratio-based system also has

advantages in being independent of absolute values of nutrients so it operates independent of

scale and allows assessing food combinations. While NPSs offer valuable insights, most don't

account for alcohol and are primarily supported by correlative research. NCS also offers an

advantage over other algorithms given its web accessibility. Future studies leveraging

smartphone access to the web-enabled algorithm could assess NCS's capacity for

personalization, generalizability across diverse populations, and causal effects on metabolic

health.

Acknowledgements

Grateful to Luke Walker for his vital assistance with coding the scoring algorithms and to Ben

Roberts for his invaluable feedback and constructive criticism on the manuscript. CJD designed

and conducted the research, drafted the paper, and was responsible for final content. CLF

performed the statistical analysis, wrote the methods section, and revised the manuscript. All

authors have read and approved the final version.

368 Data Availability

369 NHANES data used in this work is freely available from the National Center for Health Statistics.

370 Other data described in the manuscript and a code book will be made available upon request

371 pending approval.

373 Funding

374 None

376 Author Disclosures

377 C.L.F. is an Associate Editor for Journal of Nutrition and Annals of Epidemiology and on the

378 Editorial Board for Critical Reviews in Food Science and Nutrition, and works as a consultant for

379 EpidStrategies, A BlueRidge Life Sciences Company. C.J.D. is editor-in-chief at GutBites MD, a

380 not-for-profit web site aimed at making gut health research accessible. He is on the scientific

381 advisory board at Supergut, One Bio, and Oobli.

383 Declaration of Generative AI and AI-assisted technologies in the

384 writing process

385   During the preparation of this manuscript, ChatGPT was used on a limited basis to improve

386   clarity and wording of the text. The authors subsequently reviewed and revised all content. No

387   generative AI tools were used for data analysis or interpretation.

References

1. Nielsen J, Narayan KV, Cunningham SA. Incidence of obesity across adulthood in the United States, 2001-2017-a national prospective analysis. Am J Clin Nutr [Internet] Am J Clin Nutr; 2023 [cited 2024 Oct 19];117. Available from: https://pubmed.ncbi.nlm.nih.gov/36789933/

2. Hirode G, Wong RJ. Trends in the Prevalence of Metabolic Syndrome in the United States, 2011-2016. JAMA 2020;323:2526.

3. Hambleton IR, Caixeta R, Jeyaseelan SM, Luciani S, Hennis AJM. The rising burden of non-communicable diseases in the Americas and the impact of population aging: a secondary analysis of available data. Lancet Regional Health - Americas 2023;21:100483.

4. Chew NWS, Ng CH, Tan DJH, Kong G, Lin C, Chin YH, Lim WH, Huang DQ, Quek J, Fu CE, et al. The global burden of metabolic disease: Data from 2000 to 2019. Cell Metab [Internet] Cell Metab; 2023 [cited 2024 Oct 19];35. Available from: https://pubmed.ncbi.nlm.nih.gov/36889281/

5. Lane MM, Davis JA, Beattie S, Gómez-Donoso C, Loughman A, O'Neil A, Jacka F, Berk M, Page R, Marx W, et al. Ultraprocessed food and chronic noncommunicable diseases: A systematic review and meta-analysis of 43 observational studies. Obes Rev [Internet] Obes Rev; 2021 [cited 2024 Oct 19];22. Available from: https://pubmed.ncbi.nlm.nih.gov/33167080/

6. Fazzino TL, Rohde K, Sullivan DK. Hyper-palatable foods: Development of a quantitative definition and application to the US food system database. Obesity (Silver Spring) Wiley; 2019;27:1761–8.

7. Alamnia TT, Sargent GM, Kelly M. Dietary patterns and associations with metabolic risk factors for non-communicable disease. Sci Rep Nature Publishing Group; 2023;13:1–10.

8. Sadler CR, Grassby T, Hart K, Raats MM, Sokolović M, Timotijevic L. "Even We Are Confused": A Thematic Analysis of Professionals' Perceptions of Processed Foods and Challenges for Communication. Frontiers in Nutrition 2022;9:826162.

9. Vijaykumar S, McNeill A, Simpson J. Associations between conflicting nutrition information, nutrition confusion and backlash among consumers in the UK. Public Health Nutr 2021;24:914.

10. Gokani N, Grosso G. An appetite for change: shaping consumer choices through food labelling amidst global challenges. Int J Food Sci Nutr [Internet] Int J Food Sci Nutr; 2024 [cited 2024 Oct 19];75. Available from: https://pubmed.ncbi.nlm.nih.gov/38508777/

11. Pomeranz JL, Wilde P, Mozaffarian D, Micha R. Mandating Front-of-Package Food Labels in the U.S. – What are the First Amendment Obstacles? Food Policy 2019;86:101722.

12. Carb-to-Fiber Ratio Calculator for Gut Microbiome Nutrition [Internet]. Gut Bites MD. 2022 [cited 2024 Oct 19]. Available from: https://gutbites.org/carb-fiber-ratio-calculator/

13. Soutjis B. The new digital face of the consumerist mediator: the case of the "Yuka" mobile app. Journal of Cultural Economics [Internet] 2020 [cited 2024 Oct 19]; Available from: https://doi.org/10.1080/17530350.2019.1603116

14. Gut Bites MD [Internet]. Gut Bites MD. 2023 [cited 2024 Dec 21]. Available from: https://gutbites.org/nutrient-consume-score-app/

15. Labonté MÈ, Poon T, Gladanac B, Ahmed M, Franco-Arellano B, Rayner M, L'Abbé MR. Nutrient Profile Models with Applications in Government-Led Nutrition Policies Aimed at Health Promotion and Noncommunicable Disease Prevention: A Systematic Review. Adv Nutr [Internet] Adv Nutr; 2018 [cited 2024 Oct 19];9. Available from: https://pubmed.ncbi.nlm.nih.gov/30462178/

16. Greenberg D, Drewnowski A, Black R, Weststrate JA, O'Shea M. A progressive nutrient profiling system to guide improvements in nutrient density of foods and beverages. Front Nutr Frontiers Media SA; 2021;8:774409.

17. Bauner C, Rahman R. The effect of front-of-package nutrition labelling on product composition. Eur Rev Agric Econ Oxford University Press (OUP); 2024;51:482–505.

18. Martin C, Turcotte M, Cauchon J, Lachance A, Pomerleau S, Provencher V, Labonté M-È. Systematic review of nutrient profile models developed for nutrition-related policies and regulations aimed at noncommunicable disease prevention -an update. Adv Nutr Elsevier BV; 2023;14:1499–522.

19. Rey-García J, Mérida DM, Donat-Vargas C, Sandoval-Insausti H, Rodríguez-Ayala M, Banegas JR, Rodríguez-Artalejo F, Guallar-Castillón P. Less Favorable Nutri-Score Consumption Ratings Are Prospectively Associated with Abdominal Obesity in Older Adults. Nutrients Multidisciplinary Digital Publishing Institute; 2024;16:1020.

20. Egnell M, Seconda L, Neal B, Mhurchu CN, Rayner M, Jones A, Touvier M, Kesse-Guyot E, Hercberg S, Julia C. Prospective associations of the original Food Standards Agency nutrient profiling system and three variants with weight gain, overweight and obesity risk: results from the French NutriNet-Santé cohort. Br J Nutr [Internet] Br J Nutr; 2021 [cited 2024 Oct 19];125. Available from: https://pubmed.ncbi.nlm.nih.gov/32878658/

21. Poti JM, Braga B, Qin B. Ultra-processed Food Intake and Obesity: What Really Matters for Health – Processing or Nutrient Content? Curr Obes Rep 2017;6:420.

22. O'Hearn M, Erndt-Marino J, Gerber S, Lauren BN, Economos C, Wong JB, Blumberg JB, Mozaffarian D. Validation of Food Compass with a healthy diet, cardiometabolic health, and mortality among U.S. adults, 1999–2018. Nat Commun Nature Publishing Group; 2022;13:1–14.

23. Barrett EM, Shi P, Blumberg JB, O'Hearn M, Micha R, Mozaffarian D. Food Compass 2.0 is an improved nutrient profiling system to characterize healthfulness of foods and beverages.

Nature Food Nature Publishing Group; 2024;1–5.

24. Hercberg S, Touvier M, Salas-Salvado J. The Nutri-Score nutrition label. International Journal for Vitamin and Nutrition Research [Internet] Hogrefe AG; 2021 [cited 2024 Oct 19]; Available from: https://econtent.hogrefe.com/doi/10.1024/0300-9831/a000722

25. Muhammad Junaid Hasni, Mohsin Abdur Rehman, Nicolas Pontes, Muhammad Zafar Yaqub. Health Star Rating Labels: A systematic review and future research agenda. Food Qual Prefer Elsevier; 2025;122:105310.

26. Monteiro CA, Cannon G, Levy RB, Moubarac J-C, Louzada MLC, Rauber F, Khandpur N, Cediel G, Neri D, Martinez-Steele E, et al. Ultra-processed foods: what they are and how to identify them. Public Health Nutr 2019;22:936.

27. Mozaffarian D, El-Abbadi NH, O'Hearn M, Erndt-Marino J, Masters WA, Jacques P, Shi P, Blumberg JB, Micha R. Food Compass is a nutrient profiling system using expanded characteristics for assessing healthfulness of foods. Nature Food Nature Publishing Group; 2021;2:809–18.

28. Derbyshire E. Are all "ultra-processed" foods nutritional demons? A commentary and nutritional profiling analysis. Trends Food Sci Technol Elsevier BV; 2019;94:98–104.

29. Fontanelli M de M, Micha R, Sales CH, Liu J, Mozaffarian D, Fisberg RM. Application of the ≤ 10:1 carbohydrate to fiber ratio to identify healthy grain foods and its association with cardiometabolic risk factors. Eur J Nutr Springer Science and Business Media LLC; 2020;59:3269–79.

30. Grundy SM. What is the desirable ratio of saturated, polyunsaturated, and monounsaturated fatty acids in the diet? Am J Clin Nutr 1997;66:988S – 990S.

31. Vaudin A, Wambogo E, Moshfegh AJ, Sahyoun NR. Sodium and potassium intake, the sodium to potassium ratio, and associated characteristics in older adults, NHANES 2011-2016. J Acad Nutr Diet Elsevier BV; 2022;122:64–77.

32. Mendoza JA, Drewnowski A, Christakis DA. Dietary energy density is associated with obesity and the metabolic syndrome in U.S. adults. Diabetes Care American Diabetes Association; 2007;30:974–9.

33. Kelly OJ, Gilman JC, Ilich JZ. Utilizing dietary nutrient ratios in nutritional research: Expanding the concept of nutrient ratios to macronutrients. Nutrients MDPI AG; 2019;11:282.

34. Damman CJ. Perspective: Nutrition's Next Chapter – Bioactive Gaps and the Microbiome–Mitochondria Axis. Adv Nutr Elsevier; 2023;14:420–5.

35. CDC. National Health and Nutrition Examination Survey [Internet]. National Health and Nutrition Examination Survey. 2025 [cited 2025 Dec 23]. Available from: https://www.cdc.gov/nchs/nhanes/index.html?utm_source=chatgpt.com

36. NHANES Questionnaires, Datasets, and Related Documentation [Internet]. [cited 2024 Dec 17]. Available from: https://wwwn.cdc.gov/nchs/nhanes/continuousnhanes/default.aspx?BeginYear=2015

37. DMR - Food Categories : USDA ARS [Internet]. [cited 2024 Nov 7]. Available from: https://www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-group/docs/dmr-food-categories/

38. Pincus R. Aitchison, J.: The statistical analysis of compositional data. Chapman and hall, London ⬜ New York 1986, XII, 416 pp., £ 25,00. Biom J Wiley; 1988;30:794–794.

39. van den Boogaart KG. Using the R package "compositions" [Internet]. [cited 2024 Dec 17]. Available from: https://cran.r-project.org/web/packages/compositions/vignettes/UsingCompositions.pdf

40. Obesity and overweight [Internet]. [cited 2025 Jun 2]. Available from: https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight

41. Grundy SM, Brewer HB Jr, Cleeman JI, Smith SC Jr, Lenfant C. Definition of metabolic syndrome: Report of the national heart, lung, and blood institute/American heart association conference on scientific issues related to definition. Circulation Ovid Technologies (Wolters Kluwer Health); 2004;109:433–8.

42. Grundy SM, Cleeman JI, Merz CNB, Brewer HB Jr, Clark LT, Hunninghake DB, Pasternak RC, Smith SC Jr, Stone NJ, National Heart, Lung, and Blood Institute, et al. Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines. Circulation Ovid Technologies (Wolters Kluwer Health); 2004;110:227–39.

43. Petrus RR, do Amaral Sobral PJ, Tadini CC, Gonçalves CB. The NOVA classification system: A critical perspective in food science. Trends Food Sci Technol Elsevier BV; 2021;116:603–8.

44. Nielsen SJ, Kit BK, Fakhouri T, Ogden CL. Calories consumed from alcoholic beverages by U.S. adults, 2007-2010. NCHS Data Brief NCHS Data Brief; 2012;1–8.

45. Barbosa C, Dowd WN, Barnosky A, Karriker-Jaffe KJ. Alcohol consumption during the first year of the COVID-19 pandemic in the United States: Results from a nationally representative longitudinal survey. J Addict Med Ovid Technologies (Wolters Kluwer Health); 2023;17:e11–7.

46. Robinson E, Humphreys G, Jones A. Alcohol, calories, and obesity: A rapid systematic review and meta-analysis of consumer knowledge, support, and behavioral effects of energy labeling on alcoholic drinks. Obes Rev Wiley; 2021;22:e13198.

47. Golzarand M, Salari-Moghaddam A, Mirmiran P. Association between alcohol intake and overweight and obesity: a systematic review and dose-response meta-analysis of 127 observational studies. Crit Rev Food Sci Nutr Informa UK Limited; 2022;62:8078–98.

48. Moss HB. The impact of alcohol on society: a brief overview. Soc Work Public Health Informa UK Limited; 2013;28:175–7.

49. van den Akker K, Bartelet D, Brouwer L, Luijpers S, Nap T, Havermans R. The impact of the nutri-score on food choice: A choice experiment in a Dutch supermarket. Appetite Elsevier BV; 2022;168:105664.

50. Bonaccio M, Di Castelnuovo A, Ruggiero E, Costanzo S, Grosso G, De Curtis A, Cerletti C,

Donati MB, de Gaetano G, Iacoviello L, et al. Joint association of food nutritional profile by Nutri-Score front-of-pack label and ultra-processed food intake with mortality: Moli-sani prospective cohort study. BMJ BMJ; 2022;378:e070688.

51. Xu B, Fu J, Qiao Y, Cao J, Deehan EC, Li Z, Jin M, Wang X, Wang Y. Higher intake of microbiota-accessible carbohydrates and improved cardiometabolic risk factors: a meta-analysis and umbrella review of dietary management in patients with type 2 diabetes. Am J Clin Nutr Elsevier BV; 2021;113:1515–30.

52. Marcobal AM, McConnell BR, Drexler RA, Ng KM, Maldonado-Gomez MX, Conner AMS, Vierra CG, Krishnakumar N, Gerber HM, Garcia JKA, et al. Highly soluble β-glucan fiber modulates mechanisms of blood glucose regulation and intestinal permeability. Nutrients MDPI AG; 2024;16:2240.

53. Neal B, Wu Y, Feng X, Zhang R, Zhang Y, Shi J, Zhang J, Tian M, Huang L, Li Z, et al. Effect of salt substitution on cardiovascular events and death. N Engl J Med Massachusetts Medical Society; 2021;385:1067–77.

54. Ortenzi F, Kolby M, Lawrence M, Leroy F, Nordhagen S, Phillips SM, van Vliet S, Beal T. Limitations of the Food Compass Nutrient Profiling System. J Nutr Elsevier BV; 2023;153:610–4.

55. Dunford EK, Huang L, Peters SAE, Crino M, Neal BC, Ni Mhurchu C. Evaluation of Alignment between the Health Claims Nutrient Profiling Scoring Criterion (NPSC) and the Health Star Rating (HSR) Nutrient Profiling Models. Nutrients Multidisciplinary Digital Publishing Institute; 2018;10:1065.

56. Whelan K, Bancil AS, Lindsay JO, Chassaing B. Ultra-processed foods and food additives in gut health and disease. Nat Rev Gastroenterol Hepatol Springer Science and Business Media LLC; 2024;21:406–27.

57. Aguilera JM. The food matrix: implications in processing, nutrition and health. Crit Rev Food Sci Nutr Crit Rev Food Sci Nutr; 2019;59:3612–29.

58. (PDF) Nutritional Value Score rates foods based on global health priorities [Internet]. ResearchGate. [cited 2024 Nov 9]. Available from: https://www.researchgate.net/publication/375502962_Nutritional_Value_Score_rates_foods_based_on_global_health_priorities

59. Simon M-C, Sina C, Ferrario PG, Daniel H, Working Group "Personalized Nutrition" of the German Nutrition Society. Gut microbiome analysis for personalized nutrition: The state of science. Mol Nutr Food Res Wiley; 2023;67:e2200476.

60. Bernstein OM, Vegetabile BG, Salazar CR, Grill JD, Gillen DL. Adjustment for biased sampling using NHANES derived propensity weights. Health Serv Outcomes Res Methodol [Internet] Springer Science and Business Media LLC; 2022; Available from: https://link.springer.com/article/10.1007/s10742-022-00283-x

61. Va P, Dodd KW, Zhao L, Thompson-Paul AM, Mercado CI, Terry AL, Jackson SL, Wang C-Y, Loria CM, Moshfegh AJ, et al. Evaluation of measurement error in 24-hour dietary recall for assessing sodium and potassium intake among US adults - National Health and Nutrition Examination Survey (NHANES), 2014. Am J Clin Nutr Elsevier BV;

2019;109:1672–82.

62. Murakami K, Livingstone MBE. Prevalence and characteristics of misreporting of energy intake in US adults: NHANES 2003-2012. Br J Nutr Cambridge University Press (CUP); 2015;114:1294–303.

63. Tobias DK. Addressing reverse causation bias in the obesity paradox is not "one size fits all." Diabetes Care American Diabetes Association; 2017;40:1000–1.

Tables and Figures

Table 1. Study population characteristics by tertiles of energy-weighted NCS score in NHANES adults, 2015-2016

Personal characteristics	Tertile 1 (<37.08), n=1533	Tertile 2 (37.08 to 48.31), n=1533	Tertile 3 (>48.31), n=1532
Age (years)			
20 to 34	458 (33.3%)	370 (25.4%)	360 (25.4%)
35 to 49	415 (26.8%)	392 (24.8%)	339 (22.6%)
50 to 64	380 (25.2%)	434 (28.6%)	403 (27.4%)
65+	280 (14.8%)	337 (21.2%)	430 (24.7%)
Gender			
Male	764 (53.3%)	752 (47.1%)	693 (43.0%)
Female	769 (46.7%)	781 (52.9%)	839 (57.0%)
Race and non-Hispanic ethnicity			
Non-Hispanic White	589 (67.7%)	559 (68.0%)	420 (58.2%)
Non-Hispanic Black	415 (13.8%)	316 (10.4%)	220 (7.6%)
Non-Hispanic Other	116 (4.9%)	199 (8.2%)	333 (16.2%)
Hispanic	413 (13.6%)	459 (13.4%)	559 (18.0%)
Education			
<HS graduate or equivalent	293 (12.4%)	324 (12.8%)	394 (13.5%)
HS graduate or equivalent	412 (26.9%)	347 (21.6%)	264 (13.8%)
Some college or associates degree	527 (36.1%)	463 (33.4%)	389 (30.6%)
Four-year college graduate or more	301 (24.6%)	399 (32.2%)	485 (42.2%)
Poverty-to-income ratio			
<1	307 (14.7%)	265 (10.2%)	312 (13.5%)
1 to <2	399 (19.4%)	387 (20.5%)	338 (15.0%)
2 to <3	252 (18.0%)	262 (16.9%)	219 (12.8%)
≥3	460 (41.6%)	488 (45.8%)	500 (49.8%)
Missing	115 (6.3%)	131 (6.6%)	163 (8.9%)
Smoking			
Never	807 (51.6%)	869 (55.4%)	1,009 (64.1%)
Former	338 (23.7%)	360 (26.7%)	365 (25.4%)
Current, less than 20 cig/day	282 (16.0%)	242 (14.2%)	139 (8.6%)
Current, 20+ cig/day	106 (8.6%)	62 (3.7%)	19 (2.0%)
Physical activity (METS/week)			
<15	666 (38.2%)	680 (40.4%)	688 (36.1%)
15 to <75	431 (31.3%)	453 (32.7%)	510 (41.2%)
75 to <165	219 (15.8%)	195 (14.4%)	197 (13.6%)
165+	217 (14.7%)	205 (12.5%)	137 (9.0%)

Figure Legends

Figure 1. Associations of nutrient profile scores (in 10-unit values) with body mass index, waist circumference, and blood pressure. Beta-coefficients from multivariable adjusted models are displayed. Models are adjusted for age, gender, race or Hispanic ethnicity, education, smoking, poverty-income ratio, and physical activity.

Figure 2. Associations of nutrient profile scores (in 10-unit values) with obesity, abdominal obesity, and high blood pressure. Odds ratios from multivariable adjusted models are displayed. Models are adjusted for age, gender, race or Hispanic ethnicity, education, smoking, poverty-income ratio, and physical activity.

Figure 3. Associations of nutrient ratio scores (in 1-unit values) with body mass index, waist circumference, and blood pressure. Beta-coefficients from multivariable adjusted models are displayed. Models are adjusted for age, gender, race or Hispanic ethnicity, education, smoking, poverty-income ratio, and physical activity.

Figure 4. Associations of nutrient ratio scores (in 1-unit values) with obesity, abdominal obesity, and high blood pressure. Odds ratios from multivariable adjusted models are displayed. Models are adjusted for age, gender, race or Hispanic ethnicity, education, smoking, poverty-income ratio, and physical activity.

Figure 5. Heatmap of food category contributions to kcal-weighted scores. Values represented are the percentage of variation explained (R-squared*100*indicator for direction of association). Food categories that had no consumption in the study population and foods not included in a WWEIA category are not included, and only foods for which at least it explains at least 1% of variation of a score are included. Darker shading represents more variation explained, and red indicates inverse associations and blue indicates positive associations.

Supplemental Figure 1. Distribution of NCS scores for representative foods, organized by food category.

Supplemental Figure 2. Associations of nutrient profile scores (in 10-unit values) with high LDL cholesterol, low HDL cholesterol, high total cholesterol, high triglycerides, and high plasma glucose. Beta-coefficients for continuous outcomes (panel A) and odds ratios for dichotomous outcomes (panel B) from multivariable adjusted models are displayed. Models adjusted for age, gender, race or Hispanic ethnicity, education, smoking, poverty-income ratio, and physical activity.

Supplemental Figure 3. Associations of nutrient ratios (in 1-unit values) with LDL cholesterol, HDL cholesterol, total cholesterol, triglycerides, and plasma glucose. Beta-coefficients for continuous outcomes (panel A) and odds ratios for dichotomous outcomes (panel B) from multivariable adjusted models are displayed. Models are adjusted for age, gender, race or Hispanic ethnicity, education, smoking, poverty-income ratio, and physical activity.

Supplemental Figure 4. Heatmap of food category contributions to kcal-weighted scores. Values represented are the percentage of variation explained (R-squared*100*indicator for direction of association). All WWEIA food categories displayed. Darker shading represents more variation explained, and red indicates inverse associations and blue indicates positive associations.

Supplemental Table 1. Associations of nutrient profile scores (in 10-unit values) with continuous measures of anthropometry and blood pressure, and with dichotomous measures for obesity, abdominal obesity, and high blood pressure, using unweighted, weighted by grams, and weighted by kcal approaches to scores.

Multivariable regression results	NCS beta/OR (CI), p-value	Food Compass 2.0 beta/OR (CI), p-value	Health Star Rating beta/OR (CI), p-value	Nova Category beta/OR (CI), p-value	Nutri-Score beta/OR (CI), p-value
Unweighted					
Body Mass Index (kg/m2)	-0.86 (-1.18,-0.55), <0.0001	-0.73 (-1.02,-0.43), 0.0001	-0.72 (-1.05,-0.39), 0.0003	-0.97 (-1.39,-0.55), 0.0002	-0.78 (-1.13,-0.44), 0.0002
Waist Circumference (cm)	-2.08 (-2.91,-1.26), 0.0001	-1.73 (-2.48,-0.98), 0.0002	-1.71 (-2.56,-0.87), 0.0006	-2.32 (-3.41,-1.23), 0.0004	-1.91 (-2.81,-1.01), 0.0004
Systolic blood pressure (mmHg)	-0.65 (-1.28,-0.03), 0.0426	-0.69 (-1.14,-0.24), 0.0055	-0.73 (-1.53, 0.07), 0.0709	-0.66 (-1.40, 0.07), 0.0729	-0.84 (-1.66,-0.03), 0.0439
Diastolic blood pressure (mmHg)	-0.30 (-0.65, 0.05), 0.0899	-0.27 (-0.69, 0.15), 0.1915	-0.52 (-0.86,-0.19), 0.0046	-0.37 (-0.88, 0.13), 0.1382	-0.49 (-0.89,-0.08), 0.0211
Obese	0.78 (0.71, 0.85), <0.0001	0.81 (0.73, 0.89), 0.0004	0.83 (0.75, 0.91), 0.0009	0.77 (0.68, 0.87), 0.0004	0.82 (0.74, 0.91), 0.0010
Abdominal obesity	0.81 (0.72, 0.92), 0.0034	0.83 (0.74, 0.93), 0.0032	0.82 (0.73, 0.92), 0.0023	0.76 (0.66, 0.87), 0.0009	0.81 (0.72, 0.91), 0.0015
High blood pressure	0.94 (0.86, 1.03), 0.1940	0.92 (0.84, 1.02), 0.1106	0.89 (0.80, 0.99), 0.0347	0.95 (0.85, 1.07), 0.4175	0.90 (0.80, 1.02), 0.0813
Weighted by grams					
Body Mass Index (kg/m2)	-1.00 (-1.29,-0.72), <0.0001	-1.15 (-1.49,-0.80), <0.0001	-0.86 (-1.14,-0.59), <0.0001	-0.89 (-1.22,-0.57), <0.0001	-0.70 (-0.93,-0.47), <0.0001
Waist Circumference (cm)	-2.42 (-3.18,-1.66), <0.0001	-2.69 (-3.58,-1.79), <0.0001	-2.01 (-2.76,-1.26), <0.0001	-2.10 (-2.94,-1.27), 0.0001	-1.67 (-2.27,-1.07), <0.0001
Systolic blood pressure (mmHg)	-0.04 (-0.63, 0.56), 0.9005	-0.52 (-1.18, 0.15), 0.1190	-0.39 (-1.01, 0.23), 0.1963	-0.31 (-0.98, 0.36), 0.3419	-0.33 (-0.90, 0.23), 0.2244
Diastolic blood pressure (mmHg)	-0.05 (-0.35, 0.24), 0.7076	-0.50 (-0.87,-0.12), 0.0124	-0.41 (-0.69,-0.14), 0.0059	-0.35 (-0.65,-0.06), 0.0233	-0.26 (-0.50,-0.03), 0.0269
Obese	0.75 (0.69, 0.83), <0.0001	0.73 (0.64, 0.83), 0.0001	0.80 (0.73, 0.87), 0.0001	0.79 (0.70, 0.88), 0.0004	0.83 (0.78, 0.90), 0.0001
Abdominal obesity	0.76 (0.70, 0.84), <0.0001	0.74 (0.66, 0.83), <0.0001	0.79 (0.71, 0.88), 0.0003	0.79 (0.70, 0.89), 0.0009	0.82 (0.75, 0.91), 0.0008
High blood pressure	1.06 (0.98, 1.15), 0.1485	0.95 (0.85, 1.07), 0.4076	0.98 (0.90, 1.06), 0.5896	1.00 (0.92, 1.07), 0.8928	0.99 (0.92, 1.06), 0.7751
Weighted by kcal					
Body Mass Index (kg/m2)	-0.64 (-0.86,-0.41), <0.0001	-0.51 (-0.70,-0.32), <0.0001	-0.31 (-0.54,-0.08), 0.0124	-0.32 (-0.56,-0.08), 0.0128	-0.40 (-0.61,-0.19), 0.0012
Waist Circumference (cm)	-1.63 (-2.23,-1.02), <0.0001	-1.27 (-1.81,-0.72), 0.0002	-0.78 (-1.35,-0.20), 0.0114	-0.80 (-1.49,-0.11), 0.0253	-1.05 (-1.58,-0.52), 0.0007
Systolic blood pressure (mmHg)	-1.01 (-1.67,-0.35), 0.0051	-1.03 (-1.60,-0.46), 0.0015	-1.01 (-1.83,-0.19), 0.0187	-0.82 (-1.42,-0.23), 0.0100	-1.20 (-1.96,-0.43), 0.0045
Diastolic blood pressure (mmHg)	-0.56 (-0.90,-0.23), 0.0026	-0.58 (-1.04,-0.11), 0.0193	-0.75 (-1.19,-0.32), 0.0022	-0.54 (-0.99,-0.10), 0.0205	-0.80 (-1.22,-0.39), 0.0009
Obese	0.85 (0.79, 0.92), 0.0004	0.89 (0.81, 0.97), 0.0138	0.94 (0.89, 1.00), 0.0359	0.94 (0.87, 1.02), 0.1117	0.92 (0.85, 0.98), 0.0207
Abdominal obesity	0.83 (0.75, 0.91), 0.0007	0.88 (0.81, 0.95), 0.0021	0.89 (0.83, 0.95), 0.0024	0.90 (0.83, 0.98), 0.0210	0.85 (0.78, 0.92), 0.0007
High blood pressure	0.90 (0.81, 1.00), 0.0586	0.87 (0.77, 0.99), 0.0399	0.85 (0.74, 0.98), 0.0297	0.91 (0.83, 0.99), 0.0333	0.85 (0.74, 0.97), 0.0235

Outcome	Measure	Beta-coefficient	
Body mass index	NCS	-0.64	
	NCS without bioactives	-0.59	
	NCS without alcohol	-0.53	
	Food Compass 2.0	-0.51	
	Health Star Rating	-0.31	
	NOVA Category	-0.32	
	Nutri-Score	-0.40	
Waist circumference	NCS	-1.63	
	NCS without bioactives	-1.45	
	NCS without alcohol	-1.32	
	Food Compass 2.0	-1.27	
	Health Star Rating	-0.78	
	NOVA Category	-0.80	
	Nutri-Score	-1.05	
Systolic blood pressure	NCS	-1.01	
	NCS without bioactives	-0.93	
	NCS without alcohol	-1.25	
	Food Compass 2.0	-1.03	
	Health Star Rating	-1.01	
	NOVA Category	-0.82	
	Nutri-Score	-1.20	
Diastolic blood pressure	NCS	-0.56	
	NCS without bioactives	-0.53	
	NCS without alcohol	-0.80	
	Food Compass 2.0	-0.58	
	Health Star Rating	-0.75	
	NOVA Category	-0.54	
	Nutri-Score	-0.80	

Outcome	Measure	Odds Ratio	
Obesity	NCS	0.85	
	NCS without bioactives	0.86	
	NCS without alcohol	0.88	
	Food Compass 2.0	0.89	
	Health Star Rating	0.94	
	NOVA Category	0.94	
	Nutri-Score	0.92	
Abdominal Obesity	NCS	0.83	
	NCS without bioactives	0.85	
	NCS without alcohol	0.86	
	Food Compass 2.0	0.88	
	Health Star Rating	0.89	
	NOVA Category	0.90	
	Nutri-Score	0.85	
High Blood Pressure	NCS	0.90	
	NCS without bioactives	0.92	
	NCS without alcohol	0.85	
	Food Compass 2.0	0.87	
	Health Star Rating	0.85	
	NOVA Category	0.91	
	Nutri-Score	0.85	

0.71   1.0

Outcome	Measure	Beta-coefficient	
Body mass index	Energy ratio	1.75	
	Salt ratio	1.74	
	Fiber ratio	0.81	
	Fat ratio	2.24	
Waist circumference	Energy ratio	3.94	
	Salt ratio	3.90	
	Fiber ratio	1.95	
	Fat ratio	6.11	
Systolic blood pressure	Energy ratio	-0.66	
	Salt ratio	2.86	
	Fiber ratio	4.24	
	Fat ratio	-0.18	
Diastolic blood pressure	Energy ratio	-0.97	
	Salt ratio	0.24	
	Fiber ratio	1.03	
	Fat ratio	-2.87	

-5 -4 -3 -2 -1 0 1 2 3 4 5 6 7 8 9 10

Outcome	Measure	Odds Ratio	
Obesity	Energy ratio	1.58	
	Salt ratio	1.73	
	Fiber ratio	1.33	
	Fat ratio	3.65	
Abdominal Obesity	Energy ratio	1.89	
	Salt ratio	1.30	
	Fiber ratio	1.16	
	Fat ratio	3.77	
High Blood Pressure	Energy ratio	0.58	
	Salt ratio	1.20	
	Fiber ratio	1.91	
	Fat ratio	1.32	

Food category	NCS	NCS, No Bioactives	NCS, No Alcohol	Food Compass 2.0	Health Star Rating	NOVA Category	Nutri-Score
Pizza	-5.49	-6.11	-4.57	-1.47	-2.50	-3.02	-1.97
Soft drinks	-3.51	-0.99	-2.32	-3.56	-3.37	-1.05	-1.09
Frankfurter sandwiches (single code)	-3.29	-2.40	-2.95	-2.51	-0.88	-1.68	-0.90
Egg/breakfast sandwiches (single code)	-2.95	-1.28	-2.48	-0.71	-0.54	-0.45	-1.04
Tea	-1.75	-0.98	-1.92	-2.31	-2.58	-0.92	-1.85
Cookies and brownies	-1.71	-2.64	-1.04	-0.89	-0.90	-0.46	-1.15
Burgers (single code)	-1.11	-0.37	-1.09	-1.87	-0.60	-3.81	-1.28
Liquor and cocktails	-0.98	-0.44	-1.84	-0.95	-2.61	-1.71	-3.48
Biscuits, muffins, quick breads	-0.92	-1.23	-0.59	-0.15	-0.26	0.00	-0.20
French fries and other fried white potatoes	-0.90	-0.13	-0.66	-0.18	0.08	-1.54	-0.08
Sausages	-0.50	-0.20	-0.54	-0.52	-0.84	-0.68	-1.21
Salad dressings and vegetable oils	-0.49	-0.59	-0.61	-0.02	-1.19	-0.41	-0.71
Potato chips	-0.38	-0.16	-0.47	-0.22	-0.29	-1.22	-0.66
Soy-based condiments	-0.14	-0.03	-0.16	-0.11	-0.21	-1.06	-0.22
Other Mexican mixed dishes	-0.05	-0.02	-0.04	0.00	0.01	1.35	0.04
Beer	-0.02	0.01	-7.42	-4.40	-10.51	-5.96	-11.67
Wine	0.00	0.00	-0.60	-0.07	-0.76	-0.30	-1.13
Fried rice and lo/chow mein	0.01	0.00	0.01	0.02	0.42	3.04	0.52
Citrus juice	0.05	0.05	0.06	0.74	0.18	1.09	0.01
Pasta mixed dishes, excludes macaroni and cheese	0.09	0.15	0.12	0.23	0.38	0.45	1.45
Tortillas	0.41	0.61	0.44	0.19	1.21	0.08	1.32
Tomato-based condiments	0.50	0.20	0.83	0.53	0.68	0.33	1.29
Eggs and omelets	0.70	0.69	0.88	1.92	0.42	0.36	0.06
Smoothies and grain drinks	0.72	0.76	0.92	2.33	-0.05	1.64	0.00
Soups	1.05	0.88	1.27	0.81	2.05	1.15	2.03
Stir-fry and soy-based sauce mixtures	1.09	0.55	0.86	0.58	0.18	1.57	0.33
Seafood mixed dishes	1.09	0.71	1.07	1.31	0.31	0.12	0.45
Yeast breads	1.10	0.15	1.21	1.09	1.79	0.02	2.65
Bananas	1.13	0.64	1.09	1.42	0.84	1.13	1.03
Rice mixed dishes	1.96	0.90	2.10	1.24	1.71	1.25	1.60
Fish	2.52	1.23	2.34	4.60	0.68	1.63	0.66
Oatmeal	3.22	2.20	3.26	3.33	1.70	2.04	3.01
Nuts and seeds	4.53	3.40	4.27	9.81	3.70	1.67	1.14
Beans, peas, legumes	5.91	5.26	5.98	4.36	5.44	1.64	5.66
Rice	6.14	2.42	6.51	4.70	5.39	12.89	6.90