Comparing tangible retinal image characteristics with deep learning features reveals their complementarity for gene association and disease prediction

Michael J. Beyeler[1,2], Olga Trofimova[1,2], Dennis Bontempi[1,2], José Vargas Quiros[3,4], Ilenia Meloni[5,6], Bart Liefers[3,4], Adham Elwakil[5,6], Sacha Bors[1,2], Ian Quintas[1,2], Leah Böttger[1,2], Ilaria Iuliani[1,2], Sofia Ortin Vela[1,2], Federica M. Conedera[7], Mattia Tomasoni[5,6], Ciara Bergin[5], Reinier O. Schlingemann[5,8], Caroline C.W. Klaver[3,4,9,10], VascX Consortium, David M. Presby[1,2,*], Sven Bergmann[1,2,11,*]

[1]Dept. of Computational Biology, University of Lausanne, Lausanne, Switzerland
[2]Swiss Institute of Bioinformatics, Lausanne, Switzerland
[3]Dept. of Ophthalmology, Erasmus University Medical Center, Rotterdam, The Netherlands
[4]Dept. of Epidemiology, Erasmus University Medical Center, Rotterdam, The Netherlands
[5]Dept. of Ophthalmology, University of Lausanne, Fondation Asile des Aveugles, Jules Gonin Eye Hospital, Lausanne, Switzerland
[6]Platform for Research in Ocular Imaging, Fondation Asile des Aveugles, Jules Gonin Eye Hospital, Lausanne, Switzerland
[7]Department of BioMedical Research, Immunology RIA, Inselspital, University of Bern, Bern, Switzerland
[8]University Medical Centres, Amsterdam, The Netherlands
[9]Dept. of Ophthalmology, Radboud University Medical Center, Nijmegen, the Netherlands
[10]Institute of Molecular and Clinical Ophthalmology, University of Basel, Switzerland
[11]Dept. of Integrative Biomedical Sciences, University of Cape Town, Cape Town, South Africa

* joint corresponding (david.presby@unil.ch, sven.bergmann@unil.ch) and equal senior authors

Abstract

Advances in AI, including deep learning (DL), are transforming medical image analysis by enabling automated disease risk predictions. However, DL's outputs and latent space representations often lack interpretability, impeding clinical trust and biological insight. In this study, we evaluated *RETFound*, a foundation model for retinal images, by comparing its predictive performance and genetic associations to those obtained using clinically interpretable tangible image features (TIFs). Our findings revealed that fine-tuning *RETFound* to predict TIFs provides reasonable estimates for simpler TIFs, like vessel densities (R^2 = 0.91-0.93), but much less accurate approximations for more complex TIFs, like vessel tortuosities (R^2 = 0.25-0.43), highlighting *RETFound's* limitations to fully characterise the retinal vasculature. We also utilized genome wide association studies on *RETFound*'s latent space, the predicted TIFs, and their measured counterparts to better understand the physiological features that *RETFound* may be focusing on. We find that its latent space variables have many genetic associations, in particular with pathways involved in pigmentation, but only a small overlap with the significant genes identified from measured or predicted TIFs. Analysing the predictive value of the latent space variables, predicted and measured TIFs for clinical endpoints, we find that hybrid models that include all these features perform best for predicting blood pressure and body mass index, indicating that augmenting deep learning models with manually curated features may improve overall prediction capacity. Overall, this study highlights the synergistic potential of integrating deep learning with classical feature extraction, advancing our understanding of retinal biology and disease mechanisms, and paving the way toward improved diagnostic and prognostic tools in ophthalmology.

Introduction

Medical imaging has become essential for diagnosing diseases and providing presymptomatic risk prognoses. The use of computer-aided analysis, particularly through deep learning (DL), has made these processes not only more accurate but also scalable to large datasets. Recently, foundation models have emerged as a powerful DL approach generating initial models pre-trained on extensive image datasets using self-supervised learning approaches, such as masked autoencoders [1]. Such models can then be applied across a wide range of use cases as they provide a robust and efficient latent space that captures image variability in a low-dimensional form. The basic idea is to use the "encoder" of the model that transforms an image into latent variables (LVs), which can then be used as features for disease identification or prediction. In some cases, the training procedure also involves "fine-tuning" the weights of the encoder, effectively co-adapting the latent space to provide the most pertinent features for the end-point prediction.

While there are ample examples showing the power of using foundation models [1–3], they also have several disadvantages: First, the LVs are typically difficult to interpret. Although there exist techniques to elucidate which regions of an image most strongly contribute to variation in a given LV [4–6], it is generally not obvious what exact properties of the image in these regions are captured. Second, the number of LVs is usually relatively large, typically between 2^7 and 2^{10}, increasing the burden of correcting for multiple hypotheses testing. Finally, there is no guarantee that LVs from masked autoencoders, which are optimised to capture the information needed to reconstruct medical images, efficiently capture disease-relevant signals.

In contrast to DL approaches, clinicians and researchers have been examining images for decades to determine tangible image features (TIFs), such as the size or shape of previously identified anatomical objects, and for many of those there is well-established medical evidence showing their relevance for certain diseases. Recently, automated methods have been increasingly utilized for TIF extraction from medical images, facilitating large-scale analysis and enhancing diagnostic capabilities [7–13]. Among various imaging modalities, colour fundus images (CFIs), obtained via standard color fundus photography of the retina, have emerged as a particularly valuable tool due to their noninvasive nature and ability to predict a wide range of diseases. CFIs provide a cost-effective, in vivo overview of retinal morphology and some information on the underlying choroid and potential pathologies thereof [14], making them instrumental in monitoring ocular diseases such as diabetic retinopathy, macular degeneration, and glaucoma. Furthermore, they enable early risk prediction for systemic conditions, including cardiovascular and cerebrovascular disease, chronic kidney disease, and diabetes [15]. CFIs are also an attractive image modality for investigation because they support the extraction of vascular-based TIFs, facilitated by the availability of public datasets with ground-truth segmentations [16–19]. Of note, we recently published a study of 17 different morphological vascular phenotypes extracted from over 130k fundus images of close

to 72k UK Biobank participants. We found that a substantial portion of the variability in these measured TIFs (mTIFs)—features of longstanding interest to ophthalmologists, including median vessel diameter, diameter variability, main temporal angles, vascular density, central retinal equivalents, the number of bifurcations, and tortuosity—can be attributed to genetic factors, with SNP-based heritability estimates ranging from 5% to 25%. Moreover, we observed a large number of genetic association signals with genes and pathways that appear plausible to be involved in modulating the mTIFs, and many significant correlations with both ocular and systemic diseases or their risk factors [13]. However, mTIFs are inherently limited by human-driven feature selection, which dictates what is considered relevant in disease. Additionally, mTIFs may also suffer from poor generalizability in automated settings, as mathematical "one-size-fits-all" heuristics are often employed for feature extraction.

Here we investigate how mTIFs and DL-based features compare against and complement each other in the context of genetics and disease by leveraging *RETFound* [2], a recently developed transformer-based [20] foundation model for retinal images pre-trained using masked image reconstruction. We selected *RETFound* because it is a large-scale foundation model pretrained on extensive retinal imaging data, enabling extraction of generalizable latent features for multiple downstream tasks, including disease prediction and genetic association studies. This distinguishes it from task-specific models trained only for single diseases. To better understand the potential limitations of the information embedded in *RETFound*'s latent space representation, we examine how well it can predict retinal vascular mTIFs. This exploration is driven by the fact that, unlike medical endpoints, mTIFs are entirely determined by the image itself. We utilize genomic analysis to assess the underlying genetic architectures as they can inform upon what physiological aspect an LV represents within an image. To determine if TIFs outperform or complement foundation models in prediction tasks, we build models that include TIFs and LVs and assess their ability to predict disease. Lastly, as classical retinal features are established using heuristic approaches that may fail to generalize, we use *RETFound's* prediction of mTIFs to derive "deep" TIFs (dTIFs), and compare their genetic architectures as well as disease predictive capacities (see Fig. 1 for a graphical overview of our study). In summary, using *RETFound* and CFIs as an example, our study sheds light on (i) whether foundation models can predict tangible ("classical") image features, (ii) whether their latent variables can gain interpretability using genomic analyses, and (iii) whether mTIFs and deep features complement in regards to their genetic signals and disease prediction.

Results

Phenotypic data generation in terms of vascular TIFs and LVs

Using the methodology described in our previous work [12], we measured 17 TIFs from CFIs of 41 527 genotyped participants from the UK Biobank with complete measurements (see Suppl. Fig. 1 for inclusion-exclusion criteria). Each CFI was also characterised in terms of 1 024 LVs extracted with *RETFound* prior to any fine-tuning, using the weights of the network provided by the authors [2] (see Methods for details).

Figure 1 | Graphical overview of the study. (a) We selected a subset of genotyped participants from the UK Biobank who had colour fundus images (CFIs) taken, as well as available health outcome information and risk factors. (b) Top panel: Using a previously developed analysis pipeline to characterise the retinal vasculature [13], we extracted tangible image features (TIFs), such as vessel tortuosity or the number of bifurcations, from each CFI. We also used the retinal foundation model *RETFound* [2] as a DL approach to characterise CFIs both (middle) in terms of "deep TIFs" by fine-tuning *RETFound* to predict the 17 measured TIFs and (bottom) by extracting a set of 1 024 latent variables (LVs) using *RETFound's* pre-trained weights. (c) For downstream comparative analyses, we employed feature correlations, genetic analyses, and disease associations.

Modelling vascular TIFs with *RETFound*

By design, *RETFound's* LVs capture much of the information contained in CFIs, including the retinal vasculature. Thus, in principle, these LVs should also be informative of our vascular TIFs. In order to evaluate to what extent any individual TIF can be represented by a single LV, we computed all pairwise correlations between LVs and TIFs (shown as a bi-clustered cross-correlation matrix in Fig. 2a). For each TIF, we then picked the LV with the largest squared correlation, which corresponds to the explained variance of a Simple Linear Regression (SLR) model (R^2_{SLR}; top bars in Fig. 2b). Our analysis indicates that

individual LVs can represent a sizable portion of the variability of relatively simple vascular measures, like

the vascular densities $(max\ R^2_{SLR} = 0.36)$. However, this is not the case for vascular TIFs whose

measurement is more intricate, where we observe poor correlations between an individual LV and the

arterial central retinal equivalent $(max\ R^2_{SLR} = 0.09)$, tortuosity $(max\ R^2_{SLR} = 0.01)$, and tortuosity

artery/vein (A/V) ratios $(max\ R^2_{SLR} < 0.01)$. To assess the relationships between LVs, we conducted

correlation and principal component (PC) analyses. Only 0.003% of pairwise associations exhibited an R²

≥ 0.7, and 177 PCs were required to explain 95% of the cumulative variance (Suppl. Fig. 2).

Next, we investigated to what extent combining multiple LVs to predict TIFs can increase the explained

variance. As a first step, we trained Multiple Linear Regression (MLR) models with all LVs as features

using five-fold cross-validation. The corresponding out-of-sample estimates for the explained variance

R^2_{MLR} for each TIF (middle bars in Fig. 2b) are consistently higher than those from the best SLR model. We

observed similar patterns as for SLR, with simple TIFs (such as vascular densities) having more than 90%

of their variance explained and much less for more intricate TIFs, requiring explicit identification of vessels

and their type. Notably, LVs failed to predict A/V ratios for tortuosity $(R^2_{MLR} = 0.09)$ and central retinal

equivalents $(R^2_{MLR} = 0.06)$. In a second step we directly trained *RETFound* to predict TIFs, resulting in 17

fine-tuned *RETFound* models providing TIF estimates as their output, which we call "deep TIFs" (dTIFs).

For most of these dTIFs, the out-of-sample estimates R^2_{deep} for the explained variances (bottom bars in

Fig. 2b) are larger than those of the corresponding MLR models. Specifically, the fine-tuned *RETFound*

models outperformed the MLR models for 11 out of 17 TIFs, with a mean improvement in R^2 of 0.05 (95%

CI = [0.01, 0.08], t(16) = 2.45, p = 0.03). We observed the most substantial increases for the temporal

angles (arterial: $R^2_{deep} = 0.34, \Delta R^2 = 0.14$; venous: $R^2_{deep} = 0.61, \Delta R^2 = 0.22$) and arterial tortuosity

$(R^2_{deep} = 0.43, \Delta R^2 = 0.16)$, while the improvements for the other TIFs are marginal or even absent.

Gradient-weighted attention rollout maps qualitatively indicated that the model attended to expected

regions for all dTIFs (i.e., the parts of the vasculature beds that were used to heuristically determine the

corresponding mTIF via the automated pipeline; Fig. 2c).

Figure 2 | Correlations analysis. (a) Bi-clustered cross-correlation matrix between LVs (x-axis) and TIFs (y-axis). (b) Variance of TIFs explained by the best simple linear regression (SLR) models using the single best baseline LV (purple), multiple linear regression (MLR) models using all baseline LVs (green) and fine-tuning *RETFound* models (orange). Mean out-of-sample R^2 and their standard errors across the five folds are displayed. A: arterial, V: venous, eq: equivalent, std: standard deviation. (c) Gradient-weighted attention rollout maps for each dTIF, averaged across the cohort, and overlaid on a reference color fundus image. Minimum and maximum attention are calculated independently for each image.

Comparing the genetic association signals of TIFs and LVs

We previously showed that many of our TIFs have a sizable genetic component, with heritability estimates ranging from 5% for median vessel diameter to >25% for arterial tortuosity [12]. Our genome-wide association study (GWAS) further revealed many plausible genes and pathways for the vast majority of TIFs. These genetic association signals indicate that our TIFs are likely to capture some processes related to vascular maintenance that are partially modulated by genetic variants. As we do not understand what information is captured by LVs—other than that they can only partially represent TIFs— we sought to elucidate their genetic architectures, as these genetic signals can potentially shed light on the image-based features represented by the LVs. To this end, we performed GWAS, heritability estimation, gene and pathway analyses for all 1 024 LVs of the *RETFound* model (before any fine-tuning) and our 17 dTIFs.

At the SNP-level (Fig. 3a–c), the number of significantly associated variants was higher for mTIFs (91 SNPs at a Bonferroni-adjusted threshold of $5/17 \times 10^{-8}$) and dTIFs (88 SNPs at $5/17 \times 10^{-8}$) than for LVs (63 SNPs at $5/1 024 \times 10^{-8}$). Since LVs are not independent we also performed a GWAS on their 17 leading PCs, explaining 66% of their variance, which resulted in n = 28 SNPs with $p < 5/17 \cdot 10^{-8}$ (Suppl. Fig. 3). Notably, a subset of LVs showed the strongest associations, with an effect size of $\beta = 0.45$ and a nominal significance of $p < 10^{-300}$. rs12913832 at the *HERC2* locus was a key driver of this signal. In contrast, 38.8% of loci were shared between mTIFs and dTIFs, with the numbers of exclusive hits being 39 for mTIFs, 37 for dTIFs, and 53 for LVs. Gene-level aggregation using the *PascalX* tool [21,22] confirmed that the genetic architecture of dTIFs more closely resembles that of mTIFs than LVs (Fig. 3d), and while the strongest hits come from LVs (Fig. 3e), dTIFs had more independent gene discoveries (Fig. 3f). Consistently, the overlap of significantly associated genes was 50.4% between mTIFs and dTIFs, but only 12.4% between mTIFs and LVs (Fig. 3f). The most significant gene-level association was observed for the *HERC2/OCA2 locus* ($p = 3.78 \times 10^{-293}$ in LV 813), which also influences some TIFs, albeit less significantly ($p = 7.4 \times 10^{-21}$ in venous vascular density). Finally, heritability estimates indicated that most LVs have a modest genetic contribution ($h^2 < 0.1$ for 77.5% of LVs), with only one LV exhibiting a heritability $h^2 > 0.21$ (Fig. 3g). A similar pattern emerged at the gene level, with LVs showing a less polygenic architecture compared with mTIFs and dTIFs (Fig. 3h). Moreover, while genes linked to TIFs were generally specific to only one or two of the 17 phenotypes, those associated with LVs tended to influence multiple traits (Fig. 3i). Genome-wide inflation factors and *Linkage Disequilibrium (LD) Score Regression (LDSR)* intercepts were within acceptable ranges for all feature sets (Suppl. Fig. 4)

To deepen the interpretability of our genetic results, we performed gene set enrichment testing on the full MSigDB using *PascalX*. The results revealed a complete disjoint of significant pathways between TIFs and LVs (Fig. 3j), and high similarity between mTIFs and dTIFs. An in-depth look at all the significant pathways (Fig. 3k) demonstrated that significant genetic signals from both mTIFs and dTIFs predominantly come from pathways functionally implicated in vascular remodeling and endothelial biology, while genetic signals from LVs mostly point to pigmentation, neuronal, and ocular pathways.

TIF-level analyses (Fig. 4), comparing dTIFs and mTIFs in terms of heritabilities and genetic associations,

revealed that heritability estimates for dTIFs were generally comparable to those of the corresponding

mTIFs, or in some cases slightly higher (Fig. 4a). Notably, for the arterial temporal angle, the "deep" version

showed a substantially higher heritability (h^2=0.22, $\Delta h^2 = 0.14$). We made a consistent observation at the

gene level (Fig. 4b), where dTIFs and mTIFs showed similar numbers of associated genes and large

overlaps. Only for venous vascular density, the corresponding arterio-venous ratio, and venous tortuosity

the dTIFs had fewer associated genes than their corresponding mTIFs. Interestingly, "deep" arterial

tortuosity had 10 additional gene discoveries (10%), despite a slightly lower heritability ($\Delta h^2 = -0.02$),

whereas its venous counterpart lost 11 discoveries (-32%).

Figure 3 | Genetics architectures of mTIFs, dTIFs, and LVs. (a,d) Manhattan plots showing aggregated −log10(p-
values) across all traits within each feature set at the SNP level (a) and gene-score level (d). For each genomic
position, all p-values observed across traits within a given feature set are shown, enabling visualization of lead loci
for the entire feature set in a single panel rather than for a representative trait. Red horizontal lines denote genome-
wide Bonferroni significance thresholds adjusted for the number of traits within each feature set. (b,e) Corresponding
QQ plots of aggregated SNP-level (b) and gene-level (e) p-values for each feature set. Black lines denote null
expectation. (c,f,j) Venn diagrams showing overlap of genome-wide significant SNPs (c) and genes (f), and pathways
(j) across feature sets. A SNP, gene, or pathway is considered significant if it reaches genome-wide Bonferroni
significance adjusted for the number of traits within each feature set. (g,h) Distributions of heritability (g) and
polygenicity (h) across traits within each feature set. The y-axes denote trait proportions (bar values sum to 1 within
each feature set), allowing comparison of relative distributions independent of the number of traits. (i) Percentage of
genome-wide significant genes (from f) associated with an increasing number of traits within each feature set. (k)
Complete list of significant pathways, grouped by functional similarity. Asterisks indicate thresholds of statistical
significance (*: α=0.05, **: α=0.01, ***: α=0.001). Genome-wide Bonferroni significance thresholds adjusted for the
number of traits within each feature set.

228
229 **Figure 4 | Heritabilities and gene associations per TIF.** (a) SNP heritability estimates obtained via *LDSR* and the
230 difference between deep and measured equivalents. (b) Number of associated genes of mTIFs versus dTIFs,
231 including the count of genes exclusive to each (blue: exclusive to mTIFs, red: exclusive to dTIFs). Abbreviations: A
232 = Arterial; V = Venous; std = Standard Deviation; mTIFs = measured Tangible Image Features; dTIFs = Deep
233 Tangible Image Features.

Associating disease risk with measured and deep TIFs

Next, we investigated to what extent risk factors, ocular and general diseases, and clinical events can be explained by mTIFs or dTIFs. We used linear models that incorporated covariates alongside a single TIF as explanatory variables. We assessed the strength of associations using standardised effect sizes and statistical significance and quantified the total number of unique significant associations across diseases and TIFs to summarise the findings (Fig. 5). Additionally, we applied Fisher's exact test to assess whether the ratio of significant associations differed between mTIFs and dTIFs. Although not statistically significant, the largest difference in the number of significant diseases tended to be found with the venous temporal angle, with five diseases being significantly associated uniquely with dTIFs (OR = 2.74, p = 0.21). While also not significant, diabetes-related eye disease ("diabetes-eye") tended to display the largest number of differences between TIF types, with five more dTIFs than mTIFs (OR = 3.43, p = 0.17). When grouping the diseases into categories, we observed that ocular diseases had significantly more associations only with dTIFs than for mTIFs (OR = 1.75, p = 0.02). No such pattern was observed across risk factors (OR = 1.06, p = 0.84) or general diseases and events (OR = 0.92, p = 0.81).

Finally, we aimed to determine whether models that combine multiple image feature sets increase predictive power over models restricted to just covariates or one feature set at a time. We used regularised multilinear models with eight sets of explanatory features: (i) a baseline model including only covariates (Covar) that were included in each subsequent model; (ii) all 17 mTIFs; (iii) all 17 dTIFs; (iv) the 1 024 RETFound LVs; (v) all mTIFs and dTIFs; (vi) all mTIFs and the 1 024 RETFound LVs; (vii) all dTIFs and the 1 024 RETFound LVs; and (viii) all features. We found that: 1) models that incorporated TIFs with LVs performed best when predicting blood pressure and BMI (Fig. 6a and c); 2) when the outcome was an ocular disease, either there was no additional gain in predictive power over the baseline model, or the LVs tended to outperform the TIFs (Fig. 6b and Suppl. Fig. 5); and 3) both the TIFs and the LVs added to predictive power over the base model for most risk factors and diseases (Fig. 6a-c and Suppl. Fig. 5). Associated statistics for comparing feature predictive capacities can be found in Suppl. Tables 1-2.

Figure 5 | Disease associations with mTIFs and dTIFs. Each cell presents the standardised effects ("beta values") of a linear model that explains a risk factor, ocular or general disease incident with a single mTIF (lower left portion of a cell) or dTIF (upper right portion of a cell), while accounting for several covariates. Associations significant only for the dTIFs or mTIFs are indicated with 'D' or 'M', respectively, or a star (*) if significant for both. The bar graphs on the top and right indicate the total number of unique counts for each type of these associations across all diseases and TIFs, respectively. Abbreviations: A = Arterial; V = Venous; DBP = Diastolic Blood Pressure; SBP = Systolic Blood Pressure; Age BP = Age of High Blood Pressure Diagnosis; PR = Pulse Rate; PWASI = Pulse Wave Arterial Stiffness Index; HDL = High-Density Lipoprotein; LDL = Low-Density Lipoprotein; HbA1c = Glycated haemoglobin; N cig/day = self-reported number of cigarettes per day; N pack/yr = self-reported number of packs of cigarettes per year; BMI = Body Mass Index; DVT = Deep Vein Thrombosis.

Figure 6 | Predictive power of CFI features in assessing risk factors, diseases, and events. (a) Explained variance achieved by linear models built to predict risk factors. (b) Area under the receiver operating characteristic curve (AUC) achieved by logistic regression models built to predict ocular diseases. (c) AUC achieved by logistic regression models built to predict diseases or events. Feature sets include: (i) a baseline model including only covariates (see methods for list), (ii) all 17 mTIFs, (iii) all 17 dTIFs, (iv) 1 024 RETFound LVs, (v) all mTIF and 1 024 RETFound LVs, (vi) all dTIFs and 1 024 RETFound LVs, (vii) all mTIFs and dTIFs, (viii) all features combined. Abbreviations: Covar = Covariates; dTIF = Deep Tangible Image Feature; mTIF = Measured Tangible Image Feature; LVs = Latent Variables; DBP = Diastolic Blood Pressure; SBP = Systolic Blood Pressure; BMI = Body Mass Index; AUC = Area Under the Curve. Error bars reflect 95% confidence intervals.

Replication in two independent cohorts

We next assessed the reproducibility of our dTIFs, extracting them in two independent cohorts, the Rotterdam Studies (RS, N = 7 205) and the smaller *OphtalmoLaus* (N = 1 714) [23,24]. Specifically, we used the RETFound models we had fine-tuned to predict mTIFs using fundus images from the UKB to generate these "deep TIFs" (dTIFs) from images of the two replication cohorts. We found that the rank order of dTIF predictive capacity across the 17 TIFs was largely preserved in both RS (rho=0.69, p=0.002) and *OphtalmoLaus* (rho=0.66, p=0.004) (Suppl. Fig. 6). Genetically, we found that the heritabilities of dTIFs were highly correlated between the RS and the UK Biobank (r=0.82, p=6e-5; Suppl. Fig. 7). Furthermore, the signed effect sizes of our 88 independent SNP-wise discoveries (Fig. 3c) were overall very consistent in the RS (r=0.87, p=1e-26), and, while less strongly, also in OphtalmoLaus (r=0.70, p=5e-12; Suppl. Fig. 8). Applying the Benjamini-Hochberg procedure at FDR=0.05, we replicated 60 (71%) of the 84 available SNP candidates in the RS. Repeating this procedure at the gene level, we found that replication was even more complete, with 170 (81%) out of 210 gene candidates replicating at FDR=0.05 (Suppl. Fig. 9). We performed individual per-TIF analyses in both the RS and OphtalmoLaus (Suppl. Figs. 10-17), revealing strong candidate replications at both the SNP and the gene level across the majority of TIFs. Lastly, we also observed in the RS that most dTIFs had a larger heritability estimate than their corresponding mTIF (Suppl. Fig. 18; mean difference=0.066 (SE 0.0215), z=3.08, p=0.0021). We report the complete list of replicated SNPs and genes in Suppl. Tables 3-4.

Regarding diseases, we compared the R^2 of the models designed to predict systolic and diastolic blood pressure using different covariates (Fig. 6a) in the RS replication cohort. The R^2 values of the models from RS demonstrate a rank order correlation of greater than 0.9 with the corresponding models from the UKBB

301 (RS rho=0.93, p=8.63×10^{-4}; OphtalmoLaus rho=0.95, p=2.60×10^{-4}), indicating strong reproducibility of
302 these results (Suppl. Fig. 19).

Discussion

Breakthroughs in computer-aided analysis are revolutionizing the use of medical images for diagnosis and
risk assessment. These advances range from supporting physicians in segmenting anatomical structures
and extracting precise measurements, to making fully automated disease risk predictions using end-to-
end machine learning pipelines. DL algorithms typically hold internal "latent space" representations of the
images for this task, but their nonlinear entanglement makes them appear as "black boxes" lacking
straightforward explanations. This is problematic in the biomedical domain, where trust by the clinician is
essential [25–28]. Furthermore, interpretability is key for deciphering the underlying biological or
physiological mechanisms, which is pivotal for advancing treatment [29]. Visual interpretability methods,
such as attention maps [4–6] and counterfactual explanations [30], have been developed to address this
but often fall short of clarifying exactly how latent variables (LVs) drive predictions (see [28] for some recent
advances in tackling this issue).

Here we presented some novel methodological approaches aimed to shed some light on how one can
glean insights on what such LVs may present, and in particular what they fall short of. Specifically, our
case study investigated *RETFound,* a DL model using the vision transformer architecture and pre-trained
using masked image reconstruction that has recently been proposed as a foundation model for retinal
images. Our key innovation was to extract *RETFound*'s LVs for a large collection of colour fundus images
(CFIs) which we had previously characterised in terms of "tangible image features" (TIFs) of high clinical
interpretability [13]. This allowed us to characterise to what extent *RETFound*'s latent space captures those
TIFs or could be fine-tuned to predict them. Moreover, the fact that these images stem from genotyped
participants with extensive clinical phenotyping enabled us to directly compare the associations of LVs
with genotypes and disease states or risk factors with those of TIFs.

Our analysis revealed that, prior to fine-tuning, individual LVs do not strongly correlate with any of our
TIFs. Vascular densities and number of bifurcations have the closest LV representation, explaining close
to 40% of their variance, while it is less than 20% for the TIFs whose measurement is more intricate. Linear
combinations of multiple LVs have substantially higher predictive power for TIFs, in particular for the
vascular densities, explaining close to 90% of their variance. Strikingly, fine-tuning *RETFound* to predict
TIFs yielded marginal or no improvements compared to the LVs (c.f. [31–33] for similar negative results
on fine-tuning in LLMs), with the notable exception of the temporal angles between the main vascular
branches and, to a lesser degree, vessel tortuosity. These limited gains from fine-tuning may in part reflect
denoising effects or the learning of some relevant proxy trait. Nevertheless, these results suggest that
even the advanced transformer DL architecture of *RETFound* and its foundation model parameters trained
on more than 900 000 CFIs fail to provide a close approximation of well-established classical
ophthalmological parameters of the retinal vasculature. This is remarkable because, in contrast to clinical

endpoints, TIFs are fully determined by the images, i.e., they can be extracted through deterministic
algorithmic procedures based on conventional image processing.

Our analysis of genetic associations elucidates to what extent *RETFound*'s LVs are modulated genetically
and quantifies the overlap with the association signals of the TIFs. We argue that genetic associations
support physiological relevance because they point to a causal chain from a DNA difference (i.e. a genetic
variant at the level of a single nucleotide) over a molecular variation (such as differential gene expression)
to a (typically minute) phenotypic variability. Heritability can be seen as a global summary statistic of such
genetic effects, and our results indicate that TIFs predicted by *RETFound*, even when being a relatively
poor proxy of their measured counterparts, are nevertheless at least as heritable as the latter, supporting
our denoising or relevant proxy hypotheses. Gene-scoring analyses revealed that among *RETFound*'s
LVs the strongest association is observed with the *HERC2*/*OCA2* locus, a well-known, strong modulator
of pigmentation and eye color [34] that has previously been found as a top hit in a GWAS of LVs from self-
supervised deep phenotyping of CFIs [35]. Pathway analyses further supported this observation, with
latent variables showing predominant enrichment for pigmentation-related pathways. This indicates that
some of the baseline LVs predominantly capture information related to the pigmentation of the retina.

In contrast, dTIFs obtained by fine-tuning *RETFound* demonstrate a genetic architecture that more closely
mirrors that of mTIFs. Consistent with this shift, pathway analyses revealed that both mTIFs and dTIFs are
enriched for pathways related to vascular remodeling and endothelial biology, in line with their intended
representation of retinal vascular morphology. Notably, fine-tuning yielded 73 additional genetic
discoveries exclusive to dTIFs, an approximate 50% increase in the number of unique associations
observed for mTIFs. Moreover, these genetic signals showed strong replication in an independent cohort,
indicating that the fine-tuned model captures stable, transferable features rather than cohort-specific
effects. Furthermore, traits such as arterial temporal angles exhibited significantly higher heritability and a
greater number of gene discoveries compared to their measured counterparts, suggesting that fine-tuning
enhances the capacity of some TIFs to capture subtle yet physiologically pertinent vascular signals. We
thus hypothesize that directing *RETFound*'s focus toward TIFs augments to some extent their ability to
reflect the underlying vascular biology.

A complementary means of evaluating the pertinence of vascular features is in terms of their capacity to
facilitate disease prediction. In this study, we contrasted mTIFs and dTIFs by comparing their standardised
effects in logistic and linear models with diseases or risk factors as response variables. We found that the
number of significant diseases and risk factors was largely similar between mTIFs and dTIFs, but dTIFs
showed more significant associations with ocular diseases compared to mTIFs. This outcome may be
attributed to *RETFound*'s initial training on a diverse set of retinal images [2,36]. We hypothesise that its
fine-tuned versions, while directed to predict aspects of vascular morphology, might nevertheless retain
some of *RETFound*'s broader representation of retinal features. This may allow the model to capture subtle
patterns or relationships between vascular morphology and other retinal structures that are relevant to
ocular pathologies but not explicitly measured in traditional vascular assessments. The enhanced
performance in ocular disease prediction using our fine-tuned models suggests that DL-derived vascular

morphology metrics may provide additional informative features for ocular condition detection compared to conventional morphological measurements alone.

Using models that integrate multiple feature types (i.e., baseline, mTIFs, dTIFs, and LVs), we provide evidence in both discovery and replication cohorts that their integration enhances the predictive power for disease. Notably, despite being representative of the entire fundus image, including the vasculature, models that included both LVs from *RETFound* and TIFs (both measured and deep) perform best when predicting blood pressure or BMI, suggesting that TIFs provide complimentary information not fully captured by *RETFound*'s LVs. One potential reason is that the *RETFound* model may not capture the information provided by TIFs is due to its pre-training process through masked autoencoding [37], which is not designed to prescribe particular importance to vascular structures when reconstructing the image. As a result, some critical vasculature information may be excluded from its base weights. Indeed, the original authors of *RETFound* qualitatively report that the model fails to reconstruct much of the finer vasculature present in CFIs. Our results further support this, as we find that fine-tuning *RETFound* to predict vascular TIFs only captures a relatively small portion of their variance, suggesting that the model is unable to learn detailed vascular and vessel type specific information. Similar findings regarding hybrid models that combine manually- and DL-derived features have been observed in other domains, where manually engineered features capture domain-specific knowledge that DL models may overlook when sample sizes are small [38] or when training and test set distributions differ [39,40]. These results emphasize the importance of leveraging both foundation- and expert-derived features to optimize predictive performance. Moreover, future work should examine whether foundation models trained with objectives that explicitly emphasize specific retinal structures better capture and predict these features of interest.

In summary, our findings provide key insights into the interplay between DL-derived features and classical retinal image features in the context of genetics and disease prediction. While *RETFound* can predict certain TIFs, such as vessel densities, with reasonable accuracy, it provides poor estimates of more intricate features, like measures related to artery-to-vein ratios. Despite these limitations, dTIFs exhibit heritability comparable to or greater than their measured counterparts, with a notably large increase in heritability found with temporal angles of the arteries, suggesting that dTIFs may enhance physiologically relevant signal extraction. Furthermore, the genetic and clinical associations of mTIFs and dTIFs show both overlap and complementarity, reinforcing the idea that DL-derived features do not simply replicate classical measurements but may provide additional, distinct insights. Most notably, integrating LVs with TIFs leads to superior predictive performance of cardiovascular risk factors such as SBP and DBP compared to models using either feature set alone, highlighting the potential of hybrid approaches in disease modeling. Overall, this work highlights the synergy between DL and classical feature extraction for advancing our understanding of retinal biology and disease mechanisms, potentially leading to improved diagnostic and prognostic tools in ophthalmology.

Methods

Data and quality control

The UK Biobank is a population-based cohort of ~488k participants with rich, longitudinal phenotypic data, including medical history, and a median 10-year follow-up [41]. Standard retinal 45° CFIs were captured via standard color fundus photography using a Topcon 3D-OCT 1000 Mark II, resulting in a total of 173 814 CFIs from 84 813 individuals. For this study we used 41 527 of these images from the right eye, for which we were able to measure all of our 17 TIFs using our previously described analysis pipeline [13]. 96% of these images passed the quality control threshold we had used in our previous work [13] (see Suppl. Fig. 20). This effectively removed CFIs for which our algorithm failed to identify the optic disk or enough major blood vessels to measure the temporal angles. Since this happens more often for CFIs from older and diseased participants, the cohort of the present study appears to be more "healthy" with respect to its baseline risk factors (see Suppl. Table 5 for a comparison of baseline characteristics to our previous study). Since some of the LVs produced from *RETFound* encoded information specific to eye laterality (left vs. right), we opted to analyze a single eye rather than averaging across both eyes. Genotyping was performed on Axiom arrays for a total of 805 426 markers, from which ~96 million genotypes were imputed. We used the subset of 15 599 830 SNPs that had been assigned an rsID.

Measured phenotypes

The 17 mTIFs comprise tortuosity, temporal angles, vessel diameters and diameter variability, vascular density, and the number of bifurcations. With the exception of the number of bifurcations, all of these measurements were performed independently for arteries and veins. To explore asymmetries between arteries and veins, the arterio-venous ratio between diameters, tortuosities and vascular densities were also included. All mTIFs were derived using an automated image analysis pipeline described in our recent publication [13], which follows a three-step procedure: (1) deep learning–based segmentation of retinal vessels from raw color fundus images, (2) reconstruction and classification of vessel centerlines into arteries and veins, and (3) quantitative extraction of morphometric features yielding the final mTIFs.

Deep learning model

We utilised *RETFound [2]*, a novel foundation model that leverages Google's vision transformer large patch 16 architecture [42]. Briefly, *RETFound* uses a masked autoencoder (MAE) approach [37] to optimise the model towards reconstructing retinal images. For building *RETFound*, 75% of the image patches were masked and the model was trained to reconstruct the missing patches, thereby learning effective representations of the images provided as training data. We used the encoder portion of this model to extract LVs and make predictions.

Extracting Latent Variables

To extract latent variables (LVs), we utilized the encoder portion of the pre-trained *RETFound* foundation model. Based on the Vision Transformer Large (ViT-L/16) architecture, the model consists of 24 Transformer blocks with a native embedding dimension of 1 024. Input images were resized to 224×224 pixels and normalized using ImageNet-derived means and standard deviations. During extraction, unmasked 16×16 patches were projected into feature vectors and processed through the encoder's multi-

449 headed self-attention layers and multilayer perceptrons. The resulting 1 024-dimensional feature vectors
450 were then utilized as LVs for all subsequent downstream genetic and phenotypic analyses.

Predicting tangible image features

For predicting TIFs, we used the encoder portion of *RETFound* and added a multi-layer perceptron to the
head of the model. We utilised the base hyperparameters established in the original *RETFound* manuscript
[2] and trained the model over 50 epochs, with a "warm-up" in the 10 first epochs (with the learning rate r
monotonically increasing from 0 to 5×10^{-4}), followed by a cosine annealing schedule (where r decreases
from 5×10^{-4} to 1×10^{-6}). We set the batch size to 16, the dropout rate to 0.2, and the image size to
224×224 pixels. After each epoch, we evaluated the model on a validation dataset and then kept the model
that had the lowest mean absolute error across all epochs. For each of the 17 TIFs, we trained separate
models 5-fold cross-validation with an effective split of 60% for training, 20% for validation, and 20% for
testing in each fold for both RETFound fine-tuning and the multiple linear regression baselines, and report
the mean test-set r² across the five folds.

Covariate corrections

463 Genetic analysis and disease prediction models were corrected for a set of covariates known to confound
464 phenotypic variability in general, or in the eye specifically. They comprise sex, age, age-squared, sex-by-
465 age, sex-by-age-squared, spherical equivalent (defined as $spherical\ power + \frac{cylindrical\ power}{2}$), spherical
466 equivalent-squared, imaging instance, assessment centre, genotype measurement batch, and the first 20
467 genomic PCs (see Suppl. Fig. 21).

Attention maps

To characterise the model's focus during inference, we generated attribution maps for all 17 fine-tuned
RETFound models using a modified gradient-weighted attention rollout [43]. For each input image, we
registered hooks on all transformer blocks to retrieve the post-softmax attention matrices and their
corresponding gradients with respect to the model output. In each block, and for each attention head, we
computed a gradient-weighted attention tensor by taking the element-wise product between the attention
weights and their gradients, followed by a ReLU to retain only positive signal. These tensors were then
averaged across heads within each block, after which we applied rollout [44]: at every layer we added an
identity matrix to account for residual connections, and row-normalized the resulting matrix before
multiplying it into the cumulative rollout matrix. This produced a single NxN relevance matrix per image,
where N=197 denotes the number of tokens. We focused on the "CLS→patch" attention by extracting the
row corresponding to the class token [45] , yielding a 14×14 spatial attribution map once reshaped. For
each model, we computed attribution maps for all images in the cohort, averaged them to visualise
population-level patterns of model focus, and upsampled the resulting matrices to the original image
resolution using bicubic interpolation for overlay on a reference colour fundus image.

Genomic analyses

For GWAS, raw phenotypes were transformed using rank-based inverse normal transformation. GWAS
were conducted using *BGENIE [46]*. Only SNPs with minor allele frequency larger than 0.01 and

imputation quality larger than 0.8 were kept for downstream analyses. SNP-specific heritabilities and
genetic correlations between traits were calculated using *LDSR [47]* which fits an ordinary least squares
regression of SNP LD scores against their average chi-squared statistics. *LDSR* considered only common
SNPs (MAF > 0.01) with an imputation quality > 0.9. Genome-wide significant SNPs were LD-pruned using
LDlinkR::SNPclip (population = GBR; $r^2 < 0.1$) to retain an approximately independent set of variants for
reporting. Gene and pathway scores were computed using *PascalX* [21,22,48]. Both protein-coding genes
and lincRNAs were scored using the approximate "saddle" method, incorporating all SNPs within a 50 kb
window surrounding each gene. All pathways from MSigDB v7.2 [49–51] were scored using *PascalX's*
ranking mode, merging and rescoring co-occurring genes separated by less than 100 kb. Among nested
subgroups of pathways, we retained only the element with the highest significance. *PascalX* utilises LD
structure for accurate computation of gene scores, which in these analyses was derived from the UK10K
(hg19) reference panel [52]. For annotating genes in the Manhattan plots, the top ten genes for each
feature set (i.e., mTIFs, dTIFs, or LVs) were selected within a 1 000 SNP window.

Disease associations

Three distinct groups of disease traits were included in this work. First, a set of general systemic risk
factors, second, a set of ocular diseases, and third, a set of general diseases and events (see Suppl. Fig.
22). All disease data were collected from the UK Biobank (see Ref. [13] for how we mapped the data field
identifiers to each disease included in this study).

To extract beta coefficients, in the case of the continuous risk factors, linear regression was used to
estimate standardised effects. For both ocular and general diseases, time-to-event data were binarised,
and logistic regression was applied.

When examining the predictive capacity of different feature sets, linear regression was used for continuous
risk factors and logistic regression was used for disease prediction. To assess model generalization and
obtain robust estimates of predictive performance, five-fold cross-validation was employed, with an 80%
training and 20% test split in each fold. Models were trained exclusively on the 80% training portion within
each fold, and performance was evaluated on the held-out 20% test set. Reported results correspond to
averages across all test folds, such that each subject appears exactly once in a test set, reducing
dependence on any single data partition. Both linear and logistic regression models were regularized using
elastic net penalties, with optimal values for alpha and the L1 ratio determined by grid search within each
fold. We utilized the coefficient of determination (R^2) as the performance metric for continuous risk factors
and area under the receiver operating characteristic curve (AUC) as the performance metric for diseases.
To ensure that the predictive capacity of retinal features was not confounded by known risk factors present
in the covariates, we created a matched cohort for each binary disease analysis. We downsampled the
larger control group by matching each case with a unique control based on the most similar covariate
profile, a process performed using optimal bipartite matching (the Hungarian algorithm), ensuring that no
control was used more than once. This matching procedure, which resulted in a balanced sample size for
cases and controls, allows for a more direct assessment of the independent contribution of the image-
derived features. To prevent poor feature-to-sample-size ratios, diseases and events were required to
have at least 1 000 samples (cases and controls combined) to be analyzed.

For all regression analyses, traits were corrected for the covariates listed above. To determine the
significance of regression coefficients and prediction power, p-values were computed, and the Benjamini-
Hochberg procedure was applied using a false discovery rate of 0.05.

Replication in the Rotterdam Study and OphtalmoLaus

Replication was performed in two cohorts, (1) the Rotterdam Study (RS) [53], and (2) OphtalmoLaus [23],
a sub-cohort of CoLaus [54] and a member of the Global RETFound initiative [55]. Replication analyses
closely followed our previously described replication pipeline [13] including its Supplementary Methods,
with only the following minor changes. In the RS, (1) the number of subjects in the four sub-cohorts
available was 5 227 (RS-I), 2 256 (RS-II), 3 166 (RS-III), and 2 229 (RS-IV); (2) the included covariates
were age, sex, spherical equivalent, imaging device, image quality (AI score), genomic PCs 1–
10, squared terms and interactions (age squared, spherical equivalent squared, age times sex).

For the Rotterdam CFIs, the original dataset comprised 262 965 color fundus images acquired from
multiple participants using several different imaging devices. To ensure data quality and consistency, we
applied a two-stage filtering pipeline. In the first stage, we retained only macula-centered images, excluded
those with missing quality scores, applied an automated image-quality threshold of ≥0.5, and only retained
right-eye images for consistency, reducing the dataset from 262 965 to 24 941 images (90.5% reduction).
In the second stage, a single highest-quality image was selected per participant using a hierarchical
prioritization strategy. Images from the TOPCON 3D OCT-2000 FA Plus device were preferred when
available, and among these, the image with the highest quality score was selected. The same criteria were
applied across devices when the preferred model was unavailable. The final dataset comprised 8,755
images, with the following device distribution: TOPCON 3D OCT-2000 FA Plus (n=7 168; 81.9%),
TOPCON 3D OCT-1000 MARK II (n=1 030; 11.8%), TOPCON 3D OCT-1000 (n=316; 3.6%), TOPCON
DRI OCT Triton Plus (n=213; 2.4%), TOPCON TRC-50VT (n=23; 0.3%), and TOPCON DRI OCT Triton
(n=5; 0.1%). Regarding OpthalmoLaus, 6 503 images from 2 276 subjects were available, of which 4 890
images from 1 715 subjects passed QC and were used for analyses.

For feature extraction, we used our 17 *RETFound* models that were fine-tuned in the UK Biobank to
independently extract dTIFs in each of the two replication cohorts. For genetic replication, candidates at
the SNP-level were defined as independent genome-wide-significant loci passing Bonferroni correction
adjusted for both the number of independent SNPs and the number of traits (5e-8/17). Candidates at the
gene level were defined as PascalX gene scores passing the nominal Bonferroni-corrected threshold
adjusted for both the number of tested genes (25819) and the number of traits (17), p_thres=1.14e-7.

Data availability

557 GWAS summary statistics, gene and pathway scores, and all data underlying figures will be made
558 available on Zenodo upon publication. GWAS summary statistics will further be made available on the
559 GWAS Catalog. Raw UK Biobank data are protected and not open access; however, they can be obtained
560 upon project creation and acceptance via their platform (https://www.ukbiobank.ac.uk/). mTIFs [13] are
561 already available via this channel; dTIFs and baseline LVs will be added to the platform upon publication.

Code availability

Code is going to be made available on Zenodo upon publication.

Author information

VascX Consortium:

Michael Beyeler [1,2]; Ciara Bergin [3]; Sven Bergmann [1,2,4]; Dennis Bontempi [1,2]; Sacha Bors [1,2]; Leah Böttger [1,2]; Bogdan Draganski [5,6,7]; Adham Elwakil [3,8]; Györgyi V. Hamvas [9,10]; Janna Hastings [11,12]; Ilaria Iuliani [1,2]; Caroline C.W. Klaver [13,14,15,16]; Ihor Kuras [6,7]; Bart Liefers [13,14]; Ilenia Meloni [3,8]; Sofia Ortin Vela [1,2]; David Presby [1,2]; Ian Quintas [1,2]; José Vargas Quiros [13,14,]; Marc Schindewolf [9,10]; Reinier O. Schlingemann [3,17]; Mattia Tomasoni [3,8]; Olga Trofimova [1,2].

[1] *Department of Computational Biology, University of Lausanne, Lausanne, Switzerland*

[2] *Swiss Institute of Bioinformatics, Lausanne, Switzerland*

[3] *Department of Ophthalmology, University of Lausanne, Fondation Asile des Aveugles, Jules Gonin Eye Hospital, Lausanne, Switzerland.*

[4] *Department of Integrative Biomedical Sciences, University of Cape Town, Cape Town, South Africa*

[5] *Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Germany*

[6] *Insel University Hospital Bern, Switzerland*

[7] *Department of Clinical Neuroscience, Lausanne University Hospital and University of Lausanne, Switzerland*

[8] *Platform for Research in Ocular Imaging, Fondation Asile des Aveugles, Jules Gonin Eye Hospital, Lausanne, Switzerland.*

[9] *Inselspital, Bern University Hospital, University of Bern, Switzerland.*

[10] *Department for BioMedical Research, Bern University Hospital, University of Bern, Switzerland.*

[11] *Institute for Implementation Science in Health Care, Faculty of Medicine, University of Zurich, Zürich, Switzerland*

[12] *School of Medicine, University of St Gallen, St. Gallen, Switzerland*

[13] *Department of Ophthalmology, Erasmus University Medical Center, Rotterdam, The Netherlands.*

[14] *Department of Epidemiology, Erasmus University Medical Center, Rotterdam, The Netherlands.*

[15] *Department of Ophthalmology, Radboud University Medical Center, Nijmegen, the Netherlands.*

[16] *Institute of Molecular and Clinical Ophthalmology, University of Basel, Switzerland.*

[17] *University Medical Centres, Amsterdam, The Netherlands.*

References

1. Azad B, Azad R, Eskandari S, Bozorgpour A, Kazerouni A, Rekik I, et al. Foundational Models in Medical Imaging: A Comprehensive Survey and Future Vision. 2023. Available: http://arxiv.org/abs/2310.18689

2. Zhou Y, Chia MA, Wagner SK, Ayhan MS, Williamson DJ, Struyven RR, et al. A foundation model for generalizable disease detection from retinal images. Nature. 2023;622: 156–163.

3. Parres D, Paredes R. Fine-Tuning Vision Encoder–Decoder Transformers for Handwriting Text Recognition on Historical Documents. Document Analysis and Recognition - ICDAR 2023. 2023; 253–268.

4. Zhou B, Khosla A, Lapedriza A, Oliva A, Torralba A. Learning deep features for discriminative localization. 2016 IEEE Conference on Computer Vision and Pattern Recognition (CVPR). IEEE; 2016. doi:10.1109/cvpr.2016.319

5. Selvaraju RR, Cogswell M, Das A, Vedantam R, Parikh D, Batra D. Grad-CAM: Visual Explanations from Deep Networks via Gradient-based Localization. 2016 [cited 17 Dec 2024]. doi:10.1007/s11263-019-01228-7

6. Shrikumar A, Greenside P, Kundaje A. Learning important features through propagating activation differences. Precup D, Teh YW, editors. arXiv [cs.CV]. 06--11 Aug 2017. pp. 3145–3153. Available: https://proceedings.mlr.press/v70/shrikumar17a.html

7. Fhima J, Van Eijgen J, Billen Moulin-Romsée M-I, Brackenier H, Kulenovic H, Debeuf V, et al. LUNet: deep learning for the segmentation of arterioles and venules in high resolution fundus images. Physiol Meas. 2024;45. doi:10.1088/1361-6579/ad3d28

8. Quiros JV, Liefers B, van Garderen K, Vermeulen J, Center ER, Consortium S, et al. VascX Models: Model Ensembles for Retinal Vascular Analysis from Color Fundus Images. 2024. Available: http://arxiv.org/abs/2409.16016

9. Tomasoni M, Beyeler MJ, Vela SO, Mounier N, Porcu E, Corre T, et al. Genome-wide Association Studies of Retinal Vessel Tortuosity Identify Numerous Novel Loci Revealing Genes and Pathways Associated With Ocular and Cardiometabolic Diseases. Ophthalmology Science. 2023;3: 100288.

10. Zhou Y, Wagner SK, Chia MA, Zhao A, Woodward-Court P, Xu M, et al. AutoMorph: Automated Retinal Vascular Morphology Quantification Via a Deep Learning Pipeline. Trans Vis Sci Tech. 2022;11: 12–12.

11. Van Eijgen J, Fhima J, Billen Moulin-Romsée M-I, Behar JA, Christinaki E, Stalmans I. Leuven-Haifa High-Resolution Fundus Image Dataset for Retinal Blood Vessel Segmentation and Glaucoma Diagnosis. Sci Data. 2024;11: 257.

12. Ortin Vela S, Beyeler MJ, Trofimova O, Tomasoni M. Phenotypic and Genetic Characteristics of Retinal Vascular Parameters and their Association with Diseases. medRxiv. 2023. Available: https://www.medrxiv.org/content/10.1101/2023.07.07.23292368.abstract

13. Ortín Vela S, Beyeler MJ, Trofimova O, Iuliani I, Vargas Quiros JD, de Vries VA, et al. Phenotypic and genetic characteristics of retinal vascular parameters and their association with diseases. Nat Commun. 2024;15: 9593.

14. Rahimi M, Rossi A, Son T, Adejumo T, Dadzie AK, Heiferman MJ, et al. High dynamic range widefield fundus photography with transcranial illumination: a contrast agent-free method for non-mydriatic choroidal imaging. Biomed Opt Express. 2025;16: 84–96.

15. Ho H, Cheung CY, Sabanayagam C, Yip W, Ikram MK, Ong PG, et al. Retinopathy Signs Improved Prediction and Reclassification of Cardiovascular Disease Risk in Diabetes: A prospective cohort study. Sci Rep. 2017;7: 41492.

16. Staal J, Abràmoff MD, Niemeijer M, Viergever MA, van Ginneken B. Ridge-based vessel

segmentation in color images of the retina. IEEE Trans Med Imaging. 2004;23: 501–509.

17.   Fraz MM, Remagnino P, Hoppe A, Uyyanonvara B, Rudnicka AR, Owen CG, et al. An ensemble
classification-based approach applied to retinal blood vessel segmentation. IEEE Trans Biomed
Eng. 2012;59: 2538–2548.

18.   Budai A, Bock R, Maier A, Hornegger J, Michelson G. Robust vessel segmentation in fundus
images. Int J Biomed Imaging. 2013;2013: 154860.

19.   Hoover A, Kouznetsova V, Goldbaum M. Locating blood vessels in retinal images by piecewise
threshold probing of a matched filter response. IEEE Trans Med Imaging. 2000;19: 203–210.

20.   Advances in medical image analysis with vision Transformers: A comprehensive review. Medical
Image Analysis. 2024;91: 103000.

21.   Krefl D, Bergmann S. Cross-GWAS coherence test at the gene and pathway level. PLoS Comput
Biol. 2022;18: e1010517.

22.   Krefl D, Brandulas Cammarata A, Bergmann S. PascalX: a Python library for GWAS gene and
pathway enrichment tests. Bioinformatics. 2023;39. doi:10.1093/bioinformatics/btad296

23.   Meloni I, Elwakil A, Aurélia G, Flavie T, Navarro A, Bagatella L, et al. Cohort profile: OphtalmoLaus,
an extension of the CoLaus|PsyCoLaus cohort to investigate the relationships between ocular,
cardiovascular, and cognitive parameters. Research Square. 2025. doi:10.21203/rs.3.rs-7660406/v1

24.   Bergin C, Mantel I, Schlingemann RO, Tomasoni M, Wolfensberger TJ. SOIN – MI Data Lab:
Personalized Ophthalmology Through Collaborative Data Collection and Dynamic Patient Consent.
Challenges of Trustable AI and Added-Value on Health. IOS Press; 2022. pp. 281–282.

25.   Andrews B, Chang J-B, Collinson L, Li D, Lundberg E, Mahamid J, et al. Imaging cell biology.
Nature Cell Biology. 2022;24: 1180–1185.

26.   Rajpurkar P, Chen E, Banerjee O, Topol EJ. AI in health and medicine. Nat Med. 2022;28: 31–38.

27.   DeGrave AJ, Cai ZR, Janizek J, Daneshjou R, Lee S-I. Auditing the inference processes of medical-
image classifiers by leveraging generative AI and the expertise of physicians. Nature Biomedical
Engineering. 2023; 1–13.

28.   Rotem O, Schwartz T, Maor R, Tauber Y, Shapiro MT, Meseguer M, et al. Visual interpretability of
image-based classification models by generative latent space disentanglement applied to in vitro
fertilization. Nature Communications. 2024;15: 1–19.

29.   Rotem O, Zaritsky A. Visual interpretability of bioimaging deep learning models. Nat Methods.
2024;21: 1394–1397.

30.   Lang O, Gandelsman Y, Yarom M, Wald Y, Elidan G, Hassidim A, et al. Explaining in Style: Training
a GAN to explain a classifier in StyleSpace. arXiv [cs.CV]. 2021. Available:
http://arxiv.org/abs/2104.13369

31.   Fine-Tuning or Fine-Failing? Debunking Performance Myths in Large Language Models. [cited 18
Dec 2024]. Available: https://arxiv.org/html/2406.11201v1

32.   Li X, Chan S, Zhu X, Pei Y, Ma Z, Liu X, et al. Are ChatGPT and GPT-4 General-Purpose Solvers
for Financial Text Analytics? A Study on Several Typical Tasks. 2023. Available:
http://arxiv.org/abs/2305.05862

33.   Uppaal R, Hu J, Li Y. Is Fine-tuning Needed? Pre-trained Language Models Are Near Perfect for
Out-of-Domain Detection. 2023. Available: http://arxiv.org/abs/2305.13282

34.   Eiberg H, Troelsen J, Nielsen M, Mikkelsen A, Mengel-From J, Kjaer KW, et al. Blue eye color in
humans may be caused by a perfectly associated founder mutation in a regulatory element located
within the HERC2 gene inhibiting OCA2 expression. Human Genetics. 2008;123: 177–187.

35.   Xie Z, Zhang T, Kim S, Lu J, Zhang W, Lin C-H, et al. iGWAS: Image-based genome-wide

association of self-supervised deep phenotyping of retina fundus images. PLOS Genetics. 2024;20: e1011273.

36. Fu Y, Ma L, Wan S, Ge S, Yang Z. A novel clinical artificial intelligence model for disease detection via retinal imaging. Innovation (Camb). 2024;5: 100575.

37. He K, Chen X, Xie S, Li Y, Dollar P, Girshick R. Masked Autoencoders Are Scalable Vision Learners. 2022 IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR). IEEE; 2022. doi:10.1109/cvpr52688.2022.01553

38. Mavaie P, Holder L, Skinner MK. Hybrid deep learning approach to improve classification of low-volume high-dimensional data. BMC Bioinformatics. 2023;24: 1–20.

39. Lin W, Hasenstab K, Moura Cunha G, Schwartzman A. Comparison of handcrafted features and convolutional neural networks for liver MR image adequacy assessment. Scientific Reports. 2020;10: 1–11.

40. Bento N, Rebelo J, Barandas M, Carreiro AV, Campagner A, Cabitza F, et al. Comparing Handcrafted Features and Deep Neural Representations for Domain Generalization in Human Activity Recognition. Sensors (Basel, Switzerland). 2022;22: 7324.

41. Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, et al. The UK Biobank resource with deep phenotyping and genomic data. Nature. 2018;562: 203–209.

42. Dosovitskiy A, Beyer L, Kolesnikov A, Weissenborn D, Zhai X, Unterthiner T, et al. An Image is Worth 16x16 Words: Transformers for Image Recognition at Scale. 2020. Available: http://arxiv.org/abs/2010.11929

43. Chefer H, Gur S, Wolf L. Transformer Interpretability Beyond Attention Visualization. arXiv [cs.CV]. 2020. doi:10.48550/ARXIV.2012.09838

44. Abnar S, Zuidema W. Quantifying attention flow in transformers. arXiv [cs.LG]. 2020. doi:10.48550/ARXIV.2005.00928

45. Caron M, Touvron H, Misra I, Jégou H, Mairal J, Bojanowski P, et al. Emerging properties in self-supervised vision transformers. arXiv [cs.CV]. 2021. doi:10.48550/ARXIV.2104.14294

46. Band G, Marchini J. BGEN: a binary file format for imputed genotype and haplotype data. bioRxiv. 2018. p. 308296. doi:10.1101/308296

47. Bulik-Sullivan B, Loh PR, Finucane HK, Ripke S, Yang J, Patterson N, et al. LD score regression distinguishes confounding from polygenicity in genome-wide association studies. Nat Genet. 2015;47: 291–295.

48. Lamparter D, Marbach D, Rueedi R, Kutalik Z, Bergmann S. Fast and Rigorous Computation of Gene and Pathway Scores from SNP-Based Summary Statistics. 2016. doi:10.1371/journal.pcbi.1004714

49. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A. 2005;102: 15545–15550.

50. Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P. The Molecular Signatures Database (MSigDB) hallmark gene set collection. Cell Syst. 2015;1: 417–425.

51. Castanza AS, Recla JM, Eby D, Thorvaldsdóttir H, Bult CJ, Mesirov JP. Extending support for mouse data in the Molecular Signatures Database (MSigDB). Nature Methods. 2023;20: 1619–1620.

52. UK10K Consortium, Walter K, Min JL, Huang J, Crooks L, Memari Y, et al. The UK10K project identifies rare variants in health and disease. Nature. 2015;526: 82–90.

53. Ikram MA, Brusselle GGO, Murad SD, van Duijn CM, Franco OH, Goedegebure A, et al. The Rotterdam Study: 2018 update on objectives, design and main results. Eur J Epidemiol. 2017;32:

807–850.

54. Firmann M, Mayor V, Vidal PM, Bochud M, Pécoud A, Hayoz D, et al. The CoLaus study: a population-based study to investigate the epidemiology and genetic determinants of cardiovascular risk factors and metabolic syndrome. BMC Cardiovasc Disord. 2008;8: 6.

55. Tham YC, Goh JHL, Nderitu P, Zhou Y, Ran AR, Srinivasan S, et al. Building the world's first truly global medical foundation model. Nature Medicine. 2025;31: 3580–3585.

Abnar, Samira, and Willem Zuidema. 2020. "Quantifying Attention Flow in Transformers." In arXiv [cs.LG]. https://doi.org/10.48550/ARXIV.2005.00928.

"Advances in Medical Image Analysis with Vision Transformers: A Comprehensive Review." 2024. Medical Image Analysis 91 (January): 103000.

Andrews, Brenda, Jae-Byum Chang, Lucy Collinson, et al. 2022. "Imaging Cell Biology." Nature Cell Biology 24 (8): 1180–1185.

Azad, Bobby, Reza Azad, Sania Eskandari, et al. 2023. "Foundational Models in Medical Imaging: A Comprehensive Survey and Future Vision." October 28. http://arxiv.org/abs/2310.18689.

Band, Gavin, and Jonathan Marchini. 2018. "BGEN: A Binary File Format for Imputed Genotype and Haplotype Data." In bioRxiv. April 29. https://doi.org/10.1101/308296.

Bento, Nuno, Joana Rebelo, Marília Barandas, et al. 2022. "Comparing Handcrafted Features and Deep Neural Representations for Domain Generalization in Human Activity Recognition." Sensors (Basel, Switzerland) 22 (19): 7324.

Bergin, Ciara, Irmela Mantel, Reinier O. Schlingemann, Mattia Tomasoni, and Thomas J. Wolfensberger. 2022. "SOIN – MI Data Lab: Personalized Ophthalmology Through Collaborative Data Collection and Dynamic Patient Consent." In Challenges of Trustable AI and Added-Value on Health. IOS Press.

Budai, A., R. Bock, A. Maier, J. Hornegger, and G. Michelson. 2013. "Robust Vessel Segmentation in Fundus Images." International Journal of Biomedical Imaging 2013 (December): 154860.

Bulik-Sullivan, Brendan, Po Ru Loh, Hilary K. Finucane, et al. 2015. "LD Score Regression Distinguishes Confounding from Polygenicity in Genome-Wide Association Studies." Nature Genetics 47 (3): 291–295.

Bycroft, Clare, Colin Freeman, Desislava Petkova, et al. 2018. "The UK Biobank Resource with Deep Phenotyping and Genomic Data." Nature 562 (7726): 203–209.

Caron, Mathilde, Hugo Touvron, Ishan Misra, et al. 2021. "Emerging Properties in Self-Supervised Vision Transformers." In arXiv [cs.CV]. https://doi.org/10.48550/ARXIV.2104.14294.

Castanza, Anthony S., Jill M. Recla, David Eby, Helga Thorvaldsdóttir, Carol J. Bult, and Jill P. Mesirov. 2023. "Extending Support for Mouse Data in the Molecular Signatures Database (MSigDB)." Nature Methods 20 (11): 1619–1620.

Chefer, Hila, Shir Gur, and Lior Wolf. 2020. "Transformer Interpretability Beyond Attention Visualization." In arXiv [cs.CV]. https://doi.org/10.48550/ARXIV.2012.09838.

DeGrave, Alex J., Zhuo Ran Cai, Jose Janizek, Roxana Daneshjou, and Su-In Lee. 2023. "Auditing the Inference Processes of Medical-Image Classifiers by Leveraging Generative AI and the Expertise of Physicians." Nature Biomedical Engineering, December 28, 1–13.

Dosovitskiy, Alexey, Lucas Beyer, Alexander Kolesnikov, et al. 2020. "An Image Is Worth 16x16 Words: Transformers for Image Recognition at Scale." October 22. http://arxiv.org/abs/2010.11929.

Fhima, Jonathan, Jan Van Eijgen, Marie-Isaline Billen Moulin-Romsée, et al. 2024. "LUNet: Deep Learning for the Segmentation of Arterioles and Venules in High Resolution Fundus Images." Physiological Measurement 45 (5). https://doi.org/10.1088/1361-6579/ad3d28.

"Fine-Tuning or Fine-Failing? Debunking Performance Myths in Large Language Models." n.d. Accessed December 18, 2024. https://arxiv.org/html/2406.11201v1.

Firmann, Mathieu, Vladimir Mayor, Pedro Marques Vidal, et al. 2008. "The CoLaus Study: A Population-Based Study to Investigate the Epidemiology and Genetic Determinants of Cardiovascular Risk Factors and Metabolic Syndrome." BMC Cardiovascular Disorders 8 (1): 6.

Fraz, Muhammad Moazam, Paolo Remagnino, Andreas Hoppe, et al. 2012. "An Ensemble Classification-Based Approach Applied to Retinal Blood Vessel Segmentation." IEEE Transactions on Bio-Medical Engineering 59 (9): 2538–2548.

Fu, Yidian, Liang Ma, Sheng Wan, Shengfang Ge, and Zhi Yang. 2024. "A Novel Clinical Artificial Intelligence Model for Disease Detection via Retinal Imaging." Innovation (Cambridge (Mass.)) 5 (2): 100575.

He, Kaiming, Xinlei Chen, Saining Xie, Yanghao Li, Piotr Dollar, and Ross Girshick. 2022. "Masked Autoencoders Are Scalable Vision Learners." Paper presented 2022 IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR), 2022/6/18-2022/6/24, New Orleans, LA, USA. 2022 IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR), June. https://doi.org/10.1109/cvpr52688.2022.01553.

Ho, Henrietta, Carol Y. Cheung, Charumathi Sabanayagam, et al. 2017. "Retinopathy Signs Improved Prediction and Reclassification of Cardiovascular Disease Risk in Diabetes: A Prospective Cohort Study." Scientific Reports 7 (February): 41492.

Hoover, A., V. Kouznetsova, and M. Goldbaum. 2000. "Locating Blood Vessels in Retinal Images by Piecewise Threshold Probing of a Matched Filter Response." IEEE Transactions on Medical Imaging 19 (3): 203–210.

Ikram, M. Arfan, Guy G. O. Brusselle, Sarwa Darwish Murad, et al. 2017. "The Rotterdam Study: 2018 Update on Objectives, Design and Main Results." European Journal of Epidemiology 32 (9): 807–850.

Krefl, Daniel, and Sven Bergmann. 2022. "Cross-GWAS Coherence Test at the Gene and Pathway Level." PLoS Computational Biology 18 (9): e1010517.

Krefl, Daniel, Alessandro Brandulas Cammarata, and Sven Bergmann. 2023. "PascalX: A Python Library for GWAS Gene and Pathway Enrichment Tests." Bioinformatics 39 (5). https://doi.org/10.1093/bioinformatics/btad296.

Lamparter, David, Daniel Marbach, Rico Rueedi, Zoltán Kutalik, and Sven Bergmann. 2016. Fast and Rigorous Computation of Gene and Pathway Scores from SNP-Based Summary Statistics. https://doi.org/10.1371/journal.pcbi.1004714.

Lang, Oran, Yossi Gandelsman, Michal Yarom, et al. 2021. "Explaining in Style: Training a GAN to Explain a Classifier in StyleSpace." In arXiv [cs.CV]. April 27. arXiv. http://arxiv.org/abs/2104.13369.

Liberzon, Arthur, Chet Birger, Helga Thorvaldsdóttir, Mahmoud Ghandi, Jill P. Mesirov, and Pablo Tamayo. 2015. "The Molecular Signatures Database (MSigDB) Hallmark Gene Set Collection." Cell Systems 1 (6): 417–425.

Lin, Wenyi, Kyle Hasenstab, Guilherme Moura Cunha, and Armin Schwartzman. 2020. "Comparison of Handcrafted Features and Convolutional Neural Networks for Liver MR Image Adequacy Assessment." Scientific Reports 10 (1): 1–11.

Li, Xianzhi, Samuel Chan, Xiaodan Zhu, et al. 2023. "Are ChatGPT and GPT-4 General-Purpose Solvers for Financial Text Analytics? A Study on Several Typical Tasks." May 10. http://arxiv.org/abs/2305.05862.

Mavaie, Pegah, Lawrence Holder, and Michael K. Skinner. 2023. "Hybrid Deep Learning Approach to Improve Classification of Low-Volume High-Dimensional Data." BMC Bioinformatics 24 (1): 1–20.

Meloni, Ilenia, Adham Elwakil, Gryczka Aurélia, et al. 2025. "Cohort Profile: OphtalmoLaus, an Extension of the CoLaus|PsyCoLaus Cohort to Investigate the Relationships between Ocular, Cardiovascular,

and Cognitive Parameters." In Research Square. September 22. https://doi.org/10.21203/rs.3.rs-
7660406/v1.

Ortin Vela, S., M. J. Beyeler, O. Trofimova, and M. Tomasoni. 2023. "Phenotypic and Genetic
Characteristics of Retinal Vascular Parameters and Their Association with Diseases." medRxiv.
https://www.medrxiv.org/content/10.1101/2023.07.07.23292368.abstract.

Ortín Vela, Sofía, Michael J. Beyeler, Olga Trofimova, et al. 2024. "Phenotypic and Genetic
Characteristics of Retinal Vascular Parameters and Their Association with Diseases." Nature
Communications 15 (1): 9593.

Parres, Daniel, and Roberto Paredes. 2023. "Fine-Tuning Vision Encoder–Decoder Transformers for
Handwriting Text Recognition on Historical Documents." Document Analysis and Recognition -
ICDAR 2023, 253–268.

Quiros, Jose Vargas, Bart Liefers, Karin van Garderen, et al. 2024. "VascX Models: Model Ensembles
for Retinal Vascular Analysis from Color Fundus Images." September 24.
http://arxiv.org/abs/2409.16016.

Rahimi, Mojtaba, Alfa Rossi, Taeyoon Son, et al. 2025. "High Dynamic Range Widefield Fundus
Photography with Transcranial Illumination: A Contrast Agent-Free Method for Non-Mydriatic
Choroidal Imaging." Biomedical Optics Express 16 (1): 84–96.

Rajpurkar, Pranav, Emma Chen, Oishi Banerjee, and Eric J. Topol. 2022. "AI in Health and Medicine."
Nature Medicine 28 (1): 31–38.

Rotem, Oded, Tamar Schwartz, Ron Maor, et al. 2024. "Visual Interpretability of Image-Based
Classification Models by Generative Latent Space Disentanglement Applied to in Vitro Fertilization."
Nature Communications 15 (1): 1–19.

Rotem, Oded, and Assaf Zaritsky. 2024. "Visual Interpretability of Bioimaging Deep Learning Models."
Nature Methods 21 (8): 1394–1397.

Selvaraju, Ramprasaath R., Michael Cogswell, Abhishek Das, Ramakrishna Vedantam, Devi Parikh, and
Dhruv Batra. 2016. Grad-CAM: Visual Explanations from Deep Networks via Gradient-Based
Localization. October 7. https://doi.org/10.1007/s11263-019-01228-7.

Shrikumar, Avanti, Peyton Greenside, and Anshul Kundaje. 06--11 Aug 2017. "Learning Important
Features through Propagating Activation Differences." In arXiv [cs.CV], edited by Doina Precup and
Yee Whye Teh. Proceedings of Machine Learning Research. arXiv.
https://proceedings.mlr.press/v70/shrikumar17a.html.

Staal, Joes, Michael D. Abràmoff, Meindert Niemeijer, Max A. Viergever, and Bram van Ginneken. 2004.
"Ridge-Based Vessel Segmentation in Color Images of the Retina." IEEE Transactions on Medical
Imaging 23 (4): 501–509.

Subramanian, Aravind, Pablo Tamayo, Vamsi K. Mootha, et al. 2005. "Gene Set Enrichment Analysis: A
Knowledge-Based Approach for Interpreting Genome-Wide Expression Profiles." Proceedings of the
National Academy of Sciences of the United States of America 102 (43): 15545–15550.

Tham, Yih Chung, Jocelyn Hui Lin Goh, Paul Nderitu, et al. 2025. "Building the World's First Truly Global
Medical Foundation Model." Nature Medicine 31 (11): 3580–3585.

Tomasoni, Mattia, Michael Johannes Beyeler, Sofia Ortin Vela, et al. 2023. "Genome-Wide Association
Studies of Retinal Vessel Tortuosity Identify Numerous Novel Loci Revealing Genes and Pathways
Associated With Ocular and Cardiometabolic Diseases." Ophthalmology Science 3 (3): 100288.

UK10K Consortium, Klaudia Walter, Josine L. Min, et al. 2015. "The UK10K Project Identifies Rare
Variants in Health and Disease." Nature 526 (7571): 82–90.

Uppaal, Rheeya, Junjie Hu, and Yixuan Li. 2023. "Is Fine-Tuning Needed? Pre-Trained Language
Models Are Near Perfect for Out-of-Domain Detection." May 22. http://arxiv.org/abs/2305.13282.

Van Eijgen, Jan, Jonathan Fhima, Marie-Isaline Billen Moulin-Romsée, Joachim A. Behar, Eirini

Christinaki, and Ingeborg Stalmans. 2024. "Leuven-Haifa High-Resolution Fundus Image Dataset
869   for Retinal Blood Vessel Segmentation and Glaucoma Diagnosis." Scientific Data 11 (1): 257.

870 Xie, Ziqian, Tao Zhang, Sangbae Kim, et al. 2024. "iGWAS: Image-Based Genome-Wide Association of
871   Self-Supervised Deep Phenotyping of Retina Fundus Images." PLOS Genetics 20 (5): e1011273.

872 Zhou, Bolei, Aditya Khosla, Agata Lapedriza, Aude Oliva, and Antonio Torralba. 2016. "Learning Deep
873   Features for Discriminative Localization." Paper presented 2016 IEEE Conference on Computer
874   Vision and Pattern Recognition (CVPR), 2016/6/27-2016/6/30, Las Vegas, NV, USA. 2016 IEEE
875   Conference on Computer Vision and Pattern Recognition (CVPR), June.
876   https://doi.org/10.1109/cvpr.2016.319.

877 Zhou, Yukun, Mark A. Chia, Siegfried K. Wagner, et al. 2023. "A Foundation Model for Generalizable
878   Disease Detection from Retinal Images." Nature 622 (7981): 156–163.

879 Zhou, Yukun, Siegfried K. Wagner, Mark A. Chia, et al. 2022. "AutoMorph: Automated Retinal Vascular
880   Morphology Quantification Via a Deep Learning Pipeline." Translational Vision Science &
881   Technology 11 (7): 12–12.