Chronic Complications of Type 2 Diabetes Patients in Southern Ethiopia

Biniyam Demisse Andarge[1*], Desta Haftu Hayelom[1], Sayih Mehari[1], Habtamu Esubalew[1], Habtamu Wondmagegn[1], Yohannes Habtegiorgis[2], Muluken Bekele[1], Yilma Chisha[1]

[1] College of Medicine and Health Science, Arba Minch University, Arba Minch, Ethiopia

[2] Director of Quality Services, Aleta Wondo Hospital, Aleta Wondo, Sidama, Ethiopia

*Corresponding author: Biniyam Demisse Andarge, bini.demisse@gmail.com

Abstract

Background: Type 2 diabetes mellitus (T2DM) is a global health concern associated with complications that significantly impact patients' quality of life and place significant burdens on healthcare systems. While the prevalence of T2DM is rising in Ethiopia, the scope and factors contributing to its complications remain understudied. Hence, this study aimed to assess the burden and identify associated factors of chronic complications among Type 2 diabetes mellitus patients attending Wolaita Sodo University Comprehensive Specialized Hospital in Southern Ethiopia.

Methods: A facility-based cross-sectional quantitative study was conducted from July to August 2024, involving 404 systematically sampled T2DM patients. Data on sociodemographic characteristics, clinical profiles, self-care practices, and chronic complications were collected through structured interviews and medical record reviews. Descriptive statistics summarized patient characteristics, while General Poisson regression identified factors associated with chronic complications. Results were reported as Adjusted Prevalence Ratios (APRs) with 95% Confidence Intervals (CIs), and a p-value < 0.05 was considered statistically significant.

Results: Of the 404 participants, 45.54% (95% CI: 40.61 - 50.54) had at least one chronic complication. Among these, 20.05% experienced one complication, 14.60% had two, 8.42% had three, and 2.48% had four. Peripheral neuropathy (14.85%) and nephropathy (9.65%) were the most common microvascular complications, while congestive heart failure (14.11%) and cerebrovascular disorders (11.39%) were the leading macrovascular complications. Multivariable Poisson regression identified older age (APR = 2.30; 95% CI: 1.23–4.28), longer diabetes duration (APR = 1.87; 95% CI: 1.28–2.76), elevated blood pressure (APR = 2.02; 95% CI: 1.21–3.37), and higher Body Mass Index (APR = 2.63; 95% CI: 1.43–4.81) as significant factors associated with increased prevalence of chronic complications.

Conclusion: The study reveals a substantial burden of chronic complications among T2DM patients, driven by modifiable and non-modifiable risk factors such as age, diabetes duration, blood pressure, and body mass index. These findings emphasize the urgent need for targeted interventions, including improved disease management, regular monitoring, and lifestyle modifications, to mitigate complications and enhance patient outcomes.

Keywords: Type 2 diabetes mellitus, chronic complications, prevalence, risk factors, Ethiopia

Introduction

Diabetes mellitus (DM) is a chronic, heterogeneous metabolic disorder characterized by elevated blood glucose levels due to impaired insulin secretion or function. Type 2 diabetes mellitus (T2DM), the most common form of DM, primarily affects middle-aged and older adults and is associated with poor lifestyle and dietary habits (1,2). T2DM is influenced by genetic and environmental factors, including obesity, sedentary behavior, unhealthy diet, and family history, and contributes significantly to global morbidity and mortality (3).

Non-communicable diseases (NCDs) account for three-quarters of all deaths worldwide, with diabetes causing an estimated 2 million deaths annually (4). The burden is particularly high in low- and middle-income countries like Ethiopia (5). As of the 2021 Global Burden of Disease Study, 529 million people globally live with diabetes, with T2DM comprising 96% of cases (6). This prevalence is expected to reach one in ten people by 2045, placing a significant strain on healthcare resources, with current diabetes-related expenditures accounting for 11.5% (USD 966 billion) of global healthcare budgets (7,8). In Africa, 24 million people currently live with diabetes, projected to rise to 55 million by 2045 (9), and Ethiopian prevalence estimated at 2.0%-6.5% (10).

Chronic hyperglycemia in diabetes substantially raises the risk of both microvascular (retinopathy, nephropathy, neuropathy) and macrovascular complications (cardiovascular disease, stroke) (11), which can lead to severe outcomes like blindness, kidney failure, and amputations (12,13). Half of diabetes patients unaware of their disease and complications are more likely in patients with poor glucose control, whereas improved management can slow disease progression (14,15). Cardiovascular disease is the most common complication with two-thirds of diabetes patients dying from myocardial infarction or stroke (16). Additional complications include end-stage renal disease (ESRD), neuropathy, and lower-extremity amputations (17,18).

The pathogenesis of diabetes complications involves multiple risk factors, some modifiable (e.g., diet, smoking, physical inactivity) and others non-modifiable (e.g., genetics) (19,20). A high BMI is a primary risk factor globally, with other significant contributors including dietary risks, tobacco use, and low physical activity (21). In response, the World Health Organization (WHO) and the United Nations have developed global action plans and Sustainable Development Goals (SDGs) to reduce the NCD burden, aiming to decrease premature mortality from NCDs by one-third by 2030 (22,23). As a WHO member, Ethiopia has adopted these frameworks and implemented a National Strategic Plan to address NCDs.

However, despite these efforts, the national burden of diabetes and its complications continues to rise

(24,25).

In Ethiopia, diabetes care remains challenged by limited resources, and the complexity of diabetes

complications poses substantial obstacles to improving patient outcomes. As diabetic complications often

go undetected until advanced stages, there is a need for early detection and proactive management

strategies in clinical settings, especially in regions with limited diagnostic and treatment capacity.

Addressing this need, this study explores the scope and nature of complications among patients with

T2DM to improve understanding of the condition and to inform healthcare practices and policies. By

investigating the factors associated with diabetes complications, this study aims to provide healthcare

providers and policymakers with actionable insights that support the development of locally adapted

interventions to reduce complications, improve patient quality of life, and alleviate the healthcare burden

in low-resource settings.

Materials and Methods

Study setting and period

This study was conducted at Wolaita Sodo University Comprehensive Specialized Hospital, located in Sodo

Town, Wolaita Zone, Southern Ethiopia from July 1 to August 30, 2024. Sodo Town is approximately 329

km south of Addis Ababa and serves as the administrative center of the South Ethiopia Regional State.

The hospital is the largest healthcare facility in the region, with a capacity of 370 beds distributed across

four main departments: medical, gynecology/obstetrics, surgical, and pediatrics. It is staffed by 975 health

professionals and 760 administrative personnel, collectively serving a catchment population of over 5

million people. The hospital has a chronic care unit that manages over 1,350 diabetic patients, 986 of

whom are diagnosed with Type 2 diabetes mellitus (T2DM).

Study design

119 A facility-based cross-sectional study design was employed.

Population and Eligibility Criteria

The study was conducted on type-2 diabetes mellitus (T2DM) patients attending Wolaita Sodo University

Comprehensive Specialized Hospital in South Ethiopia during the study period. Eligible participants were

adult patients aged 18 years and older with a clinical diagnosis of T2DM, who were willing to provide

information and had attended at least three follow-up visits at the hospital. Exclusion criteria were applied

to patients with incomplete medical records, missing essential data such as laboratory results, or lacking

documented information on diabetes-related complications. Additionally, patients with Type 1 diabetes

mellitus or gestational diabetes were excluded from the study.

Sample size determination

129 The required sample size was calculated with a 43% estimated proportion of complications among

130 diabetic patients (26), a 5% margin of error, and a 95% confidence level, yielding a sample size of 376

131 patients. Using these parameters, the sample size (n) was determined with the following formula:

132
$$\text{n} = \frac{(Za/2)^2 \times P(1-P)}{d^2} \qquad = \frac{(1.96)^2 \times 0.43(1-0.43)}{0.05^2}$$

133 Accounting for a 10% non-response rate, the final sample size was 414 patients. Further analysis validated

134 the adequacy of the sample size.

Sampling Technique and Procedures

This study used a systematic random sampling method to select participants from T2DM patients

attending routine follow-up visits. Before sampling, the total list of eligible patients receiving care at the

diabetic clinic during the study period was obtained. The sampling interval (k≈2) was determined by

dividing the total number of eligible patients (986) by the desired sample size of 414. During the study

period, the research team approached patients who visited the clinic, explained the study purpose, and

obtained informed consent. Participants who agreed to participate had their clinical records reviewed,

and data was collected accordingly.

Data Collection Tools and Procedure

Data collection was conducted using a pre-tested, structured questionnaire and checklist developed from

a thorough review of relevant literature. The questionnaire gathered information on socio-demographic

characteristics, clinical characteristics, self-care practices, and complications. Socio-demographic

characteristics included age, sex, marital status, education level, occupation, place of residence, and

family history of diabetes mellitus (DM). Clinical characteristics focused on the duration of diabetes,

glycemic control, blood pressure status, body mass index (BMI), current medication regimen, and

biochemical data such as HbA1c levels and lipid profiles. Self-care practices assessed lifestyle habits,

including dietary practices (e.g., salt intake), physical activity, smoking, alcohol consumption, foot care,

and the use of a glucometer for blood glucose monitoring. Complication assessment documented the

presence of both microvascular complications (e.g., peripheral neuropathy, nephropathy, retinopathy,

and sexual dysfunction) and macrovascular complications (e.g., hypertension, angina, myocardial

infarction, stroke, transient ischemic attack, heart failure, and foot-related issues like ulcers and
amputations). Complications were identified based on both self-reports and clinical documentation.

The data collection process was conducted by two trained and experienced nurses under the supervision
of the principal investigator and a medical doctor. Training was provided to ensure consistency and
accuracy in administering the tools. Eligible patients provided written informed consent before
participating in the study. The questionnaire was administered through face-to-face interviews conducted
in private settings to ensure confidentiality. Interviews were conducted in the local language to facilitate
comprehension. Following the interviews, medical records were reviewed to gather additional clinical
data and verify documented complications. Daily reviews of completed questionnaires were carried out
to ensure completeness and accuracy, with discrepancies resolved promptly.

Operational Definitions

Chronic complications of diabetes mellitus: Chronic complications are defined as the presence of one or
more vascular complications in a diabetic patient as diagnosed by a physician. These complications are
classified as either microvascular or macrovascular. Microvascular complications include neuropathy,
nephropathy, sexual dysfunction, and retinopathy. Macrovascular complications include hypertension,
coronary artery disease (angina and myocardial infarction), cerebrovascular disorders (stroke, and
transient ischemic attack), congestive heart failure, and peripheral artery disease (foot ulcers, and leg
amputations). Patients without any documented chronic complications are categorized as having no
complications (27).

Glycemic level: Glycemic levels are determined based on the three blood sugar measurements. These
were classified into two categories: optimal glycemic control (average fasting blood sugar of 80–130
mg/dL), and suboptimal glycemic control (average FBS of <70 mg/dL and >130 mg/dL) (2).

Blood pressure category: Blood pressure measurements were taken from three observations, and the
average of the second and third readings was recorded for analysis. Raised BP was defined as systolic
blood pressure (SBP) ≥ 140 mmHg and/or diastolic blood pressure (DBP) ≥ 90 mmHg (28).

Anthropometric measurement: Body mass index (BMI) is calculated by dividing the participant's weight
(kg) by height squared (m²) after data collection. Classified as; Normal: BMI of 18.5-24.9 kg/m²,
Overweight: BMI of 25-29.9 kg/m², and Obese: BMI of ≥30 kg/m² (28).

Self-care practices: refer to self-reported behaviors and actions taken by diabetes patients to manage
their condition. This includes regular blood glucose monitoring, dietary salt intake, physical activity, foot
care, and avoidance of harmful behaviors like smoking and alcohol consumption.

Data Entry and Analysis

Data were entered into Epi-Data Manager version 4.2 and analyzed in R software version 4.4.1. Descriptive
statistics including frequencies, percentages, means, and standard deviations were calculated to
summarize the study variables. Chronic complications were computed as count data to give the number
of complications developed per patient, and yield values ranged from zero to four. Preliminary analysis
showed the data were found to be under-dispersed (0.53), and count models were compared using the
Akaike Information Criterion (AIC) and Bayesian Information Criterion (BIC). The General Poisson
regression model was selected for final analysis based on the lowest AIC and BIC values. Adjusted
Prevalence Ratio (APR) with 95% confidence intervals (CIs) were reported and p-values < 0.05 were
considered statistically significant. Model fitness was assessed using the Hosmer-Lemeshow goodness-of-
fit test ($p = 0.59$), and multicollinearity was assessed using the Variance Inflation Factor (VIF), with all
variables with VIF values below 5.

Data Quality Assurance

A one-day training session was provided to data collectors and supervisors covering study objectives, data
collection instruments, ethical guidelines, and specific medical record review procedures. A pre-test
involving 5% of the sample size was conducted at Dilfana Primary Hospital to evaluate the data collection
tools and procedures. During data collection, the principal investigator and supervisors maintained
continuous oversight, conducting daily spot-checks of completed questionnaires to promptly address
discrepancies. Forms were reviewed each day to verify completeness and accuracy, and double data entry
was performed on a subset of questionnaires to identify and resolve any inconsistencies. Finally,
biostatisticians and subject matter experts reviewed the statistical analysis to validate the study findings.

Results

Sociodemographic Factors

The study included a total of 404 diabetes mellitus patients with a response rate of 97.58%. The mean age of respondents was 44.80 ± 14.1 years, where 48.17% were between 40-64 years, and 57.67% were male. Most participants were married (79.21%), had at least secondary education (57.18%), and resided in urban areas (59.16%). The majority were having an insurance (65.59%) and reported moderate social support (53.96%). Additionally, 43.56% had a family history of diabetes (Table 1).

Table 1: Sociodemographic Characteristics of Patients with Type-2 Diabetes Mellitus in South Ethiopia, 2024 (n=404).

Variable	Categories	Frequency	Proportion
Age	Mean (SD)= 44.80 ± 14.10		
	18-39	140	34.65
	40-64	235	58.17
	65+	29	7.18
Sex	Male	233	57.67
	Female	171	42.33
Marital status	Married	320	79.21
	Single	77	19.06
	Others*	7	1.73
Educational level	No formal education	87	21.53
	Primary (Grade 1–8)	86	21.29
	Secondary (Grade 9–12)	116	28.71
	Tertiary (12+)	115	28.47
Occupation	Paid employee	95	23.51
	Merchant	67	16.58
	Farmer	63	15.59
	Daily laborer	14	3.47
	House-wife	85	21.04
	Non-employee	17	4.21
	Students	63	15.59
Residence	Urban	239	59.16
	Rural	165	40.84
Health Insurance	Yes	265	65.59
	No	139	34.41
Social support	Low	130	32.18
	Moderate	218	53.96
	High	56	13.86
Family History of DM	Yes	176	43.56
	No	228	56.44

Others=Widowed, Divorced*

Clinical Characteristics

The clinical characteristics of the patients revealed that 58.91% had diabetes for less than 5 years, with 46.04% on insulin monotherapy and 64.85% having suboptimal glycemic control. Additionally, 51.49% had raised blood pressure, 64.60% were either overweight or obese, 39.61% had high triglycerides (≥200 mg/dL), and 45.55% had high cholesterol levels (≥240mg/dL) (Table 2).

Table 2: Clinical Characteristics of Patients with Type-2 Diabetes Mellitus in South Ethiopia, 2024 (n=404).

Variable	Value	Frequency	Proportion
DM duration, years	<5 years	238	58.91
	≥5 years	166	41.09
Current regimen	Insulin monotherapy	186	46.04
	Insulin + OGAs	104	25.74
	OGA monotherapy	30	7.43
	OGA combination therapy	84	20.79
Glycemic control	Optimal	142	35.15
	Suboptimal	262	64.85
Blood Pressure	Normal	196	48.51
	Raised	208	51.49
BMI	Normal	143	35.40
	Overweight	186	46.04
	Obese	75	18.56
Triglycerides	Normal (150–199mg/dL)	104	25.74
	Border line (150–199mg/dL)	140	34.65
	High & very high (≥200mg/dL)	160	39.61
Total cholesterol	Normal (<200mg/dL)	105	25.99
	Border line (200–239mg/dL)	115	28.46
	High (≥240mg/dL)	184	45.55

Burden and Patterns of Chronic Complications

Among diabetes mellitus (DM) patients, 45.54% (95% CI: 40.61-50.54) developed at least one complication. Among those who developed complications, 20.05% had one, 14.60% had two, 8.42% had three, and 2.48% had developed four complications (Figure 1).

Figure 1: Number of Complications Developed Among Patients with Type-2 Diabetes Mellitus in South Ethiopia, 2024. (n=404)

In this study, microvascular complications were prevalent with peripheral neuropathy affecting 14.85% of patients, nephropathy 9.65%, retinopathy 4.21%, and sexual dysfunction 0.50%. In terms of macrovascular complications, congestive heart failure (CHF) affected 14.11% of patients, peripheral artery disease (foot problems) affected 9.90%, cerebrovascular disorders (stroke and transient ischemic attack) affected 11.39%, and coronary artery disease (angina or myocardial infarction) affected 4.70% of patients (Figure 2).

Figure 2: Types of Complications Developed Among Patients with Type-2 Diabetes Mellitus in South Ethiopia, 2024. (n=404)

Factors Associated with Chronic Complications

On bivariate Poisson regression analysis age, marital status, educational level, residence, duration of diabetes mellites, glycemic control, blood pressure, body mass index, and hypertriglyceridemia were significantly associated with chronic complications, then exported to the multivariable regression model for further analysis.

In the multivariable Poisson regression analysis, four factors were identified as significantly associated with the prevalence of chronic complications among diabetes patients. Older age was associated with an increased prevalence of complications. Patients aged 40–64 years had a significantly higher prevalence of complications (APR = 1.78; 95% CI: 1.05, 3.01), while those aged 65 years and older showed an even higher prevalence (APR = 2.30; 95% CI: 1.23, 4.28) compared to those aged 18–39 years.

The duration of diabetes was also a significant factor. Patients with diabetes for five or more years had a 1.87 times higher prevalence of complications (APR = 1.87; 95% CI: 1.28, 2.76) compared to those with a shorter duration of the disease. Raised blood pressure significantly increased the prevalence of complications, with hypertensive patients having twice the prevalence of complications (APR = 2.02; 95% CI: 1.21, 3.37) compared to those with normal blood pressure.

Both overweight and obesity were significantly associated with the prevalence of complications. Overweight patients had a 1.82 times higher prevalence of complications (APR = 1.82; 95% CI: 1.05, 3.35), while obese patients had a 2.63 times higher prevalence (APR = 2.63; 95% CI: 1.43, 4.81) compared to patients with a normal BMI (Table 3).

Table 3: Factors Associated with Chronic Complications Among Patients with Type-2 Diabetes Mellitus in South Ethiopia, 2024 (n=404).

Variable	Categories	Chronic Complications	
		CPR (95% CI)	APR (95% CI)
Age	18-39	1	1
	40-64	3.63 (1.59, 8.27)	**1.78 (1.05, 3.01)****
	65+	4.43 (1.91, 10.29)	**2.30 (1.23, 4.28)****
Marital status	Married	1	1
	Single	0.12(0.07,0.23)	1.17(0.58,2.15)
	Others	1.36(0.75,2.48)	1.02(0.52,1.82)
Educational level	No formal education	1	1
	Primary (Grade 1–8)	0.63(0.49,0.81)	0.96(0.72, 1.28)
	Secondary (Grade 9–12)	0.38(0.29,0.50)	0.97(0.69,1.33)
	Tertiary (12+)	0.35(0.26,0.46)	1.06(0.66,1.71)
Residence	Urban	1	1
	Rural	0.70(0.57,0.87)	1.05(0.84,1.32)
DM duration, years	<5 years	1	1
	≥5 years	3.36 (2.57, 4.38)	**1.87 (1.28,2.76)****
Glycemic control	Optimal	1	1
	Suboptimal	2.32(1.78,3.02)	1.11(0.85,1.48)

Blood Pressure	Normal	1	1
	Raised	3.78 (2.46, 5.82)	**2.02 (1.21, 3.37)****
BMI	Normal	1	1
	Overweight	2.98 (1.72, 5.18)	**1.82 (1.05, 3.35)***
	Obese	4.96 (3.00, 8.19)	**2.63 (1.43, 4.81)*****
Triglycerides	Normal (<150mg/dL)	1	1
	Border line (150–199mg/dL)	2.12 (1.19, 3.75)	1.01 (0.56, 1.87)
	High (≥200mg/dL)	3.36 (2.57, 4.38)	1.28 (0.77, 2.26)

*p<0.05, **p<0.01, ***p<0.001; 1=reference

Discussion

This study revealed a significant burden of chronic complications among type-2 diabetes mellitus patients, with 45.54% developed at least one complication, and one in five patients presented with a multimorbidity. Among those, 20.05% had developed one complication, 14.60% had two, 8.42% had three, and 2.48% had developed four complications. These findings are consistent with other studies conducted in Ethiopia (26,29,30), and a slightly higher prevalence was observed in this study compared to other studies in the country (31,32). This suggests an ongoing burden of chronic complications among diabetes patients in the country, emphasizing the need to monitor blood glucose levels and manage risk factors to reduce the overall burden. Notably, the prevalence observed in this study is lower than that reported in studies from high and middle-income countries (33–36). This discrepancy may be due to differences in healthcare infrastructure, access to advanced healthcare services, early diagnosis, and management practices that might result in more accurate detection and reporting of complications.

Microvascular complications such as peripheral neuropathy (14.85%) and nephropathy (9.65%) were prevalent. Macrovascular complications, such as congestive heart failure (14.10%) and cerebrovascular disorders (11.39%) were also common, reflecting ongoing challenges in glycemic control and a pattern consistent with global findings where vascular problems dominate (16,37). The varied distribution of complications, with many patients suffering from multiple issues, shows a challenge in managing diabetes mellitus patients within the study area and may reflect the gaps in diabetes management strategies.

In this study, age emerged as a significant factor in the prevalence of complications, with patients aged 40–64 and those 65 years and older experiencing a higher rate compared to younger patients. This finding aligns with previous research indicating that aging is a critical risk factor for diabetes complications due

to the cumulative effects of prolonged hyperglycemia and age-related metabolic changes (31,34).

Similarly, the duration of diabetes was strongly associated with complications, as patients living with

diabetes for five or more years had significantly higher prevalence rates. This underscores the importance

of early diagnosis and consistent management of diabetes to prevent complications and reduce long-term

health risks (38) (39).

Raised blood pressure was another risk factor associated with a twofold increase in the prevalence of

complications. This finding is consistent with evidence indicating that elevated blood pressure

exacerbates the risk of both microvascular and macrovascular complications among individuals with

diabetes (40). This finding highlights the need for regular blood pressure monitoring and effective

management as integral components of diabetes care to prevent the progression of complications (40).

Similarly, Body mass index (BMI) was also a significant factor, with obesity and overweight strongly linked

to a higher prevalence of complications. This association reflects the well-documented link between an

increased body weight, increased insulin resistance, and the progression of type 2 diabetes (41). With the

rising prevalence of obesity globally and in Ethiopia, public health interventions that focus on promoting

healthy eating, physical activity, and lifestyle changes are urgently needed to mitigate the risks associated

with obesity and its contribution to diabetes complications (42,43).

Limitations of the study

311   The study's cross-sectional design limits the ability to establish causal relationships between the identified

312   factors and the complications of diabetes mellitus, and it does not provide the progression of

313   complications over time. Additionally, the reliance on self-reported data, particularly regarding self-care

314   practices, introduces potential biases such as recall bias or social desirability bias. Another significant

315   limitation is the possibility of missing undiagnosed complications.

Conclusion

This study highlights a significant burden of chronic complications among Type 2 diabetes mellitus patients

in the study area. Factors including older age, longer diabetes duration, elevated blood pressure, and

higher body mass index, were significantly associated with an increased prevalence of complications.

These findings underscore the critical need for targeted interventions, including early diagnosis, enhanced

glycemic and blood pressure control, and lifestyle modifications to mitigate risk factors. Strengthening

healthcare policies and practices to address these determinants can play a pivotal role in improving

patient outcomes and reducing the prevalence of diabetes-related complications.

Abbreviations

APR: Adjusted Prevalence Ratio; BMI: Body Mass Index; DALYs: Disability-Adjusted Life Years; DM:

Diabetes Mellitus; ESRD: End-Stage Renal Disease; NCDs: Non-Communicable Diseases; SDGs: Sustainable

Development Goals; T2DM: Type 2 Diabetes Mellitus; USD: United States Dollar; WHO: World Health

Organization

Ethical Consideration

Ethical approval was granted by the Institutional Review Board of Arba Minch University College of

Medicine and Health Sciences (IRB/23166/2024). Consent was obtained from each participant after

explaining the study objectives. Confidentiality was maintained by omitting personal identifiers from the

questionnaire, and all data were securely stored and accessible only to the research team.

Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author

upon reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

The author declares that no financial support was received for research, authorship, and publication.

Author Contributions

BDA contributed to the study's conception, design, data collection, and analysis, interpreted the results,

and drafted the manuscript. YC and MB contributed to the study's conception, design, and interpretation

of results and critically reviewed the manuscript. DHH, SM, HE, HW and YH contributed to the

interpretation of results and critically reviewed the manuscript. All authors have reviewed and approved

the final version of the manuscript and agree to be accountable for all aspects of the work, ensuring the

accuracy and integrity of the research.

Acknowledgments

The authors wish to extend their heartfelt gratitude to Wolaita Sodo University Comprehensive

Specialized Hospital for providing support and access to conduct this study. We are deeply thankful to all

the patients who participated in this study. Special thanks also go to the dedicated data collectors for their

hard work and commitment. This research would not have been possible without their invaluable

contributions.

References

1. American Diabetes Association; Diagnosis and Classification of Diabetes Mellitus. Diabetes Care. 2014 Jan 1;37(Supplement_1):S81–90. Available from: https://dx.doi.org/10.2337/dc14-S081

2. Introduction: Standards of Medical Care in Diabetes—2018. Diabetes Care. 2018 Jan 1; 41(Supplement_1):S1–2. Available from: https://dx.doi.org/10.2337/dc18-Sint01

3. Henson J, Anyiam O, Vishnubala D. Type 2 Diabetes. Exercise Management for Referred Medical Conditions. 2023 Jun 23;223–52. Available from: https://www.ncbi.nlm.nih.gov/books/NBK513253/

4. Piovani D, Nikolopoulos GK, Bonovas S. Non-Communicable Diseases: The Invisible Epidemic. J Clin Med. 2022 Oct 1;11(19):5939.

5. Bigna JJ, Noubiap JJ. The rising burden of non-communicable diseases in sub-Saharan Africa. Lancet Glob Health. 2019 Oct 1;7(10):e1295–6.

6. Ong KL, Stafford LK, McLaughlin SA, Boyko EJ, Vollset SE, Smith AE, et al. Global, regional, and national burden of diabetes from 1990 to 2021, with projections of prevalence to 2050: a systematic analysis for the Global Burden of Disease Study 2021. The Lancet. 2023 Jul 15;402(10397):203–34.

7. Hossain MJ, Al-Mamun M, Islam MR. Diabetes mellitus, the fastest growing global public health concern: Early detection should be focused. Health Sci Rep. 2024 Mar 1;7(3):e2004.

8. Williams R, Karuranga S, Malanda B, Saeedi P, Basit A, Besançon S, et al. Global and regional estimates and projections of diabetes-related health expenditure: Results from the International Diabetes Federation Diabetes Atlas, 9th edition. Diabetes Res Clin Pract. 2020 Apr 1;162.

9. Magliano DJBEJ committee, IDA 10th edition scientific. Diabetes by region. In: IDF Diabetes Atlas. 10th edition. Brussels: International Diabetes Federation; 2021. Available from: https://www.ncbi.nlm.nih.gov/books/NBK581937/

10. Bishu KG, Jenkins C, Yebyo HG, Atsbha M, Wubayehu T, Gebregziabher M. Diabetes in Ethiopia: A systematic review of prevalence, risk factors, complications, and cost. Obes Med. 2019 Sep 1;15:100132.

11. Turner R. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). Lancet. 1998 Sep 12;352(9131):837–53.

12. Sapra A, Bhandari P. Diabetes. StatPearls. 2023 Jun 21; Available from: https://www.ncbi.nlm.nih.gov/books/NBK551501/

13. Unger RH, Orci L. Paracrinology of islets and the paracrinopathy of diabetes. Proc Natl Acad Sci U S A. 2010 Sep 9;107(37):16009.

14.   Perreault L, Pan Q, Mather KJ, Watson KE, Hamman RF, Kahn SE. Effect of regression from
prediabetes to normal glucose regulation on long-term reduction in diabetes risk: Results from
the Diabetes Prevention Program Outcomes Study. The Lancet. 2012 Jun 16;379(9833):2243–51.

15.   Papatheodorou K, Banach M, Bekiari E, Rizzo M, Edmonds M. Complications of Diabetes 2017. J
Diabetes Res. 2018;2018.

16.   Wannamethee SG, Shaper AG, Whincup PH, Lennon L, Sattar N. Impact of Diabetes on
Cardiovascular Disease Risk and All-Cause Mortality in Older Men: Influence of Age at Onset,
Diabetes Duration, and Established and Novel Risk Factors. Arch Intern Med. 2011 Mar 14;
171(5):404–10.

17.   Deshpande AD, Harris-Hayes M, Schootman M. Epidemiology of diabetes and diabetes-related
complications. Phys Ther. 2008 Nov;88(11):1254–64.

18.   Forbes JM, Cooper ME. Mechanisms of diabetic complications. Physiol Rev. 2013 Jan 1;
93(1):137–88.

19.   Pinchevsky Y, Butkow N, Raal FJ, Chirwa T, Rothberg A. Demographic and Clinical Factors
Associated with Development of Type 2 Diabetes: A Review of the Literature. Int J Gen Med.
2020; 13:121.

20.   Kyrou I, Tsigos C, Mavrogianni C, Cardon G, Van Stappen V, Latomme J, et al. Sociodemographic
and lifestyle-related risk factors for identifying vulnerable groups for type 2 diabetes: a narrative
review with emphasis on data from Europe. BMC Endocr Disord. 2020 Mar 12;20(Suppl 1).

21.   Lin X, Xu Y, Pan X, Xu J, Ding Y, Sun X, et al. Global, regional, and national burden and trend of
diabetes in 195 countries and territories: an analysis from 1990 to 2025. Scientific Reports 2020
10:1. 2020 Sep 8;10(1):1–11.

22.   Hyder AA, Rylance S, Al Saegh A, Feigin VL, Kataria I, Laatikainen T, et al. Strengthening evidence
to inform health systems: opportunities for the WHO and partners to accelerate progress on non-
communicable diseases. BMJ Glob Health. 2023 Nov 19;8(11):e013994.

23.   Singh Thakur J, Nangia R, Singh S. Progress and challenges in achieving noncommunicable
diseases targets for the sustainable development goals. FASEB Bioadv. 2021 Aug 1; 3(8):563.

24.   Misganaw A, Haregu TN, Deribe K, Tessema GA, Deribew A, Melaku YA, et al. National mortality
burden due to communicable, non-communicable, and other diseases in Ethiopia, 1990-2015:
Findings from the Global Burden of Disease Study 2015. Popul Health Metr. 2017 Jul 21;15(1):1–
17.

25.   Awedew AF, Tezera *, Berheto M, Dheresa M, Tadesse S, Hailemariam A, et al. The Burden of
Non-Communicable Diseases and Its Implications for Sustainable Development Goals Across
Regions in Ethiopia. Ethiopian Journal of Health Development. 2023 Oct 20;37(2).

26.   Letta S, Aga F, Yadeta TA, Geda B, Dessie Y. Poor Self-Care Practices and Being Urban Resident
Strongly Predict Chronic Complications Among Patients with Type 2 Diabetes in Eastern Ethiopia:
A Hospital-Based Cross-Sectional Study. Diabetes Metab Syndr Obes. 2022;15:2095.

27. Kosiborod M, Gomes MB, Nicolucci A, Pocock S, Rathmann W, Shestakova M V., et al. Vascular
complications in patients with type 2 diabetes: prevalence and associated factors in 38 countries
(the DISCOVER study program). Cardiovasc Diabetol. 2018 Nov 28;17(1):1.

28. World Health Organization. The WHO STEPwise approach to noncommunicable disease risk
factor surveillance manual (2017); https://cdn.who.int/media/docs/default-source/ncds/ncd-
surveillance/steps/steps-manual.pdf?sfvrsn=c281673d_10

29. YimamAhmed M, Ejigu SH, Zeleke AZ, Hassen MY. Glycemic Control, Diabetes Complications and
Their Determinants Among Ambulatory Diabetes Mellitus Patients in Southwest Ethiopia: A
Prospective Cross-Sectional Study. Diabetes Metab Syndr Obes. 2020;13:1089.

30. Berhe KK, Mselle LT, Gebru HB. The magnitude of chronic diabetes complications and its
associated factors among diabetic patients attending the general hospitals in Tigray region,
Northern Ethiopia. PLoS One. 2023 Aug 1;18(8).

31. Seid MA, Akalu Y, Gela YY, Belsti Y, Diress M, Fekadu SA, et al. Microvascular complications and
its predictors among type 2 diabetes mellitus patients at Dessie town hospitals, Ethiopia.
Diabetol Metab Syndr. 2021 Dec 1;13(1):86.

32. Sheleme T, Mamo G, Melaku T, Sahilu T. Prevalence, Patterns and Predictors of Chronic
Complications of Diabetes Mellitus at a Large Referral Hospital in Ethiopia: A Prospective
Observational Study. Diabetes Metab Syndr Obes. 2020;13:4909.

33. Bruce MSP, Mallika MCV. Prevalence of complications of diabetes among patients with diabetes
mellitus attending a tertiary care centre in Tamil Nadu. Int J Community Med Public Health. 2019
Mar 27;6(4):1452–6.

34. Li J, Chattopadhyay K, Xu M, Chen Y, Hu F, Chu J, et al. Prevalence and associated factors of
vascular complications among inpatients with type 2 diabetes: A retrospective database study at
a tertiary care department, Ningbo, China. PLoS One. 2020 Jun 1;15(6).

35. Pedersen ML, Jacobsen JL, Lynge AR. Micro- and macrovascular complications among
Greenlanders and Danes with type 2 diabetes mellitus in Nuuk, Greenland. Int J Circumpolar
Health. 2010;69(2):195–207.

36. Faselis C, Katsimardou A, Imprialos K, Deligkaris P, Kallistratos M, Dimitriadis K. Microvascular
Complications of Type 2 Diabetes Mellitus. Curr Vasc Pharmacol. 2020 May 3;18(2):117–24.

37. Chen J, Yin D, Dou K. Intensified glycemic control by HbA1c for patients with coronary heart
disease and Type 2 diabetes: a review of findings and conclusions. Cardiovascular Diabetology
2023 22:1. 2023 Jun 22;22(1):1–16.

38. Yan Z, Cai M, Han X, Chen Q, Lu H. The Interaction Between Age and Risk Factors for Diabetes and
Prediabetes: A Community-Based Cross-Sectional Study. Diabetes, Metabolic Syndrome and
Obesity. 2023;16:85.

39. Antar SA, Ashour NA, Sharaky M, Khattab M, Ashour NA, Zaid RT, et al. Diabetes mellitus:
Classification, mediators, and complications; A gate to identify potential targets for the

development of new effective treatments. Biomedicine & Pharmacotherapy. 2023 Dec
1;168:115734.

40. Bretzel RG. Comorbidity of diabetes mellitus and hypertension in the clinical setting: A review of
prevalence, pathophysiology, and treatment perspectives. Clin Ther. 2007 Jan 1;29(SUPPL.
B):S35–43.

41. Chen Y, Zhang XP, Yuan J, Cai B, Wang XL, Wu XL, et al. Association of body mass index and age
with incident diabetes in Chinese adults: a population-based cohort study. BMJ Open. 2018
;8(9):21768.

42. Vetrivel Venkatasamy V, Pericherla S, Manthuruthil S, Mishra S, Hanno R. Effect of Physical
activity on Insulin Resistance, Inflammation and Oxidative Stress in Diabetes Mellitus. J Clin Diagn
Res. 2013 Aug 1;7(8):1764.

43. Syeda USA, Battillo D, Visaria A, Malin SK. The importance of exercise for glycemic control in type
2 diabetes. American Journal of Medicine Open. 2023 Jun 1;9:100031.