Title: Early pregnancy health behaviors, adverse pregnancy outcomes, and maternal blood pressure after pregnancy

Short title: Health behaviors, APOs, and future maternal BP

Andrea C Kozai, PhD, CSCS: Department of Epidemiology, University of Pittsburgh

Bethany Barone Gibbs, PhD, FAHA: Department of Epidemiology & Biostatistics, West Virginia University School of Public Health

Lisa D Levine, MD, MSCE: Department of Obstetrics & Gynecology, University of Pennsylvania Perelman School of Medicine

Abbi D Lane, PhD, FAHA: Department of Applied Exercise Science, University of Michigan

Mitali Ray, PhD, RN: Department of Health Promotion & Development, University of Pittsburgh School of Nursing

William Grobman, MD, MBA: Department of Obstetrics & Gynecology, Brown University School of Medicine

Philip Greenland, MD: Departments of Medicine (Cardiology) and Preventive Medicine, Northwestern University Feinberg School of Medicine

Matthew K Hoffman, MD, MPH: Department of Obstetrics & Gynecology, ChristianaCare

C Noel Bairey Merz, MD, FAHA: Department of Cardiology, Smidt Heart Institute, Cedars Sinai Medical Center

Phyllis C Zee, MD, PhD: Department of Neurology, Northwestern University Feinberg School of Medicine

Lisa Mims, MD: Department of Obstetrics & Gynecology, Indiana University School of Medicine

Natalie A Cameron, MD, MPH: Departments of General Internal Medicine and Preventive Medicine, Northwestern University Feinberg School of Medicine

George Saade, MD: Department of Obstetrics & Gynecology, Eastern Virginia Medical School

Robert M Silver, MD: Department of Obstetrics & Gynecology, University of Utah

Natalie A Bello, MD: Department of Cardiology, Cedars Sinai Medical Center

Eliza C Miller, MD, MS: Department of Neurology, University of Pittsburgh

Uma M Reddy, MD, MPH: Department of Obstetrics & Gynecology, Columbia University

Judith H Chung, MD, PhD: Department of Obstetrics & Gynecology, University of California, Irvine

Kara M Whitaker, PhD, FAHA: Departments of Health, Sport, & Human Physiology and Epidemiology, University of Iowa

Janet M Catov, PhD: Departments of Obstetrics, Gynecology & Reproductive Sciences, Epidemiology, and Clinical and Translational Sciences, University of Pittsburgh.

Corresponding author:

     Andrea C. Kozai, andrea.kozai@pitt.edu

     130 N. Bellefield Ave

     Pittsburgh, PA, 15213

Keywords:

Life's Essential 8, adverse pregnancy outcomes, maternal cardiovascular health, nuMoM2b-HHS

Disclosures

The authors have no financial disclosures relevant to this work.

Abstract

Background: Individual health behaviors are associated with pregnancy outcomes, but their joint effects are rarely considered and their associations with health beyond pregnancy are uncertain. We aimed to examine associations between combinations of first trimester health behaviors with adverse pregnancy outcomes (APOs) and blood pressure 2-7 years after delivery, and to estimate the proportion of associations with later blood pressure that were mediated by APOs.

Methods: Participants in the nuMoM2b and follow-up Heart Health Study prospective cohort were included. Physical activity, diet, sleep duration, and smoking were scored using the Life's Essential 8 framework. APOs were prospectively abstracted from medical records and included hypertensive disorders of pregnancy, preterm birth, gestational diabetes, small-for-gestational-age birth, or stillbirth. Latent profiles of health behaviors were constructed with structural equation modelling. Models were built using first trimester cardiovascular health scores on a scale of 0-100 for each of the four health behaviors. Risk of APOs and incident hypertension, and differences in continuous blood pressure 2-7 years after delivery based on behavioral profiles were assessed with Poisson or linear regression. Mediation analysis examined the proportion of associations between behavioral profiles and blood pressure mediated by APOs.

Results: Among 8,700 nulliparas, four behavioral profiles were identified: Healthiest Behaviors (36%), Healthy Activity/Sleep with Poor Diet/Smoking (21%), Healthy Sleep Only (32%), and Least Healthy Behaviors (11%). Forty-nine percent (4,497) returned for blood pressure assessment 2-7 years after delivery. Adjusted risk of APOs was 12-23% higher across the less healthy behavioral profiles compared to the Healthiest Behaviors profile. Further, adjusted systolic and diastolic blood pressures were 0.86 and 1.04 mmHg higher, respectively, 2-7 years after delivery in those with less healthy behavioral patterns compared to the Healthiest Behaviors profile. Eleven percent of the association between behavioral profiles and diastolic blood pressure was mediated by APOs. Rates of incident hypertension were not significantly different across behavioral profiles.

Conclusions: Less healthy combined health behavior profiles in early pregnancy were associated with APOs and higher blood pressure 2-7 years after delivery. The observed associations with later blood pressure were predominately direct effects, with limited mediation through APOs.

Non-standard Abbreviations and Acronyms:

APO: adverse pregnancy outcome

BMI: body mass index

CVH: cardiovascular health

HEI: Healthy Eating Index

HELLP: hemolysis, elevated liver enzymes, and low platelets

MVPA: moderate-to-vigorous physical activity

nuMoM2b-HHS: Nulliparous Pregnancy Outcomes: Monitoring Mothers-to-be Heart Health Study

Introduction

Pregnancy is considered a cardiovascular stress test that can provide an early warning for future risk of cardiovascular disease.[1] Adverse pregnancy outcomes (APOs), including hypertensive disorders of pregnancy, preterm birth, or gestational diabetes are associated with a more than two-fold higher risk of developing hypertension within seven years of delivery[2–7], suggesting that shared antecedents may exist. However, accurately identifying those who will develop APOs remains elusive, and prevention strategies are limited.

The American Heart Association's Life's Essential 8 framework for cardiovascular health (CVH)[8] includes health factors (blood glucose, lipids, body mass index [BMI], and blood pressure) and health behaviors (physical activity, diet quality, sleep duration, and nonsmoking) that can be assessed in clinical settings and provides a metric to help identify those at higher risk of APOs and later cardiovascular disease.[1] An early-pregnancy atherogenic profile derived from the health factor components has been associated with both APOs and incident hypertension 2-7 years after delivery in nulliparous individuals.[9] Separate modifiable health behavior components also impact pregnancy health[10–16], but importantly, health behaviors do not exist in isolation and adherence to multiple healthy behaviors can have additive beneficial effects.[17–19] Co-occurring profiles of behavioral patterns during pregnancy have yet to be evaluated as precursors of APOs, and it is unknown whether combined health behaviors in early pregnancy are related to long-term cardiovascular outcomes such as hypertension.

Furthermore, there is a growing body of evidence demonstrating that, while APOs are associated both with poor early-pregnancy CVH and later cardiovascular disease, early pregnancy health factors themselves are related to future CVD risk even among those without APOs.[5,9,20] While this has been demonstrated for clinical health factors such as BMI, blood pressure, lipids, and glucose, the contributions of health behaviors to these associations are uncertain. The Nulliparous Pregnancy Outcomes: Monitoring Mothers-to-be prospective cohort and its follow-up Heart Health Study (nuMoM2b-HHS) includes early-pregnancy assessment of health behaviors, prospective collection of rigorously defined APO diagnoses, and postpartum follow-up of cardiovascular risk metrics. Therefore, the objectives of this secondary analysis were: 1) to identify profiles of health behaviors in early pregnancy, 2) examine the associations of these profiles with incident APOs and maternal blood pressure several years after delivery, and 3) determine whether APOs mediated associations between behavioral profiles and later maternal blood pressure.

Methods

Study Design and Population

Study protocols for the nuMoM2b-HHS prospective cohort have been described previously.[21,22] Briefly, 10,038 nulliparous pregnant individuals were recruited between 2010-2013. Participants were enrolled in the first trimester and followed through delivery at eight clinical sites across the United States. Participants were eligible if they were 6 weeks+

0 days to 13 weeks+ 6 days pregnant with a singleton gestation and had no prior pregnancy that lasted past 20 weeks. In-person postpartum follow-up occurred 2-7 years following the index pregnancy (n=4,508). Ethical approval was obtained at each clinical site and participants provided written informed consent. Participants included in this analysis delivered at or after 20 weeks+ 0 days of gestation; those included in the APO analysis had complete pregnancy outcomes data, while those included in the analysis of future blood pressure returned for an in-person follow-up assessment and had complete blood pressure data. See **Figure 1** for the study flow diagram.

Health Behaviors

Health behaviors were self-reported at the first trimester assessment. Leisure-time physical activity type, duration, and frequency were assessed using a questionnaire adapted from the Behavior Risk Factor Surveillance System[23,24], converted to metabolic equivalents of the task using the Compendium of Physical Activities, 2nd edition[25], and quantified as minutes per week of moderate-to-vigorous physical activity (MVPA). Smoking history and exposure to secondhand smoke in the home were self-reported by questionnaire. Nightly hours of sleep duration were self-reported separately for weekdays and weekends; the mean was computed to calculate average nightly sleep duration over a 1-week period. Diet quality was assessed using a modified Block Food Frequency Questionnaire[26,27] and converted to a Healthy Eating Index (HEI) score. Physical activity, smoking, and sleep questions referenced the first trimester, while the diet questionnaire referenced the three months immediately preceding conception.

Each behavioral component was assigned a score from 0-100 using the American Heart Association's Life's Essential 8 algorithm (**Table S1**). Scores ≥80 indicate "ideal" health, while scores of 50-79 indicate "moderate" health and scores <50 indicate "poor" health.[28] Component scores for physical activity ranged from 0 (no weekly MVPA) to 100 (≥150 minutes/week MVPA) with intermediate scores for values between 0-150 minutes/week. HEI score percentiles were used to calculate the diet score. For example, 0 points corresponded to HEI <25th percentile, while 100 points corresponded to HEI ≥95th percentile. Sleep scores accounted for the U-shaped association between sleep duration and cardiovascular health, in which both low and high sleep durations are adverse.[29] A sleep score of 100 corresponded to 7-9 hours of sleep, with lower scores for fewer or greater hours of nightly sleep. Finally, the smoking score was adapted for pregnancy from the standard scoring algorithm. A score of 0 indicated current smoking during pregnancy, while a score of 100 indicated never smoking tobacco; intermediate scores were assigned for former smokers. In accordance with the standard scoring algorithm, twenty points were subtracted in cases of exposure to secondhand smoke in the home for all except current smokers.[8]

Adverse Pregnancy Outcomes

Pregnancy outcomes were determined through medical record abstraction. We used a binary composite APO variable, defined as presence of any of the following: hypertensive disorders of pregnancy (>140 mmHg systolic or >90 mmHg diastolic at or after 20 gestational weeks with or without proteinuria, end-organ involvement, and/or

HELLP syndrome), gestational diabetes, preterm birth (<37 weeks of gestation at delivery), small-for-gestational-age birth (<5th percentile using Alexander curves), or stillbirth.

Blood Pressure 2-7 Years After Delivery

Resting blood pressure was measured at the in-person nuMoM2b-HHS1 visit, occurring 2-7 years following the index pregnancy, using a standardized protocol.[22] Following a seated rest, blood pressure was measured three times using an automated device (Omron Healthcare Incorporated, Lake Forest, IL); the last two measurements were averaged and used for analysis. Incident hypertension was defined as new onset of stage 1 hypertension (≥130 mmHg systolic or ≥80 mmHg diastolic) or the use of blood pressure lowering medication at follow-up. Those with measured first trimester stage 1 hypertension were excluded from incident hypertension analyses. Finally, we examined associations with continuous values of systolic and diastolic blood pressure at follow-up.

Covariates and Demographic Characteristics

Questionnaires and interviews assessed participant age at study enrollment, income, education, insurance, perceived racial discrimination (Krieger Discrimination Scale[30], categorized as none, moderate, or high), and the social construct of self-reported racial and ethnic identity. For those self-reporting anti-hypertensive medication usage at the follow-up visit (n=117), we added 10/5 mmHg to measured systolic/diastolic blood pressures, respectively.[31] Height and weight were measured at the first trimester study visit using standardized protocols and were converted into body mass index (BMI) in kg/m^2. Self-reported pre-pregnancy diabetes mellitus was assessed at the first trimester visit by

interview. Resting blood pressure was measured at the first trimester study visit using the same protocol described above.

Statistical Analyses

Latent (unobserved) classes of combinations of the four health behaviors were constructed empirically using the "gsem" command in Stata, which can compute the models using continuous values. Models were built using first trimester CVH scores on a scale of 0-100 for each of the four health behaviors (physical activity, diet, sleep duration, and smoking) based on the full nuMoM2b sample (N=10,038), and the best-fitting model was selected based on values of Akaike/Bayesian Information Criterion, sample size in each class, and model interpretability; posterior probabilities were also assessed.

We assessed the risk of having an APO among behavioral latent profiles using Poisson regression with robust error variance. Crude models were first adjusted for participant characteristics including age, insurance status, income, education, and perceived racial discrimination; a final model also included first trimester blood pressure and BMI, which are health factors that may both mediate and confound the association between health behaviors and our outcomes.[32,33]

Next, we estimated the risk of incident stage 1 hypertension 2-7 years following delivery associated with early pregnancy behavioral profiles using Poisson regression with robust error variance. In addition, we tested the association between behavioral profiles and continuous blood pressure 2-7 years after delivery; separate models were tested for systolic and diastolic blood pressure using linear regression. Individuals with early-

pregnancy stage 1 hypertension (n=598, 13%) were excluded from the incident

hypertension models but were included in analyses of APOs and continuous blood

pressure values. Crude associations for both incident hypertension and continuous blood

pressure values were first adjusted for the time elapsed between delivery and the follow-up

assessment and first-trimester participant sociodemographic characteristics. A final

model added covariate adjustment for first trimester blood pressure and BMI.

Next, we conducted a mediation analysis to estimate the proportion of any

significant associations between early-pregnancy health behaviors and blood pressure

outcomes 2-7 years after delivery that were mediated by APOs. We used a multiplicative

model with an interaction term between behavioral profile and APO status (modeled as a

count outcome using Poisson regression). Decomposition of the total effects included

estimations of how much early pregnancy health behaviors directly affected later blood

pressure (natural direct effect) and how much of the effect was explained by APOs (natural

indirect effect); we also estimated the proportions of the total effect mediated by APOs.

Crude models were adjusted using the framework depicted in **Figure S1**.

Finally, we conducted the following sensitivity analyses: first, we evaluated

associations between early-pregnancy health behaviors and the APO and continuous

blood pressure outcomes when excluding those with early-pregnancy stage 1 hypertension

or diabetes. Second, the clinical cutoff for hypertension in pregnancy is ≥140 mmHg

systolic and/or ≥90 mmHg diastolic and the pregnancy measurements were taken prior to

the implementation of revised blood pressure categorizations[34], so we examined whether

associations with incident hypertension were different using incidence of stage 2

hypertension. All analyses were conducted using Stata version 18 (StataCorp, College

Station, TX) and significance was accepted at p<0.05.

Results

In total, 8,700 participants (mean ± SD 26.9 ± 5.6 years, 26.2 ± 6.3 kg/m^2 BMI, 40%

underrepresented race or ethnicity) with complete pregnancy outcomes data were

included in the APO analysis (**Table 1**). Model fit criteria for latent classes of early-

pregnancy health behaviors found a four-class solution to be the most appropriate (AIC =

343338.1, BIC = 343504.0, Entropy = 0.85; **Table S2**). Participants were categorized as

engaging in Healthiest Behaviors (n=3,158, 36.3%), Healthy Activity/Sleep with Poor

Diet/Smoking (n=1,807, 20.8%), Healthy Sleep Only (n=2,804, 32.2%), or Least Healthy

Behaviors (n=931, 10.7%) (**Figure 2**). Participants in the Healthiest Behaviors profile were

more likely to be older, have a lower BMI, have higher education and income, self-identify

as non-Hispanic White, have lower systolic and diastolic blood pressure in the first

trimester of pregnancy, and have a lower prevalence of stage 1 hypertension at baseline

compared to those in the other behavioral profiles.

Early pregnancy behavioral profiles were significantly associated with risk of APOs

(**Table 2**), with those in less healthy profiles having a 21-34% higher unadjusted risk

compared to those in the Healthiest Behaviors profile [overall Chi-squared (3) = 33.28,

p<0.001]. These significant associations remained following adjustment for

sociodemographic characteristics, ranging from a 12-23% higher risk. These associations

persisted in the Healthy Activity/Sleep with Poor Diet/Smoking (15% higher, 95%

confidence interval 2-30%) and Least Healthy Behaviors (18% higher, 95% confidence

interval 1-38%) profiles upon further adjustment for early pregnancy blood pressure and

BMI. A full summary of individual APO diagnoses by behavioral profiles is shown in **Table**

S3.

Of the original cohort, 4,497 returned for post-delivery follow-up at 3.2 ± 0.9 years

(range, 2-7 years) following the index pregnancy and had complete blood pressure data.

Given the substantial overlap in APO risk among the three less favorable behavioral

profiles, we dichotomized the groups as Healthiest Behaviors and Less Healthy Behaviors

(including the Healthy Activity/Sleep with Poor Diet/Smoking, Healthy Sleep Only, and

Least Healthy Behaviors profiles) for analyses relating to maternal blood pressure 2-7 years

following delivery. Incident stage 1 hypertension was found in 18.1% of the sample who

participated in the follow-up visit and did not have early-pregnancy stage 1 hypertension

(688 of 3,793 participants), with no significant differences in incidence between early-

pregnancy behavioral profiles (Healthiest Behaviors profile: 247 of 1,429, 17.3%; other

three Less Healthy Behaviors profiles combined: 441 of 1,923, 18.7%, IRR = 1.08 [95%

confidence interval 0.94 – 1.24, p=0.290], **Table 3**).

Crude linear regression results for continuous blood pressure found differences that

were retained after adjustment for visit date and sociodemographic factors (**Table 3**),

wherein participants in the combined less favorable behavior profiles had 0.86 mmHg

higher systolic blood pressure (p=0.035) and 1.04 mmHg higher diastolic blood pressure

(p=0.004) compared to the Healthiest Behaviors profile. After additional adjustment for first trimester blood pressure and BMI, these associations were no longer significant.

In analyses examining the mediation by APOs of the association between early pregnancy behavioral profiles and long-term blood pressure, significant total, natural direct, and natural indirect effects were found in crude models for both diastolic and systolic blood pressure (**Table 4**) which were retained upon adjustment for visit date and sociodemographic factors. When the models were additionally adjusted for early pregnancy blood pressure and BMI, total and natural direct effects remained significant for diastolic blood pressure while associations with systolic blood pressure were no longer significant. The proportion of the association mediated through APOs was 11% for diastolic blood pressure in Model 2 (p=0.034) and was nonsignificant for systolic blood pressure.

Sensitivity analyses excluding participants with pre-pregnancy stage 1 hypertension and/or diabetes from the models estimating APO risk and associations with continuous blood pressure 2-7 years after delivery did not change the direction or interpretation of our results (**Tables S4 and S5**). Similarly, rates of incident stage 2 hypertension were not different between those with healthy versus less healthy behaviors (**Table S6**).

Discussion

In this diverse prospective cohort of individuals enrolled during a first pregnancy we found four distinct first-trimester health behavior profiles that were associated with risk of subsequent APOs. Those with less healthy behavioral profiles were 12-23% more likely to

experience an APO than those with the healthiest behavioral profile. Further, less healthy early-pregnancy behavioral profiles were associated with higher blood pressure 2-7 years following delivery. Mediation analysis suggested that these effects were predominantly direct, rather than mediated through APOs. Our findings indicate that healthy behavior combinations in early pregnancy exhibit statistically significant associations with APOs and that behaviors in early pregnancy may be important for long-term cardiovascular risk whether or not an APO occurs.

By utilizing the Life's Essential 8 framework, we were able to construct latent profiles of healthy behaviors (physical activity, diet quality, sleep duration, and nonsmoking) in early pregnancy. This allowed us to leverage naturally occurring covariance in healthy behaviors during pregnancy, rather than forcing disaggregation. We found that lower adherence to healthy behaviors was associated with greater risk of APOs. This was particularly evident in the Least Healthy Behaviors profile (comprised of individuals with "poor" adherence to physical activity, diet quality, and sleep duration behaviors and "moderate" adherence to nonsmoking behavior), in which the adjusted risk ratio was highest. These results suggest that healthy behaviors in early pregnancy may be additive, and that "poor" adherence to at least two healthy behaviors in early pregnancy is related to elevated risk of APOs. Our findings are in contrast to those from a small cohort (n=202)[15] in which individual health behaviors, but not a composite score, were associated with features of placental health. This discrepancy may be attributable to cohort differences, distinct scoring algorithms, or study outcomes.

In addition, associations between behavioral profiles and APOs were independent of first trimester blood pressure and BMI. These findings suggest that engaging in healthy behaviors is an important contributor to pregnancy outcomes for all pregnant individuals, regardless of these clinical health factors in early pregnancy. Our results support existing evidence that healthy behaviors are associated with more favorable pregnancy outcomes[10–16,35], and add important knowledge about the combined effects of multiple healthy behaviors. Furthermore, these results emphasize the importance of healthy behaviors even for those with favorable early-pregnancy blood pressure and BMI.

It is also important to note sociodemographic differences among participants in each behavioral profile in our study. Those in profiles with the lowest adherence to healthy behaviors and the highest risk of APOs were likely impacted by structural factors that may have influenced the ability to adhere to and the interplay between the healthy behaviors (e.g., neighborhood factors that may influence access to both physical activity and healthy diet options). While we adjusted for sociodemographic factors, future research should investigate the drivers of these behavioral profiles and identify areas for intervention, such as enhancement of the built environment.

We found no significant association between behavioral profiles and incident hypertension 2-7 years after delivery. However, when examining associations among behavioral profiles and continuous blood pressure over this time, we found that those with less healthy behavioral profiles had 0.86 mmHg higher systolic blood pressure and 1.04 mmHg higher diastolic blood pressure 2-7 years following delivery than those in the Healthiest Behaviors profile. Our mediation analysis suggests that health behaviors may

play a direct role in longer-term continuous blood pressure levels in the years following

delivery, with a small but significant 11% of the effect on diastolic blood pressure mediated

through APOs. This is similar to two prior studies in the nuMoM2b-HHS cohort. In the first,

the association of an early-pregnancy atherogenic profile of poor lipids, C-reactive protein,

and insulin with postpartum incident hypertension was partially mediated by APOs.[9] In the

second, hypertensive disorders of pregnancy mediated a small but statistically significant

proportion of the association between early pregnancy obesity and incident hypertension.[20]

Taken together, the results of our study and this prior work suggest that early pregnancy

CVH components are robustly associated with future maternal cardiovascular risk, with a

small proportion of these associations mediated by APO occurrence (7-14% across these

studies). Our findings point to APOs as possible contributors to higher future blood

pressure, while behavioral profiles in early pregnancy are important whether or not an APO

occurs.

While 0.86-1.04 mmHg higher blood pressure over a mean follow-up of three years

is not a clinically large difference, small increases in blood pressure in the reproductive

years can compound cardiovascular risk over time.[36,37] We found that adjusting for first

trimester blood pressure and BMI attenuated our results, suggesting that these clinical

factors may themselves mediate the association between health behaviors in early

pregnancy and later blood pressure. There are multiple possible explanations of this

finding. First, it is possible that early pregnancy blood pressure confounded the association

with blood pressure in the years following delivery (i.e., those with higher blood pressure

following delivery simply started with higher blood pressure in early pregnancy). Second, it

is likely that blood pressure and BMI were in the causal pathway between early pregnancy health behaviors and blood pressure 2-7 years after delivery (i.e., those with healthier behaviors had more favorable early-pregnancy blood pressure and BMI due to the engagement in those healthy behaviors).[32,33] Given this potential mediating effect, we chose to report Model 2 as our primary results, which did not include early-pregnancy blood pressure and BMI. In either case, these findings emphasize the potential importance of maximizing pre-conception and early pregnancy health in order to limit the development of CVD risk factors in this critical peripartum window[1], and suggest that engaging in healthy behaviors may be a strategy to achieve this. Future interventional research is needed to determine whether improving engagement in healthy behaviors in the prenatal period is an effective means to improve long-term maternal health.

Pregnancy is a time of near-universal healthcare access, and motivation may be high to engage in behavior change.[38,39] Our results suggest that modifiable health behaviors group together in distinct ways, and that risks of multiple unhealthy behaviors compound one another and may be related to health beyond pregnancy. Notably, we did not see large differences in risk between our three less healthy behavioral profile groups, but each group with two or more behaviors classified as "poor" were at higher risk. This suggests that behavior change interventions may employ strategies to improve adherence to whichever health behavior is the most easily modifiable for the individual rather than targeting the adoption of a particular behavioral profile. Early pregnancy may be an ideal time to assess behavioral profiles and engage individuals in behavior change.

Strengths, Limitations, and Future Directions

This study has several strengths. We utilized a longitudinal approach to the reproductive years by examining early-pregnancy behavioral profile exposures and followed participants through pregnancy and several postpartum years. Our large, diverse prospective cohort of people experiencing a first birth captured well-documented APOs. In addition, we were able to identify four latent health behavior profiles, demonstrating for the first time how different behavioral combinations in early pregnancy are related to perinatal and long-term maternal health. Those who returned for a postpartum visit 2-7 years after delivery were demographically similar to the full pregnancy cohort[2], suggesting these results may be generalizable to our sample at large.

A limitation of this study included the self-report of health behaviors. However, given that these behaviors were reported prior to the outcomes occurring, recall bias or other measurement error would likely be random and therefore biased toward the null. In addition, the diet questionnaire referenced the 3 months prior to conception, which may not have accurately captured first trimester dietary patterns. Despite these limitations, our strong posterior probabilities of latent class membership suggest that the behavioral profiles are distinct and any bias is likely nondifferential. Furthermore, our activity questionnaire was limited to leisure-time physical activities. Future investigations should examine whether behavioral patterns are different among those with differing levels of occupational activity. Finally, health behaviors may have shifted between delivery and the postpartum follow-up assessment; future research should examine whether behavioral profiles are consistent after childbirth and whether more recent behavioral profiles explain additional variability in cardiovascular risk measures.

Conclusions

Our results suggest that co-occurring combinations of modifiable health behaviors are important and are related to pregnancy outcomes and blood pressure years following delivery. These findings add to the growing body of literature indicating that health in early pregnancy is a strong contributor to both pregnancy health and CVH following pregnancy. Importantly, health behaviors also had direct associations with blood pressure in the years following delivery and thus are important for those with and without pregnancy complications. Improving health behaviors in the prenatal context shows promise for reducing maternal risk during and possibly beyond pregnancy.

Acknowledgements

Grant funding from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD): U10 HD063036; U10 HD063072; U10 HD063047; U10 HD063037; U10 HD063041; U10 HD063020; U10 HD063046; U10 HD063048; and U10 HD063053. In addition, support was provided by Clinical and Translational Science Institutes: UL1TR001108 and UL1TR000153. Cooperative agreement funding from the National Heart, Lung, and Blood Institute and the Eunice Kennedy Shriver National Institute of Child Health and Human Development: U10- HL119991; U10-HL119989; U10-HL120034; U10-HL119990; U10-HL120006; U10-HL119992; U10-HL120019; U10-HL119993; U10-HL120018, and U01HL145358; and the National Center for Advancing Translational Sciences through UL-1- TR000124, UL-1-TR000153, UL-1-TR000439, and UL-1- TR001108; and the Barbra Streisand Women's Cardiovascular Research and Education Program, and the Erika J. Glazer Women's Heart Research Initiative, Cedars-Sinai Medical Center, Los Angeles. The Sedentary Behavior and Cardiovascular Health in Young Women ancillary study is supported by NHLBI R01HL158652. ACK is supported under NHLBI T32HL083825.

References

1. Khan SS, Cameron NA, Lindley KJ. Pregnancy as an Early Cardiovascular Moment: Peripartum Cardiovascular Health. *Circulation Research*. 2023;132(12):1584–1606.
2. Haas DM, Parker CB, Marsh DJ, Grobman WA, Ehrenthal DB, Greenland P, Bairey Merz CN, Pemberton VL, Silver RM, Barnes S, McNeil RB, Cleary K, Reddy UM, Chung JH, Parry S, et al. Association of Adverse Pregnancy Outcomes With Hypertension 2 to 7 Years Postpartum. *Journal of the American Heart Association*. 2019;8(19):e013092.
3. Sukmanee J, Liabsuetrakul T. Risk of future cardiovascular diseases in different years postpartum after hypertensive disorders of pregnancy: A systematic review and meta-analysis. *Medicine*. 2022;101(30):e29646.
4. Tassi A, Sala A, Mazzera I, Restaino S, Vizzielli G, Driul L. Long-term outcomes of patients with preeclampsia, a review of the literature. *Hypertension in Pregnancy*. 2023;42(1):2217448.
5. Fraser A, Catov JM. Placental syndromes and long-term risk of hypertension. *Journal of Human Hypertension*. 2023:1–4.
6. Giorgione V, Jansen G, Kitt J, Ghossein-Doha C, Leeson P, Thilaganathan B. Peripartum and Long-Term Maternal Cardiovascular Health After Preeclampsia. *Hypertension*. 2023;80(2):231–241.
7. Boucheron P, Lailler G, Moutengou E, Regnault N, Gabet A, Deneux-Tharaux C, Kretz S, Grave C, Mounier-Vehier C, Tsatsaris V, Plu-Bureau G, Blacher J, Olié V. Hypertensive disorders of pregnancy and onset of chronic hypertension in France: the nationwide CONCEPTION study. *European Heart Journal*. 2022;43(35):3352–3361.
8. Lloyd-Jones DM, Allen NB, Anderson CAM, Black T, Brewer LC, Foraker RE, Grandner MA, Lavretsky H, Perak AM, Sharma G, Rosamond W, on behalf of the American Heart Association. Life's Essential 8: Updating and Enhancing the American Heart Association's Construct of Cardiovascular Health: A Presidential Advisory From the American Heart Association. *Circulation*. 2022:101161CIR0000000000001078.
9. Catov JM, McNeil RB, Marsh DJ, Mercer BM, Merz CNB, Parker CB, Pemberton VL, Saade GR, Chen Y (Ida), Chung JH, Ehrenthal DB, Grobman WA, Haas DM, Parry S, Polito L, et al. Early Pregnancy Atherogenic Profile in a First Pregnancy and Hypertension Risk 2 to 7 Years After Delivery. *Journal of the American Heart Association*. 2020;10(5):e017216.
10. Davenport MH, Ruchat S-M, Poitras VJ, Jaramillo Garcia A, Gray CE, Barrowman N, Skow RJ, Meah VL, Riske L, Sobierajski F, James M, Kathol AJ, Nuspl M, Marchand A-A, Nagpal TS, et al. Prenatal exercise for the prevention of gestational diabetes mellitus and hypertensive disorders of pregnancy: a systematic review and meta-analysis. *British Journal of Sports Medicine*. 2018;52(21):1367–1375.
11. Bernardo D, Carvalho C, Mota J, Ferreira M, Santos P, Karger S. The Influence of Pregestational Body Mass Index and Physical Activity Patterns on Maternal, Delivery, and Newborn Outcomes in a Sample of Portuguese Pregnant Women: A Retrospective Cohort Study. *Portuguese Journal of Public Health*. 2023.
12. Miller C, Boushey C, Benny P, Ma S, Huang J, Lim E, Lee M-J. Diet quality predicts hypertensive disorders of pregnancy in Asian and Pacific Islander Cohort. *Nutrition and Health*. 2022:026010602211096.

13. Makarem N, Chau K, Miller EC, Gyamfi-Bannerman C, Tous I, Booker W, Catov JM, Haas DM, Grobman WA, Levine LD, McNeil R, Bairey Merz CN, Reddy U, Wapner RJ, Wong MS, et al. Association of a Mediterranean Diet Pattern With Adverse Pregnancy Outcomes Among US Women. *JAMA Network Open*. 2022;5(12):e2248165.

14. Lu Q, Zhang X, Wang Y, Li J, Xu Y, Song X, Su S, Zhu X, Vitiello MV, Shi J, Bao Y, Lu L. Sleep disturbances during pregnancy and adverse maternal and fetal outcomes: A systematic review and meta-analysis. *Sleep Medicine Reviews*. 2021;58:101436.

15. Teo SM, Murrin CM, Mehegan J, Douglass A, Hébert JR, Segurado R, Kelleher CC, Phillips CM. Associations between the maternal healthy lifestyle score and its individual components during early pregnancy with placental outcomes. *Placenta*. 2023;139:75–84.

16. Wang J, Yang W, Xiao W, Cao S. The association between smoking during pregnancy and hypertensive disorders of pregnancy: A systematic review and meta-analysis. *International Journal of Gynecology & Obstetrics*. 2022;157(1):31–41.

17. Reeves MJ, Rafferty AP. Healthy Lifestyle Characteristics Among Adults in the United States, 2000. *Archives of Internal Medicine*. 2005;165(8):854–857.

18. Loef M, Walach H. The combined effects of healthy lifestyle behaviors on all cause mortality: A systematic review and meta-analysis. *Preventive Medicine*. 2012;55(3):163–170.

19. Loprinzi PD, Fitzgerald EM, Woekel E, Cardinal BJ. Association of Physical Activity and Sedentary Behavior with Biological Markers Among U.S. Pregnant Women. *Journal of Women's Health*. 2013;22(11):953–958.

20. Khan SS, Petito LC, Huang X, Harrington K, McNeil RB, Bello NA, Bairey Merz CN, Miller EC, Ravi R, Scifres C, Catov JM, Pemberton VL, Varagic J, Zee PC, Yee LM, et al. Body Mass Index, Adverse Pregnancy Outcomes, and Cardiovascular Disease Risk. *Circulation Research*. 2023;133(9):725–735.

21. Haas DM, Parker CB, Wing DA, Parry S, Grobman WA, Mercer BM, Simhan HN, Hoffman MK, Silver RM, Wadhwa P, Iams JD, Koch MA, Caritis SN, Wapner RJ, Esplin MS, et al. A description of the methods of the Nulliparous Pregnancy Outcomes Study: monitoring mothers-to-be (nuMoM2b). *American Journal of Obstetrics and Gynecology*. 2015;212(4):539.e1-539.e24.

22. Haas DM, Ehrenthal DB, Koch MA, Catov JM, Barnes SE, Facco F, Parker CB, Mercer BM, Bairey-Merz CN, Silver RM, Wapner RJ, Simhan HN, Hoffman MK, Grobman WA, Greenland P, et al. Pregnancy as a Window to Future Cardiovascular Health: Design and Implementation of the nuMoM2b Heart Health Study. *American Journal of Epidemiology*. 2016;183(6):519–530.

23. Centers for Disease Control and Prevention, Division of Nutrition, Physical Activity, and Obesity. A Data Users Guide to the BRFSS Physical Activity Questions. 2011. https://www.cdc.gov/brfss/data_documentation/pdf/pa-rotatingcore_brfssguide_508comp_07252013final.pdf

24. Shea S, Stein AD, Lantigua R, Basch CE. Reliability of the behavioral risk factor survey in a triethnic population. *American Journal of Epidemiology*. 1991;133(5):489–500.

25. Shephard RJ. 2011 Compendium of Physical Activities: A Second Update of Codes and MET Values. *Yearbook of Sports Medicine*. 2012;2012:126–127.

26. Block G, Coyle LM, Hartman AM, Scoppa SM. Revision of Dietary Analysis Software for the Health Habits and History Questionnaire. *American Journal of Epidemiology*. 1994;139(12):1190–1196.

27. Parrott MS, Bodnar LM, Simhan HN, Harger G, Markovic N, Roberts JM. Maternal cereal consumption and adequacy of micronutrient intake in the periconceptional period. *Public Health Nutrition*. 2009;12(8):1276–1283.

28. U.S. Department of Health and Human Services. *Physical Activity Guidelines for Americans, 2nd edition*. Washington, DC: U.S. Department of Health and Human Services; 2018.

29. Yin J, Jin X, Shan Z, Li S, Huang H, Li P, Peng X, Peng Z, Yu K, Bao W, Yang W, Chen X, Liu L. Relationship of Sleep Duration With All-Cause Mortality and Cardiovascular Events: A Systematic Review and Dose-Response Meta-Analysis of Prospective Cohort Studies. *Journal of the American Heart Association*. 2017;6(9):e005947.

30. Krieger N, Sidney S. Racial discrimination and blood pressure: the CARDIA Study of young black and white adults. *American Journal of Public Health*. 1996;86(10):1370–1378.

31. Cheng S, Xanthakis V, Sullivan LM, Vasan RS. Blood Pressure Tracking Over the Adult Life Course. *Hypertension*. 2012;60(6):1393–1399.

32. Stuart JJ, Tanz LJ, Rimm EB, Spiegelman D, Missmer SA, Mukamal KJ, Rexrode KM, Rich - Edwards Janet W. Cardiovascular Risk Factors Mediate the Long-Term Maternal Risk Associated With Hypertensive Disorders of Pregnancy. *Journal of the American College of Cardiology*. 2022;79(19):1901–1913.

33. Schoenaker DA, Soedamah-Muthu SS, Mishra GD. Quantifying the mediating effect of body mass index on the relation between a Mediterranean diet and development of maternal pregnancy complications: the Australian Longitudinal Study on Women's Health12. *The American Journal of Clinical Nutrition*. 2016;104(3):638–645.

34. Whelton PK, Carey RM, Aronow WS, Casey DE, Collins KJ, Dennison Himmelfarb C, DePalma SM, Gidding S, Jamerson KA, Jones DW, MacLaughlin EJ, Muntner P, Ovbiagele B, Smith SC, Spencer CC, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018;71(6):e13–e115.

35. Al-Huda F, Shapiro GD, Davenport MH, Bertagnolli M, Dayan N. Association between Cardiorespiratory Fitness and Hypertensive Disorders of Pregnancy: A Systematic Review and Meta-Analysis. *Journal of Clinical Medicine*. 2022;11(15):4364.

36. Ji H, Kim A, Ebinger JE, Niiranen TJ, Claggett BL, Bairey Merz CN, Cheng S. Sex Differences in Blood Pressure Trajectories Over the Life Course. *JAMA Cardiology*. 2020;5(3):255–262.

37. Kishi S, Teixido -Tura Gisela, Ning H, Venkatesh BA, Wu C, Almeida A, Choi E-Y, Gjesdal O, Jacobs DR, Schreiner PJ, Gidding SS, Liu K, Lima JAC. Cumulative Blood Pressure in Early Adulthood and Cardiac Dysfunction in Middle Age. *Journal of the American College of Cardiology*. 2015;65(25):2679–2687.

38. Zinsser LA, Stoll K, Wieber F, Pehlke-Milde J, Gross MM. Changing behaviour in pregnant women: A scoping review. *Midwifery*. 2020;85:102680.

39. Lindqvist M, Lindkvist M, Eurenius E, Persson M, Mogren I. Change of lifestyle habits – Motivation and ability reported by pregnant women in northern Sweden. *Sexual & Reproductive Healthcare*. 2017;13:83–90.

Table 1. Participant characteristics by health behavior latent profiles.

	Healthiest Behaviors (n=3,158; 36%)	Healthy Activity/Sleep with Poor Diet/Smoking (n=1,807; 21%)	Healthy Sleep Only (n=2,804; 32%)	Least Healthy Behaviors (n=931; 11%)	P
Maternal age (years)	29.0 ± 4.9	26.2 ± 5.7	25.8 ± 5.6	24.4 ± 5.5	<0.001
Body mass index (kg/m²)	25.1 ± 5.1	27.1 ± 6.9	26.8 ± 6.5	27.1 ± 7.3	<0.001
Maternal Race					<0.001
American Indian	2 (<1%)	2 (<1%)	0 (0%)	3 (<1%)	
Asian	167 (5%)	51 (3%)	107 (4%)	28 (3%)	
Black non-Hispanic	183 (6%)	298 (16%)	481 (17%)	203 (22%)	
Hispanic	338 (11%)	275 (15%)	659 (24%)	226 (24%)	
Multiracial	103 (3%)	87 (5%)	111 (4%)	48 (5%)	
Native Hawaiian	9 (<1%)	11 (<1%)	8 (<1%)	5 (<1%)	
White non-Hispanic	2,334 (74%)	1,074 (59%)	1,422 (51%)	415 (45%)	
Not available	22 (<1%)	9 (<1%)	16 (<1%)	3 (<1%)	
Educational Attainment					<0.001
Less than high school	58 (2%)	186 (10%)	314 (11%)	142 (15%)	
Completed high school or GED	98 (3%)	254 (14%)	455 (16%)	209 (22%)	
Some college	362 (11%)	429 (24%)	670 (24%)	243 (26%)	
Associate or technical degree	261 (8%)	198 (11%)	324 (12%)	99 (11%)	
Completed college	1156 (37%)	448 (25%)	641 (23%)	146 (16%)	
Degree work beyond college	1219 (39%)	292 (16%)	394 (14%)	92 (10%)	
Percentage of Federal Poverty Level					<0.001
<100%	177 (6%)	284 (20%)	452 (22%)	194 (31%)	
100-200%	263 (9%)	250 (18%)	387 (18%)	115 (18%)	
>200%	2459 (85%)	887 (62%)	1253 (60%)	318 (51%)	
Insurance Payor					<0.001
Commercial	2000 (64%)	892 (50%)	1371 (49%)	334 (36%)	
Medicare/Medicaid	296 (9%)	583 (32%)	990 (36%)	448 (48%)	

Military/Other	39 (1%)	44 (2%)	42 (2%)	24 (3%)	
Two or more sources	730 (23%)	219 (12%)	296 (11%)	91 (10%)	
None (personal income)	80 (3%)	56 (3%)	75 (3%)	27 (3%)	
1st Trimester Systolic BP (mmHg)	108.6 ± 10.4	110.0 ± 11.0	109.2 ± 11.4	108.8 ± 11.5	<0.001
1st Trimester Diastolic BP (mmHg)	67.0 ± 8.1	67.5 ± 8.5	67.0 ± 8.5	66.4 ± 8.7	0.014
1st Trimester Stage 1 Hypertension	332 (10.7%)	257 (14.5%)	369 (13.5%)	122 (13.4%)	<0.001
1st Trimester Stage 2 Hypertension	35 (1.1%)	36 (2.0%)	47 (1.7%)	18 (2.0%)	0.058
Composite APO	882 (28%)	635 (35%)	948 (34%)	349 (37%)	<0.001

Reported as mean ± standard deviation or frequency and percentage, as appropriate. Differences among behavioral latent classes tested using one-way analysis of variance for continuous variables or Chi-squared tests for categorical variables. APO: adverse pregnancy outcomes; BP: blood pressure; GED: general education diploma.

Table 2. Incident rate ratios (IRR) of adverse pregnancy outcomes by behavioral profiles.

Group	Model 1 IRR (95% CI)	P	Model 2 IRR (95% CI)	P	Model 3 IRR (95% CI)	P
Healthiest Behaviors	1.00 (ref)	-	1.00 (ref)	-	1.00 (ref)	-
Healthy Activity/Sleep with Poor Diet/Smoking	**1.26 (1.16, 1.37)**	**<0.001**	**1.20 (1.09, 1.32)**	**<0.001**	**1.15 (1.04, 1.27)**	**0.004**
Healthy Sleep Only	**1.21 (1.12, 1.31)**	**<0.001**	**1.12 (1.02, 1.22)**	**0.016**	1.09 (0.99, 1.19)	0.068
Least Healthy Behaviors	**1.34 (1.21, 1.48)**	**<0.001**	**1.23 (1.09, 1.39)**	**0.001**	**1.18 (1.04, 1.34)**	**0.010**

Model 1: unadjusted; Model 2: adjusted for maternal age, education, insurance, income, and perceived racial discrimination; Model 3: additionally adjusted for first trimester blood pressure and body mass index. Estimates modeled with Poisson regression with robust error variance. Bold font indicates significant difference (p<0.05).

Table 3. Association of incident hypertension and continuous blood pressure by behavioral profiles.

Group	Model 1	P	Model 2	P	Model 3	P
Incident Hypertension [IRR (95% CI)]						
Healthiest Behaviors	1.00 (ref)	-	1.00 (ref)	-	1.00 (ref)	-
Less Healthy Behaviors	1.08 (0.94, 1.24)	0.290	1.01 (0.85, 1.19)	0.906	0.94 (0.80, 1.10)	0.442
Systolic Blood Pressure, mmHg [ß coefficient (95% CI)]						
Healthiest Behaviors	Reference	-	Reference	-	Reference	-
Less Healthy Behaviors	**1.42 (0.72, 2.12)**	**<0.001**	**0.86 (0.06, 1.66)**	**0.035**	0.54 (-0.23, 1.32)	0.170
Diastolic Blood Pressure, mmHg [ß coefficient (95% CI)]						
Healthiest Behaviors	Reference	-	Reference	-	Reference	-
Less Healthy Behaviors	**1.49 (0.87, 2.10)**	**<0.001**	**1.04 (0.33, 1.74)**	**0.004**	0.54 (-0.14, 1.22)	0.118

Model 1: unadjusted; Model 2: adjusted for time elapsed between visits, maternal age, education, insurance, income, and perceived racial discrimination; Model 3: additionally adjusted for first trimester blood pressure and body mass index. Incident hypertension modeled with Poisson regression with robust error variance; continuous blood pressure estimates modeled with linear regression. Bold font indicates significant difference (p<0.05). IRR: incidence rate ratio.

Table 4. Effect decomposition of the association between behavioral profiles and future blood pressure mediated by adverse pregnancy outcomes.

	Effect Decomposition			
	Natural Indirect Effect, *P*	Natural Direct Effect, *P*	Total Effect, *P*	Proportion mediated, *P*
Systolic Blood Pressure (mmHg)				
Model 1	**0.23, p<0.001**	**1.40, p<0.001**	**1.62, p<0.001**	**14%, p=0.002**
Model 2	**0.16, p=0.023**	**0.97, p=0.018**	**1.13, p=0.007**	14%, p=0.054
Model 3	0.09, p=0.110	0.69, p=0.084	0.78, p=0.055	11%, p=0.183
Diastolic Blood Pressure (mmHg)				
Model 1	**0.21, p<0.001**	**1.52, p<0.001**	**1.74, p<0.001**	**12%, p=0.001**
Model 2	**0.15, p=0.022**	**1.24, p=0.001**	**1.38, p<0.001**	**11%, p=0.034**
Model 3	0.07, p=0.113	**0.80, p=0.024**	**0.87, p=0.015**	8%, p=0.154

Model 1: unadjusted; Model 2: adjusted for maternal age, education, insurance, income, and perceived racial discrimination; Model 3: additionally adjusted for first trimester blood pressure and body mass index. Bold font indicates significant difference (p<0.05).

Figure 1. Study flow diagram. APO: adverse pregnancy outcome; nuMoM2b-HHS1: Nulliparous Pregnancy Outcomes: Monitoring Mothers-to-be Heart Health Study.

Figure 2. Health behavior cardiovascular health component scores by behavioral profiles. Mean (SD) cardiovascular health (CVH) scores for each health behavior component within each behavioral profile. Higher score indicates more ideal CVH; scores <50 are "poor" CVH; 50-79 are "moderate" CVH, and scores ≥80 are "high" CVH as indicated by markers on the y-axis.

Supplemental Tables

Table S1. Cardiovascular Health Scoring algorithm for health behaviors in pregnancy.
MVPA: moderate-to-vigorous physical activity; HEI: Healthy Eating Index.

	Value	Cardiovascular Health Score
Physical Activity (minutes MVPA/week)	0	0
	1-29	20
	30-59	40
	60-89	60
	90-119	80
	120-149	90
	150+	100
Diet (HEI percentile)	<25th	0
	25th-49th	25
	50th-74th	50
	75th-94th	80
	≥95th	100
Sleep Duration (mean hours/night)	<4	0
	4 - <5	20
	5 - <6 or ≥10	40
	6 - <7	70
	9 - <10	90
	7 - <9	100
Smoking	Current smoking	0
	Smoked within 3 months of conception	25
	Former smoker, quit prior to 3 months before conception	50
	Never smoker	100
	Lives with current smoker	Subtract 20 points unless score already = 0

Table S2. Latent Class metrics.

Solution comparisons	AIC	BIC
1 Class Solution	355988.6	356046.3
2 Class Solution	353844.9	353938.7
3 Class Solution	345209.8	345339.6
4 Class Solution	343338.1	343504.0
5 Class Solution	343348.1	343550.1
4 Class Solution Characteristics	**N (%)**	**Posterior probability**
Healthiest behaviors	3,158 (36.5%)	0.78
Healthy activity/sleep with poor diet/smoking	1,807 (20.6%)	0.73
Healthy sleep only	2,804 (32.3%)	0.95
Least healthy behaviors	931 (10.7%)	0.95
Overall	8,700 (100%)	0.85

AIC: Akaike's information criterion; BIC: Bayesian information criterion.

Table S3. Individual adverse pregnancy outcome diagnoses by behavioral profiles

	Healthiest Behaviors (n=3,158)	Healthy Activity/Sleep with Poor Diet/Smoking (n=1,807)	Healthy Sleep Only (n=2,804)	Least Healthy Behaviors (n=931)
Composite APO (n, %)	882 (30%)	635 (35%)	948 (34%)	349 (37%)
Preterm Birth	237 (8%)	192 (11%)	292 (10%)	108 (12%)
Early Preterm Birth	65 (2%)	57 (3%)	109 (4%)	38 (4%)
Hypertensive Disorders of Pregnancy	389 (12%)	290 (16%)	406 (14%)	159 (17%)
Gestational Diabetes	117 (4%)	84 (5%)	148 (5%)	47 (5%)
Small for Gestational Age	327 (10%)	216 (12%)	353 (13%)	122 (13%)
Stillbirth	12 (0.4%)	16 (0.9%)	20 (0.7%)	6 (0.6%)

APO: adverse pregnancy outcome. Early preterm birth defined as prior to 34 weeks of gestation at birth.

Table S4. Incident rate ratios (IRR) of adverse pregnancy outcomes by behavioral profiles, excluding those with pre-pregnancy hypertension or diabetes.

Group	Model 1 IRR (95% CI)	P	Model 2 IRR (95% CI)	P	Model 3 IRR (95% CI)	P
Healthiest Behaviors	1.00 (ref)	-	1.00 (ref)	-	1.00 (ref)	-
Healthy Activity/Sleep with Poor Diet/Smoking	**1.23 (1.12, 1.36)**	**<0.001**	**1.18 (1.06, 1.32)**	**0.003**	**1.15 (1.03, 1.28)**	**0.015**
Healthy Sleep Only	**1.18 (1.08, 1.29)**	**<0.001**	1.09 (0.98, 1.21)	0.108	1.07 (0.96, 1.19)	0.209
Least Healthy Behaviors	**1.33 (1.19, 1.49)**	**<0.001**	**1.22 (1.06, 1.41)**	**0.006**	**1.21 (1.05, 1.39)**	**0.010**

Model 1: unadjusted; Model 2: adjusted for maternal age, education, insurance, income, and perceived racial discrimination; Model 3: additionally adjusted for first trimester blood pressure and body mass index. Estimates modeled with Poisson regression with robust error variance. Bold font indicates significant difference (p<0.05).

Table S5. Association of incident hypertension and continuous blood pressure by behavioral profiles, excluding those with pre-pregnancy diabetes (incident hypertension) or pre-pregnancy diabetes or hypertension (continuous blood pressure).

Group	Model 1	*P*	Model 2	*P*	Model 3	*P*
Incident Stage Hypertension [IRR (95% CI)]						
Healthiest Behaviors	1.00 (ref)	-	1.00 (ref)	-	1.00 (ref)	-
Less Healthy Behaviors	1.07 (0.93, 1.23)	0.363	1.01 (0.85, 1.19)	0.938	0.93 (0.79, 1.10)	0.412
Systolic Blood Pressure, mmHg [ß coefficient (95% CI)]						
Healthiest Behaviors	Reference	-	Reference	-	Reference	-
Less Healthy Behaviors	**1.20 (0.50, 1.89)**	**0.001**	**0.83 (0.02, 1.65)**	**0.045**	0.56 (-0.24, 1.36)	0.169
Diastolic Blood Pressure, mmHg [ß coefficient (95% CI)]						
Healthiest Behaviors	Reference	-	Reference	-	Reference	-
Less Healthy Behaviors	**1.29 (0.66, 1.92)**	**<0.001**	**0.94 (0.21, 1.67)**	**0.012**	0.56 (-0.15, 1.27)	0.120

Model 1: unadjusted; Model 2: adjusted for time elapsed between visits, maternal age, education, insurance, income, and perceived racial discrimination; Model 3: additionally adjusted for first trimester blood pressure and body mass index. Bold font indicates significant difference (p<0.05). IRR: incidence rate ratio.

Table S6. Association of incident stage 2 hypertension by behavioral profiles.

Group	Model 1	P	Model 2	P	Model 3	P
Incident Stage Hypertension [IRR (95% CI)]						
Healthiest Behaviors	1.00 (ref)	-	1.00 (ref)	-	1.00 (ref)	-
Less Healthy Behaviors	1.25 (0.94, 1.66)	0.129	1.12 (0.79, 1.59)	0.522	0.96 (0.69, 1.35)	0.818

Model 1: unadjusted; Model 2: adjusted for time elapsed between visits, maternal age, education, insurance, income, and perceived racial discrimination; Model 3: additionally adjusted for first trimester blood pressure and body mass index. Bold font indicates significant difference (p<0.05). IRR: incidence rate ratio.

Figure S1. Schematic of mediation analysis to determine the mediating effect of adverse pregnancy outcomes (APO) on the association between early pregnancy behavioral profiles and systolic and diastolic blood pressure 2-7 years following delivery. BMI: body mass index; E: exposure; M: mediator; O: outcome.