Title: The Effect of Three Key Administrative Errors on Patient Trust in Physicians: Prescription Errors, Confidentiality Breaches, and Appointment Errors

Tetsuro Aita, MD, PhD[1,2#]; Yoshia Miyawaki, MD, PhD, MPH[3#]; Yu Katayama, MD, PhD[3]; Kosuke Sakurai, PharmD, PhD[4]; Nao Oguro, MD, PhD[2,5,6]; Takafumi Wakita, PhD[7]; Nobuyuki Yajima, MD, PhD, MPH[2,5,8]; Ashwin B. Gupta, MD[9,10*]; Noriaki Kurita, MD, PhD[2,5,11*]

#Dr. Aita and Dr. Miyawaki share co-first authorship.

*Dr. Kurita and Dr. Gupta share co-last authorship.

[1]Department of General Internal Medicine and Family Medicine, Fukushima Medical University, Fukushima City, Fukushima, Japan

[2]Department of Clinical Epidemiology, Graduate School of Medicine, Fukushima Medical University, Fukushima City, Fukushima, Japan

[3]Department of Nephrology, Rheumatology, Endocrinology and Metabolism, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama-city, Okayama, Japan

[4]Department of Pharmacy, Showa University Koto Toyosu Hospital, Koto-ku,, Tokyo, Japan

[5]Division of Rheumatology, Department of Medicine, Showa University School of Medicine, Shinagawa-ku, Tokyo, Japan

[6]Department of Healthcare Epidemiology, School of Public Health in the Graduate

School of Medicine, Kyoto University, Kyoto-city, Kyoto, Japan

[7]Department of Sociology, Kansai University, Suita-city, Osaka, Japan

[8]Center for Innovative Research for Communities and Clinical Excellence, Fukushima

Medical University, Fukushima City, Fukushima, Japan

[9]Medicine Service, VA Ann Arbor Healthcare System, Ann Arbor, Michigan

[10]Division of Hospital Medicine, Department of Internal Medicine, University of

Michigan Medical School, Ann Arbor, Michigan

[11]Department of Innovative Research and Education for Clinicians and Trainees

(DiRECT), Fukushima Medical University Hospital, Fukushima City, Fukushima,

Japan

Corresponding author details:

Noriaki Kurita, MD, PhD, FACP

Department of Clinical Epidemiology, Graduate School of Medicine,

Fukushima Medical University

1 Hikarigaoka, Fukushima City

Fukushima 960-1295, Japan Postal address

kuritanoriaki@gmail.com

Abstract

Importance

Understanding how administrative errors, such as prescription errors, appointment

scheduling errors, and patient confidentiality breaches impact trust in physicians is

crucial for improving patient-physician relationships and healthcare outcomes.

Objectives

To investigate the association between administrative errors, general trust in physicians

and interpersonal trust in a physician.

Design

Cross-sectional study.

Setting

An online panel survey targeting adults across Japan.

Participants

Adults aged ≥20 years who had received treatment at least twice for non-communicable

diseases within the past 6 months.

Exposure

Past experiences with prescription errors, confidentiality breaches, and appointment

scheduling errors by personal physicians treating their non-communicable diseases.

Primary Outcomes and Measurements

General trust in physicians and interpersonal trust in a physician were measured using

the Japanese version of the Wake Forest Physician Trust Scale.

Results

Among the 661 participants, nearly 14% reported experiencing at least one type of

administrative error. Prescription errors were associated with a significant decrease in

general trust in physicians (–9.78 points, 95% confidence interval (CI): –13.74 to –5.81).

Confidentiality breaches had the most significant negative impact on interpersonal trust

(–14.09 points, 95% CI: –24.35 to –3.83), followed by appointment scheduling errors (–

13.56 points, 95% CI: –22.48 to –4.65). Mediation analysis revealed that the association

between prescription errors and reduced general trust was partially mediated by

decreased trust in personal physicians.

Conclusions and Relevance

Administrative errors during care for non-communicable diseases significantly

undermine patients' trust in physicians. Physicians should prioritise improving their

practices, particularly regarding prescription errors, as these errors harm interpersonal

relationships and have broader implications for the public's perception of physicians.

key messages

What is already known on this topic

Patient trust in physicians has been conceptualised as two constructs: trust in personal

physicians and trust in physicians in general. While administrative errors, including

medication prescription errors, can cause physical harm to patients, the impact of such

errors, along with issues like appointment scheduling errors and breaches of

confidentiality, on trust in physicians, has not been fully explored.

What this study adds

This study found that prescription errors were associated with a decline in general trust

in physicians. Appointment scheduling errors and breaches of confidentiality were

shown to undermine interpersonal trust in personal physicians. Additionally, loss of

trust in personal physicians partially mediated the association between prescription

errors and loss of general trust in physicians.

How this study might affect research, practice or policy

Enhancing administrative practices, such as ensuring accuracy in medication prescribing,

improving appointment scheduling, and safeguarding patient confidentiality, could

strengthen the physician-patient relationship on a personal level and pave the way for

better future interactions with other physicians.

Introduction

Patient trust in physicians is central to medical care and plays an important role

in medication adherence, adoption of recommended preventive behaviours, and

ensuring the continuity of care. [1–3] Trust in physicians can be classified into two

constructs: interpersonal trust, which refers to trust in a specific physician, and general

trust, which pertains to trust in physicians as a collective group. [4,5] General trust is

particularly important for enabling physicians to gain confidence with new patients who

have limited prior interactions. [4] Although numerous studies have shown that medical

errors causing physical harm result in diminished trust in the involved physician or

institution, [6–8] limited understanding of how administrative errors in outpatient

services—errors that do not necessarily involve physical harm—affect both

interpersonal and general trust exist.

Prescription errors, for instance, can lead to the delayed recognition of adverse

events, causing physical and emotional distress for patients and their families. [8,9]

These errors, whether executive actions (e.g. incorrect prescriptions) or omissions (e.g.

failure to prescribe), are known to impact patient well-being; however, their influence

on trust in physicians remains unclear. [9] Similarly, appointment scheduling errors

represent a common administrative omission, yet their impact on trust has not been

extensively studied. [10] In addition, patient confidentiality breaches, a cornerstone of

the doctor-patient relationship, pose another critical issue. While confidentiality

breaches may not cause direct physical harm, they can erode trust, result in patient

humiliation, and lead to broad disadvantages (i.e., legal sanctions). [11,12] While it is

intuitive that confidentiality violations damage interpersonal trust, their potential to

diminish general trust in the healthcare system has not been quantitatively examined.

[12] This distinction is critical, as general trust may be impacted differently to these

errors than interpersonal trust. [4,5]

Our study aimed to examine the impact of administrative errors—such as

prescription errors, patient confidentiality breaches, and appointment scheduling

errors—on both interpersonal trust in their physicians and general trust in physicians as

a group among adults with non-communicable diseases. Furthermore, we explored

whether interpersonal trust in their physicians mediates the relationship between these

errors and general trust, offering insights into the importance of different dimensions of

trust in healthcare management.

Methods

Study Design and Subjects

This cross-sectional study used an online panel survey conducted through Cross

Marketing, a web-based company located in Shinjuku-ku, Tokyo, to recruit Japanese

participants with non-communicable diseases. The survey was conducted on April 27

and 28, 2020, targeting individuals aged 20 years and older. Eligible participants were

recruited consecutively and their responses were collected simultaneously. The study

protocol was approved by the Ethics Review Board of Okayama University Hospital

(No. ken-2405-041). Participants were presented with an online consent statement at the

beginning of the survey, and only those who completed the survey were included in the

study. This study adhered to the Strengthening the Reporting of Observational Studies

in Epidemiology (STROBE) guidelines. [13]

Screening Process and Data Collection

Quality assurance measures to mitigate random variability and enhance data

reliability have been described previously. [5,14] Briefly, the survey employed multiple

screening items to ensure the validity of responses. Participants were initially asked to

select any non-communicable diseases for which they had received treatment at least

twice within the past 6 months, with the option of selecting multiple conditions. They

were then asked to identify the most burdensome diseases based on their selections.

Any inconsistencies between the diseases reported in these two items led to the

exclusion of the participant's data. Furthermore, respondents were asked to list the

medications prescribed for their most burdensome diseases. Any discrepancies between

the most burdensome disease and appropriate medications for that disease resulted in

exclusion. Two researchers independently verified the listed medications against an

online database of label information and resolved any discrepancies by consensus to

ensure consistency. An exception was made for respondents who selected cancer and

reported "none" for prescribed medications, as the watch-and-wait approach is

considered appropriate for certain cancer treatments. Additionally, a response time cut-

off of 300 s, determined through prior pilot testing, was applied to exclude overly quick

responses that could indicate random or non-serious answers.

Outcome

The primary outcomes were general trust in physicians and interpersonal trust in personal physicians. Trust was measured using the Japanese version of the 5-item Wake Forest Physician Trust Scale: *Trust in Doctors Generally* and *Interpersonal Trust in Physician*. [5,15] Each item was rated on a 5-point Likert scale ranging from 1 (*totally disagree*) to 5 (*totally agree*). One negatively worded item was reverse coded, and the total score for each scale was normalised to a range of 0 to 100.

Before answering the *Interpersonal Trust in Physician* scale, respondents were instructed as follows: "Please think of the doctor who cares for your [the most troublesome disease chosen by the participants was automatically displayed here] when you answer these questions. He or she will be considered your doctor for this survey. For the next question, we are interested in your honest opinion about your doctor. Please choose the answer that best matches your opinion for each question."

For the *Trust in Doctors Generally* scale, respondents received the following instructions: "The following questions may seem similar to the previous ones. However, they are not about your doctors, however, doctors in general. No need to be concerned if these issues have not been considered previously. No correct or incorrect responses were obtained. Please choose the answer that best matches your thoughts on doctors."

Exposures

Exposure included three types of administrative errors: prescription errors,

appointment scheduling failures, and confidentiality breaches. Respondents were first

provided with the following instructions: "Please select the response that best describes

your opinion regarding the following statements about your doctor who cares for your

[the most troublesome disease chosen by the participants was automatically displayed

here]". The following items were then presented: "Has your doctor ever failed to

prescribe your medication, or mistakenly prescribed wrong types or dosages of your

medication?", "Has your doctor previously forgotten an outpatient or exam

appointment?", "Has your doctor ever previously discussed your illness or personal

information with you in a place where someone you did not know could hear you?" For

each item, respondents were asked to select either (1) (*have had*) or (2) (*have not*).

Covariates

Potential confounders were identified based on prior literature and clinical

expertise. These included age, [1,16] sex, [16] final education, household income, [16]

reported disease, duration of the patient-physician relationship, [1] and patient's general

level of interpersonal traits. The patients' general level of interpersonal trust was

assessed using the 6-item General Trust Scale, [17] rated on a 5-point Likert scale

ranging from *strongly disagree* to *strongly agree*. The total score was computed by

summing the responses to all items.

Statistical Analysis

Demographic and clinical variables are summarised as medians with interquartile ranges

for continuous variables and frequencies with percentages for categorical variables. We

used general linear models to estimate the associations between the three types of

administrative errors and both interpersonal trust in a physician and general trust in

physicians. Cluster-robust variance estimation was used to account for the within-

prefecture clustering. [18] To assess whether trust in personal physicians mediates the

relationship between specific administrative errors and general trust in physicians, we

conducted a mediation analysis. A series of mediation models were fitted to examine

the hypothesised relationships between administrative errors, interpersonal trust

(mediator), and general trust while adjusting for the previously described covariates.

The mediation analysis was performed using the user-written Stata command

"sgmediation2". [19] All statistical analyses were performed using Stata version 18

(StataCorp).

Results

Participant characteristics

A total of 964 participants met the inclusion criteria; however, 247 were

excluded due to inconsistent responses regarding their most troubling disease and

medication, and 46 were excluded for responding too quickly to provide logical answers.

[5] Of the remaining 671 eligible participants, 661 (mean age: 63 years; 26.5% female)

were included in the primary analyses and 10 were excluded due to missing data on

covariates. The baseline characteristics are shown in Table 1. The most common

diseases were cancer (n = 255; 38.6%), diabetes mellitus (n = 191; 28.9%), and

depression (n = 127; 19.2%). Among the participants, 92 (13.9%) reported experiencing

at least one administrative error, 72 (10.9%) experienced a prescription error, 18 (2.7%)

experienced a confidentiality breach, and 26 (3.9%) reported an appointment reservation

error.

Association between trust in physicians and three types of administrative errors

After adjusting for covariates, all three types of errors were associated with decreased

interpersonal trust in their physicians. As shown in Table 2 and Fig. 1, confidentiality

breaches had the most significant negative impact on interpersonal trust scores

(coefficient estimate: –14.09 points [95% confidence interval (CI): –24.35 to –3.83; p =

0.008), followed by appointment reservation errors (–13.56 points [95% CI: –22.48 to –

4.65; p = 0.004). Medication prescription errors also significantly reduced interpersonal

trust (–5.56 points [95% CI: –10.70 to –0.41; p = 0.035). In contrast, general trust in

physicians was primarily associated with medication prescription errors, which led to a

reduction of 9.78 points [95% CI: –13.74 to –5.81; p < 0.001). No significant

associations were found between general trust in physicians and appointment

reservation errors (–6.12 points [95% CI: –12.59 to 0.35; p = 0.063) or confidentiality

breaches (–5.23 points [95% CI: –11.33 to 0.88; p = 0.091).

The mediating role of trust in their own physician in the association between medication

prescription error and trust in general physicians

Medication prescription errors were the only type of administrative error significantly

associated with decreased general trust in physicians. Both direct effects (69% of the

total effect) and indirect effects (31% of the total effect) of medication prescription

errors on general trust in physicians were observed. Mediation analysis revealed that a

decline in interpersonal trust in their physicians partially mediated the relationship

between medication prescription errors and the overall decline in general trust in

physicians (Sobel test, $p = 0.031$; see Fig. 2).

Discussion

This study underscores the significant impact of administrative errors on patient trust in

physicians among individuals with non-communicable diseases such as cancer, diabetes,

and depression. Prescription errors were associated with reduced trust in both personal

physicians and physicians in general, with the loss of trust in personal physicians

partially mediating the loss of trust in general physicians.

Confidentiality breaches and missed appointment scheduling resulted in the greatest

reduction in interpersonal trust scores, highlighting that ensuring reliable appointment

scheduling is as crucial as maintaining patient confidentiality.

The reported incidence of prescription errors in primary care outpatient services

affiliated with an academic medical centre in the United States is 7.6%. [20] In

comparison, the prevalence of prescription errors in this study was 10.9%, focusing

specifically on missed prescriptions or errors in type or dosage. One reason our

percentage was slightly higher than that of the previous study is that our study relied on

patient-reported reflections over a longer recall period, whereas the prior study used

short-term prospective verification of prescription records. [20] Few studies have

comprehensively examined the loss of trust in physicians owing to prescription errors,

including their effects on general trust. Prior research quantified trust in physicians

using vignettes involving adverse medication events; however, that study assessed trust

based on disclosure practices rather than the presence of the error itself. [21] In this

study, the reduction in trust in one's physician due to prescription errors (5.6 points)

was comparable to the impact of misdiagnosis by a different physician (4.3 points). [14]

This finding suggests that prescription errors, which may be harmless, [9] are often an

unexpected experience, causing negative and persistent psychological consequences.

Additionally, our findings on the loss of trust in general physicians stemming from

prescription errors by their personal physicians support the argument that trust in

general physicians is influenced to some extent by individual experiences with their

personal physicians. [4,5] In other words, prescription errors can erode trust not only in

specific physicians but also in the broader healthcare system. Furthermore, the results

highlighted that the loss of trust in general physicians due to prescription errors was

partially mediated by diminished trust in one's own physicians. On the other hand, the

direct loss of trust in general physicians attributable to prescription errors may be

shaped by the societal perceptions of physicians conveyed through the media and other

social channels. [4,5,22] Particularly, exaggerated portrayals of medical errors in

television dramas may contribute to negative public perceptions of healthcare systems.

[22] For example, an analysis of medical errors depicted in six episodes from each of

eight popular U.S. medical television series (1994–2018) revealed that medical errors

occurred at a rate of 6.4 times per hour, with medication errors being the third most

common. [22]

The findings indicate that, among patients with non-communicable diseases

requiring ongoing management, the loss of interpersonal trust resulting from physicians

discussing patient illnesses or personal information in a way that can be overheard by

others, is greater than that caused by prescription errors. The issue of healthcare

professionals discussing patient information in institutional spaces with limited privacy

has been studied in many countries. For instance, a study conducted in five U.S.

hospitals found that approximately 14% of elevator rides involved inappropriate

comments from healthcare professionals. The majority of confidentiality violations,

including graphic discussions of patient identifiers and treatment plans, were attributed

to physicians. [23] However, breaches of confidentiality stemmed from patients'

perceptions of insufficient auditory privacy during interactions with their physicians.

This issue can be attributed not only to physicians' attitudes toward confidentiality but

also to the design of consultation rooms, a problem that may be specific to Japan.

Consultation rooms in Japan are often not fully walled off on all four sides. In particular,

the side opposite the patient entrance is frequently left open to facilitate the movement

of medical staff between consultation rooms. This design flaw allows conversations to

be overheard by patients in adjacent rooms. Similar issues have been reported in a U.S.

emergency room study where patients often overheard discussions about themselves

and others due to insufficient space and inadequate walls on all sides. [24] Another potential source of confidentiality breaches, one not unique to Japan, may involve ward rounds during past hospitalisations, where sensitive patient information was overheard by others in neighbouring beds. [11] Unlike prescription errors, breaches of confidentiality did not significantly affect general trust in physicians in this study. This finding suggests that such breaches are viewed by patients as isolated incidents attributable to specific physicians rather than reflective of the healthcare system as a whole. [12]

The loss of trust in physicians resulting from missed appointment scheduling may be a unique issue in Japan. While follow-up appointments are commonly scheduled through electronic healthcare records in many countries, [25] this task is typically performed by the physicians themselves at the end of consultations in Japan. In contrast, in the U.S. and other countries, appointment scheduling is often handled by a hospital administrators. [10] Although clerical errors leading to missed appointments are occasionally reported globally, the administrative burden on Japanese physicians may contribute to oversights of appointment reservations. Such errors are often discovered only when patients inform their physicians. The magnitude of the loss of trust in personal physicians associated with missed appointments, comparable to that of confidentiality breaches, suggests that patients attribute these administrative errors to their personal physician rather than to systemic issues.

This study has several strengths. First, we quantified the impact of specific

categories of administrative errors on trust in both one's personal physician and general

physicians, offering valuable insights into the dynamics of patient trust within

healthcare settings. Second, by adjusting for patients' general tendency to trust others as

a covariate, the analysis isolated the impact of these errors on trust in physicians

independently of general trustworthiness.

However, this study has certain limitations, which must be acknowledged. First,

the classification of administrative errors relied on participants' self-reports, which may

have introduced misclassification. For instance, confidentiality breaches may have been

underestimated due to the reliance on participants' recollections of past interactions.

Second, the study did not examine the specifics of medical errors, such as the type of

medication error (e.g. dosage exceeding label instructions or misprescription of a

medication with a different indication) or the consequences of these errors (e.g. the

presence, or absence, or severity of adverse events).

In conclusion, this study provides unique insights into how administrative errors,

such as prescription errors, missed appointment scheduling, and confidentiality breaches,

directly and negatively impact the quality of physician-patient interactions. Addressing

these issues is crucial for fostering strong physician-patient relationships and improving

outcomes, particularly in the long-term management of Japanese adults with non-

communicable diseases.

Figure Legend

Fig. 1. Association between Trust in physicians and administrative errors

The general linear model described in the footnotes of Table 2 was fitted to each outcome. Red circles indicate the adjusted mean difference (coefficient) in Trust in Doctors Generally scores associated with administrative errors. The blue square indicates the adjusted mean difference (coefficient) of Interpersonal Trust in Physician scores associated with administrative errors. Error bars indicate 95% confidence intervals.

Fig. 2. Mediating Role of Interpersonal Trust in the Association Between Medication Prescription Errors and General Trust in Doctors

This figure shows the direct and indirect paths between medication prescription errors and general trust in doctors, mediated by interpersonal trust in a physician. Direct effect: –6.75, indirect effect: –3.02 (Sobel test, $p = 0.031$). The total effect of medication errors on general trust is –9.78, with 30.9% of this effect mediated through interpersonal trust in the attending physician.

Ethical Approval: The study protocol was approved by the Ethics Review Board of Okayama University Hospital (No. ken-2405-041).

Author Contributions: Aita, Miyawaki, and Kurita had full access to all data in the study and took responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Aita, Kurita.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Aita, Miyawaki, Kurita.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Aita, Miyawaki, Kurita.

Obtained funding: Kurita

Administrative, technical, or material support: Wakita, Yajima

Supervision: Yajima, Wakita, Gupta

Conflict of Interest Disclosures: Dr. Kurita reported receiving grants from the Japan Society for the Promotion of Science; consulting fees from GlaxoSmithKline K.K.; and payments for speaking at and participating in educational events from Chugai Pharmaceutical Co. Ltd., Sanofi K.K., Mitsubishi Tanabe Pharma Corporation, and Japan College of Rheumatology.

Funding/Support: This study was supported by JSPS KAKENHI, Grant Numbers 19KT0021 (N. K.) and 22K19690 (N. K.).

Role of the Funder/Sponsor: The funder had no role in the study design, analyses,

interpretation of data, writing of the manuscript, or decision to submit the manuscript
for publication.

Data Sharing Statement: The data supporting the findings of this study are available
from the corresponding author upon request.

Additional Contributions: TA, YK, YM, TF, NY, AG, and NK designed and
conceived the study. RS, KS, and NO collected data. TA, YK, YM, AG, and NK
analysed and interpreted the results and drafted the manuscript. All the authors have
read and approved the final version of the manuscript.

Reference

1 Thom DH, Ribisl KM, Stewart AL, et al. Further validation and reliability
testing of the Trust in Physician Scale. The Stanford Trust study physicians. *Med
Care* 1999;37:510–7. doi: 10.1097/00005650-199905000-00010.

2 Thom DH, Kravitz RL, Bell RA, et al. Patient trust in the physician: relationship
to patient requests. *Fam Pract* 2002;19:476–83. doi: 10.1093/fampra/19.5.476.

3 Kurita N, Oguro N, Miyawaki Y, et al. Trust in the attending rheumatologist,
health-related hope and medication adherence among Japanese systemic lupus
erythematosus patients. *Rheumatol (Oxf Engl)* 2023;62:2147–53. doi:
10.1093/rheumatology/keac565.

4 Hall MA, Camacho F, Dugan E, et al. Trust in the medical profession:
conceptual and measurement issues. *Health Serv Res* 2002;37:1419–39. doi:

10.1111/1475-6773.01070.

5 Oguro N, Suzuki R, Yajima N, et al. The impact that family members' health

care experiences have on patients' trust in physicians. *BMC Health Serv Res*

2021;21:1122. doi: 10.1186/s12913-021-07172-y.

6 Southwick FS, Cranley NM, Hallisy JA. A patient-initiated voluntary online

survey of adverse medical events: the perspective of 696 injured patients and

families. *BMJ Qual Saf* 2015;24:620–9. doi: 10.1136/bmjqs-2015-003980.

7 Elder NC, Jacobson CJ, Zink T, et al. How experiencing preventable medical

problems changed patients' interactions with primary health care. *Ann Fam Med*

2005;3:537–44. doi: 10.1370/afm.346.

8 Iedema R, Sorensen R, Manias E, et al. Patients' and family members'

experiences of open disclosure following adverse events. *Int J Qual Health Care*

2008;20:421–32. doi: 10.1093/intqhc/mzn043.

9 Wittich CM, Burkle CM, Lanier WL. Medication errors: an overview for

clinicians. *Mayo Clin Proc* 2014;89:1116–25. doi: 10.1016/j.mayocp.2014.05.007.

10 Lacy NL, Paulman A, Reuter MD, et al. Why we don't come: patient

perceptions on no-shows. *Ann Fam Med* 2004;2:541–5. doi: 10.1370/afm.123.

11 Bourke J, Wessely S. Confidentiality. *BMJ* 2008;336:888–91. doi:

10.1136/bmj.39521.357731.BE.

12 Beltran-Aroca CM, Girela-Lopez E, Collazo-Chao E, et al. Confidentiality

breaches in clinical practice: what happens in hospitals? *BMC Med Ethics*

2016;17:52. doi: 10.1186/s12910-016-0136-y.

13 von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of

Observational Studies in Epidemiology (STROBE) statement: guidelines for

reporting observational studies. *Ann Intern Med* 2007;147:573–7. doi:

10.7326/0003-4819-147-8-200710160-00010.

14 Suzuki R, Yajima N, Sakurai K, et al. Association of patients' past misdiagnosis

experiences with trust in their current physician among Japanese adults. *J Gen*

Intern Med 2022;37:1115–21. doi: 10.1007/s11606-021-06950-y.

15 Dugan E, Trachtenberg F, Hall MA. Development of abbreviated measures to

assess patient trust in a physician, a health insurer, and the medical profession.

BMC Health Serv Res 2005;5:64. doi: 10.1186/1472-6963-5-64.

16 Croker JE, Swancutt DR, Roberts MJ, et al. Factors affecting patients' trust and

confidence in GPs: evidence from the English national GP patient survey. *BMJ*

Open 2013;3:e002762. doi: 10.1136/bmjopen-2013-002762.

17 Yamagishi T, Yamagishi M. Trust and commitment in the United States and

Japan. *Motiv Emot* 1994;18:129–66. doi: 10.1007/BF02249397.

18 Colin Cameron A, Miller DL. A practitioner's guide to cluster-robust inference.

J Hum Resour 2015;50:317–72. doi: 10.3368/jhr.50.2.317.

19 Mize TD. sgmediation2 – Sobel-Goodman tests of mediation in Stata.

https://www.trentonmize.com/software/sgmediation2 (Accessed 8 Aug 2023).

20 Gandhi TK, Weingart SN, Seger AC, et al. Outpatient prescribing errors and the

impact of computerized prescribing. *J Gen Intern Med* 2005;20:837–41. doi:

10.1111/j.1525-1497.2005.0194.x.

21 Mazor KM, Reed GW, Yood RA, et al. Disclosure of medical errors: what

factors influence how patients respond? *J Gen Intern Med* 2006;21:704–10. doi:

10.1111/j.1525-1497.2006.00465.x.

22 Carney M, King TS, Yumen A, et al. The depiction of medical errors in a

sample of medical television shows. *Cureus* 2020;12:e11994. doi:

10.7759/cureus.11994.

23 Ubel PA, Zell MM, Miller DJ, et al. Elevator talk: observational study of

inappropriate comments in a public space. *Am J Med* 1995;99:190–4. doi:

10.1016/s0002-9343(99)80139-9.

24 Olsen JC, Cutcliffe B, O'Brien BC. Emergency department design and patient

perceptions of privacy and confidentiality. *J Emerg Med* 2008;35:317–20. doi:

10.1016/j.jemermed.2007.10.029.

25 Alawadhi A, Palin V, van Staa T. Investigating the reasons for missing an

outpatient appointment in Royal Hospital, Sultanate and Oman: perspectives of

patients and medical staff in a survey. *Health Sci Rep* 2022;5:e470. doi:

10.1002/hsr2.470.

Table 1. Participants' characteristics

Baseline Characteristics	Total (N = 661)
Age, years	63.0 (56.0–70.0)
Male sex	486 (73.5%)
Education	
Junior high school	19 (2.9%)
High school	209 (31.6%)
Junior college	65 (9.8%)
University	325 (49.2%)
Graduate school	30 (4.5%)
Not answered	13 (2.0%)
Household income	
<1,000,000 yen	40 (6.1%)
1,000,000 – < 3,000,000 yen	157 (23.8%)
3,000,000 – < 5,000,000 yen	203 (30.7%)
5,000,000 – < 10,000,000 yen	205 (31.0%)
≥ 10,000,000 yen	56 (8.5%)
Patient-physician relationship duration	
<1 years	60 (9.1%)
1 – < 3 years	212 (32.1%)
≥ 3 years	389 (58.9%)
General level of interpersonal trust	54.2 (45.8–70.8)
Reported diseases	
Arrhythmia	37 (5.6%)
Ischemic heart disease	119 (18.0%)
Heart failure	15 (2.3%)
Diabetes mellitus	191 (28.9%)
Rheumatoid arthritis/Systemic lupus erythematosus	17 (2.6%)
Cancer	255 (38.6%)
Depression	127 (19.2%)
The most troublesome disease	
Arrhythmia	17 (2.6%)
Ischemic heart disease	89 (13.5%)
Heart failure	8 (1.2%)
Diabetes mellitus	175 (26.5%)
Rheumatoid arthritis/Systemic lupus erythematosus	13 (2.0%)

Cancer	242 (36.6%)
Depression	117 (17.7%)

83 Note*:* Values are presented as medians (IQR) for continuous variables and n (%) for categorical variables.

Table 2. Association of administrative errors with trust in doctors generally and interpersonal trust in a physician

	Interpersonal trust in a physician		Trust in doctors generally	
	Coefficient (95% CI)	P-value	Coefficient (95% CI)	P-value
Prescription error	–5.56 (–10.70 to –0.41)	0.035	–9.78 (–13.74 to –5.81)	<0.001
Appointment reservation error	–13.56 (–22.48 to –4.65)	0.004	–6.12 (–12.59 to 0.35)	0.063
Confidentiality breach	–14.09 (–24.35 to –3.83)	0.008	–5.23 (–11.33 to 0.88)	0.091

Note*:* The models were adjusted for baseline characteristics such as age, sex, final education, household income, duration of the patient-physician relationship, reported comorbidities, and patient's general interpersonal trust level. Standard errors were estimated using cluster-robust variance, assuming that the prefectures were cluster units. Abbreviations: CI, confidence interval.