Stimulant medication use and development of the dopamine system: a naturalistic long-term follow-up of boys and men with ADHD

Zarah van der Pal[1]* MSc, Liesbeth Reneman[1] PhD, Henk JMM Mutsaerts[2] MD PhD, Antonia Kaiser[3,4] PhD, Marco A Bottelier[5] MD PhD, Hilde M Geurts[6] PhD, Anouk Schrantee[1] PhD

1. Department of Radiology & Nuclear Medicine, Amsterdam UMC location University of Amsterdam, Amsterdam, Netherlands
2. Department of Radiology & Nuclear Medicine, Amsterdam UMC location Vrije Universiteit, Amsterdam, Netherlands
3. CIBM Center for Biomedical Imaging, Lausanne, Switzerland
4. Animal Imaging and Technology, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland
5. Accare, Centre for Academic Child and Adolescent Psychiatry, UMC Groningen, Groningen, Netherlands
6. Dutch Autism and ADHD Research Center, Amsterdam Brain and Cognition, University of Amsterdam, Amsterdam, Netherlands

* Corresponding author: address: Meibergdreef 9 (Room Z0-174), 1105 AZ, Amsterdam, The Netherlands; phone: +31205661644; email: z.vanderpal@amsterdamumc.nl

Competing interests: The authors have nothing to disclose.

Abstract

Stimulant medication is commonly used by children and adolescents with attention-deficit/hyperactivity disorder (ADHD), however its long-lasting effects on the developing brain remain unclear. In our previous ePOD-methylphenidate randomised controlled trial (RCT) we found that short-term stimulant treatment influences development of the brain's dopamine system in an age-dependent manner, in line with animal studies suggesting persisting effects on brain development into adulthood. In this 4-year naturalistic follow-up of the initial ePOD-methylphenidate RCT, we investigated the long-term age-dependent effects of stimulant treatment on development of the dopamine system in children and adults with ADHD (n=56; adolescents aged 10-17 years, adults aged 23-43 years). At baseline and 4-year follow-up, we assessed estimates of dopamine-reactivity using pharmacological MRI. We estimated relative CBF (rCBF) before a single-dose methylphenidate-challenge (resting rCBF) and the rCBF-response to a single-dose methylphenidate-challenge. Linear mixed models were constructed to evaluate the effect of stimulant medication use, age and visit on resting rCBF and rCBF-response. We found no evidence for long-term age-dependent effects of stimulant treatment on development of the dopamine system, suggesting that our previously identified short-term effects may be transient. We did identify age-dependent associations between estimates of dopamine function in the medial prefrontal cortex and stimulant treatment. This was unrelated to ADHD symptom severity and may hold predictive value for (extent of) stimulant medication use after ADHD diagnosis in children and adolescents.

Introduction

Stimulant medication, such as methylphenidate, belongs to the largest class of psychotropic medications prescribed for the treatment of attention-deficit/hyperactivity disorder (ADHD). Methylphenidate blocks the dopamine and noradrenaline transporters in the brain, leading to an increase in extracellular dopamine and noradrenaline [1, 2]. Methylphenidate's short-term safety has been documented in many studies and its efficacy is among the highest of psychiatric medications. However, despite its widespread and prolonged use, surprisingly little is known on the long-term effects of stimulant medication on the human brain in development.

During adolescence, the brain undergoes rapid development, showcasing specific sensitive periods [3] which render the brain particularly susceptible to interventions, like stimulant medications, which might alter these crucial developmental processes. For example, the ratio of excitatory D1/5 receptors and inhibitory D2/3/4 receptor expression changes throughout development [4-6]. Indeed, animal studies suggest that exposure to stimulant medication at different developmental stages affects the outgrowth of the dopamine system, with long-lasting consequences [7, 8]. As such, administering methylphenidate during crucial developmental periods may imprint enduring effects on brain maturation ('the neurochemical imprinting hypothesis' [9]), significantly manifesting as individuals approach adulthood.

In humans, studies investigating the effect of stimulant medication on the development of the dopamine and noradrenaline systems remain sparse, and are hampered by the in-accessibility of these systems without using invasive (i.e. radioactive) imaging tracers. In addition, these studies exhibit a lack of convergent findings [10-14], accompanied by methodological challenges such as the absence of longitudinal treatment study designs. To this purpose, we previously validated the non-invasive method 'pharmacological MRI' to evaluate estimates of dopamine function, in which alterations in resting cerebral blood flow (CBF) in response to an acute challenge with methylphenidate are assessed (i.e. *dopamine-reactivity*) [15-17]. We subsequently applied this technique in a prospective longitudinal randomised controlled trial (RCT) aimed at probing the effects of methylphenidate treatment on the development of the dopamine and noradrenaline systems in the brain. In this RCT ("effects of Psychotropic drugs On the Developing brain" (ePOD) Methylphenidate project, NTR3103), we showed that 4-month treatment with methylphenidate, compared to placebo, decreased the CBF-response to an acute challenge with methylphenidate after one-week washout in stimulant treatment-naive children, but not adults, with ADHD [18]. These findings suggest *short-term* lasting age-dependent effects of methylphenidate treatment on dopamine-reactivity. However, in a related cross-sectional study exploring the *long-term* effects of age-of-first-stimulant treatment on the brain's response to methylphenidate, we observed no differences in dopamine-reactivity between participants that initiated stimulant treatment during childhood and those that initiated stimulant treatment during adulthood, and stimulant treatment-naive participants [19]. However, the group exposed to stimulants during childhood exhibited lower baseline CBF in the anterior cingulate cortex, compared with the group that had not received stimulant treatment. This discrepancy could signify early-induced alterations by stimulant medication on the developing dopamine and noradrenaline systems.

To further investigate the *long-term* effects of stimulant treatment, we here conducted a 4-year naturalistic follow-up of participants from the initial ePOD-methylphenidate RCT, assessing alterations in resting CBF and response to an acute challenge with methylphenidate (dopamine-reactivity). We hypothesised that stimulant treatment would induce persistent long-term age-dependent effects on development of the dopamine and noradrenaline systems in the brain. Specifically, we hypothesised that higher stimulant medication use during the naturalistic follow-up would be associated with lower dopamine-reactivity in adolescents, but not adults. Moreover, we expected that the age-dependent effects previously identified in our RCT [18] could still be identified after 4 years; in line with the neurochemical imprinting hypothesis which states that neurochemical imprinting effects increase with age in preclinical studies [9, 20, 21].

Methods

Study design and participants

The present study constitutes the naturalistic 4-year follow-up assessment of the ePOD-methylphenidate RCT (2016-2019 [22]). The initial RCT was a 16-week double-blind, randomised, placebo-controlled trial of methylphenidate treatment (2011-2015) in stimulant treatment-naive participants with ADHD [18, 23]. The ePOD-methylphenidate RCT protocol was approved by the Central Committee on Research Involving Human Subjects (identifier NL34509.000.10). The 4-year follow-up assessment was approved by the local medical ethical committee of the Academic Medical Center (NL54972.018.15). All participants and parents or legal representatives of the children provided written informed consent.

In the original RCT, 50 boys (aged 10-12 years) and 49 men (aged 23-30 years) with ADHD were included. An experienced psychiatrist diagnosed all participants according to the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV, 4th edition [24]) using a structured interview (Diagnostic Interview Schedule for Children fourth edition, DISC-IV [25]) in children or in parents and the Diagnostic Interview for Adult ADHD [26] in adults (for details on recruitment and exclusion criteria, see Supplementary Materials). For the 4-year naturalistic follow-up assessment, participants were contacted by phone and/or email to ask if they wanted to participate in the follow-up assessment. Exclusion criteria were contraindications to MRI.

The present study reports on the primary outcome of the long-term ePOD-methylphenidate project, namely stimulant medication effects on dopamine-reactivity assessed using pharmacological MRI (i.e. CBF-response to an acute challenge with methylphenidate). At baseline and 4-year follow-up, MR scanning was performed before and 90 minutes after a single-dose methylphenidate-challenge (0.5 mg/kg, with a maximum of 20 mg for adolescents and a maximum of 40 mg for adults) (Figure 1A). To exclude possible acute effects of stimulant medication use (half-life: 2-10 hours), participants were asked to stop using stimulant medication one week prior to the follow-up assessment (one-week washout period). A total of 33 adolescents and 25 adults returned for the 4-year follow-up assessment, for details see the results section.

Figure 1. Study design and regions of interest (ROIs). A) At baseline and 4-year follow-up assessment, participants underwent MR scanning before and 90 minutes after a single-dose short-acting methylphenidate-challenge (0.5mg/kg; max. 20 mg for adolescents, max. 40 mg for adults) to evaluate dopamine-reactivity. Stimulant medication use between baseline and follow-up (cumulative dose/exposure duration) was calculated from medication received during the randomised controlled trial (RCT) and pharmacy prescription information from the naturalistic follow-up. B) ROIs: anterior cingulate cortex (ACC), medial prefrontal cortex (mPFC), striatum and thalamus.

Stimulant medication use

We calculated stimulant medication use per participant based on medication received during the initial trial and medication prescription information between the trial and 4-year follow-up assessment obtained from participant's pharmacies. The medication use variables cumulative dose (mg) and exposure duration (months) were extracted. Moreover, we determined age at start of medication use (years), stimulant medication use at follow-up (yes/no) and stimulant treatment-naivety at follow-up (yes/no) (for details, see Supplementary Materials).

ADHD symptom severity

At baseline and 4-year follow-up, self-reported ADHD symptom scores were assessed using the inattentive and hyperactivity/impulsivity subscales of the Disruptive Behavior Disorder Rating-Scale (DBD-RS [27]) in adolescents, and the ADHD-Rating Scale (ADHD-RS [26]) in adults. Although these scales are primarily used as continuous measures of ADHD symptom severity

and are insufficient for establishing a clinical diagnosis, we here include the cut-off values to provide a broader context for interpretation. Clinically significant symptoms were defined as scores >15 on the DBD-RS inattentive and hyperactive/impulsive subscales for adolescents [28] and scores >10 on the ADHD-RS for adults [26].

MRI - acquisition

MR data were acquired using a 3T MR scanner (baseline: Intera or Achieva, follow-up: Ingenia; Philips Medical Systems, Best, The Netherlands) using an 8-channel receive-only head coil. A pseudo-continuous arterial spin labelling (PCASL) sequence with a gradient-echo echo-planar imaging readout was acquired before (pre-MPH) and after (post-MPH) the methylphenidate-challenge (parameters: TR/TE=4000/14 ms; post-label delay range=1525-2143 ms; label duration=1650 ms; FOV=240x240x119 mm; 75 dynamics; voxel size=3x3x7 mm; 17 contiguous slices; no background suppression). Moreover, we acquired a 3D T1-weighted scan (parameters: TR/TE=9.8/4.6 ms; FOV=256x256x120; voxel size=0.875x0.875x1.2 mm; 120 slices) for registration purposes. Heart rate was measured using a peripheral pulse unit.

MRI - processing

Image processing and CBF quantification were performed using the ExploreASL pipeline (v1.12.0 Beta [29]) in Matlab (vR2022b). Briefly, the T1-weighted images were segmented into grey matter (GM), white matter and cerebrospinal fluid using the Computational Anatomy Toolbox 12 [30]. Next, for the ASL time series, motion artefacts were detected using motion estimation, and a fixed spike exclusion threshold (1mm) was applied for outlier exclusion. CBF (mL/min/100g) was calculated with the recommended single-compartment model from the perfusion-weighted and M0 images [31]. Since no background suppression was used, the mean of the control images was used as M0 image. Rigid-body registration to the corresponding T1-weighted image was performed for all intermediate and final images, followed by nonlinear registration to MNI space. Finally, visual quality control was performed [29].

For two adolescents, background suppression was used for the ASL sequences at baseline. For four adolescents, at 4-year follow-up, the ASL MR data were exported as perfusion images instead of raw unsubtracted ASL images. Therefore, image processing was adjusted for these six participants (for details, see Supplementary Materials).

Median CBF (mL/min/100g) was estimated for global GM and four predefined regions of interest (ROIs): anterior cingulate cortex (ACC), medial prefrontal cortex (mPFC), striatum and thalamus (Figure 1B; for the Brainnetome parcellation numbers, see Supplementary Table 1). The striatum was selected because it is rich in dopamine transporters, which are targeted by stimulant medication such as methylphenidate. The choice for the ACC, mPFC, and thalamus was based on animal literature demonstrating the largest age-dependent effects of methylphenidate in these important projections of the striatum [7, 8]. Moreover, these four ROIs were assessed in previous short-term studies in the same sample [18, 32].

All analyses were conducted using R v4.4.1 (R Development Core Team, 2011). We calculated CBF for each ROI relative to global CBF (i.e. relative CBF, rCBF), to account for large physiological, vascular, and scanner variability in global CBF (Chen et al., 2011). The rCBF-response to an acute methylphenidate-challenge was defined as *(post-MPH - pre-MPH)/pre-MPH*.

Linear mixed effect models (LMMs; *lme4* package version 1.1-35-5) were used to assess the effect of stimulant medication use (cumulative dose, exposure duration), visit (baseline, follow-up), age group (adolescents, adults) on i) rCBF before a single-dose methylphenidate-challenge (resting rCBF), and ii) the rCBF-response to a single-dose methylphenidate-challenge (rCBF-response). Moreover, we evaluated whether the short-term effects identified in the initial RCT were still present at follow-up using LMMs including the treatment group during the RCT (methylphenidate, placebo) instead of continuous medication use variables as predictors. In all models, we applied multiple comparison correction (FDR=5%) to adjust for the four ROIs evaluated.

The following variables were included as covariates in all models: MR scanner at baseline (Intera/Achieva), standardised scan interval (offset 48 months, corresponding to a 4-year follow-up), and demeaned age at baseline (per age group; to correct for the larger age range at baseline among adults than adolescents). Moreover, mean motion (mm; in the pre-MPH rCBF analysis) and change in mean motion (mm, calculated as *post-MPH - pre-MPH*; in the rCBF-response analysis) were included as covariates in all models. In addition, we explored associations between rCBF, motion and heart rate. Finally, for ROIs showing significant age-dependent medication effects on resting rCBF and rCBF-response, we explored correlations between rCBF and ADHD symptom severity.

Exploratory whole-brain analyses

To evaluate whether stimulant treatment impacted additional brain regions beyond the four ROIs evaluated in the main analysis, we performed an exploratory whole-brain analysis evaluating the relation between stimulant medication use and (change in) CBF-response to methylphenidate. In addition, we explored the association between the whole-brain CBF-response and the spatial distribution of dopamine and noradrenaline receptors to identify which receptor systems may be mediating the observed response (for details, see Supplementary Materials).

Results

Participant characteristics

Of the 50 children and 49 adults who participated in the initial trial, 33 children and 25 adults returned for the 4-year follow-up assessment. Baseline characteristics were comparable for participants that returned and did not return for follow-up (Supplementary Table 2). One adolescent and one adult were excluded from analysis due to incomplete MRI data and undisclosed stimulant medication use before baseline, respectively. The final sample consisted of 32 adolescents (aged 11.2±0.9 years at baseline) and 24 adults (aged 29.9±5.0 years at

baseline) (Table 1). Post-MPH MRI data was missing for 3 adolescents and 7 MRI data points were excluded from analysis due to insufficient data quality or high motion (>75 excluded frames). This resulted in inclusion of 53 participants at baseline (30 adolescents, 23 adults) and 54 participants at follow-up (30 adolescents, 24 adults) for the resting rCBF analysis, and 50 participants at baseline (27 adolescents, 23 adults) and 52 participants at follow-up (28 adolescents, 24 adults) for the rCBF-response analysis.

Exposure duration to stimulant medication was higher in adolescents compared with adults (W=531, P=.02), whereas cumulative dose did not differ between age groups (W=456, P=.24; Table 1). Two adult participants were prescribed atomoxetine, a non-stimulant medication for treatment of ADHD, in addition to stimulant medication.

<u>Stimulant treatment was not significantly associated with change in rCBF over time</u>
In contrast to our hypothesis, we found no evidence for an association between stimulant medication use and change in resting rCBF or rCBF-response over time, in either age group or any of the ROIs (*age*medication*visit interaction*; all P>.10; Table 2). The rCBF-response did not change significantly between baseline and follow-up in either age group or any ROI (*age*visit interaction*; all P>.10) (Table 2B).

Resting rCBF in the mPFC was consistently higher in adults than adolescents at both visits (*age effect*; all P<.046). Moreover, thalamic resting rCBF showed a decrease between baseline and follow-up, in both age groups (*visit effect*; all P<.019) (Table 2A).

<u>Extent of stimulant use was significantly associated with resting rCBF and rCBF-response in an age-dependent manner</u>
Across visits, thalamic resting rCBF was significantly associated with stimulant medication use in an age-dependent manner (cumulative dose: $F(1,51)=7.7$, P=.030; exposure duration: $F(1,51)=10.1$, P=.010) (*age*medication interaction*; Table 2A). Post-hoc analysis revealed that adults (but not adolescents) taking more stimulant medication, and having a longer exposure duration, showed lower thalamic resting rCBF (Figure 2A). Moreover, across age groups and visits, we found a significant association between stimulant medication use and resting rCBF in the other ROIs (*medication effect*; ACC, mPFC, striatum) (Table 2A), such that participants taking more stimulant medication, and having a longer exposure duration, showed higher resting rCBF.

Additionally, rCBF-response in the mPFC was significantly associated with stimulant medication use in an age-dependent manner (cumulative dose: $F(1,52)=10.9$, P=.007; exposure duration: $F(1,52)=11.2$, P=.006) (*age*medication interaction*; Table 2B), such that adolescents (but not adults) taking more stimulant medication, and having a longer exposure duration, showed a lower rCBF-response to an acute methylphenidate-challenge (Figure 2B). No such effects were observed in the other ROIs (Table 2B).

Figure 2. Age*medication interaction effects on resting rCBF and rCBF-response to a methylphenidate-challenge. A) Thalamic resting rCBF was associated with medication use in adults, but not in adolescents. B) mPFC rCBF-response was associated with medication use in adolescents, but not in adults. The plots show the average rCBF values across visits, as the age-by-medication interaction effects are independent of time. mPFC=medial prefrontal cortex; rCBF=relative cerebral blood flow.

<u>Sensitivity analyses reveal robustness of the findings</u>
We found no evidence that initial treatment allocation during the ePOD-methylphenidate RCT (i.e.
methylphenidate or placebo arm) had long-lasting effects on resting rCBF or rCBF-response to a
methylphenidate-challenge (Supplementary Table 3). Absolute CBF values were higher in
adolescents than adults and reduced between baseline and follow-up in adolescents only
(Supplementary Table 4). Moreover, we found no associations between (change in) ADHD
symptom severity and (change in) resting rCBF or rCBF-response (Supplementary Table 5; linear
Pearson correlations). Heart rate increased after the methylphenidate-challenge, but this was not
associated with rCBF-response (Supplementary Tables 6-7). Motion was higher in adolescents
than adults, and reduced after a single-dose methylphenidate-challenge in adolescents only
(Supplementary Table 8). (Change in) motion was a significant predictor in the thalamic resting
rCBF and mPFC rCBF-response models (Table 2), but results were robust with and without
motion in these models. We found no associations between rCBF and motion in any of the other
models (Supplementary Table 9).
15
<u>Whole-brain CBF-response was significantly associated with spatial distribution of noradrenaline</u>
<u>transporters</u>
We conducted an exploratory whole-brain analysis to evaluate whether stimulant treatment
impacted additional brain regions beyond the four ROIs evaluated in the main analysis. In both
age groups, we found no additional associations between stimulant medication use and (change
in) CBF-response to methylphenidate. Furthermore, exploratory analysis of the association
between the whole-brain CBF-response and the spatial distribution of dopamine and
noradrenaline receptors revealed that the CBF-response maps were significantly associated with
noradrenaline transporter (NAT) distribution in both adolescents and adults. Although the
dopamine transporter (DAT) and dopamine D1 and D2/3 receptors are also targets of
methylphenidate, we found no evidence for associations of these receptors' distribution with
whole-brain CBF-response (Table 3).
28
29 **Discussion**
30
In this 4-year naturalistic follow-up study using pharmacological MRI, we found no evidence for
long-term age-dependent effects of stimulant treatment on development of resting CBF and CBF-
response to an acute challenge with methylphenidate (dopamine-reactivity) in adolescents when
compared to adults. This contrasts with our previous short-term findings in the initial RCT [18].
Nevertheless, our findings revealed that thalamic resting rCBF and the mPFC rCBF-response to
a methylphenidate-challenge were associated with the extent of stimulant treatment during the 4-
year follow-up period in an age-dependent manner. These associations were already present
before the commencement of stimulant treatment, but were unrelated to ADHD symptom severity.
Finally, we found that CBF-response was associated with NAT distributions, but not dopamine
receptor/transporter distributions.
41
42
43

No evidence for long-term age-dependent effects of stimulant treatment on development of the dopamine system

Guided by animal studies and the neurochemical imprinting hypothesis, which suggests that stimulant administration during development has lasting effects that increase with age [9, 20, 21], we hypothesised that the effects we identified in the initial ePOD-methylphenidate RCT [18] would persist long-term. In our previous RCT in the same study population, children receiving 4-month methylphenidate treatment showed reduced CBF responsiveness to a methylphenidate-challenge in the thalamus and striatum, unlike adults or placebo groups. However, our current long-term follow-up data from the same cohort reveal no age-dependent changes over time in CBF-responses, nor dose-dependent stimulant treatment effects.

The discrepancy between our short-term findings in the initial RCT and this long-term follow-up may have several explanations. First, our previously identified short-term effects may be transient due to neuroplasticity - the brain's ability to adapt and reorganise in response to internal and external stimuli [33]. Second, heterogeneous medication use and treatment adherence during the naturalistic follow-up, in contrast to the RCT, may have resulted in more subtle effects than we could detect [34, 35]. Additionally, the reduced sample size due to participant attrition may have limited our statistical power to identify such differences. Alternatively, potential stimulant medication effects might become apparent after a longer follow-up period, when the adolescents have reached young adulthood and the neurochemical imprinting effects are suggested to be most pronounced.

Notably, the existing literature presents mixed findings regarding long-term effects of stimulant treatment on brain development. Some studies reported that participants taking stimulant medication were similar to neurotypical controls regarding cortical thickness [36] and brain functioning (at rest [37] or during an attentional task [38]), whereas other studies reported no associations between stimulant treatment and brain development [22, 39, 40] or inconclusive results [14]. Besides methodological heterogeneity and reporting issues, the lack of convergence between these studies may be related to factors like biological factors/genetic predispositions, sample age and behavioural characteristics (for an extensive discussion, see [14, 41]).

Age-dependent associations between stimulant treatment and rCBF

Exploring the age-dependent associations between rCBF at baseline and subsequent stimulant medication use, we found that this relation was present in adults only for thalamic resting rCBF, and in adolescents only for the mPFC rCBF-response to a methylphenidate-challenge. Notably, these associations remained present also at 4-year follow-up.

Adults with higher thalamic resting rCBF were prescribed less stimulant medication for shorter durations in the years following treatment initiation. That we found this relation solely in adults may be related to the gradual decline in dopamine D2/3 receptors in the thalamus in adulthood [42, 43]. The thalamus plays an important role in attention and cognitive functioning, processes that are commonly affected in ADHD and can effectively be improved with stimulant treatment [38, 44, 45]. Individual differences in thalamic function (e.g. perfusion) may become apparent in adulthood, which may be associated with efficacy of stimulant treatment, and subsequently with the extent of medication use.

Adolescents that exhibited a higher (more positive) rCBF-response to the methylphenidate-challenge at baseline were prescribed less medication for shorter durations,

whereas adolescents with lower responsiveness, or even a negative rCBF-response, were prescribed more medication for a longer duration. This finding aligns with the significant maturation of the prefrontal cortex during adolescence, where dopamine D1 and D2 receptor densities peak and then decline into adulthood [5, 46, 47]. The heightened rCBF-response observed in adolescents compared to adults could be attributed to the relatively high density of D1 receptors in the mPFC [6, 48]. Notably, D1 receptors are activated only at higher dopamine levels, and the relatively high dose of methylphenidate used during the challenge may have specifically enhanced D1 receptor activity in the mPFC.

Both the thalamus and mPFC also exhibit a high density of NAT [48], another important target of methylphenidate. Interestingly, we observed that the whole-brain CBF-response was associated with NAT distribution, but this relation appeared to be similar across age groups. While this whole-brain approach provides valuable neurobiological insights into the mechanisms of methylphenidate action, it does not capture inter-individual variability in receptor distributions. Understanding how receptor distributions vary across individuals—and how this variability influences brain function—could provide crucial insights into the observed differences in medication responses and usage patterns.

Different adolescent neural trajectories may also be associated with varying ADHD phenotypes [49]. In our study, the variation in mPFC dopamine-reactivity was not *linearly* associated with (changes in) self-reported severity of ADHD symptoms. Nonetheless, ADHD symptoms have been shown to follow *non-linear* trajectories across the lifespan; for example, both persistent, remittent and subthreshold symptom trajectories have been described [50, 51]. This may explain the lack of correlation between ADHD symptoms and mPFC dopamine-reactivity in adolescents, but also the differences observed between the adolescent and adult group. Given our study's design, in which both adolescents and adults were stimulant treatment-naive at baseline, it is possible that individuals diagnosed with ADHD in adulthood may have experienced subthreshold ADHD symptoms during childhood and adolescence, progressing to diagnosis in adulthood, thus constituting an ADHD subpopulation [50]. Moreover, reported distinct genetic variations and risk factors between childhood and adult ADHD diagnoses may contribute to diverse patterns of mPFC function [52].

Methodological considerations

A critical strength of this study is its longitudinal design with stimulant treatment-naive participants at baseline, ruling out the influence of prior stimulant treatment on brain development. Additionally, the 4-year follow-up period addresses the need to investigate the potentially long-lasting effects of stimulant treatment. Stimulant medication is commonly prescribed to children and adolescents for extended periods; however, neuroimaging studies with long-term follow-up are scarce, and those that do exist often do not include stimulant treatment-naive participants (e.g. [36] and [37]). Furthermore, the use of pharmacological MRI provided insights into how stimulant medication affects dopamine function over time and presents an interesting technique for subtyping and stratification, contributing to a better understanding of the neural mechanisms that influence treatment patterns.

Our findings should be interpreted in light of several methodological considerations. First, although pharmacy prescription data provides more reliable medication use information during the naturalistic follow-up compared with participant self-report, we assumed complete prescription

adherence for all participants (aligning with e.g. [53-55]). However, medication adherence rates can vary substantially [34, 35], necessitating the inclusion of reliable treatment adherence measures in future research [56]. Moreover, while this study focused on males with ADHD [23], future studies should investigate medication effects in females as well [57]. Other limitations include the restricted sample size at follow-up due to attrition and the use of different MR scanners at baseline and follow-up. Still, we were able to identify an age-dependent association between mPFC dopamine-reactivity and stimulant treatment, which may hold predictive value for optimisation of treatment approaches for ADHD. Future studies should explore whether long-term effects of stimulant treatment on brain development become evident in young adulthood, when such effects are suggested to be most pronounced [9].

Conclusion

In conclusion, this study found no evidence for long-lasting effects of stimulant treatment on development of the dopamine system. We did identify an age-dependent association between estimates of dopamine function in the mPFC and extent of stimulant treatment, which may hold predictive value for (extent of) stimulant medication use after ADHD diagnosis in children and adolescents. Our findings highlight the value of including stimulant treatment-naive participants and using pharmacological MRI to investigate functioning of the dopamine system in the brain. Large-scale longitudinal studies are encouraged to further explore the potential of pharmacological MRI for optimisation of treatment approaches in ADHD, and disentangle the underlying (neurobiological) processes.

Data availability statement

References

1. Hannestad, J., et al., *Clinically relevant doses of methylphenidate significantly occupy norepinephrine transporters in humans in vivo.* Biol Psychiatry, 2010. **68**(9): p. 854-60.
2. Volkow, N.D., et al., *Mechanism of action of methylphenidate: insights from PET imaging studies.* J Atten Disord, 2002. **6 Suppl 1**: p. S31-43.
3. Wahlstrom, D., T. White, and M. Luciana, *Neurobehavioral evidence for changes in dopamine system activity during adolescence.* Neurosci Biobehav Rev, 2010. **34**(5): p. 631-48.
4. Chen, Y.I., et al., *Pharmacologic neuroimaging of the ontogeny of dopamine receptor function.* Dev Neurosci, 2010. **32**(2): p. 125-38.
5. Jucaite, A., et al., *Age-related reduction in dopamine D1 receptors in the human brain: from late childhood to adulthood, a positron emission tomography study.* Neuroscience, 2010. **167**(1): p. 104-10.
6. Manza, P., et al., *Cortical D1 and D2 dopamine receptor availability modulate methylphenidate-induced changes in brain activity and functional connectivity.* Commun Biol, 2022. **5**(1): p. 514.
7. Andersen, S.L., *Stimulants and the developing brain.* Trends Pharmacol Sci, 2005. **26**(5): p. 237-43.

8. Andersen, S.L., et al., *Juvenile methylphenidate modulates reward-related behaviors and cerebral blood flow by decreasing cortical D3 receptors.* Eur J Neurosci, 2008. **27**(11): p. 2962-72.

9. Andersen, S.L. and C.P. Navalta, *Altering the course of neurodevelopment: a framework for understanding the enduring effects of psychotropic drugs.* Int J Dev Neurosci, 2004. **22**(5-6): p. 423-40.

10. Szobot, C.M., et al., *The acute effect of methylphenidate on cerebral blood flow in boys with attention-deficit/hyperactivity disorder.* Eur J Nucl Med Mol Imaging, 2003. **30**(3): p. 423-6.

11. Langleben, D.D., et al., *Effects of methylphenidate discontinuation on cerebral blood flow in prepubescent boys with attention deficit hyperactivity disorder.* J Nucl Med, 2002. **43**(12): p. 1624-9.

12. Schweitzer, J.B., et al., *A positron emission tomography study of methylphenidate in adults with ADHD: alterations in resting blood flow and predicting treatment response.* Neuropsychopharmacology, 2003. **28**(5): p. 967-73.

13. Ulrich, M., et al., *Methylphenidate Differentially Affects Intrinsic Functional Connectivity of the Salience Network in Adult ADHD Treatment Responders and Non-Responders.* Biology (Basel), 2022. **11**(9).

14. Pereira-Sanchez, V., et al., *Systematic Review: Medication Effects on Brain Intrinsic Functional Connectivity in Patients With Attention-Deficit/Hyperactivity Disorder.* J Am Acad Child Adolesc Psychiatry, 2021. **60**(2): p. 222-235.

15. Jenkins, B.G., *Pharmacologic magnetic resonance imaging (phMRI): imaging drug action in the brain.* Neuroimage, 2012. **62**(2): p. 1072-85.

16. Schrantee, A., et al., *Dopaminergic system dysfunction in recreational dexamphetamine users.* Neuropsychopharmacology, 2015. **40**(5): p. 1172-80.

17. Reneman, L., et al., *Imaging of the dopamine system with focus on pharmacological MRI and neuromelanin imaging.* Eur J Radiol, 2021. **140**: p. 109752.

18. Schrantee, A., et al., *Age-Dependent Effects of Methylphenidate on the Human Dopaminergic System in Young vs Adult Patients With Attention-Deficit/Hyperactivity Disorder: A Randomized Clinical Trial.* JAMA Psychiatry, 2016. **73**(9): p. 955-62.

19. Schrantee, A., et al., *Long-term effects of stimulant exposure on cerebral blood flow response to methylphenidate and behavior in attention-deficit hyperactivity disorder.* Brain Imaging Behav, 2018. **12**(2): p. 402-410.

20. Moll, G.H., et al., *Early methylphenidate administration to young rats causes a persistent reduction in the density of striatal dopamine transporters.* J Child Adolesc Psychopharmacol, 2001. **11**(1): p. 15-24.

21. Urban, K.R., B.D. Waterhouse, and W.-J. Gao, *Distinct Age-Dependent Effects of Methylphenidate on Developing and Adult Prefrontal Neurons.* Biological Psychiatry, 2012. **72**(10): p. 880-888.

22. van der Pal, Z., et al., *Stimulant medication use and apparent cortical thickness development in attention-deficit/hyperactivity disorder: a prospective longitudinal study.* Front Psychiatry, 2024. **15**: p. 1365159.

23. Bottelier, M.A., et al., *The effects of Psychotropic drugs On Developing brain (ePOD) study: methods and design.* BMC Psychiatry, 2014. **14**: p. 48.

24. American Psychiatric Association, *Diagnostic and statistical manual of mental disorders, 4th ed.* Diagnostic and statistical manual of mental disorders, 4th ed. 1994, Arlington, VA, US: American Psychiatric Publishing, Inc.

25. Ferdinand, R. and J.v.d. Ende, *DISC-IV Diagnostic Interview Schedule for Children [Dutch Translation NIMH-DISC-IV]. Afdeling Kinder- en Jeugdpsychiatrie, Sophia Kinderziekenhuis/Academisch Ziekenhuis Rotterdam, Rotterdam, The Netherlands.* 1998.

26. Kooij, J.J.S., *Adult ADHD: Diagnostic Assessment and Treatment*, in *Adult ADHD*. 2013, Springer London: London. p. 97-99.

27. Pelham, W.E., Jr., et al., *Teacher ratings of DSM-III-R symptoms for the disruptive behavior disorders.* J Am Acad Child Adolesc Psychiatry, 1992. **31**(2): p. 210-8.

28. Oosterlaan, J., et al., *VvGK6-16: Vragenlijst voor Gedragsproblemen bij Kinderen.* 2008: Pearson.

29. Mutsaerts, H., et al., *ExploreASL: An image processing pipeline for multi-center ASL perfusion MRI studies.* Neuroimage, 2020. **219**: p. 117031.

30. Gaser, C., *Partial Volume Segmentation with Adaptive Maximum A Posteriori (MAP) Approach.* NeuroImage, 2009. **47**: p. S121.

31. Alsop, D.C., et al., *Recommended implementation of arterial spin-labeled perfusion MRI for clinical applications: A consensus of the ISMRM perfusion study group and the European consortium for ASL in dementia.* Magn Reson Med, 2015. **73**(1): p. 102-16.

32. Schrantee, A., et al., *The age-dependent effects of a single-dose methylphenidate challenge on cerebral perfusion in patients with attention-deficit/hyperactivity disorder.* Neuroimage Clin, 2017. **13**: p. 123-129.

33. Martin, S.J., P.D. Grimwood, and R.G. Morris, *Synaptic plasticity and memory: an evaluation of the hypothesis.* Annu Rev Neurosci, 2000. **23**: p. 649-711.

34. Ahmed, R. and P. Aslani, *Attention-deficit/hyperactivity disorder: an update on medication adherence and persistence in children, adolescents and adults.* Expert Review of Pharmacoeconomics \& Outcomes Research, 2013. **13**(6): p. 791--815.

35. Charach, A., et al., *Documenting adherence to psychostimulants in children with ADHD.* J Can Acad Child Adolesc Psychiatry, 2008. **17**(3): p. 131-6.

36. Shaw, P., et al., *Psychostimulant treatment and the developing cortex in attention deficit hyperactivity disorder.* Am J Psychiatry, 2009. **166**(1): p. 58-63.

37. Kaminski, A., et al., *Change in striatal functional connectivity networks across 2 years due to stimulant exposure in childhood ADHD: results from the ABCD sample.* Translational Psychiatry, 2024. **14**(1): p. 463.

38. Hart, H., et al., *Meta-analysis of functional magnetic resonance imaging studies of inhibition and attention in attention-deficit/hyperactivity disorder: exploring task-specific, stimulant medication, and age effects.* JAMA Psychiatry, 2013. **70**(2): p. 185-98.

39. Lizanne, J.S.S., et al., *Thinner Medial Temporal Cortex in Adolescents With Attention-Deficit/Hyperactivity Disorder and the Effects of Stimulants.* Journal of the American Academy of Child & Adolescent Psychiatry, 2015. **54**(8): p. 660-667.

40. Shaw, P., et al., *Mapping the development of the basal ganglia in children with attention-deficit/hyperactivity disorder.* J Am Acad Child Adolesc Psychiatry, 2014. **53**(7): p. 780-9.e11.

41. Lizanne, J.S.S., Patrick, and D. Sarah, *MR imaging of the effects of methylphenidate on brain structure and function in Attention-Deficit/Hyperactivity Disorder.* European Neuropsychopharmacology, 2013. **23**(10): p. 1151-1164.

42. Kaasinen, V., et al., *Age-related dopamine D2/D3 receptor loss in extrastriatal regions of the human brain.* Neurobiol Aging, 2000. **21**(5): p. 683-8.

43. Seaman, K.L., et al., *Differential regional decline in dopamine receptor availability across adulthood: Linear and nonlinear effects of age.* Hum Brain Mapp, 2019. **40**(10): p. 3125-3138.

44. Garcia-Cabezas, M.A., et al., *Distribution of the dopamine innervation in the macaque and human thalamus.* Neuroimage, 2007. **34**(3): p. 965-84.

45. Kasparek, T., P. Theiner, and A. Filova, *Neurobiology of ADHD From Childhood to Adulthood: Findings of Imaging Methods.* J Atten Disord, 2015. **19**(11): p. 931-43.

46. Andersen, S.L. and M.H. Teicher, *Sex differences in dopamine receptors and their relevance to ADHD.* Neurosci Biobehav Rev, 2000. **24**(1): p. 137-41.

47. Weickert, C.S., et al., *Postnatal alterations in dopaminergic markers in the human prefrontal cortex.* Neuroscience, 2007. **144**(3): p. 1109-19.

48. Udo de Haes, J.I., et al., *Methylphenidate-induced activation of the anterior cingulate but not the striatum: a [15O]H2O PET study in healthy volunteers.* Hum Brain Mapp, 2007. **28**(7): p. 625-35.

49. Shaw, P. and G. Sudre, *Adolescent Attention-Deficit/Hyperactivity Disorder: Understanding Teenage Symptom Trajectories.* Biol Psychiatry, 2021. **89**(2): p. 152-161.

50. Franke, B., et al., *Live fast, die young? A review on the developmental trajectories of ADHD across the lifespan.* Eur Neuropsychopharmacol, 2018. **28**(10): p. 1059-1088.

51. Norman, L.J., et al., *Longitudinal trajectories of childhood and adolescent attention deficit hyperactivity disorder diagnoses in three cohorts.* EClinicalMedicine, 2023. **60**: p. 102021.

52. Palladino, V.S., et al., *Genetic risk factors and gene-environment interactions in adult and childhood attention-deficit/hyperactivity disorder.* Psychiatr Genet, 2019. **29**(3): p. 63-78.

53. Gajria, K., et al., *Adherence, persistence, and medication discontinuation in patients with attention-deficit/hyperactivity disorder - a systematic literature review.* Neuropsychiatr Dis Treat, 2014. **10**: p. 1543-69.

54. Hodgkins, P., R. Sasane, and W.M. Meijer, *Pharmacologic treatment of attention-deficit/hyperactivity disorder in children: incidence, prevalence, and treatment patterns in the Netherlands.* Clin Ther, 2011. **33**(2): p. 188-203.

55. Winterstein, A.G., et al., *Utilization of pharmacologic treatment in youths with attention deficit/hyperactivity disorder in Medicaid database.* Ann Pharmacother, 2008. **42**(1): p. 24-31.

56. Thompson, K., J. Kulkarni, and A.A. Sergejew, *Reliability and validity of a new Medication Adherence Rating Scale (MARS) for the psychoses.* Schizophrenia Research, 2000. **42**(3): p. 241-247.

57. Rapoport, I.L. and A.P. Groenman, *A Review of Sex and Gender Factors in Stimulant Treatment for Attention-Deficit/Hyperactivity Disorder: Knowledge Gaps and Future Directions.* PsyArChiv Preprints, 2024.

Acknowledgements

We would like to thank all participants and their parents for their contribution to this study and all students that helped with collection and analysis of the data. A preprint version of this manuscript is available at https://doi.org/10.1101/2024.12.30.24319766.

Author contributions

Conceptualisation: ZP, LR, HMG, AS. Formal analysis: ZP. Funding acquisition: LR. Investigation: AK, MAB, AS. Methodology: HJMM. Resources: LR. Supervision: LR, HMG, AS. Writing - original draft: ZP, LR, AS. Writing - review & editing: ZP, LR, HJJM, AK, MAB, HMG, AS.

Funding

This study was funded by the Dutch non-profit organisations Kiddy GoodPills and Suffigium. The ePOD-MPH RCT (baseline data) was funded by a personal research grant awarded to LR by the Academic Medical Center, University of Amsterdam, and 11.32050.26ERA-NET PRIOMEDCHILD FP 6 (EU).

Competing interests

The authors have nothing to disclose.

1 **Figure and Table legends**
2
Figure 1. Study design and regions of interest (ROIs). A) At baseline and 4-year follow-up
assessment, participants underwent MR scanning before and 90 minutes after a single-dose
short-acting methylphenidate-challenge (0.5mg/kg; max. 20 mg for adolescents, max. 40 mg for
adults). Stimulant medication use between baseline and follow-up (cumulative dose/exposure
duration) was calculated from medication received during the randomised controlled trial (RCT)
and pharmacy prescription information from the naturalistic follow-up. B) Predefined ROIs:
anterior cingulate cortex (ACC), medial prefrontal cortex (mPFC), striatum and thalamus.
10
11 **Table 1. Participant characteristics of the study participants.** Data are presented as mean
12 (standard deviation; SD), median (interquartile range; IQR), fraction (yes/no, MPH/placebo) or
13 percentage (%). ADHD=attention-deficit/hyperactivity disorder; BL=baseline; rCBF=relative
14 cerebral blood flow; FU=follow-up; IQ=intelligence quotient; MPH=methylphenidate; RCT =
15 randomised controlled trial.
16
Figure 2. Age*medication interaction effects on resting rCBF and rCBF-response to a
methylphenidate-challenge. A) Thalamic resting rCBF was associated with medication use in
adults, but not in adolescents. B) mPFC rCBF-response was associated with medication use in
adolescents, but not in adults. The plots show the average rCBF values across visits, as the age-
by-medication interaction effects are independent of time. mPFC=medial prefrontal cortex;
rCBF=relative cerebral blood flow.
23
Table 2. Linear mixed effect model results. Bold values represent significant effects after
multiple comparison correction (FDR=5%, 4 comparisons). Separate models were used per
medication variable (cumulative dose, exposure duration). The variables mean or change in
motion, MR scanner at baseline, baseline age, and scan interval were included as covariates in
all models. For readability covariates were excluded from the table, except for motion, which
showed a significant effect in some models. ACC=anterior cingulate cortex; mPFC=medial
prefrontal cortex; P_{FDR}=P-value after FDR multiple comparison correction.
31
32 **Table 3. Associations between whole-brain CBF-response to a methylphenidate challenge**
33 **and dopamine and noradrenaline receptor distributions.** Correlation coefficients were
34 obtained using spin tests. Bold values represent significant associations. BL=baseline;
35 CBF=cerebral blood flow; FU=follow-up.

Table 1. Participant characteristics of the study participants.

	Adolescents n = 32		Adults n = 24		
	BL	FU	BL	FU	Statistics[1]
Demographics					
Age (years, *mean (SD)*)	11.2 (0.9)	15.3 (1.4)	29.9 (5.0)	34.4 (4.6)	
Estimated IQ[2] (*mean (SD)*)	104.5 (17.7)		106.7 (5.0)		
Handedness (*% right-handed*)	93.8		91.7		
ADHD symptom severity (*mean (SD)*)[3]					
DBD-RS inattentive	22.8 (3.4)	11.8 (5.2)			
DBD-RS hyperactive/impulsive	15.9 (5.6)	7.2 (4.4)			
ADHD-RS			33.1 (9.9)	24.5 (8.5)	
Clinically significant symptoms[4] (*%*)					
DBD-RS inattentive	96.9	26.7			
DBD-RS hyperactive/impulsive	53.1	6.7			
ADHD-RS			100.0	94.7	
Medication					
Age at start (years, *mean (SD)*)		11.8 (1.4)		29.2 (4.0)	
Cumulative dose[5] (mg, *median (IQR)*)		9455 (3533 - 23630)		4778 (750 - 13228)	W= 456, p = .24
Exposure duration[6] (months, *median (IQR)*)		16 (6 - 25)		4 (1 - 16)	W = 531, p = .02, adolescents > adults
Medication use at FU (*yes/no*)		16/13		2/17	
Stimulant-treatment naive at FU (*yes/no*)		3/29		5/19	
ePOD-MPH RCT treatment group (*MPH/placebo*)		11/21		11/13	
Magnetic resonance imaging					
Resting rCBF[7] (*mean (SD)*)					
ACC	1.27 (0.11)	1.27 (0.09)	1.32 (0.09)	1.33 (0.07)	
mPFC	1.11 (0.07)	1.16 (0.09)	1.13 (0.05)	1.17 (0.05)	
Striatum	0.98 (0.10)	1.01 (0.10)	1.00 (0.06)	1.01 (0.08)	
Thalamus	0.83 (0.09)	0.80 (0.090	0.97 (0.08)	0.91 (0.09)	
rCBF-response to a MPH-challenge[7] (*mean (SD)*)					
ACC	-0.01 (0.10)	-0.02 (0.08)	-0.04 (0.06)	-0.02 (0.07)	
mPFC	0.04 (0.08)	0.003 (0.07)	0.003 (0.05)	0.01 (0.05)	
Striatum	0.02 (0.07)	0.01 (0.09)	0.04 (0.05)	0.04 (0.08)	
Thalamus	0.02 (0.13)	0.05 (0.13)	0.03 (0.08)	0.04 (0.11)	
Scan interval (months, *mean (SD)*)		48.6 (14.2)		53.3 (9.1)	t(54) = -1.41, p = .16

[1] Two-sample t-test or Mann Whitney U test.
[2] For adolescents: subtest Wechsler Intelligence Scale for Children (WISC); for adults: National Adult Reading Test (NART, Dutch translation).
[3] For adolescents: Disruptive behavior disorder rating scale (DBD-RS); for adults: Attention Deficit Hyperactivity Disorder-Rating Scale (ADHD-RS).
[4] Thresholds indicating clinically significant symptoms were >15 for DBD-RS inattention and hyperactive/impulsive subscales (adolescents) and >10 for ADHD-RS (adults).
[5] Converted to methylphenidate-equivalents.
[6] Calculated with a 30-day permissible gap.
[7] rCBF calculated as *CBF region of interest/CBF total grey matter*. rCBF-response calculated as *(post-MPH - pre-MPH)/pre-MPH*.

Table 2. Linear mixed effect model results.

A) Three-way linear mixed effects model with pre-MPH rCBF	ACC	mPFC	Striatum	Thalamus
Cumulative dose				
Age	$F(1,103)=4.3$, $P=.041$, $P_{FDR}=.055$	**$F(1,58)=5.4$, $P=.023$, $P_{FDR}=.046$**	$F(1,57)=0.5$, $P=.47$, $P_{FDR}=.47$	**$F(1,55)=26.1$, $P<.001$, $P_{FDR}<.001$**
Visit	$F(1,103)=0.02$, $P=.88$, $P_{FDR}=.88$	$F(1,55)=4.8$, $P=.033$, $P_{FDR}=.066$	$F(1,50)=1.9$, $P=.18$, $P_{FDR}=.24$	**$F(1,52)=10.7$, $P=.002$, $P_{FDR}=.008$**
Medication	$F(1,103)=4.2$, $P=.042$, $P_{FDR}=.056$	**$F(1,53)=6.4$, $P=.014$, $P_{FDR}=.028$**	**$F(1,51)=6.8$, $P=.012$, $P_{FDR}=.028$**	$F(1,51)<0.01$, $P=.99$, $P_{FDR}=.99$
Age * Visit	$F(1,103)=0.07$, $P=.79$, $P_{FDR}=.99$	$F(1,55)<0.01$, $P=.99$, $P_{FDR}=.99$	$F(1,50)=0.5$, $P=.50$, $P_{FDR}=.99$	$F(1,52)=0.2$, $P=.67$, $P_{FDR}=.99$
Age * Medication	$F(1,103)=0.2$, $P=.70$, $P_{FDR}=.70$	$F(1,53)=1.4$, $P=.23$, $P_{FDR}=.31$	$F(1,51)=1.6$, $P=.22$, $P_{FDR}=.31$	**$F(1,51)=7.7$, $P=.008$, $P_{FDR}=.030$**
Visit * Medication	$F(1,103)=0.06$, $P=.81$, $P_{FDR}=.81$	$F(1,54)=0.7$, $P=.42$, $P_{FDR}=.79$	$F(1,48)=0.3$, $P=.59$, $P_{FDR}=.79$	$F(1,50)=1.3$, $P=.26$, $P_{FDR}=.79$
Age * Visit * Medication	$F(1,103)=0.01$, $P=.90$, $P_{FDR}=.90$	$F(1,54)=0.3$, $P=.62$, $P_{FDR}=.90$	$F(1,48)=0.1$, $P=.71$, $P_{FDR}=.90$	$F(1,50)=0.9$, $P=.33$, $P_{FDR}=.94$
Mean Motion	$F(1,103)=0.02$, $P=.90$, $P_{FDR}=.90$	$F(1,98)=1.5$, $P=.23$, $P_{FDR}=.46$	$F(1,103)=0.7$, $P=.39$, $P_{FDR}=.53$	**$F(1,101)=7.0$, $P=.009$, $P_{FDR}=.037$**
Exposure duration				
Age	$F(1,103)=3.8$, $P=.054$, $P_{FDR}=.072$	**$F(1,54)=5.6$, $P=.022$, $P_{FDR}=.043$**	$F(1,54)=0.5$, $P=.49$, $P_{FDR}=.49$	**$F(1,52)=27.3$, $P<.001$, $P_{FDR}<.001$**
Visit	$F(1,103)<0.01$, $P=.98$, $P_{FDR}=.98$	$F(1,54)=3.2$, $P=.08$, $P_{FDR}=.16$	$F(1,50)=1.2$, $P=.28$, $P_{FDR}=.38$	**$F(1,51)=8.8$, $P=.005$, $P_{FDR}=.019$**
Medication	**$F(1,103)=5.6$, $P=.020$, $P_{FDR}=.027$**	**$F(1,53)=8.7$, $P=.005$, $P_{FDR}=.015$**	**$F(1,51)=7.7$, $P=.007$, $P_{FDR}=.015$**	$F(1,50)=1.7$, $P=.20$, $P_{FDR}=.20$
Age * Visit	$F(1,103)=0.2$, $P=.64$, $P_{FDR}=.82$	$F(1,54)=0.3$, $P=.58$, $P_{FDR}=.82$	$F(1,50)=0.1$, $P=.82$, $P_{FDR}=.82$	$F(1,51)=0.3$, $P=.59$, $P_{FDR}=.82$
Age * Medication	$F(1,103)=0.8$, $P=.36$, $P_{FDR}=.56$	$F(1,53)=0.4$, $P=.54$, $P_{FDR}=.56$	$F(1,51)=0.3$, $P=.56$, $P_{FDR}=.56$	**$F(1,51)=10.1$, $P=.003$, $P_{FDR}=.010$**
Visit * Medication	$F(1,103)=0.02$, $P=.89$, $P_{FDR}=.89$	$F(1,53)=0.9$, $P=.34$, $P_{FDR}=.89$	$F(1,49)=0.1$, $P=.79$, $P_{FDR}=.89$	$F(1,50)=0.4$, $P=.53$, $P_{FDR}=.89$
Age * Visit * Medication	$F(1,103)=0.2$, $P=.67$, $P_{FDR}=.78$	$F(1,53)=1.1$, $P=.29$, $P_{FDR}=.69$	$F(1,49)=0.1$, $P=.78$, $P_{FDR}=.78$	$F(1,50)=0.9$, $P=.34$, $P_{FDR}=.69$
Mean Motion	$F(1,103)<0.01$, $P=.98$, $P_{FDR}=.98$	$F(1,97)=1.0$, $P=.33$, $P_{FDR}=.44$	$F(1,103)=1.5$, $P=.22$, $P_{FDR}=.44$	**$F(1,99)=8.2$, $P=.005$, $P_{FDR}=.021$**

B) Three-way linear mixed effects model with rCBF-response to a MPH-challenge	ACC	mPFC	Striatum	Thalamus
Cumulative dose				
Age	$F(1,53)=0.9$, $P=.34$, $P_{FDR}=.43$	**$F(1,53)=10.6$, $P=.002$, $P_{FDR}=.008$**	$F(1,55)=0.9$, $P=.34$, $P_{FDR}=.43$	$F(1,40)=0.6$, $P=.43$, $P_{FDR}=.43$
Visit	$F(1,53)=0.4$, $P=.54$, $P_{FDR}=.54$	$F(1,51)=1.4$, $P=.25$, $P_{FDR}=.54$	$F(1,55)=0.7$, $P=.40$, $P_{FDR}=.54$	$F(1,39)=1.1$, $P=.29$, $P_{FDR}=.54$
Medication	$F(1,53)=2.4$, $P=.13$, $P_{FDR}=.17$	**$F(1,53)=8.8$, $P=.004$, $P_{FDR}=.018$**	$F(1,55)=2.7$, $P=.11$, $P_{FDR}=.17$	$F(1,40)=0.1$, $P=.82$, $P_{FDR}=.82$
Age * Visit	$F(1,53)=0.4$, $P=.53$, $P_{FDR}=.53$	$F(1,51)=3.7$, $P=.059$, $P_{FDR}=.23$	$F(1,55)=0.8$, $P=.37$, $P_{FDR}=.53$	$F(1,39)=0.5$, $P=.50$, $P_{FDR}=.53$
Age * Medication	$F(1,53)=0.5$, $P=.47$, $P_{FDR}=.47$	**$F(1,52)=10.9$, $P=.002$, $P_{FDR}=.007$**	$F(1,55)=5.0$, $P=.030$, $P_{FDR}=.059$	$F(1,40)=3.0$, $P=.09$, $P_{FDR}=.12$
Visit * Medication	$F(1,53)=0.4$, $P=.54$, $P_{FDR}=.71$	$F(1,51)=0.1$, $P=.71$, $P_{FDR}=.71$	$F(1,55)=2.7$, $P=.11$, $P_{FDR}=.42$	$F(1,40)=0.4$, $P=.51$, $P_{FDR}=.71$
Age * Visit * Medication	$F(1,52)=3.1$, $P=.09$, $P_{FDR}=.33$	$F(1,51)=0.1$, $P=.80$, $P_{FDR}=.80$	$F(1,55)=2.0$, $P=.16$, $P_{FDR}=.33$	$F(1,39)=0.2$, $P=.62$, $P_{FDR}=.80$
Change in Motion	$F(1,90)=1.1$, $P=.29$, $P_{FDR}=.58$	**$F(1,94)=8.5$, $P=.005$, $P_{FDR}=.018$**	$F(1,88)<0.01$, $P=.92$, $P_{FDR}=.92$	$F(1,83)=0.1$, $P=.73$, $P_{FDR}=.92$
Exposure duration				
Age	$F(1,53)=0.5$, $P=.47$, $P_{FDR}=.75$	**$F(1,53)=13.3$, $P<.001$, $P_{FDR}=.002$**	$F(1,57)=0.3$, $P=.57$, $P_{FDR}=.75$	$F(1,42)=0.1$, $P=.75$, $P_{FDR}=.75$
Visit	$F(1,54)=1.4$, $P=.24$, $P_{FDR}=.25$	$F(1,53)=1.3$, $P=.25$, $P_{FDR}=.25$	$F(1,58)=1.5$, $P=.22$, $P_{FDR}=.25$	$F(1,42)=2.3$, $P=.14$, $P_{FDR}=.25$
Medication	$F(1,51)=0.6$, $P=.45$, $P_{FDR}=.45$	$F(1,52)=6.2$, $P=.016$, $P_{FDR}=.065$	$F(1,55)=1.7$, $P=.19$, $P_{FDR}=.38$	$F(1,40)=0.7$, $P=.41$, $P_{FDR}=.45$
Age * Visit	$F(1,54)=1.4$, $P=.23$, $P_{FDR}=.23$	$F(1,52)=4.7$, $P=.036$, $P_{FDR}=.14$	$F(1,58)=1.5$, $P=.23$, $P_{FDR}=.23$	$F(1,42)=2.0$, $P=.16$, $P_{FDR}=.23$
Age * Medication	$F(1,52)=0.1$, $P=.74$, $P_{FDR}=.74$	**$F(1,52)=11.2$, $P=.002$, $P_{FDR}=.006$**	$F(1,55)=1.6$, $P=.22$, $P_{FDR}=.34$	$F(1,40)=1.3$, $P=.25$, $P_{FDR}=.34$
Visit * Medication	$F(1,52)=0.9$, $P=.33$, $P_{FDR}=.45$	$F(1,52)=0.03$, $P=.86$, $P_{FDR}=.86$	$F(1,56)=3.6$, $P=.07$, $P_{FDR}=.26$	$F(1,41)=1.0$, $P=.31$, $P_{FDR}=.45$
Age * Visit * Medication	$F(1,51)=5.2$, $P=.027$, $P_{FDR}=.11$	$F(1,51)=0.4$, $P=.55$, $P_{FDR}=.55$	$F(1,56)=2.0$, $P=.17$, $P_{FDR}=.26$	$F(1,41)=1.7$, $P=.20$, $P_{FDR}=.26$
Change in Motion	$F(1,91)=1.7$, $P=.20$, $P_{FDR}=.39$	$F(1,91)=5.6$, $P=.018$, $P_{FDR}=.070$	$F(1,88)=0.02$, $P=.89$, $P_{FDR}=.89$	$F(1,84)=0.2$, $P=.66$, $P_{FDR}=.88$

Table 3. Associations between whole-brain CBF-response to a methylphenidate challenge and dopamine and noradrenaline receptor distributions. BL=baseline; CBF=cerebral blood flow; FU=follow-up.

	Adolescents		Adults	
	BL	FU	BL	FU
Dopamine transporter	r = .03, p = .80	r = -<.01, p = .99	r = .02, p = .83	r = -.01, p = .93
Dopamine D1 receptor	r = .28, p = .059	r = .09, p = .55	r = -<.01, p = .99	r = .04, p = .75
Dopamine D2/D3 receptor	r = .17, p = .28	r = -.08, p = .64	r = .05, p = .76	r = .04, p = .81
Noradrenaline transporter	**r = -.39, p = .020**	r = -.24, p = .19	**r = -.38, p = .020**	**r = -.37, p = .030**