AI Perspectives on the Present and Future of Antidepressant Pharmaceutical Treatment Based on Anti-inflammatory Strategies: A Scoping Review of Randomised Controlled Clinical Trials

Yan Bo[1*]

[1]Department of Medicine, Northwest Minzu University, Lanzhou 73000, China

* Correspondence:
Corresponding Author
Yan Bo thoseformylovedpeople@outlook.com

Keywords: depression, anti-inflammatory strategy, randomized controlled trials,
pharmaceutical treatment, scoping review.

Abstract

Background

Depression remains an unresolved issue on a global scale. Recently, a novel concept of 'anti-
inflammatory-based pharmacotherapy' has been developed. Despite the role of inflammation in
depression having been discussed in many reviews at various levels, the prevalence of this new
concept in randomised controlled clinical studies and its implications remain elusive. The aim of this
scoping review was precisely to explore in depth the current status of inflammation in randomised
controlled clinical trial studies of depression.

Methods

PubMed was systematically searched from inception to December 11 2024. Studies that researches
on the treatment of depression based on anti-inflammatory strategies were included. Study
characteristics and outcomes were extracted and organized thematically.

Findings

11 reports of randomised controlled clinical trials were included, which accumulated 907 depressed
patients. All studies found that there is a connection between the effects of anti-inflammatory drugs
in treating depression and a large decrease in the levels of inflammatory markers in the blood of
depressed patients compared to before treatment. Three inflammatory markers, CRP, IL-6 and TNF-
alpha, were the most frequently mentioned. The current strategy of anti-inflammatory drug
administration did not differ fundamentally from the previous strategy of traditional antidepressant
drugs combined with psychotherapy.

Interpretation

At present, the use of anti-inflammatory strategies for the pharmacological treatment of depression
has limited research value and poor feasibility. The future direction of the new concept of anti-
inflammatory strategies for the treatment of depression, proposed in the context of the association
between inflammation and depression, is that psychiatrists, researchers, and psychotherapists should

shift their future focus from pharmacological treatments based on anti-inflammatory strategies to
non-pharmacological treatments of anti-inflammatory strategies, such as positive thinking, exercise,
and so on. The popularity of purely clinical randomised controlled studies in the depression
population is extremely low when considering the financial investment in research and the benefits of
translating the results. In the future, public interest studies, low research costs, and research protocols
with mass generalisability will be more likely to stimulate the depression community's interest in
participating in research. The potential value and feasibility of future research lies in the application
of an integrated AI platform to assist pharmacological treatment of depression based on anti-
inflammatory strategies.

Registration DOI: https://doi.org/10.17605/OSF.IO/A64GC.

1   Research in context

1.1   Evidence before this study

Depression remains an unresolved issue on a global scale. Recently, a novel concept of 'anti-
inflammatory-based pharmacotherapy' has been developed. Despite the role of inflammation in
depression having been discussed in many reviews at various levels, the prevalence of this new
concept in randomised controlled clinical studies and its implications remain elusive. My previous
preliminary search of the PubMed database revealed no extant systematic reviews publishing insights
on how to view anti-inflammatory strategies based on the treatment of patients with depression-
associated inflammation.

1.2   Added value of this study

To the best of our knowledge, this is the first scoping review that investigates treatment of patients
with depression-related inflammation based on anti-inflammatory strategies. It uncovers the
connection between anti-inflammatory drugs, reduced inflammatory markers, and treatment effects.
It also points out the limitations of current anti-inflammatory strategies in treating depression,
suggesting a shift to non-pharmacological approaches. Additionally, it proposes the potential of AI
in improving treatment, providing new directions for future research.

1.3   Implications of all the available evidence

Current evidence shows anti-inflammatory strategies for depression treatment have limitations.
Future research should focus on non-pharmacological approaches, AI-assisted treatment, and
clarifying inflammation-depression links for better patient care.

2   Introduction

In recent years, the role of inflammation in depression and the effects of anti-inflammatory drugs on
depression have emerged as significant areas of mental health research. A growing body of evidence
strongly indicates that inflammatory processes are closely associated to the pathogenesis of
depression (1). Currently, the anti-inflammatory drugs commonly used to treat depression in the
clinic are non-steroidal anti-inflammatory drugs (NSAIDs) and Omega-3 fatty acids. NSAIDs, such
as aspirin and ibuprofen, are considered to have an antidepressant effect because they can lower
inflammation in the body and thus improve depressive symptoms (2). A substantial body of clinical
research has demonstrated significant improvements in mood and overall mental health in depressed
patients treated with NSAIDs. Omega-3 fatty acids, believed to possess anti-inflammatory properties,

have also exhibited favourable outcomes in the management of depression. Studies have indicated that a diet abundant in Omega-3s is associated with a reduced risk of depression. These fatty acids may modulate inflammatory responses and neurotransmitter levels, thereby contributing to their antidepressant effect (3). In addition to the two aforementioned classes of drugs, other substances with anti-inflammatory bioactive effects, such as steroids, have demonstrated antidepressant effects by suppressing the inflammatory response, thereby potentially alleviating depressive symptoms. Patients who have used steroids for a prolonged period have exhibited a substantial reduction in depressive symptoms (4).

Nevertheless, the question of whether these findings can modify the prevailing status quo for depressed patients, who require either long-term or short-term cognitive therapy, remains unresolved (5). This shift in the management of depression emphasises the transition of long-term fixed cognitive therapy into a therapeutic process involving the regular administration of medication. This transformation in the treatment process is undoubtedly more appealing to patient groups requiring short, rapid, and convenient treatments for depression, such as working adults and school students. The present moment is characterised by a crisis in the translation of research findings into clinical practice in the field of depression. This crisis results in the failure to fully implement significant discoveries concerning the underlying mechanisms of depression in clinical treatment. The reasons for this phenomenon are not clear-cut, and there is a pressing need for novel approaches to enhance the return on investment in depression research.

This study proposes that reducing inflammation in the body should be a goal in treating depression. This idea is backed by the new area of research on developing anti-inflammatory drug treatments for depression. Concurrently, the field of artificial intelligence is experiencing rapid growth and gradual infiltration into all aspects of self-health assessment and teletherapy (6,7). In this context, the question arises as to how artificial intelligence can be utilised to assist in the pharmacological treatment of depression under effective human management.

The objective of this scoping review is to examine the current state of inflammation in randomised controlled clinical trial studies of depression. The identification of current trends in this area and the collation of existing frameworks that may be used to guide future research is also intended, with a view to providing new ideas and approaches to the treatment of depression.

3 Literature Review

3.1 Overview

Neuroinflammation, defined as an immune response within the central nervous system, involves a series of processes including the activation of glial cells and the release of cytokines and chemokines. In the development of depression, abnormal neuroinflammation interferes with neurotransmitter metabolism, affects neuroplasticity, and in turn triggers depressive symptoms. Relatedly, studies using radiomics to predict remission in Cushing's disease have highlighted the role of inflammation in neuroendocrine contexts, offering insights into similar processes in depression(11).

Conversely, signalling pathways represent a series of intracellular molecular interactions, including the NLRP3 (12), NF-κB (13) and P2X7R (14), which play a pivotal role in the transmission and regulation of neuroinflammation and depression. These signalling pathways are of central interest in the study of the pathogenesis and therapeutic targets of depression. Advanced AI techniques, such as deep learning for segmenting pituitary adenomas, could be adapted to study these signaling pathways in depression (15).

In 2017, Miller and Raison published a image entitled 'Transmitting stress-induced inflammatory
signals to the brain'. This image was edited to reflect the adjustment of factors that trigger
inflammation-associated depression from stress to environmental factors, heritage factors, and
personal habits **(Figure 1)**. This adjustment is more consistent with existing evidence describing the
inflammatory pathogenesis of depression (16). **Figure 1-A** depicts the pathological process by which
depression susceptibility factors stimulate inflammation through humoral, neural and cellular
pathways. Depression susceptibility factors include environmental factors, genetic factors, and
lifestyle habits. The environmental factors referred to here include both human social and family
environments and other scenarios where human interaction occurs, as well as natural environments
that can trigger microbial infections in humans. Microorganisms are mentioned because patients with
depression often have positive lipolyaccharide (LPS) results or signs of infection in their laboratory
tests. The strong action of depression susceptibility factors activates the sympathetic nervous system
to release catecholamines, and this results in the production of myeloid cells by the bone marrow and
their gradual release into the peripheral circulation. Here we describe the inflammatory response to
depression initiated by monocytes in a cellular pathway using monocytes in myeloid cells as an
example. The monocyte in the peripheral blood circulation will try to fulfil its mission of capture,
phagocytosis. Examples include bacterial metabolites or other substances that stimulate damage-
associated molecular patterns (DMAPs). The most frequent site in the human body where monocytes
are confronted with bacteria and associated metabolites is the gut, where monocytes take full
advantage of recognising microbial-associated molecular patterns (MAMPs) and thus clearing
microbes. However, DMAPs and MAMPs are optimal mediators for the activation of inflammatory
signalling pathways, such that a range of expression occurs within monocytes, such as nuclear factor-
κ b (NF-κ b) and pyrin structural domain protein 3 (NLRP3) inflammatory vesicles. This expression
is a cascade response that can also interact with each other. For example, stimulation of NLRP3 in
turn activates caspase 1, leading to the production of mature interleukin-1β (IL-1β) and IL-18, as well
as splitting of the glucocorticoid receptor, leading to glucocorticoid resistance. This expression
results in the recruitment of large amounts of cytokines such as IL-1β and IL-18, which in turn
recruits other immune cells against the cytokine chemical gradient to come and participate in the
immune process. As immune mobilisation continues, the number of cytokines produced gradually
increases, and these cytokines do not stop at specific sites of the immune response, but travel with
monocytes to the brain via humoral and neural pathways. The immune cells present in the brain are
partly called astrocytes and partly called microglia. These microglia are activated by a large number
of cytokines to an M1 pro-inflammatory phenotype which releases CC-chemokine ligand 2 (CCL2),
which in turn attracts activated monocytes to the brain via the cellular pathway. Once the monocytes
enter the brain, the peripheral immune response will spread to the centre, which may result in central
inflammation and exacerbate the manifestations of depression. **Figure 1-B** highlights how nerve cells
in the inflammatory response affect human behaviour through cellular pathways. If the inflammatory
response is present in the brain, then there is no doubt that neurotransmitter transmission and
metabolism will certainly be disrupted. The effects we are most interested in are depressive
symptoms, such as lack of motivation and lack of pleasure in behaviour. Here elaborate on
interferons (IFNs), IL-1β and tumour necrosis factor (TNF) as representatives of cytokines. These
three cytokines reduce the availability of monoamine-serotonin (5-HT), dopamine (DA) and
norepinephrine (NE) - through activation of the mitogen-activated protein kinase (MAPK) pathway.
With cytokine-induced reduction in astrocyte glutamate reuptake and stimulation of astrocyte
glutamate release. Glutamate (Glu) is an excitatory amino acid neurotransmitter. Excess Glu, when
bound to extrasynaptic N-methyl-D-aspartate receptor (NMDAR), leads to decreased brain-derived
neurotrophic factor (BDNF) and excitotoxicity. Low levels of DA and BDNF levels of
neurotransmitters passed between nerve cells and neuronal cells are then not consistent with

concentration levels in a non-inflammatory state, a result that can lead to a lack of reward motivation
and a lack of rapture in cortical striatal circuits.

Figure 1 Inflammatory pathogenesis of depression. (A) Humoral, netural and cellular route. (B)
Neurotransmitters. ASC, apoptosis-associated speck-like protein containing a CARD; HMGB1, high
mobility group box 1; LPS, lipopolysaccharide; TLR, Toll-like receptor. EAAT2, excitatory amino
acid transporter 2; NF-κB, nuclear factor-κB; Glu, Glutamate; HSP, heat shock protein; NMDAR, N-
methyl-D-aspartate receptor; BDNF, brain-derived neurotrophic factor; Glu, Glutamate; MAPK,
mitogen-activated protein kinase; DA, dopamine; NE, norepinephrine; 5-HT monoamine-serotonin;
IFNs, interferons; TNF, tumour necrosis factor; CCL2, CC-chemokine ligand 2; IL-1β, interleukin-
1β; pyrin structural domain protein 3 (NLRP3); NF-κ b ,nuclear factor-κ b; MAMPs, microbial-
associated molecular patterns; DMAPs, damage-associated molecular patterns; LPS, lipolyaccharide.

The present review covers 31 animal studies focusing on neuroinflammatory target-based therapeutic
strategies for depression, involving both pharmacological interventions (including SGLT2 inhibitors,
vitamin D analogues, natural extracts and other compounds) and non-pharmacological interventions
(e.g. These include acupuncture, exercise, and light therapy. The present review comprehensively
incorporates the results of the animal model studies, with the aim of presenting an all-rounded picture
of the research in this field and to provide a comprehensive overview of the clinical translation of the
scoping review. It aims to present a comprehensive picture of the research in this field and to provide
an animal-level basis for a scoping review to explore clinical translation.

In the actual population of depressed patients, neuroinflammation does not exist in isolation, but
rather involves the interaction of multiple systems, including the neurological, endocrine, and
immune systems. The lifestyle and environmental factors of each depressed patient have been shown

to indirectly affect neural circuits, thereby complicating the manifestation of neuroinflammation. It
has been demonstrated that both the patient's personal factors and the complex neuroinflammation
can have a significant impact on the onset, development, and treatment response of depression. The
present study has identified a variety of novel anti-inflammatory drugs and non-pharmacological
treatments in animal experiments that modulate the three main signalling pathways of
neuroinflammation, namely NLRP3 (12), NF-κB (13) and P2X7R (14). This modulation has been
shown to result in an improvement of depressive symptoms and neuroinflammation. In terms of
pharmacological interventions, Empagliflozin has been shown to regulate autophagy or inflammation
and neurogenesis in the hippocampus through activation of AMPK (19). Calcitriol has been
demonstrated to inhibit the P2X7R/NLRP3/caspase-1 signalling pathway, thereby reducing
neuroinflammation (20). Among non-pharmacological interventions, acupuncture (14) and exercise
(21) have been shown to modulate relevant inflammatory and apoptotic pathways, improve
neuroplasticity and reduce neuroinflammation, respectively. The primary debate has focused on
single-target versus multi-target modulation strategies and pharmacological versus non-
pharmacological intervention modalities. Single-target modulation emphasises precise inhibition of
specific key signalling pathways, whereas multi-target modulation argues for integrated modulation
of multiple associated signalling pathways to cope with disease complexity. The majority of extant
studies are in the short-term observation and small-scale research stage. At this stage, animal studies
have not yet been able to clearly explain how these drugs act on specific neuroinflammatory targets
according to the dynamic changes in animals, and effectively regulate the functional activities of
neural circuits in the hippocampus, prefrontal cortex, and other brain regions that are closely related
to depression, so as to achieve a stable and long-lasting antidepressant effect.

Although the main focus has centred around therapeutic strategies for depression based on
neuroinflammatory targets, the question of how different pharmacological and non-pharmacological
treatments can precisely and dynamically modulate the interaction between neuroinflammation and
depression-associated neural circuits in a complex human physiological and pathological
environment has not been thoroughly investigated.

3.2   Development

In the early research on depression, the focus was primarily on the traditional theory of monoamine
neurotransmitter imbalance, with a relative paucity of understanding of the role of neuroinflammation
in depression (22). However, as research progressed, it was found that the inflammatory response
was associated with depression. Nonetheless, at this stage, research remained at the level of
observational studies, with no exploration of the specific mechanisms of neuroinflammation-related
signalling pathways and effective intervention strategies.

In recent years, however, there has been a shift in focus towards the exploration of the role of
neuroinflammation-associated signalling pathways, such as NLRP3 (12), NF-κB (13) and P2X7R
(14), in the development of depression. These studies have identified pharmacological and non-
pharmacological intervention options with therapeutic potential to modulate these signalling
pathways and thereby improve depressive symptoms. However, the majority of these studies are in
the animal experimentation phase and have yet to demonstrate sufficient clinical relevance.The
findings from these animal experiments are not yet representative of a comprehensive understanding
and effective solution to the problem of achieving long-term stable modulation of
neuroinflammation-related signalling pathways in depression and avoiding adverse effects in order to
achieve a sustained antidepressant effect. This requires a significant volume of in-depth research.

3.3 Research status

Pharmacological interventions

A study by Muhammad et al demonstrated that Empagliflozin can regulate the balance of autophagy or inflammatory dynamics and PKCζ-mediated neurogenesis in the hippocampus through activation of AMPK, inhibit the NLRP3 inflammatory vesicle pathway, and ameliorate depressive symptoms. In the reserpine-induced rat model of depression, Empagliflozin has been shown to restore monoamines and autophagy or inflammatory balance in the hippocampus, and to promote neuroplasticity. However, no in-depth study has yet been conducted on whether its sustained effects on the signalling pathways are affected by adaptive modulation of the organism after long-term use (19).

A study by Ma et al suggests that the combination of brexpiprazole and fluoxetine produces a rapid antidepressant effect in an inflammatory model of depression, possibly through stimulation of the BDNF-TrkB signalling pathway as well as modulation of dendritic spine density. Consequently, it is hypothesised that brexpiprazole-assisted SSRI treatment is expected to produce rapid antidepressant effects in patients with depression (22).

A study by Bao et al revealed that chronic stress activates endogenous retroviruses (ERVs) in a mouse model, prompting an immune-inflammatory response in microglia in the basolateral amygdala (BLA), which ultimately triggers negative emotional behaviours. Antiretroviral therapy, inhibition of reverse transcriptase, and knockdown of p53, a gene that regulates the transcription of ERVs, have been shown to be effective in suppressing chronic stress-induced immunoinflammatory responses in microglia and ameliorating negative emotional behaviours in mice. This remarkable effect relies on reversing microglial cell morphology and biological immune inflammation. This particular use of antipsychotics may generate new therapeutic inspiration for psychiatrists (23).

Wang et al. demonstrated that vitamin D analogues could attenuate neuroinflammation, alleviate LPS-induced depressive-like behaviours and cognitive deficits, and have protective effects on neurons through inhibition of the P2X7R/NLRP3/caspase-1 signalling pathway. However, the potential adverse effects of long-term use require further investigation. In particular, the indirect effects of abnormal calcium metabolism on neuroinflammation-related signalling pathways need to be further explored (20).

A study by Rostevanov et al. demonstrates that leukotrienes (LTs) enhance depression-like symptoms in mice, exhibiting a non-sex-selective effect. However, this study also reveals that sex selectivity is present in inflammatory homeostasis, as evidenced by the increased levels of IL-6 and TNF-alpha in brain regions in males and decreased levels in females. The clinical and translational implications of this sex-selective correlation in inflammatory homeostasis remain to be elucidated (24).

A study by Li et al. examined the effects of liver-sparing granules (SGKL) on the gut, brain, and behaviour of chronic restraint stress (CRS) rats, and found that SGKL could improve depression. The study also found that SGKL could improve depression-like behaviour by altering the intestinal microbiota and metabolites targeting the PI3K/Akt/mTOR pathway (25).

Progesterone alleviated depressive-like behavior in mice. It did so by reducing neuroinflammatory responses and oxidative stress(26).

A study by Casaril et al. demonstrated that the selenium-containing compound 3-((4-
chlorophenyl)selanyl)-1-methyl-1H-indole (CMI) was able to reverse acute restraint stress-induced
depressive-like behaviour in mice by regulating oxidative stress and neuroinflammation, and by
restoring corticosterone levels and related gene expression. This selenium-enriched compound,
endowed with oxidative and anti-inflammatory bioactivities, may represent a promising candidate for
future dietary interventions in the prevention of depression (27).

A total of three studies on nutritional supplements have reported that propolis, n-3 polyunsaturated
fatty acids (PUFAs) and skeletal muscle glycine all improve depression-like behaviour. This
mechanism is associated with anti-inflammatory and antioxidant properties 32−34.

A total of fourteen studies on natural extracts, including Fustin, Berberine, and Cinnamic Acid, were
identified as modulating oxidative stress, neuroinflammatory markers, and associated signalling
pathways in animal models to enhance depression- and anxiety-like behaviours. However, the
majority of these studies were of a short-term nature, and there was an absence of longitudinal
research on the stability of the signalling pathways and the potential challenges associated with
tolerance during extended use. Furthermore, there is a paucity of long-term follow-up studies.The
differences between these studies are summarised in **Table 1**.

Table 1. Comparison of 14 research reports on natural extracts. The abbreviations in the table
are explained below. BDNF: brain-derived neurotrophic factor; TrkB: tropomyosin receptor kinase
B; PI3K: phosphatidylinositol 3-kinase; Akt: Protein kinase B; NLRP3: NLR family pyrin domain
containing 3; caspase-1: Cysteinyl aspartate specific proteinase-1; ASC: apoptosis-associated speck-
like protein containing a CARD; IL-1β: interleukin-1β; NF-κB: nuclear factor kappa-B; TLR4: toll-
like receptor 4; FoxO1: forkhead box O1; CREB: cyclic AMP response element-binding protein; NR:
not report.

Plant extract	Pathway or mechanism	Effect
Loganin in Corni Fructus(33)	BDNF-TrkB; PI3K/Akt	Improve depression-like behavior, promote hippocampal neuroplasticity, and inhibit neuroinflammation
Berberine(12)	NLRP3/caspase-1/ASC/IL-1β	Improves depression-like behavior and reverses neuroplasticity damage
Aerva javanica(31)	Regulates neuroinflammation, oxidative stress and monoaminergic pathways	Improve depression-like behavior
Nootkatone(41)	NF-κB/NLRP3 inflammasome pathway	Improve depression-like behavior, inhibit proinflammatory cytokine expression, and inhibit neuroinflammation
Cinnamic acid(42)	Regulates inflammation and oxidative stress	Improve depression-like behavior, improve inflammatory cytokines and oxidative stress parameters, and regulate BDNF levels
Thymoquinone(39)	Regulates inflammatory, cytokines and BDNF	Improve depression-like behavior, reduce inflammatory cytokine mRNA expression in hippocampus and amygdala, and regulate BDNF and NeuN mRNA levels
Baicalein(43)	BDNF, TrkB, CREB	Improve depression-like behavior
Baicalin(34)	PI3K/AKT/FoxO1/TLR4	Improve depression-like behavior
Sinomenine(38)	Regulates neuroinflammation and neuroplasticity	Improve depression-like behavior
Zanthoxylum alatum(32)	Regulates hippocampal neuroinflammation and monoamine neurotransmitters	Improve depression-like behavior

Salvianolic acid B(36)	Regulates autophagy and NLRP3 inflammasome	Improve behavioral deficits and neuroinflammatory responses
Vitex megapotamica(35)	NR	Improved depression-like behavior, analgesic effect
Clove essential oil(40)	Anti-inflammatory and antioxidant	Improves depression-like behavior and cognitive memory deficits
Fustin(13)	Regulates oxidative stress, neuroinflammatory markers, NF-κB, caspase-3, BDNF	Improve depression-like behavior

Non-pharmacological interventions

Four studies of acupuncture have indicated that acupuncture modulates oxidative stress,
neuroinflammation, and apoptosis-related pathways and attenuates chronic stress-induced depressive-
like behaviours and cognitive deficits, but the mechanism of maintenance of the long-term effects of
acupuncture stimulation and how individual differences in sensitivity to acupuncture affect the
modulation of neuroinflammation-related signalling pathways have not yet been conclusively
demonstrated (14,44–46).

A study by Xie et al. from Renmin Hospital of Wuhan University, China, demonstrated that
swimming exercise increases the expression of proteins related to neuroplasticity in the
hippocampus, reduces neuroinflammation, and improves depressive behaviours. However, further
systematic studies are required to investigate the long-term regulation of neuroinflammatory
signalling pathways by different exercise intensities, frequencies, and durations, and the differential
effects of exercise among individuals of different ages, genders, and health conditions. Further
systematic studies are needed to investigate the differential effects of exercise in individuals of
different age groups, genders and health conditions (21).

A study by Universitat de Barcelona, Spain, examined the effects of brain-gut photobiomodulation
(PBM) on mice with chronic stress. Chronic stress is an important risk factor for depression and is
associated with gut microbiota dysbiosis. Existing treatments for depression are deficient, and PBM
has potential as a non-invasive therapy. Utilising an experimental model, the study ascertained that
chronic stress instigates alterations in behaviour and hippocampal protein expression in mice.
Moreover, the study determined that PBM modulates the gut microbiota, restores hippocampal Sirt1
levels, improves cognition and reduces neuroinflammation. This novel physical light therapy
approach to treat stress-related depression provides a novel direction for further research (47).

3.4 Perspective

The choice of single target versus multi-target regulation is a major point of debate in the field.Some
studies have argued in favour of the search for efficient inhibitors targeting specific key signalling
pathways. These studies suggest that precise blockade of the pathway can effectively control
neuroinflammation and thus improve depressive symptoms. For example, certain natural compounds
have been studied with a focus on their inhibitory effects on NLRP3 inflammatory vesicle activation
and have achieved some antidepressant effects (12,36,41). Conversely, it has been proposed that
depression is a multifactorial and intricate condition, and that single-target modulation may not be
sufficiently comprehensive in addressing the multifaceted nature of the disease and the heterogeneity
of individual patients. The argument is made that a more effective approach would be the
comprehensive modulation of multiple interconnected signalling pathways, such as the concurrent

modulation of autophagy, inflammation, and oxidative stress-related pathways, to achieve a more
enduring antidepressant effect. Empagliflozin's ability to simultaneously regulate multiple signalling
pathways is considered to be a significant mechanism contributing to its antidepressant effects (19).

A further point of contention for achieving clinical translational value is the comparison of the
advantages of pharmacological versus non-pharmacological interventions. Pharmacological
interventionists emphasise the precision and actionability of drugs in modulating neuroinflammatory
signalling pathways, which can act directly on the target through specific molecular mechanisms and
are easy to standardise. Non-pharmacological interventionists counter by highlighting the advantages
of natural, holistic regulation and reduced adverse effects that non-pharmacological methods, such as
acupuncture and exercise, offer. These methods can indirectly regulate neuroinflammation by
improving the physiological and psychological state of the body. From a holistic standpoint, these
approaches may be more suited to long-term maintenance therapy. However, the mechanisms
underlying non-pharmacological interventions are intricate and challenging to delineate with
precision, and the intricacies of their regulation of neuroinflammation-related signalling pathways are
yet to be thoroughly investigated.

The present study offers a plethora of novel concepts and methodologies for the management of
depression. Nevertheless, considerable progress is yet to be made before the comprehensive
resolution of the long-term stable regulation of neuroinflammation-related signalling pathways and
the circumvention of adverse effects can be achieved. Future research should focus on the following
fuzzy general directions: first, to carry out long-term follow-up studies to observe the long-term
effects and potential adverse effects of different interventions in animal models, and to analyse in
depth the dynamic changes of the signalling pathways; second, to strengthen the research on
individual difference factors (e.g., gene polymorphisms , lifestyle, intestinal flora, etc.) and to explore
the personalised treatment options; thirdly, to further explore the different interventions between
theThirdly, to further explore the joint application mode and synergistic mechanism between
different interventions, to give full play to the advantages of pharmacological and non-
pharmacological interventions, and to improve the effect and stability of depression treatment.

It is the contention of this team that the progress made thus far in animal studies is not transferable to
a clinical context. With regard to research methodology, there is a necessity for animal models to be
further optimised in order to more accurately replicate the pathophysiological processes of human
depression. While the development of new animal models for inducing depression is a more arduous
task, the enhancement of the design and implementation of clinical studies is a more attainable goal,
thereby improving the extrapolation and utility of findings. While acknowledging the value of animal
studies in identifying interventions that promote or inhibit key factors, it is crucial to recognise the
limitations in assessing the extent to which these interventions occur in actual organisms. It is highly
probable that moderation conclusions are minor or insignificant in real-world settings. To enhance
the credibility of the research, the authors' team should consider publishing the specific numbers in
addition to the published data and images, rather than relying solely on the P-value significance. The
bias caused by the credibility of the interpretation of results by the mere independent presence of P
values has been previously reported (48). In addition to the continued exploration of new drug targets
and compounds, attention must be paid to the safety and tolerability of drugs, especially with regard
to potential risks during long-term use. It is evident that discovering a promising new drug is a very
difficult process.

In the context of non-pharmacological interventions, a comprehensive analysis of the mechanisms
underlying acupuncture, exercise and other therapeutic modalities is imperative. This analysis should

encompass the specific sites and modalities of their regulation of neuroinflammation-related
signalling pathways. The objective is to formulate a more scientific and standardised intervention
protocol. However, it should be noted that acupuncture is a technique that falls under the umbrella of
traditional Chinese medicine (TCM). TCM is an empirical medicine, and the deliberate
administration of acupuncture using scientific protocols may potentially diminish the original
therapeutic effect. Interdisciplinary cooperation is crucial for comprehensive analysis of the
pathogenesis and therapeutic principles of depression, integrating multidisciplinary knowledge such
as neuroscience, immunology, psychology, genetics, etc. Unfortunately, in-depth studies from these
perspectives can erode research teams' confidence. The current ability and methods to study from the
subcellular organelle level are limited, and compared to the cost of research, cognitive therapy and
exhortation to daily exercise may be the most effective option at this time.

4   Methods

My previous initial search of the PubMed database revealed no extant systematic reviews publishing
insights on how to view anti-inflammatory strategies based on the treatment of patients with
depression-related inflammation. Therefore, I conducted a scoping review of extant clinical
randomised controlled studies based on anti-inflammatory strategies for the treatment of patients with
depression-related inflammation. The purpose of the scoping review was to sort out existing
treatment protocols, assess the rationale and consistency of the selection of inflammatory biomarkers,
evaluate the value to clinical practice, and make implementable recommendations. This scoping
review was previously registered on the OSF platform at (17,18). To ensure sufficient transparency
and detail, I followed a standard scoping review framework, first defined by Arksey and
O'Malley (49), then later expanded by Levac et al.(50). This review is reported in accordance with
the PRISMA extension for scoping reviews (51).

4.1   Search strategy and selection criteria

As clinical randomised controlled studies are the highest level of research evidence that constitutes
guidance for clinical practice, all strategies involving inflammation-based strategies whether
pharmacological or non-pharmacological regimens, but applied to patients with depression-related
inflammation, should be included. This is similar to Fei Zhao's emphasis on the inclusion of
randomised controlled studies in evidence-based medical research (5). In this scoping review,
depression was recognised by the DSM-V specification(52). Depression-associated inflammation, on
the other hand, is not nowadays recognised as a diagnostic term, and therefore inflammation was
recognised according to the Mesh entry specification in the Pubmed database (53).

I excluded non-randomised controlled studies, as well as studies that did not diagnose depression
purely as depression or co-morbidity. This is because the idea of treating depression based on
inflammatory strategies is not yet commonly used in clinical practice and remains in theoretical
discussion, and expanding the scope of the disease is not conducive to synthesising the evidence. In
addition, I excluded literature published in non-English languages, editorials, conferences, books,
advocacy or secondary research that could not be accessed in sufficient detail, which is similar to
Lorna's idea (54).

On 12 November 2024 I performed a systematic search of the PubMed database. This study was
designed as a scoping review rather than a systematic review. Therefore, Embase, the Cochrane
Library, and Web of Science were not searched due to concerns about the speed and efficiency of the
scoping review, despite the fact that multiple databases were searched more broadly and
comprehensively.This is similar to the point made by Johannes(55). I did not search Embase, the

Cochrane Library, or Web of Science. In my searches, no date constraints were applied in order to be
as comprehensive as possible. The complete search strategy is shown below:

1#: ((((("Depression"[Mesh]) OR (Depressive Symptoms[Title/Abstract])) OR (Depressive
Symptom[Title/Abstract])) OR (Symptom, Depressive[Title/Abstract])) OR (Emotional
Depression[Title/Abstract])) OR (Depression, Emotional[Title/Abstract])

2#: ((((("Inflammation"[Mesh]) OR (Inflammations[Title/Abstract])) OR (Innate Inflammatory
Response[Title/Abstract])) OR (Inflammatory Response, Innate[Title/Abstract])) OR (Innate
Inflammatory Responses[Title/Abstract])

3#: ((((("Therapeutics"[Mesh]) OR (Therapeutic[Title/Abstract])) OR (Therapy[Title/Abstract])) OR
(Therapies[Title/Abstract])) OR (Treatment[Title/Abstract])) OR (Treatments[Title/Abstract])

4#: 1# AND 2# AND 3#

Once the search was completed, I would then import the literature into Zotero Literature Manager for
screening. As the scoping review was done independently by me, I screened the literature a total of 3
times in order to avoid screening bias and reviewed the results of each screening, which were in
almost perfect agreement. The use of kappa coefficients for measuring the confidence of those who
screened the literature is not reported here as it does not involve a second person who screened the
literature.

4.2 Data charting

Data were extracted for the final included literature and the detailed programme was referred to the
research protocol(17).

(1) Basic information such as study authors, year, country, journal / publication source, financial
support and type of study.

(2) Patient participation information such as sample size, study subgroups, level of depression,
diagnostic criteria, age and gender of test and control groups.

(3) Details of pharmacological or non-pharmacological treatment regimens based on anti-
inflammatory strategies.

(4) Study results and lessons learnt.

4.3 Data synthesis

448 This scoping review primarily uses narrative synthesis to superimpose evidence and logical reasoning
449 to create a map of the inflammatory pathology of depression (**Figure 1**). In case of any implausibility
450 or lack of rigour in the research, I will use logical retrospective reasoning and add appropriate
451 solutions or recommendations.

4.4 Role of the funding source

453 No funding supported this research. The funders of the study had no role in study design, data
454 collection, data analysis, data interpretation, or writing of the article.

5   Results

5.1   Search and study characteristics

Included articles were published between 2008 and 2024 **(Table 2)**. Studies were conducted predominantly in the United Kingdom (n=3) and the China (n=3). Studies pertaining to RCT (n=12) methods were identified **(Figure 2)**.

Figure 2 PRISMA flow diagram. A total of 1057 articles were screened. Thirty-two articles met eligibility criteria, including 11 articles. This is represented in the PRISMA flow diagram. Study characteristics were extracted from primary studies only, and all 11 articles were included in the content analysis.

Table 2. Study characteristics of included articles (n=11 articles).

Study author	Year	Country	Journal/source of publication	Funding received	Study type
Gillian M. Mackay(56)	2008	United Kingdom	Clinical and Experimental Pharmacology and Physiology	No	RCT
Charles L. Raison(57)	2012	United States	JAMA psychiatry	Yes	RCT
JeremyF.Weinberger(58)	2014	United States	Brain,Behavior,andImmunity	Yes	RCT
Zhang Xiaoling(59)	2018	China	Pakistan journal of pharmaceutical sciences	No	RCT
Chen Mu-Hong(60)	2018	Taiwan (China)	Psychiatry Research	Yes	RCT
Zhi-Yong Cao(61)	2020	China	Journal of Affective Disorders	Yes	RCT
Nastaran Hariri(62)	2020	Iran	Phytotherapy research: PTR	Yes	RCT

Maria Antonietta Nettis(63)	2021	United Kingdom	Neuropsychopharmacology	Yes	RCT
Bernhard T. Baune(64)	2021	Australia	European Neuropsychopharmacology	Yes	RCT
Maria Antonietta Nettis(65)	2023	United Kingdom	Journal of Psychopharmacology	Yes	RCT
Emma Sampson(66)	2024	Germany	International Journal of Neuropsychopharmacology	Yes	RCT

5.2 Patient engagement characteristics and identification

The eleven studies included in this review collectively encompassed 907 patients diagnosed with depression, with major depression being the most prevalent form of depression (n=8). The primary diagnostic criterion employed for depression was the DSM-IV (n=7).The majority of the depressed populations studied were middle-aged. The anti-inflammatory drugs employed in the treatment of these patients included Fluoxetine (n=1), Infliximab (n=2), ketamine (n=1), Minocycline (n=2), Votioxetine + celecoxib (n=2), selective serotonin reuptake inhibitors (n=1), Lacquer Tree (n=1). The dispersion in the distribution of these drugs was large **(Table 3)**.

Table 3. Patient engagement characteristics of included studies (n=11). T indicates experimental treatment method in the study subgroup. C indicates placebo method or other common treatment. MDD indicates major depressive disorder. Sex takes the form of male/female ratios or percentages are shown. *Marked places are for median and interquartile representation. Places marked with # denote minimum and maximum values.NR denotes not reported. DSM-IV and DSM-V denote the fourth and fifth editions, respectively, of the American Psychiatric Association's Diagnostic and Statistical Manual of Mental Disorders.

Study author	Sample size	Research grouping	Degree of depression	Diagnostic criteria	Age T	Age C	Sex T	Sex C
Gillian M. Mackay(56)	51	T Fluoxetine C Psychological counseling	Moderate/MDD	DSM-IV	39.0 ± 2.4	37.5 ± 2.1	5/23	8/15
Gillian M. Mackay(56)	35	T Fluoxetine+T3 C Psychological counseling	Moderate/MDD	DSM-IV	44.3 ± 4.1	37.5 ± 2.1	4/8	8/15
Charles L. Raison(57)	60	T Infliximab C Placebo	MDD	DSM-IV	42.5±8.2	44.3±9.4	67%	67%
JeremyF.Weinberger(58)	60	T Infliximab C Placebo	MDD	DSM-IV	41.1±8.0	46.0±9.8	7/12	4/13
Zhang Xiaoling(59)	130	T C	NR	NR	36.2±5.4	35.7±6.2	30/35	29/36
Chen Mu-Hong(60)	NR	T 0.5mg/kg ketamine C Placebo	MDD	DSM-IV	48.46±11.01	48.63±8.12	12.50%	37.50%
Chen Mu-Hong(60)	NR	T 0.2mg/kg ketamine C Placebo	MDD	DSM-IV	44.96±12.31	48.63±8.12	26.10%	37.50%
Zhi-Yong Cao(61)	56	T SSRI + GZA C SSRI + PBO	MDD	DSM-V	28.50(21.00, 33.25)	23.00 (21.00, 28.00)	14/12	23/7
Nastaran Hariri(62)	60	T Lacquer tree C Placebo	Mild/moderate	DSM-IV	42.19±8.44	44.03±11.80	0	0
Maria Antonietta Nettis(63)	44	T Minocycline C placebo	MDD	DSM-V	47.0±10.0	43.7±10.7	44.4%	42.90%
Bernhard T. Baune(64)	200	T Votioxetine + celecoxib C votioxetine + placebo	MDD	DSM-IV	45(33.5, 45)	48(30.5, 48)	46%	38%
Maria Antonietta Nettis(65)	44	T Minocycline C placebo	MDD	DSM-V	25-60	25-60	NR	

| Emma Sampson(66) | 119 | T Votioxetine + celecoxib
C votioxetine + placebo | MDD | DSM-IV | 49(39.5, 57) | 47(29, 54) | 15/28 | 14/31 |

5.3 Time and inflammatory index

In the eleven studies that incorporated drug treatment cycles ranging from two to eighteen weeks, the vast majority of inflammatory markers examined cumulatively were undetectable in depressed patients. The inflammatory markers that were reasonably detectable in patients with depression in the clinical studies included CRP, hs-CRP, sTNFR1, sTNFR2, IL-2, IL-6, and TNF-α. The utilisation of hs-CRP as an inflammation indicator could be approached via a stratified strategy, e.g., by establishing cut-off levels for hsCRP at 3 mg/L. The benefits of this stratification remain to be elucidated.The merits of such stratified analyses are yet to be demonstrated. A thorough review of the extant literature reveals that the concept of inflammation remains ambiguous in these studies, with the methodology for assessing the dynamic recovery of depression still relying on traditional depression rating scales. Furthermore, there is a paucity of investigation into the concept of inflammation and the indicators of inflammation mentioned in the study's methodology **(Table 4)**.

Table 4. Time and inflammatory index. The following section provides an explanation of the abbreviated terms. IL-2: Interleukin-2; CRP: C-reactive protein; hs-CRP: High-sensitivity C-reactive protein; sTNFR1: Soluble Tumor Necrosis Factor Receptor 1; sTNFR2: Soluble Tumor Necrosis Factor Receptor 2; IL-6: Interleukin-6; TNF-α: Tumor Necrosis Factor-α; IFN-γ: Interferon-γ; IL-1β: Interleukin-1β; IL-4: Interleukin-4; IL-8: Interleukin-8; IL-10: Interleukin-10; IL-12p70: Interleukin-12p70; IL-13: Interleukin-13.

Study author	Time	Inflammatory index
Gillian M. Mackay(56)	18W	IL-2, CRP
Charles L. Raison(57)	12W	hs-CRP
JeremyF.Weinberger(58)	8W	hs-CRP, sTNFR1, sTNFR2
Zhang Xiaoling(59)	6W	IL-2, IL-6, TNF-α
Chen Mu-Hong(60)	2W	CRP, IL-6, TNF-α
Zhi-Yong Cao(61)	4W	CRP, IL-6, TNF-α, IL - 1β, IFN-γ, cortisol
Nastaran Hariri(62)	12W	CRP, IL-6, TNF-α
Maria Antonietta Nettis(63)	4W	hs-CRP, IL-6
Bernhard T. Baune(64)	6W	hs-CRP
Maria Antonietta Nettis(65)	4W	CRP, hs-CRP, IFN-γ, TNF-α, IL-1β, IL-2, IL-4, IL-6, IL-8, IL-10, IL-12p70, IL-13
Emma Sampson(66)	6W	hs-CRP

5.4 Benefits and challenges of patient engagement

Anti-inflammatory strategies have been shown to have the potential to improve symptoms in the treatment of depression. Studies using glycopyrrolate (GZA) as an adjunctive treatment found that

patients treated with an SSRI + GZA had more pronounced remission of depressive symptoms and
higher rates of treatment response and remission at 2 and 4 weeks of treatment compared to the
placebo group. This finding suggests that by reducing inflammation, it may enhance the mood
symptoms of depressed patients and improve treatment outcomes. Anti-inflammatory treatments help
to regulate the inflammatory state in depressed patients. A study of overweight or obese depressed
women found that supplementation with lacuna (Rhus coriaria L.) in combination with a calorie-
restricted diet significantly reduced the patients' body weight, BMI, body fat, visceral fat levels, and
levels of MDA, an indicator of oxidative stress. Additionally, serum levels of IL-6 and TNF-alpha
(although not to a significant level) and hs-CRP levels were significantly reduced. This finding
suggests that anti-inflammatory strategies may have a positive effect on the inflammatory state of the
body in patients with depression by regulating the levels of inflammatory factors, thereby improving
the condition.

A significant challenge for precision treatment is the existence of substantial inter-patient variability
in response to anti-inflammatory treatments, as evidenced by a study of minocycline in patients with
treatment-resistant depression. While no substantial difference was observed in terms of
improvement in depressive symptoms between the minocycline and placebo groups, further stratified
analyses revealed that patients with baseline hsCRP levels ≥3 mg/L exhibited a more favourable
response to minocycline, while those with <3 mg/L demonstrated a less favourable response. This
finding underscores the notion that a patient's baseline inflammation level exerts a significant
influence on their response to anti-inflammatory therapy. Consequently, there is an imperative for the
development of more precise biomarkers to facilitate the screening of patients who may benefit from
personalised treatment regimens. A further critical concern pertains to the safety of anti-inflammatory
drugs, which is a pivotal consideration in clinical practice. Some studies have identified potential
risks associated with certain drugs. For instance, celecoxib, when studied as an adjunct to the
antidepressant vortioxetine, was found to be non-inferior to a placebo in the treatment of depression.
However, the drug itself carries a risk of side effects. Therefore, when selecting anti-inflammatory
drugs for depression treatment, it is crucial to carefully weigh efficacy against safety to ensure that
the benefits outweigh the risks for patients during treatment.

Inflammatory marker levels might be able to predict how a patient will respond to anti-inflammatory
treatment. For example, the levels of certain markers like CRP and TNF at the start of treatment have
been linked to how well a drug called infliximab works in treating severe depression. Patients with
higher starting levels of these markers tend to do better with infliximab, while those with lower CRP
levels do better with a placebo. This finding underscores the potential of inflammatory markers as
significant predictors of treatment response, thereby aiding in the informed decision-making process
for clinical practitioners.

I still don't fully understand how inflammatory markers affect the treatment of depression, and more
research is needed to figure out what they mean and how important they are. One study looked at
fluoxetine and escitalopram for treating depression. It found that levels of inflammatory factors like
IL - 2, IL - 6, and TNF - α went down a lot in both groups after treatment. But the drop was bigger in
the escitalopram group, and this was related to how much the depressive symptoms improved. This
finding suggests that distinct inflammatory factors may contribute differently to the pathophysiology
of depression and respond variably to treatment.

Further research is necessary to elucidate their mechanisms of action. Anti-inflammatory strategies
for treating depression have shown benefits only in cases where patients exhibit high inflammatory
factors. However, these strategies also face challenges, such as individual differences and drug

safety. Inflammation-related biomarkers have been found to be strongly associated with treatment
efficacy. As outlined in **Table 3**, the cost of testing for these inflammatory indicators in a standard
hospital in China ranges from 1000 to 1500 RMB, while traditional therapies, based on the
experience of our group's common discussion that 1000 RMB is sufficient to complete a course of
depression treatment in any hospital in China. It is evident that the drug regimens employed in
conjunction with anti-inflammatory strategies or those intended to calm inflammation are less
generalisable.

Future research should address the following questions:

(1) How can we find out if a depressed patient has a strong inflammatory response?

(2) Should the goal of treating depression with anti-inflammatory methods be to stop the
inflammation or just to make the depression better?

(3) If a patient takes both regular drugs and anti-inflammatory drugs, how do we classify these
drugs?

(4) Do psychiatrists have a clear list of which anti-inflammatory drug combinations to use?

(5) Can using anti-inflammatory drugs help depressed patients take their medicine more regularly and
get better faster?

The above five questions should be thoroughly investigated in the development of anti-inflammatory-
based treatments for depression, and we call on psychiatrists, neurological surgeons, and
psychotherapists to be open and transparent in their research and to publish their findings.

6   Discussion

A scoping review was conducted to identify examples of medications based on anti-inflammatory
strategies for the treatment of depression. This review yielded a substantial amount of information for
future research in this field. While the prevalence of anti-inflammatory strategy-based medications
for depression is currently low, the included studies accommodated 907 patients in total, with
treatment based on anti-inflammatory strategies exhibiting significant variation.

The findings of the present literature review section revealed that the immune mechanisms of
inflammation do not apply to depression. In cellular and animal studies, experimental models and
intervention protocols are fixed, which facilitates the placement of confounding factors within a
manageable range. The role of immune inflammation in depression is convincing without taking into
account the multitude of factors that contribute to the pathogenesis of depression, and is limited to
the specific way in which the experimental model is modelled. A particular experimental model of
depression itself stimulates depression through inflammatory immune mechanisms (67,68). However,
the conditions under which humans acquire depression are uncertain and could be inheritance,
environmental changes, acute event stress, etc. (69,70). We hypothesise that some specific triggers of
depression are consistent with inflammatory immune mechanisms. Yet, triggers of depression are not
usually considered in clinical guidelines for the treatment of depression. Clinical experience within
our team has revealed that depression is treated with psychotherapy (the specific type of which
depends on the patient's preferences) when available, and in cases where psychotherapy is ineffective
or of low benefit, with antidepressant medication. For patients with depression accompanied by
underlying diseases, priority is given to treating the underlying diseases, with psychiatry providing

assistance in the synergistic treatment of depression. In the context of depression accompanied by
underlying diseases, the hypothesis that immunoinflammation alone can explain the pathological
mechanisms of depression is not substantiated. The emergence of anti-inflammatory strategies for the
treatment of depression appears to be an endeavour to circumvent the conventional treatment
pathway for depression by exploring an alternative inflammatory-immune etiological pathway. A
comparison of the potential targets mentioned in the literature review section with the treatment
protocols in the randomised controlled studies suggests that this subversive translation is inadequate.
Secondly, the new approach to treating depression is much more expensive than the traditional
methods. This is because the new pathway may involve more costly drugs, additional tests to monitor
inflammation levels, or longer treatment periods, which all contribute to the higher overall cost.

As demonstrated in the literature review section of this paper, the feasibility of treating depression
based on anti-inflammatory strategies is derived from inflammatory vesicles (as hypothesised). The
hypothesis is that stress activates the NLRP3 inflammasome, which prompts caspase-1 to cleave
precursor forms of interleukin-1β (pro-IL-1β) and interleukin-18 (pro-IL-18). The resultant process
of NLRP3 inflammasome activation produces the mature inflammatory cytokines IL-1β and IL-18
(71). These cytokines then activate microglia and astrocytes, which, in turn, release interleukin-6 (IL-
6), tumour necrosis factor-α (TNF-α), and other molecules (72). This process of increased release of
inflammatory mediators (73) could explain why inflammatory indicators in clinical studies frequently
refer to IL-1β, IL-6 and TNF-α. However, one of the clinical studies reported that the vast majority of
the cytokines were below the detected levels (65), which allows two inferences to be drawn.

The initial hypothesis is that inflammatory vesicles are inadequate in explaining the inflammatory
pathogenesis of depression. The secondary hypothesis is that the inflammation indicator is not
reasonably set. In order to comprehensively discuss the potential clinical application of the
inflammatory pathological mechanism of depression, it is hypothesized that the inflammation
indicator is not set up rationally. A salient point is that, according to the activation process of
inflammatory vesicles, the reduction of these vesicles in the blood of depressed patients to normal
range levels by anti-inflammatory treatment is indicative of the therapeutic endpoint being reached
(this set of assumptions is not mentioned in the 11 studies).

It is recommended, based on the assumptions outlined above, that intravenous nano-micro AI robots
be used for patients diagnosed with depression, thus addressing the current research deficit in the
field. This recommendation is made on the condition that patients are fully informed of the
advantages and disadvantages of this intervention, and that the treatment protocol for intravenous
nano-mini-AI robots has been approved by an ethics committee following a thorough review. The
present state of AI development is constrained by the utilisation of machine learning algorithms that
employ vast informational databases stored in remote servers. This technological framework is
inadequate for the comprehension of the intricacies inherent in the human body.Nevertheless, the
prevailing strategy for addressing depression, which is predicated on anti-inflammatory
methodologies, exhibits a dearth of research endeavours concerning the analysis of diverse
inflammatory markers within the human organism and the extrapolation of depression treatment
outcomes. The integration of an AI robot, capable of being implanted into the human body via
intravenous infusion, signifies a potential paradigm shift, promising a future replete with
possibilities.

It is imperative to elucidate the following stages of AI development:(1) The production of all
inflammatory mediators or cytokines during the activation of inflammatory vesicles is detected using
transcriptomics or single-cell sequencing. All potential inflammatory markers detected are then

analysed for differences using a healthy adult population and a depressed adult population. This
process results in the identification of the first potentially usable inflammatory markers. (2)
Considering the rapidity of clinical application, it is necessary to analyse the first batch of
inflammation indicators in the blood level again to obtain the second batch of inflammation
indicators. (3) Considering the difficulty for clinicians to interpret the data of the inflammation
indicators, it is necessary to establish the second batch of inflammation indicators in the healthy adult
population to establish a constant model, so that we can get the minimum and maximum values of
each inflammation indicator in the healthy adult population. (4) Inflammation indicators that
establish a norm in the healthy adult population can be used as the final inflammation indicators. (5)
At this stage, there is no accurate and objective assessment method for the diagnosis and evaluation
of depression in the medical field, and we need to rely on the subjective judgement of psychiatrists
and psychologists. The question of whether the final inflammation index can become the gold
standard for assessing depression needs to be addressed by psychiatrists and psychologists rather than
psychological rating scales (the protocol for assessing the dynamic treatment effect of depression
mentioned in 11 studies is the change in the score of psychological rating scales).

In the context of contemporary clinical randomised controlled studies, the methodology employed for
the selection of inflammatory indicators remains largely opaque, thereby engendering a sense of
distrust among researchers and practitioners who engage with the extant literature. This opacity
hinders the accumulation and advancement of knowledge in the field of clinical translational research
on the pathological mechanisms of inflammation in depression. To address these concerns, we
propose a systematic approach to the organisation of evidence, commencing from the initial phase
delineated above. This approach involves the integration of artificial intelligence (AI) technologies to
enhance the transparency and reliability of the research process.

It is hypothesised that the AI could assist researchers, psychiatrists and patients with depression to
work together to develop new conceptual therapies based on anti-inflammatory strategies. For
example, AI has been successfully applied to improve diagnosis and treatment planning in pituitary
adenoma cases, suggesting a model for depression therapy development(74,75).

This hypothesis is based on past experience in developing a shared health needs assessment for the
AI integration platform (6). Patients can take information from different test results they get at the
hospital and put it into the AI system for analysis(76). The AI system will then carefully look at all
the information and keep it up to date as new reports come in. This way, it creates a record of the
patient's health that is focused on the patient, similar to what a hospital would have(77).

The AI integration platform uses closed-source computing on patient data, drawing from a database
of all user data on the master server and Internet data (78). This allows patients to easily access their
depression health assessment reports without training. It also provides personalized report
interpretations. If the report links inflammation to depression, patients can choose anti-inflammatory
medication, strengthening treatment beliefs. During treatment, patients can submit medical records to
the platform, which projects medication and health changes. Early warning results are sent to doctors,
enhancing treatment effectiveness. AI technology improves patient care, reduces medical waste, and
helps doctors quickly identify and treat abnormal health data, enabling more precise decisions and
better treatment plans.

A research team at Harbin Institute of Technology has developed a micro-robot(79). It can mimic
white blood cells, moving through cell gaps to reach blood vessel lumens. The robot can carry
medications for targeted treatment, thanks to its internal structure that enables signal transmission for

real-time feedback on its internal environment. This real - time feedback is similar to that from
imaging and blood tests. The micro - robot has the potential to reduce the economic burden of
monitoring inflammation in depressed patients. As the technology advances, it may interact with AI
in the future**(Figure 3)**. Wang et al.'s research also supports the proposed AI machine therapy
hypothesis(80).

Figure 3. A prospective micro-robot application for interaction with depressed patients. AI
microbots treating depression. After patients give consent, microbots are administered into the body.
They start a 24-hour continuous body examination, uploading data hourly to the healthcare provider's
computer. When an inflamed area is detected, AI compares it with normal data. If abnormal,
microbots dispense medication. If not, they notify the provider and help with treatment until the
patient's data is normal. After recovery, with consent, microbots in suspension re-enter the body.
They resume 24-hour, hourly-updated body checks, continuously monitoring the patient.

In addition, existing molecular pathological explanations of depression suggest that abnormalities in
the neurochemical metabolism of 5-hydroxytryptamine and dopamine are the primary biological
pathogenesis of depression. Such lesions within the brain are usually not easily detectable, even when
imaging techniques are used. Here we make use of other brain lesions and difficult to detect diseases
(pituitary adenomas). This allows a side-by-side assessment of the need for AI and avoids the bias of
depression assessment alone. Zhu et al. reviewed 14 cases of ectopic pituitary adenomas and noted
that ectopic pituitary adenomas in the cavernous sinus and suprasellar space were easily
misdiagnosed as suspicious intrasellar masses on imaging. Apparently, it is difficult to accurately
diagnose ectopic pituitary adenomas even with imaging data. Such a clinical assessment dilemma is a
great challenge for clinicians (81). Li et al. reviewed previously treated thyrotropin-secreting
pituitary adenomas and proposed the new idea of 'partial remission of pituitary adenomas alone'.
However, the clinical presentation of pituitary adenoma patients is insidious and difficult to track,
especially with the suspicion of a co-secretory tumour (82). How can clinicians input data from
pituitary adenoma patients into an AI platform that assesses the disease progression and insidious
clinical presentation of patients with pituitary adenomas, and what are the outcomes? It is possible
that the poor prognosis of the current pituitary adenoma patient will be known sooner than the human

physician. Previously treated suprasellar pituitary adenomas were reviewed by Zhu et al. Nine
patients were treated with extended endoscopic transsphenoidal surgery and only one tumour
recurred without serious postoperative complications. The implication of this retrospective study is
that extended endoscopic transsphenoidal surgery may be a safe and effective method of resecting
these tumours. In type II Suprasellar pituitary adenomas, the pituitary stalk should be carefully
protected intraoperatively. This protection of the pituitary stalk will be easier to achieve if an AI
platform is available to assist the surgeon during the operation (83). Li et al. reviewed 232 previous
cases of endocrine pituitary tumour disease in children at Peking Union Medical College Hospital.
This would have reduced the presentation of cystic changes or haemorrhage by approximately 25% if
the diagnosis had been made early and treated aggressively (84). Based on these four papers, we
believe that the current dilemma of how to accurately identify hard-to-detect lesions in the brain
cannot be overcome in a short period of time without the aid of AI to assist the judgement of human
doctors.

7   Conclusion

At present, the use of anti-inflammatory strategies for the pharmacological treatment of depression
has limited research value and poor feasibility. The future direction of the new concept of anti-
inflammatory strategies for the treatment of depression, proposed in the context of the association
between inflammation and depression, is that psychiatrists, researchers, and psychotherapists should
shift their future focus from pharmacological treatments based on anti-inflammatory strategies to
non-pharmacological treatments of anti-inflammatory strategies, such as positive thinking, exercise,
and so on. The popularity of purely clinical randomised controlled studies in the depression
population is extremely low when considering the financial investment in research and the benefits of
translating the results. In the future, public interest studies, low research costs, and research protocols
with mass generalisability will be more likely to stimulate the depression community's interest in
participating in research.

8   Conflict of Interest

734 The authors declare that the research was conducted in the absence of any commercial or financial
735 relationships that could be construed as a potential conflict of interest.

9   Acknowledgments

This scoping review was previously registered on the OSF platform. Pre-established research team
members can be accessed in the registration link (https://doi.org/10.17605/OSF.IO/A64GC).

To ensure that authentic authorship qualifications meet authorship standards
(https://www.nature.com/nature-portfolio/editorial-policies/authorship). Here, I would like to thank
those who have helped in the past in the preliminary stages of this research.

I would like to thank Miss Fengjiao Zhao (Shantou University Medical College), Miss Jingyi Li
(North China University of Science and Technology) and Miss Yuanyin Deng (Zhejiang University)
for providing assistance in document delivery. And I would like to thank Mr. Yifei Chen (The
Affiliated Hospital of Yangzhou University), Hsu Yi Liang (Monash University), Haoran Dai
(Nanjing Medical University) and Wenjie Jiang (Northwest Minzu University) for providing
assistance in images drawing. Moreover, I would like to thank Miss Ren Sha (Inner Mongolia
Chaopai New Materials Co., Ltd.) and Youwei Wang (Beihua University) for providing assistance in

management and budget support. What's more, I would like to thank Mr. Ming Wu (Gongli Hospital
affiliated to Second Military Medical University) for providing assistance.

I make sure that the contributions of the collaborators mentioned in this acknowledgement are
genuine and valid.

10   Reference

1.   Beurel E, Toups M, Nemeroff CB. The Bidirectional Relationship of Depression and
Inflammation: Double Trouble. *Neuron* (2020) 107:234–256. doi:
10.1016/j.neuron.2020.06.002

2.   Andrade C. Anti-Inflammatory Treatments for Depression: Perspectives on How to Read a
Meta-Analysis Critically. *J Clin Psychiatry* (2019) 80:19f12907. doi: 10.4088/JCP.19f12907

3.   Bai S, Guo W, Feng Y, Deng H, Li G, Nie H, Guo G, Yu H, Ma Y, Wang J, et al. Efficacy and
safety of anti-inflammatory agents for the treatment of major depressive disorder: a systematic
review and meta-analysis of randomised controlled trials. *J Neurol Neurosurg Psychiatry*
(2020) 91:21–32. doi: 10.1136/jnnp-2019-320912

4.   Köhler O, Benros ME, Nordentoft M, Farkouh ME, Iyengar RL, Mors O, Krogh J. Effect of
anti-inflammatory treatment on depression, depressive symptoms, and adverse effects: a
systematic review and meta-analysis of randomized clinical trials. *JAMA Psychiatry* (2014)
71:1381–1391. doi: 10.1001/jamapsychiatry.2014.1611

5.   Zhao F. Is cognitive behavioral therapy an efficacious treatment for psychological interventions
in body dysmorphic disorders? A meta-analysis based on current evidence from randomized
controlled trials. *J Affect Disord* (2024)

6.   Miao C, Bo Y, Xu J, Wu Y, Shen J, Fan J, Wang Y, Li WT. Health Needs Assessment:
Comparison of Applications of All-in-One AI Platform during the COVID-19 Pandemic
between Mainland China and Hong Kong. *Am J Health Behav* (2024)

7.   Liang Q. Effects of home-based telemedicine and mHealth interventions on blood pressure in
stroke patients: a systematic evaluation and meta-analysis of randomized controlled trials. *J
Stroke Cerebrovasc Dis* (2024)

8.   Becher B, Spath S, Goverman J. Cytokine networks in neuroinflammation. *Nat Rev Immunol*
(2017) 17:49–59. doi: 10.1038/nri.2016.123

9.   Brenton JN, Banwell B, Bergqvist AGC, Lehner-Gulotta D, Gampper L, Leytham E, Coleman
R, Goldman MD. Pilot study of a ketogenic diet in relapsing-remitting MS. *Neurol
Neuroimmunol Neuroinflammation* (2019) 6:e565. doi: 10.1212/NXI.0000000000000565

10.   Cheng Y, Pardo M, Armini R de S, Martinez A, Mouhsine H, Zagury J-F, Jope RS, Beurel E.
Stress-induced neuroinflammation is mediated by GSK3-dependent TLR4 signaling that
promotes susceptibility to depression-like behavior. *Brain Behav Immun* (2016) 53:207–222.
doi: 10.1016/j.bbi.2015.12.012

11.   Zhang W, Zhang D, Liu S, Wang H, Liu X, Dai C, Fang Y, Fan Y, Wei Z, Feng M, et al.
Predicting delayed remission in Cushing's disease using radiomics models: a multi-center study.
Front Oncol (2024) 13:1218897. doi: 10.3389/fonc.2023.1218897

12.   Qin Z, Shi D-D, Li W, Cheng D, Zhang Y-D, Zhang S, Tsoi B, Zhao J, Wang Z, Zhang Z-J.
Berberine ameliorates depression-like behaviors in mice via inhibiting NLRP3 inflammasome-
mediated neuroinflammation and preventing neuroplasticity disruption. *J Neuroinflammation*
(2023) 20:54. doi: 10.1186/s12974-023-02744-7

13.   Bawadood AS, Al-Abbasi FA, Alghamdi AM, Alqurashi MM, Sheikh RA, Alzarea SI, Sayyed
N, Kazmi I. Fustin alleviates lipopolysaccharide-induced anxiety-depression-like performances
by modulation of oxidative stress/neuroinflammatory markers/ NF-κB/caspase-3/BDNF
expression in rodents. *Eur Rev Med Pharmacol Sci* (2024) 28:419–432. doi:
10.26355/eurrev_202401_34931

14.   Chen W, Chen Y, Cheng W, Li P, Shen J, Tong T, Lai L, Yan S, Huang Z, Li J, et al.
Acupuncture exerts preventive effects in rats of chronic unpredictable mild stress: The
involvement of inflammation in amygdala and brain-spleen axis. *Biochem Biophys Res*
Commun (2023) 646:86–95. doi: 10.1016/j.bbrc.2023.01.046

15.   Wang H, Zhang W, Li S, Fan Y, Feng M, Wang R. Development and Evaluation of Deep
Learning-based Automated Segmentation of Pituitary Adenoma in Clinical Task. *J Clin*
Endocrinol Metab (2021) 106:2535–2546. doi: 10.1210/clinem/dgab371

16.   Miller AH, Raison CL. The role of inflammation in depression: from evolutionary imperative to
modern treatment target. *Nat Rev Immunol* (2016) 16:22–34. doi: 10.1038/nri.2015.5

17.   Bo Y. AI perspectives on the present and future of antidepressant pharmaceutical treatment
based on anti-inflammatory strategies: A scoping review protocol of randomised controlled
clinical trials. (2024) osf.io/k9pue

18.   Riegelman A, Topor MK, van Veggel N, Yeung SK, Call M, Mellor DT. Coles5, Katherine S.
Corker6, Gilad Feldman7, David Moreau8, Thomas Nordström4, Jade S. Pickering9,.
https://doi.org/g5fj

19.   Muhammad RN, Albahairy MA, Abd El Fattah MA, Ibrahim WW. Empagliflozin-activated
AMPK elicits neuroprotective properties in reserpine-induced depression via regulating
dynamics of hippocampal autophagy/inflammation and PKCζ-mediated neurogenesis.
Psychopharmacology (Berl) (2024) 241:2565–2584. doi: 10.1007/s00213-024-06663-0

20.   Wang C, Cui C, Xie X, Chen B, Feng L, Jiang P. Calcitriol attenuates lipopolysaccharide-
induced neuroinflammation and depressive-like behaviors by suppressing the
P2X7R/NLRP3/caspase-1 pathway. *Psychopharmacology (Berl)* (2024) 241:1329–1343. doi:
10.1007/s00213-024-06565-1

21.   Xie Y, Wu Z, Sun L, Zhou L, Xiao L, Wang H, Wang G. Swimming exercise reverses chronic
unpredictable mild stress-induced depression-like behaviors and alleviates neuroinflammation
and collapsing response mediator protein-2-mediated neuroplasticity injury in adult male mice.
Neuroreport (2022) 33:272–282. doi: 10.1097/WNR.0000000000001779

22. Ma M, Ren Q, Yang C, Zhang J-C, Yao W, Dong C, Ohgi Y, Futamura T, Hashimoto K.
Antidepressant effects of combination of brexpiprazole and fluoxetine on depression-like
behavior and dendritic changes in mice after inflammation. *Psychopharmacology (Berl)* (2017)
234:525–533. doi: 10.1007/s00213-016-4483-7

23. Bao H, Yan J, Huang J, Deng W, Zhang C, Liu C, Huang A, Zhang Q, Xiong Y, Wang Q, et al.
Activation of endogenous retrovirus triggers microglial immuno-inflammation and contributes
to negative emotional behaviors in mice with chronic stress. *J Neuroinflammation* (2023) 20:37.
doi: 10.1186/s12974-023-02724-x

24. Rostevanov IS, Betesh-Abay B, Nassar A, Rubin E, Uzzan S, Kaplanski J, Biton L, Azab AN.
Montelukast induces beneficial behavioral outcomes and reduces inflammation in male and
female rats. *Front Immunol* (2022) 13:981440. doi: 10.3389/fimmu.2022.981440

25. Li J, Li Y, Duan W, Zhao Z, Yang L, Wei W, Li J, Li Y, Yu Y, Dai B, et al. Shugan granule
contributes to the improvement of depression-like behaviors in chronic restraint stress-
stimulated rats by altering gut microbiota. *CNS Neurosci Ther* (2022) 28:1409–1424. doi:
10.1111/cns.13881

26. Nouri A, Hashemzadeh F, Soltani A, Saghaei E, Amini-Khoei H. Progesterone exerts
antidepressant-like effect in a mouse model of maternal separation stress through mitigation of
neuroinflammatory response and oxidative stress. *Pharm Biol* (2020) 58:64–71. doi:
10.1080/13880209.2019.1702704

27. Casaril AM, Domingues M, Bampi SR, de Andrade Lourenço D, Padilha NB, Lenardão EJ,
Sonego M, Seixas FK, Collares T, Nogueira CW, et al. The selenium-containing compound 3-
((4-chlorophenyl)selanyl)-1-methyl-1H-indole reverses depressive-like behavior induced by
acute restraint stress in mice: modulation of oxido-nitrosative stress and inflammatory pathway.
Psychopharmacology (Berl) (2019) 236:2867–2880. doi: 10.1007/s00213-018-5151-x

28. Celebi G, Gocmez SS, Ozer C, Duruksu G, Yazır Y, Utkan T. Propolis prevents vascular
endothelial dysfunction by attenuating inflammation and oxidative damage in the chronic
unpredictable stress model of depression in rats. *J Pharm Pharmacol* (2023) 75:1418–1429.
doi: 10.1093/jpp/rgad071

29. Hu L, Zeng X, Yang K, Peng H, Chen J. n-3 polyunsaturated fatty acids improve depression-
like behavior by inhibiting hippocampal neuroinflammation in mice via reducing TLR4
expression. *Immun Inflamm Dis* (2022) 10:e707. doi: 10.1002/iid3.707

30. Huang N, Wang Y, Zhan G, Yu F, Li S, Hua D, Jiang R, Li S, Wu Y, Yang L, et al.
Contribution of skeletal muscular glycine to rapid antidepressant effects of ketamine in an
inflammation-induced mouse model of depression. *Psychopharmacology (Berl)* (2019)
236:3513–3523. doi: 10.1007/s00213-019-05319-8

31. Arshad HM, Ahmad F-U-D, Lodhi AH. Methanolic Extract of Aerva javanica Leaves Prevents
LPS-Induced Depressive Like Behavior in Experimental Mice. *Drug Des Devel Ther* (2022)
16:4179–4204. doi: 10.2147/DDDT.S383054

32. Barua CC, Haloi P, Saikia B, Sulakhiya K, Pathak DC, Tamuli S, Rizavi H, Ren X.
Zanthoxylum alatum abrogates lipopolysaccharide-induced depression-like behaviours in mice

by modulating neuroinflammation and monoamine neurotransmitters in the hippocampus.
Pharm Biol (2018) 56:245–252. doi: 10.1080/13880209.2017.1391298

33.   Gong M, Wang J, Song L, Wu X, Wang Y, Li B, Zhang Y, Qin L, Duan Y, Long B. Role of
BDNF-TrkB signaling in the antidepressant-like actions of loganin, the main active compound
of Corni Fructus. *CNS Neurosci Ther* (2023) 29:3842–3853. doi: 10.1111/cns.14305

34.   Guo L-T, Wang S-Q, Su J, Xu L-X, Ji Z-Y, Zhang R-Y, Zhao Q-W, Ma Z-Q, Deng X-Y, Ma S-
P. Baicalin ameliorates neuroinflammation-induced depressive-like behavior through inhibition
of toll-like receptor 4 expression via the PI3K/AKT/FoxO1 pathway. *J Neuroinflammation*
(2019) 16:95. doi: 10.1186/s12974-019-1474-8

35.   Hamann FR, Zago AM, Rossato MF, Beck VR, Mello CF, de Brum TF, de Carvalho LM,
Faccin H, Oliveira SM, Rubin MA. Antinociceptive and antidepressant-like effects of the crude
extract of Vitex megapotamica in rats. *J Ethnopharmacol* (2016) 192:210–216. doi:
10.1016/j.jep.2016.07.045

36.   Jiang P, Guo Y, Dang R, Yang M, Liao D, Li H, Sun Z, Feng Q, Xu P. Salvianolic acid B
protects against lipopolysaccharide-induced behavioral deficits and neuroinflammatory
response: involvement of autophagy and NLRP3 inflammasome. *J Neuroinflammation* (2017)
14:239. doi: 10.1186/s12974-017-1013-4

37.   Li P, Zhang F, Li Y, Zhang C, Yang Z, Zhang Y, Song C. Isoginkgetin treatment attenuated
lipopolysaccharide-induced monoamine neurotransmitter deficiency and depression-like
behaviors through downregulating p38/NF-κB signaling pathway and suppressing microglia-
induced apoptosis. *J Psychopharmacol Oxf Engl* (2021) 35:1285–1299. doi:
10.1177/02698811211032473

38.   Liu S, Xu S, Wang Z, Guo Y, Pan W, Shen Z. Anti-Depressant-Like Effect of Sinomenine on
Chronic Unpredictable Mild Stress-Induced Depression in a Mouse Model. *Med Sci Monit Int*
Med J Exp Clin Res (2018) 24:7646–7653. doi: 10.12659/MSM.908422

39.   Nazir S, Farooq RK, Nasir S, Hanif R, Javed A. Therapeutic effect of Thymoquinone on
behavioural response to UCMS and neuroinflammation in hippocampus and amygdala in
BALB/c mice model. *Psychopharmacology (Berl)* (2022) 239:47–58. doi: 10.1007/s00213-021-
06038-9

40.   Sung W-W, Yeh T-M, Shih W-L. Additive effect of clove essential oil combined with hydrogen
inhalation improves psychological harm caused by lipopolysaccharide in mice. *BMC*
Complement Med Ther (2024) 24:399. doi: 10.1186/s12906-024-04682-0

41.   Zhao X-H, An N, Xia M-H, Liu W-P, Wang Q-Q, Bao J-Z. Nootkatone Improves Chronic
Unpredictable Mild Stress-Induced Depressive-Like Behaviors by Repressing NF-κB/NLRP3-
Mediated Neuroinflammation. *Chin J Integr Med* (2023) 29:37–43. doi: 10.1007/s11655-022-
3725-2

42.   Zhuo R, Cheng X, Luo L, Yang L, Zhao Y, Zhou Y, Peng L, Jin X, Cui L, Liu F, et al.
Cinnamic Acid Improved Lipopolysaccharide-Induced Depressive-Like Behaviors by Inhibiting
Neuroinflammation and Oxidative Stress in Mice. *Pharmacology* (2022) 107:281–289. doi:
10.1159/000520990

43. Zhao X, Kong D, Zhou Q, Wei G, Song J, Liang Y, Du G. Baicalein alleviates depression-like
behavior in rotenone- induced Parkinson's disease model in mice through activating the
BDNF/TrkB/CREB pathway. *Biomed Pharmacother Biomedecine Pharmacother* (2021)
140:111556. doi: 10.1016/j.biopha.2021.111556

44. Jia Z, Yu W, Li X, Dong T, Wang X, Li J, Yang J, Liu Y. Du-moxibustion ameliorates
depression-like behavior and neuroinflammation in chronic unpredictable mild stress-induced
mice. *J Affect Disord* (2024) 358:211–221. doi: 10.1016/j.jad.2024.05.025

45. Shen J, Hao C, Yuan S, Chen W, Tong T, Chen Y, Shahzad Aslam M, Yan S, Li J, Zeng J, et al.
Acupuncture alleviates CUMS-induced depression-like behaviors of rats by regulating oxidative
stress, neuroinflammation and ferroptosis. *Brain Res* (2024) 1826:148715. doi:
10.1016/j.brainres.2023.148715

46. Tong T, Hao C, Shen J, Liu S, Yan S, Aslam MS, Chen Y, Chen W, Li J, Li Y, et al.
Electroacupuncture ameliorates chronic unpredictable mild stress-induced depression-like
behavior and cognitive impairment through suppressing oxidative stress and
neuroinflammation in rats. *Brain Res Bull* (2024) 206:110838. doi:
10.1016/j.brainresbull.2023.110838

47. Sancho-Balsells A. Brain–gut photobiomodulation restores cognitive alterations in chronically
stressed mice through the regulation of Sirt1 and neuroinflammation. *J Affect Disord* (2024)

48. Harrington D, D'Agostino RBS, Gatsonis C, Hogan JW, Hunter DJ, Normand S-LT, Drazen
JM, Hamel MB. New Guidelines for Statistical Reporting in the Journal. *N Engl J Med* (2019)
381:285–286. doi: 10.1056/NEJMe1906559

49. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res*
Methodol (2005) 8:19–32. doi: 10.1080/1364557032000119616

50. Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement*
Sci (2010) 5:69. doi: 10.1186/1748-5908-5-69

51. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, Moher D, Peters MDJ,
Horsley T, Weeks L, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist
and Explanation. *Ann Intern Med* (2018) 169:467–473. doi: 10.7326/M18-0850

52. Diagnostic and statistical manual of mental disorders: DSM-5™, 5th ed. Arlington, VA, US:
American Psychiatric Publishing, Inc. (2013). xliv, 947 p.

53. Inflammation. *Natl Libr Med* https://www.ncbi.nlm.nih.gov/mesh/68007249 [Accessed January
3, 2025]

54. Bo L, Van Der Miesen AIR, Klomp SE, Williams ZJ, Szatmari P, Lai M-C. The missing
clinical guidance: a scoping review of care for autistic transgender and gender-diverse people.
eClinicalMedicine (2024) 76:102849. doi: 10.1016/j.eclinm.2024.102849

55. Schneider-Thoma J, Chalkou K, Dörries C, Bighelli I, Ceraso A, Huhn M, Siafis S, Davis JM,
Cipriani A, Furukawa TA, et al. Comparative efficacy and tolerability of 32 oral and long-acting
injectable antipsychotics for the maintenance treatment of adults with schizophrenia: a

systematic review and network meta-analysis. *Lancet Lond Engl* (2022) 399:824–836. doi:
10.1016/S0140-6736(21)01997-8

56. Mackay GM, Forrest CM, Christofides J, Bridel MA, Mitchell S, Cowlard R, Stone TW,
Darlington LG. Kynurenine metabolites and inflammation markers in depressed patients treated
with fluoxetine or counselling. *Clin Exp Pharmacol Physiol* (2009) 36:425–435. doi:
10.1111/j.1440-1681.2008.05077.x

57. Raison CL, Rutherford RE, Woolwine BJ, Shuo C, Schettler P, Drake DF, Haroon E, Miller
AH. A randomized controlled trial of the tumor necrosis factor antagonist infliximab for
treatment-resistant depression: the role of baseline inflammatory biomarkers. *JAMA Psychiatry*
(2013) 70:31–41. doi: 10.1001/2013.jamapsychiatry.4

58. Weinberger JF, Raison CL, Rye DB, Montague AR, Woolwine BJ, Felger JC, Haroon E, Miller
AH. Inhibition of tumor necrosis factor improves sleep continuity in patients with treatment
resistant depression and high inflammation. *Brain Behav Immun* (2015) 47:193–200. doi:
10.1016/j.bbi.2014.12.016

59. Xiaoling Z, Yunping H, Yingdong L. Analysis of curative effect of fluoxetine and escitalopram
in the depression treatment based on clinical observation. *Pak J Pharm Sci* (2018) 31:1115–
1118.

60. Chen M-H, Li C-T, Lin W-C, Hong C-J, Tu P-C, Bai Y-M, Cheng C-M, Su T-P. Rapid
inflammation modulation and antidepressant efficacy of a low-dose ketamine infusion in
treatment-resistant depression: A randomized, double-blind control study. *Psychiatry Res*
(2018) 269:207–211. doi: 10.1016/j.psychres.2018.08.078

61. Cao Z-Y, Liu Y-Z, Li J-M, Ruan Y-M, Yan W-J, Zhong S-Y, Zhang T, Liu L-L, Wu R, Wang
B, et al. Glycyrrhizic acid as an adjunctive treatment for depression through anti-inflammation:
A randomized placebo-controlled clinical trial. *J Affect Disord* (2020) 265:247–254. doi:
10.1016/j.jad.2020.01.048

62. Hariri N, Darafshi Ghahroudi S, Jahangiri S, Borumandnia N, Narmaki E, Saidpour A. The
beneficial effects of sumac (Rhus coriaria L.) supplementation along with restricted calorie diet
on anthropometric indices, oxidative stress, and inflammation in overweight or obese women
with depression: A randomized clinical trial. *Phytother Res PTR* (2020) 34:3041–3051. doi:
10.1002/ptr.6737

63. Nettis MA, Lombardo G, Hastings C, Zajkowska Z, Mariani N, Nikkheslat N, Worrell C,
Enache D, McLaughlin A, Kose M, et al. Augmentation therapy with minocycline in treatment-
resistant depression patients with low-grade peripheral inflammation: results from a double-
blind randomised clinical trial. *Neuropsychopharmacol Off Publ Am Coll*
Neuropsychopharmacol (2021) 46:939–948. doi: 10.1038/s41386-020-00948-6

64. Baune BT, Sampson E, Louise J, Hori H, Schubert KO, Clark SR, Mills NT, Fourrier C. No
evidence for clinical efficacy of adjunctive celecoxib with vortioxetine in the treatment of
depression: A 6-week double-blind placebo controlled randomized trial. *Eur*
Neuropsychopharmacol J Eur Coll Neuropsychopharmacol (2021) 53:34–46. doi:
10.1016/j.euroneuro.2021.07.092

65.   Nettis MA, Lombardo G, Hastings C, Zajkowska Z, Mariani N, Nikkheslat N, Sforzini L,
Worrell C, Begum A, Brown M, et al. The interaction between kynurenine pathway, suicidal
ideation and augmentation therapy with minocycline in patients with treatment-resistant
depression. *J Psychopharmacol Oxf Engl* (2023) 37:531–538. doi:
10.1177/02698811231173588

66.   Sampson E, Kavakbasi E, Mills NT, Hori H, Schubert KO, Fourrier C, Baune BT. Emotional
Blunting in Depression in the PREDDICT Clinical Trial: Inflammation-Stratified Augmentation
of Vortioxetine With Celecoxib. *Int J Neuropsychopharmacol* (2024) 27: doi:
10.1093/ijnp/pyad066

67.   Zhang Y, Liu L, Liu Y-Z, Shen X-L, Wu T-Y, Zhang T, Wang W, Wang Y-X, Jiang C-L.
NLRP3 Inflammasome Mediates Chronic Mild Stress-Induced Depression in Mice via
Neuroinflammation. *Int J Neuropsychopharmacol* (2015) 18: doi: 10.1093/ijnp/pyv006

68.   Alcocer-Gómez E, Ulecia-Morón C, Marín-Aguilar F, Rybkina T, Casas-Barquero N, Ruiz-
Cabello J, Ryffel B, Apetoh L, Ghiringhelli F, Bullón P, et al. Stress-Induced Depressive
Behaviors Require a Functional NLRP3 Inflammasome. *Mol Neurobiol* (2016) 53:4874–4882.
doi: 10.1007/s12035-015-9408-7

69.   Belmaker R.H., Agam Galila. Major Depressive Disorder. *N Engl J Med* 358:55–68. doi:
10.1056/NEJMra073096

70.   Remes O, Mendes JF, Templeton P. Biological, Psychological, and Social Determinants of
Depression: A Review of Recent Literature. *Brain Sci* (2021) 11: doi:
10.3390/brainsci11121633

71.   Roy S, Arif Ansari M, Choudhary K, Singh S. NLRP3 inflammasome in depression: A review.
Int Immunopharmacol (2023) 117:109916. doi: https://doi.org/10.1016/j.intimp.2023.109916

72.   Schroder K, Tschopp J. The inflammasomes. *Cell* (2010) 140:821–832. doi:
10.1016/j.cell.2010.01.040

73.   Dinarello CA. Interleukin-1 in the pathogenesis and treatment of inflammatory diseases. *Blood*
(2011) 117:3720–3732. doi: 10.1182/blood-2010-07-273417

74.   Zhang W, Sun M, Fan Y, Wang H, Feng M, Zhou S, Wang R. Machine Learning in
Preoperative Prediction of Postoperative Immediate Remission of Histology-Positive Cushing's
Disease. *Front Endocrinol* (2021) 12:635795. doi: 10.3389/fendo.2021.635795

75.   Zhang W, Wu X, Wang H, Wu R, Deng C, Xu Q, Liu X, Bai X, Yang S, Li X, et al. Federated
Learning for Predicting Postoperative Remission of Patients with Acromegaly: A Multicentered
Study. *World Neurosurg* (2025) 193:1036–1046. doi: 10.1016/j.wneu.2024.10.091

76.   He Z, Zhang R, Diallo G, Huang Z, Glicksberg BS. Editorial: Explainable artificial intelligence
for critical healthcare applications. *Front Artif Intell* (2023) 6: doi: 10.3389/frai.2023.1282800

77.   Aittokallio T, Fang EF. Editorial overview–Artificial intelligence methodologies in structural
biology: Bridging the gap to medical applications. *Curr Opin Struct Biol* (2024) 87:102862. doi:
10.1016/j.sbi.2024.102862

78.   Gao G, Tian D, Li R, Hao X. Application and Development of Artificial Intelligence in Medical
Field. In: Lee G, editor. *2018 5TH INTERNATIONAL SYMPOSIUM ON COMPUTER,*
COMMUNICATION, CONTROL AND AUTOMATION (3CA 2018). Newark: Information
Engineering Research Inst, USA (2018). p. 215–219 doi: 10.5729/3ca.2018.1.215

79.   Shen Y, Jin D, Fu M, Liu S, Xu Z, Cao Q, Wang B, Li G, Chen W, Liu S, et al. Reactive
wetting enabled anchoring of non-wettable iron oxide in liquid metal for miniature soft robot.
Nat Commun (2023) 14:6276. doi: 10.1038/s41467-023-41920-4

80.   Wang WS, Mendonca R, Kording K, Avery M, Johnson MJ. Towards Data-Driven
Autonomous Robot-Assisted Physical Rehabilitation Therapy. *IEEE Int Conf Rehabil Robot*
Proc (2019) 2019:34–39. doi: 10.1109/ICORR.2019.8779555

81.   Zhu J, Wang Z, Zhang Y, Li X, Liu J, Deng K, Lu L, Pan H, Wang R, Yao Y, et al. Ectopic
pituitary adenomas: clinical features, diagnostic challenges and management. *Pituitary* (2020)
23:648–664. doi: 10.1007/s11102-020-01071-x

82.   Li X, Lynch L, Xing H, Wang Z, Zhu J, Deng K, Wang R, Yao Y, Lian X. Cosecreting
TSH/GH pituitary adenomas-an 8-year experience in a single tertiary center. *Pituitary* (2020)
23:573–581. doi: 10.1007/s11102-020-01064-w

83.   Zhu J, Wang Z, Zhang Y, Liu J, Li X, Deng K, Lu L, Yao Y. Suprasellar pituitary adenomas: a
10-year experience in a single tertiary medical center and a literature review. *Pituitary* (2020)
23:367–380. doi: 10.1007/s11102-020-01043-1

84.   Li X, Deng K, Zhang Y, Feng M, Xing B, Lian W, Yao Y. Pediatric pituitary neuroendocrine
tumors-a 13-year experience in a tertiary center. *Front Oncol* (2023) 13:1270958. doi:
10.3389/fonc.2023.1270958

1042

1043

1044

1045