Development and Validation of a Nutrient Profiling Model for Shopping Baskets: The Grocery Basket Score (GBS) Methodology

Paul Windisch[1*], Sandro Marcon[1], Javier Orts[1], David Faeh[1,2,3], Richard Hurrell[1,4], André Naef[1]

[1] dacadoo AG, Zurich, Switzerland

[2] Epidemiology, Biostatistics and Prevention Institute, Division of Chronic Disease Epidemiology, University of Zurich, Zurich, Switzerland

[3] Health Department, Bern University of Applied Sciences, Bern, Switzerland

[4] Professor emeritus, Institute of Food, Nutrition and Health, ETH Zurich, Switzerland

* Correspondence:

Paul Windisch, MD

dacadoo AG

Othmarstrasse 8

8008 Zurich

Tel: +41 44 251 23 23

ORCID-ID: 0000-0003-1040-4888

paul.windisch@dacadoo.com

Disclosures:

P.W., S.M., J.O., D.F., R.H., and A.N. and. report employment at or consulting for dacadoo

AG, a technology company that has business activity with retailers, including grocery stores.

dacadoo intends to file 'GBS' as a trademark and commercialize GBS^{TM} software and/or

consulting services. The other authors declare no additional conflict of interest.

Author contributions: Conceptualization, P.W., A.N.; methodology, P.W, S.M., A.N.; formal

analysis, P.W., S.M., J.O., A.N.; data curation, S.M., J.O.; writing—original draft preparation,

P.W.; writing—review and editing, S.M., J.O., D.F., R.H., A.N.; supervision, A.N.; project

administration, P.W., A.N.;

Data availability:

This paper contains information from the Open Food Facts (https://world.openfoodfacts.org/)

and OFF-GBS (https://github.com/windisch-paul/OFF-GBS) databases, which are made

available under the Open Database License (ODbL,

https://opendatacommons.org/licenses/odbl/1-0/).

Ethical approval:

Not applicable (publicly available, anonymized data)

Funding:

No external funding was received for this project.

Word count (excluding abstract, figures, tables, references): 4488

48 Abstract

Background: Rating a person's diet as a whole, as opposed to rating individual food items,

could help better inform consumers about the health value of their diet. Our goal was to

develop an automated health rating of grocery shopping baskets, based on the nutrient

composition of all the food items contained within the basket, and without knowing how

many people consume the food items over what period of time. It is envisaged that the

rating, or score, can be deployed by retailers on top of existing loyalty or digital shopping

basket programs.

Methods: We developed a novel model for calculating Grocery Basket Scores (GBS) that

uses nutrient energy densities rather than absolute quantities. It was based on self-reported

daily food intake from the National Health and Nutrition Examination Survey (NHANES) as

well as mortality follow-up data. We conducted a validation against the Alternate Healthy

Eating Index (AHEI) as well as the Nutri-Score.

Results: The nutrient energy density (GBS) model penalized consuming calories from sugar

and saturated fats, high salt intake, as well as consuming calories from beverages.

Furthermore, a penalty was applied to foods that have a low ratio of protein, vitamin C, or

iron relative to the total number of calories. Fiber consumption was rewarded. The model

showed a high degree of correlation with the AHEI (absolute Pearson correlation coefficient:

0.62 for the AHEI without the protective effect of moderate alcohol consumption and 0.60

with it) and the Nutri-Score (absolute Pearson correlation coefficient: 0.60).

Conclusions: The proposed nutrient energy density model is aligned with the

recommendations formulated in nutritional guidelines, as indicated by a high degree of

correlation with the AHEI and Nutri-Score. Long term use of a GBS from a single or multiple

grocery stores could help consumers adhere to dietary guidelines.

Introduction

Although the impact of diet on non-communicable diseases and mortality has been investigated in many studies (1), what constitutes an optimal diet remains subject to discussion. As a consequence, nutrition profiling and rating systems have been developed to inform consumers and nudge them towards healthier shopping choices (2).

One example is the Nutri-Score, which has been endorsed by several countries and is found on a variety of products (3). Profiling systems such as the Nutri-Score focus on rating individual foods within a given group, which allows for printing scores on the products and enables the consumer to make direct comparisons between different items from the same category (4). While this is useful, focusing on individual foods also comes with tradeoffs. For example, one could compose an entire shopping basket consisting exclusively of items that received an A grading by the Nutri-Score due to their low-calorie density and high fiber content while containing little protein, even though the Nutri-Score rewards protein. As protein deficiency has now become an issue in the elderly in industrialized nations (5) it illustrates that assessing nutrition as a whole is more complex than summing up or even averaging individual food ratings. This is also relevant to consumers, as they must decide what share of poorly rated products in their basket is still acceptable.

Nutritional guidelines such as the Alternate Healthy Eating Index (AHEI) tend to provide recommendations in quantities, such as servings, per day (6,7). The AHEI was designed to predict chronic disease risk and has also been validated for predicting mortality (8). While this approach can help consumers understand how they should compose their diet, it can still be challenging to translate this information into actionable shopping decisions as a consumer must still assess how many servings are contained in a single product and keep track of the number of products they need for a given timeframe.

Our goal was to develop a nutrient energy density model that avoids serving sizes and can

be used directly in supermarkets to translate recommendations from nutritional guidelines

into actionable decisions and, at the same time, will complement product-level rating

systems such as Nutri-Score. To this end, we set out to develop a model to rate shopping

baskets for their health properties, assuming that these baskets account for a substantial

amount of a consumer's total nutrition. Although a single basket of food products may have

little relevance, long term purchases from one retailer, or from a group of retailers who have

accepted to share information, would be meaningful.

Additionally, we did not want to develop a model that required a manual ingredient

annotation but instead wanted a model that could be automatically deployed by retailers on

top of existing loyalty or digital shopping basket program. Finally, we sought a data-based

model that avoided strong reliance on opinionated expert judgment and based our nutrient

energy density model on the link between nutrition and eventual mortality. We conducted a

validation by comparing the outputs of the final GBS model with the AHEI and the Nutri-

Score to ensure seamless integration into existing guidance.

The hypothesis was that while there would be differences with the Nutri-Score due to the

factors mentioned above, GBS and Nutri-Score would be in agreement for the majority of

items. For the AHEI, we hypothesized a better agreement with GBS than with Nutri-Score,

due to the alignment of rating the consumer's food purchases as a whole (i.e. dietary

patterns) as opposed to rating individual items.

Methods

Data

The data used in this study is a subset of the continuous National Health and Nutrition Examination Survey (NHANES) (9) from 2001 to 2018, which has been linked with the latest mortality data for the participants from 2019 (10). The dietary data thus does not constitute data derived from retail purchases, but is self-reported complete daily food intake data from each participant. As part of the NHANES nutritional assessment, participants of the respective cohort completed a 24-hour dietary recall interview in person with a trained dietary interviewer and participated in a second phone-based interview 3 to 10 days later. The food intake obtained from these recalls was then linked through codes to the Food and Nutrient Database for Dietary Studies (FNDDS) (11),and was used to calculate the respective daily nutrient intakes for each participant in each recall.

Individuals younger than 17, older than 90, those with missing nutritional information, and those who reported no calorie intake, were excluded from the analysis. This resulted in a dataset containing the daily nutrient intake of 35'462 individuals. Data from 25% of the individuals was randomly assigned to the test set, with data from the remaining individuals being used as the training set to model eventual mortality from dietary signals.

Considerations for Potential Dietary Signals

We generated two types of dietary signals, namely "nutrient energy density signals" and "low nutrient-energy-ratio signals". As one cannot know how long a shopping basket will last in a household, or how many people consume the food items in the basket, we calculated dietary

signals based on energy density, that is energy or weight from a given nutrient relative to the

total basket energy.

Nutrient energy density signals that were evaluated over the course of the project were

calories from carbohydrates divided by total calories, calories from sugar divided by total

calories, calories from proteins divided by total calories, calories from fat divided by total

calories, calories from saturated fats divided by total calories, calories from unsaturated fats

divided by total calories, calories from alcohol divided by total calories, amount of fiber

divided by total calories, amount of sodium divided by total calories, and calories from

beverages divided by the weight of the beverages.

A further set of signals were calculated for proteins and certain vitamins and minerals as

indicators of low dietary intake. These "low nutrient-energy-ratio signals" indicate that a

person is consuming a low amount of a given nutrient relative to their total calorie

consumption. In order to calculate the nutrient-energy-ratio signal, the daily recommended

intake of the nutrient was divided by the daily recommended calories, resulting in a

recommended density of the nutrient per calorie. The low nutrient-energy-ratio signal was

included in the model if the nutrient per calorie density was below the recommended density.

The signal becomes stronger the lower the density gets. The nutrients which were evaluated

as low nutrient-energy-ratio signals and the respective recommendations that were used for

the calculation are provided in Supplementary Table 1. Nutrient-energy-ratio signals were

generated for vitamin C, folate, potassium, calcium protein, and iron.

All dietary signals were then evaluated as potential inputs to Cox proportional hazard models

for predicting mortality. As a requirement for all dietary signals, there should be a plausible

biological mechanism linking each signal to mortality. Such a link should be supported by a

consistent correlation of the respective signal with mortality in NHANES (9).

Another consideration was that the signals should be computable for any retailer without

additional manual labeling of foods being required. Therefore, we focused on nutrients that

are either part of the labeling requirements in many countries or that could be reliably

imputed, e.g., from the name of the food and its categories. Lastly, we wanted the model to

be based on different, heterogeneous signals to reduce the likelihood of single items

achieving very high scores by performing well across all signals, as a shopping basket

consisting only of one item is unlikely to be conducive to healthy nutrition (low dietary

diversity). In other words, we sought a mix of signals that provides a reasonable fingerprint

of grocery shopping behavior.

182   Modeling

A schema of the modeling process is presented in Figure 1. For the modeling and validation,

we considered the items from the daily dietary recall as a person's shopping basket. Due to

high degrees of correlation between different potential signals, different combinations of

signals were evaluated while omitting others. The goal was to minimize redundancy, identify

combinations of signals that fulfill all requirements mentioned above, avoid multicollinearity

as a source of model misspecification, and ultimately to use the signals as inputs to a Cox

proportional hazard model. Avoiding multicollinearity was particularly important as many

potential signals were highly correlated (e.g. calories from sugar divided by total calories,

calories from added sugar divided by total calories, and calories from carbohydrates divided

by total calories).

If highly correlated signals are used as independent variables in a regression, the model

cannot reliably separate them, resulting in wild coefficient swings with small changes in the

data. This increases the variance of the estimates and leads to wider confidence intervals.

Moreover, as multicollinear variables tend to "cancel out", they can lead to coefficients that

are not bio-plausible when viewed in isolation, thus making the interpretation of the model

difficult, as expected relationships between dietary patterns and mortality risk disappear.

In addition, we evaluated a joint model that used the same signals for both foods and

beverages as well as separate models with different signals for foods and beverages

respectively.

Standardization of the signals was performed for interpretability reasons, by subtracting the

mean and dividing by the standard deviation, such that the resulting model coefficients are

on the same scale. Signals were also clipped to the limits of 1.5 times the interquartile range

(IQR) rule for outlier detection (Tukey's fences).

Since the model's goal was to predict a time-to-event endpoint and the NHANES data is

censored, Cox proportional hazard models were used to estimate the effect of different

combinations of the dietary signals (defined above) on mortality. The models were adjusted

for age, sex, body-mass index (BMI) and the poverty-income ratio (i.e. household income

divided by the poverty threshold) by including them as model coefficients and setting them to

zero at inference time. The Cox model which included a comprehensive set of dietary

signals and showed good performance at predicting mortality, while including a

comprehensive set of signals, was chosen for further validation. This nutrient energy density

model was then used as the basis for the Grocery Basket Score (GBS).

The output that is obtained by multiplying the regression coefficient with the strength of the

respective dietary signal is the logarithm of the hazard ratio. Afterwards, the hazards were

transformed in order to achieve a score resembling a normal distribution ranging from zero

(worst) to one thousand (best) which is how the score was supposed to be presented to

users.

To this end, the obtained values were clipped beyond -0.8 and 1.5, normalized between 0

and 1, and subtracted from 1. Values were then raised to a power of 1.5. and multiplied by

1'000.The separate levels of adjustment for the core and beverages model are available in

Supplementary Tables 2 and 3, respectively. Unless specified otherwise, data analysis was

performed using Python (v. 3.8.10) as well as the numpy (v. 1.23.5), pandas (v. 1.3.5) and

lifelines (v. 0.27.7) packages.

Validation

For the validation against the AHEI, we computed the AHEI scores for 4'382 subjects with

dietary recalls from NHANES 2017-18, computed the GBS using the same recalls and

calculated the correlation coefficient between the GBS and AHEI, both with and without the

protective effect of moderate alcohol consumption. The AHEI was computed in R (v. 4.3.2)

using the dietary index package (v. 1.0.3). In addition, we also computed the Nutri-Score

2017 using the pyNutriScore package (v.0.0.2) and calculated a total Nutri-Score for each

participant's nutritional information as a calorie-weighted average of the Nutri-Scores of the

individual food items they consumed. We then calculated the correlation between the

calorie-weighted Nutri-Scores and the AHEI.

For the validation against the Nutri-Score, we retrieved food items from the Open Food Facts

database (12). In the preprocessing, we ignored items with a missing product code or name.

Furthermore, we filtered items with missing energy information, missing Nutri-Score 2023, or

where the macronutrients summed to more than 101 grams per 100 grams of food, items

with saturated fat contents larger than total fat, items with more sugar than carbohydrates,

items where one of the nutrients is larger than the 99[th] percentile and items missing

information regarding any of protein, fat, carbohydrate, or salt content. The remaining

913'298 items were scored using the GBS model and compared to the provided Nutri-

Scores by computing the Pearson correlation coefficient.

Due to occasionally missing labeling information in the Open Food Facts database, we also

developed an imputation model to complete the required label information. Specifically, our

model is a deep neural network that implements a sequence model operating on the product

categories and name using custom embeddings to impute fiber, iron, and vitamin C. A

database with the imputed label information is available at https://github.com/windisch-

paul/OFF-GBS.

In addition to computing the correlation coefficient for the whole dataset, clustering was

performed to allow for a cluster-by-cluster correlation and comparison. Items were clustered

into different food clusters based on their assigned tags in the Open Food Facts database.

Due to the high number of tags in the database and each item potentially being assigned

multiple tags, it was deemed infeasible to manually define disjointed food clusters. A

clustering approach based on context frequency of tags was used instead.

As a first step, we filtered out tags that appeared fewer than 10'000 times in the whole

corpus. This left 78 tags. As a second step, the term frequency-inverse document frequency

matrix (tf-idf) was calculated. The idea was to vectorize tags based on their frequency and

relevance. The tf-idf matrix was then reduced using singular-value decomposition (SVD) by

keeping the 19 most relevant components. Finally, the K-Means clustering algorithm was

applied to find 14 food clusters, with the 15th cluster manually assigned to beverages. The

rationale of performing SVD is that K-Means would perform better on the reduced

eigenvector space rather than on the original tf-idf matrix. The number of singular values and

clusters was selected empirically using the "elbow-rule".

In addition to computing correlation coefficients, we calculated the absolute number and the

percentage of items in a food cluster for which the difference between the z-scores of the

GBS and the Nutri-Score was greater than 1, as well as the number and percentages of

items with a difference greater than 2. This was done to highlight the items that caused the

biggest disagreement.

Data Source
NHANES 2001-2018 with mortality data through 2019
n = 35,462 (75% training, 25% testing)

Signal Engineering
Considerations for signals: Biological link to mortality, availability
of nutrient information, reduced risk of multicollinearity

Nutrient Energy Density Signals
e.g., calories from sugar / total calories

Low Nutrient-Energy-Ratio Signals
e.g., protein intake below recommended density

Signal Preprocessing
Standardization (z-score) and outlier clipping (Tukey's fences, 1.5*IQR)

Model Training
Multiple Cox proportional hazard models with different signal combinations
Time-to-event analysis with mortality as endpoint
Adjusted for age, sex, BMI, and poverty-income ratio

Final Model Selection
Based on biologically plausible coefficients and
signals covering diverse aspects of nutrition

Figure 1: Schema of the Grocery Basket Score (GBS) Modeling Process. Conceptual framework

for the development of the Grocery Basket Score using NHANES 2001-2018 data linked with mortality

data (n=35,462). Following data preprocessing and standardization, two categories of dietary signals

were generated: nutrient energy density signals and low nutrient-energy-ratio signals. Signal

construction was informed by established biological mechanisms, nutrient information availability, and

multicollinearity considerations. Multiple Cox proportional hazard models, adjusted for demographic

and anthropometric covariates, were evaluated. The final GBS model was selected based on

coefficient plausibility and comprehensive representation of diverse nutritional aspects.

Results

Modeling

During follow-up, 14.6% (n = 5'167) of participants died (training set: 3'839 deaths out of 26'598 participants = 14.4%; test set: 1'328 deaths out of 8'874 participants = 15.0%). After several iterations, an architecture obtained by scoring the combination of foods and beverages using a core model, and only the beverages using a dedicated beverages model then, combining the results as a calorie-weighted average of the respective scores, demonstrated areas under the receiver operating characteristic curve (ROC-AUCs) ranging from 0.854 for estimating mortality over a 1-year, and 0.915 over a 15-year time frame (Figure 2).

The dietary signals and respective regression coefficients of the model are presented in Table 1. The regression coefficients indicate that the model penalizes consuming calories from sugar, saturated fats, and beverages as well as high salt intake. Fiber consumption is rewarded. Furthermore, a penalty is applied to diets with a low ratio of protein, vitamin C, or iron consumption relative to the total calories.

As the direction of these coefficients was deemed to be in line with current nutritional research and the model offered the best compromise between predictive performance, comprehensiveness of the signals, and signal plausibility, it was selected as the first version of the Grocery Basket Score (GBS) model. Note that this selection represents a judgment call as selecting whichever combination of signals achieved the best performance could have yielded implausible coefficients due to multicollinearity.

A schema of performing inference with the GBS model is provided in Figure 3. In short, inference is performed by calculating the dietary signals, multiplying them with the respective regression coefficients, weighting core and beverage model by calorie contribution, and then performing the power shift as outlined in the methodology section.

In addition, we have calculated the GBS for ten random NHANES participants using their

dietary recalls with scores ranging from 262 to 709 and present this information in Table 2.

The respective food items that the people consumed are provided in Supplementary Tables

4a and 4b.

To derive additional insights into possible correlations between signals, we also created a

correlation matrix between different parameters based on items in the Open Food Facts

database, which is presented in Figure 4.

Validation

The correlation coefficients between the GBS and the AHEI for 4'382 participants in

NHANES 2017-2018 were 0.62 for the AHEI without the protective effect of moderate

alcohol consumption and 0.60 with it. The correlation coefficients between the calorie-

weighted Nutri-Scores and the AHEI were -0.48 and -0.47, respectively. The corresponding

scatterplots are presented in Figure 5.

The overall correlation coefficient between the GBS and the Nutri-Score on single food items

in the whole Open Food Facts database was -0.60. This inverse correlation is due to healthy

items receiving low numeric values by the Nutri-Score but high numeric values by the GBS.

The clustering of items from the Open Food Facts database resulted in 15 clusters, and the

correlation coefficients per cluster ranged from -0.31 to -0.74. The clusters and respective

correlation coefficients are provided in Table 3. The weakest correlation coefficient was

found in cluster 7 with a correlation coefficient of -0.31, which was due to items being scored

more favorably by the Nutri-Score than by the GBS. In all other clusters, correlation

coefficients were stronger than -0.46.

Figure 6 shows the percentage of items per food cluster where the z-scores between the

GBS and the Nutri-Score differ by more than 1 and 2 as well as the direction of the

disagreement (i.e., which model rated more favorably). Samples of items scored differently

by the GBS compared to the Nutri-Score are provided in Supplementary Tables 5 - 8.

351 **Figure 2. Performance of the Grocery Basket Score (GBS) Model**. Areas under the receiver

352 operating characteristic curve (ROC-AUCs) demonstrating the performance of the GBS Model when

353 predicting mortality for NHANES participants on the test set over 1- to 15-year timeframes.

Figure 3. Schema of the Grocery Basket Score (GBS) model at inference. The calories, nutrients, and weights in the basket are summarized for foods and beverages respectively. The data for beverages are passed to the beverages model which only computes one dietary signal (calories from beverages / weight of beverages) and multiplies it with the respective regression coefficient. The data for both foods and beverages are passed to the core model, which computes seven dietary signals (calories from saturated fats / total calories, low protein-energy-ratio, calories from sugar / total calories, amount of fiber / total calories, amount of sodium / total calories, low vitamin-C-energy-ratio, low iron-energy-ratio) and multiplies them with the respective regression coefficients. The regression coefficients are presented in Table 1.

The outputs of the core and beverage models are weighted by the ratio of the total calories handled

369   by the respective models. If a basket's calories come almost exclusively from food, the output of the

370   core model will dictate the total score. If a basket's calories come almost exclusively from beverages,

371   the output of the beverage model will dictate the total score. The power shift from the resulting

372   logarithm of the hazard ratio to the 0 to 1'000 scale is performed according to the description in the

373   methods section.

Core model			
Dietary signal	**Regression coefficient**	**Mean prior to standardization**	**Standard deviation prior to standardization**
Calories from saturated fats [kcal] / total calories [kcal]	0.059	0.1208	0.0357
Low protein-energy-ratio	0.020	0.0001	0.0009
Calories from sugar [kcal] / total calories [kcal]	0.056	0.1662	0.0772
Amount of fiber [g] / total calories [kcal]	- 0.081	0.090	0.0039
Amount of sodium [mg] / total calories [kcal]	0.064	1.8112	0.4730
Low vitamin-C-energy-ratio	0.153	0.0171	0.0147
Low iron-energy-ratio	0.084	0.0005	0.0011
Beverage model			
Dietary Signal	**Regression coefficient**	**Mean prior to standardization**	**Standard deviation prior to standardization**
Calories from beverages [kcal] / weight of beverages [g]	0.071	0.1726	0.1451

Table 1. Dietary signals and regression coefficients of the core and beverage models of the

Grocery Basket Score (GBS). Positive regression coefficients indicate increased mortality risk, while

negative regression coefficients indicate decreased mortality risk. The low nutrient-energy-ratio

signals were obtained by reviewing nutrition guidelines and dividing the daily recommended intake of

the nutrient by the daily recommended calories from the same source, resulting in a recommended

density of the nutrient per calorie. The low nutrient-energy-ratio signal is considered if the nutrient per

calorie density in the basket is below the recommended density. The signal becomes stronger as the

ratio becomes lower. To calculate the GBS for a person's basket, each dietary signal is multiplied with

the respective regression coefficient. The resulting output is the logarithm of the hazard ratio and

387 undergoes the power shift described in the methodology section to create the final GBS on a scale of

388 0 to 1'000.

	54539	24231	47699	79039	99176	26285	85578	56160	61920	60290
Age range	56-60	21-25	71-75	30-35	61-65	36-40	61-65	76-80	16-20	41-45
Sex	F	M	F	F	F	M	F	F	M	F
Calories from saturated fats [kcal] / total calories [kcal]	0.162	0.104	0.0855	0.081	0.0722	0.1273	0.0787	0.0992	0.1539	0.0807
Low protein-energy-ratio	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Calories from sugar [kcal] / total calories [kcal]	0.1091	0.1146	0.1685	0.1025	0.0881	0.1603	0.1288	0.1768	0.2163	0.3104
Amount of fiber [g] / total calories [kcal]	0.0061	0.0058	0.0061	0.0071	0.0149	0.0134	0.0106	0.0107	0.0094	0.0182
Amount of sodium [mg] / total calories [kcal]	2.3562	1.8412	2.3944	3.2855	2.1745	1.875	2.0381	1.3206	1.6508	1.5614
Low vitamin-C-energy-ratio	0.012	0.036	0.04	0.051	0.0	0.016	0.0	0.02	0.003	0.0
Low iron-energy-ratio	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.002
Calories from beverages [kcal] / weight of beverages [g]	0.1073	0.3293	0.3046	0.1834	0.0015	0.2899	0.0039	0.0194	0.4108	0.1845
GBS	373.0	324.0	287.0	262.0	709.0	461.0	626.0	504.0	480.0	554.0

390 **Table 2. Grocery Basket Score (GBS) calculation for 10 random participants from NHANES**

391 **using their answers to dietary recalls.** The table presents the strength of the respective signals and

392 the resulting scores. The consumed foods including the respective quantities are presented in

393 Supplementary Table 4a (first five patients) and 4b (remaining patients). *F = female, M = male.*

Cluster #	Open Food Facts Tags	Pearson correlation coefficient
0	seafood; fishes-and-their-products; fishes; fatty-fishes; canned-foods	-0.60
1	fruits-and-vegetables-based-foods; plant-based-foods-and-beverages; plant-based-foods; vegetables-based-foods; canned-foods	-0.65
2	biscuits-and-cakes; sweet-snacks; biscuits; snacks; cakes	-0.74
3	meats-and-their-products; meats; prepared-meats; sausages; hams	-0.73
4	dairies; fermented-foods; fermented-milk-products; cheeses; desserts	-0.46
5	cereals-and-potatoes; plant-based-foods-and-beverages; plant-based-foods; cereals-and-their-products; breads	-0.68
6	condiments; sauces; groceries; plant-based-foods-and-beverages	-0.59
7	salty-snacks; appetizers; milks; chips-and-fries; crisps	-0.31
8	meals; pasta-dishes; pizzas-pies-and-quiches; soups; meals-with-meat	-0.53
9	frozen-foods; desserts; frozen-desserts; ice-creams-and-sorbets; dairy-desserts	-0.58
10	confectioneries; sweet-snacks; snacks; cocoa-and-its-products; chocolate-candies	-0.53
11	snacks; sweet-snacks; confectioneries; biscuits-and-cakes; biscuits	-0.74
12	spreads; breakfasts; sweet-spreads; plant-based-spreads; fruit-and-vegetable-preserves	-0.66
13	fats; vegetable-fats; vegetable-oils; plant-based-foods-and-beverages; plant-based-foods	-0.72
14	beverages	-0.60
Overall		**-0.60**

398 **Table 3. Pearson correlation coefficients by food clusters between the Grocery Basket Score**

399 **(GBS) and the Nutri-Score.** Greater absolute numbers indicate stronger correlations.

Figure 4. Pearson correlation coefficients between different parameters based on items in the Open Food Facts database. The NOVA Score tries to approximate the degree of processing a food item has undergone with higher scores indicating a higher degree.

Figure 5. Scatterplot of the Grocery Basket Score (GBS - top) and calorie-weighted Nutri-Score

(bottom) with Alternate Healthy Eating Index (AHEI) ratings. Each dot represents one of 4'382

NHANES 2017 - 2018 participants. Results of using the AHEI without the protective effect of

moderate alcohol consumption are presented on the left and results with the protective effect of

moderate alcohol consumption are displayed on the right. A steeper slope indicates a stronger

correlation. *PCC = Pearson correlation coefficient.*

Figure 6. Discrepancies in Grocery Basket Score (GBS) and Nutri-Score ratings by food clusters. Percentage of items per food cluster where the z-scores between the GBS and the Nutri-Score differ by more than 1 (top) and 2 (bottom). Items that were rated more favorably by the GBS than the Nutri-Score are presented on the left, while items rated more favorably by Nutri-Score can be found on the right. The contents of the clusters can be found in Table 3.

Discussion

The main findings of this study are that the novel nutrient energy density model tested,

predicted healthy diet scores that, like Nutri-score and AHEI, are in alignment with the

nutrition guidelines issued by the U.S. Departments of Agriculture and of Health and Human

Services or the World Health Organization (13,14). This new nutrient energy density model

therefore can now be used as a Grocery Basket Score (GBS) to automatically score foods

and beverages in grocery shopping baskets according to their health properties based on

their link to mortality. Its final score is computed as a calorie-weighted average of two

separate models, one scoring a combination of foods and beverages and the other once

scoring only beverages.

The core model's characteristics can be derived from the regression coefficients assigned to

its nutrient energy density signals: It penalizes consuming calories from sugar and saturated

fats, as well as high salt intake. Furthermore, a penalty is applied to baskets with a low ratio

of protein, vitamin C, or iron, relative to their total calories. Fiber consumption is rewarded.

For the beverages model, calorie density is penalized.

Even though the GBS was designed to be deployed on whole shopping baskets and not for

assessing individual products, the high degree of correlation with the Nutri-Score, indicated

by a correlation coefficient of -0.60 on items from the Open Food Facts database,

demonstrates the considerable overlap in principles such as rewarding the presence of fiber

and penalizing the presence of saturated fats, sugar, and salt. This should reduce the risk of

confusing consumers should the GBS be deployed in regions where the Nutri-Score is also

used for individual products.

The design differences to the Nutri-Score become evident when looking at the food clusters

with lower correlation coefficients and the items for which the scores disagreed the most.

The differences in the scoring of items in the cluster with the lowest absolute correlation

coefficient (i.e., cluster 7) appear to be mainly driven by (low-fat) milk, which is rated more

negatively by the GBS than by the Nutri-Score. The same applies to cluster 4 where the

differences appear to be driven by dairy products such as cottage cheese or yoghurt. This is

due to the lack of vitamin C, iron, and fiber, which is penalized by our model as a balanced

shopping basket should contain these nutrients. However, milk and egg whites are obviously

not supposed to be sources of fiber. When part of a more comprehensive shopping basket,

they would have a more positive effect due to improving a low protein-energy-ratio and

increasing the denominator (i.e., the total calories), thereby improving the sugar and

saturated fat divided by total calorie signals.

Cluster 10 contains sweet snacks, such as chocolate, which tend to be scored more

positively by the GBS than by the Nutri-Score. This seems to be mainly because the Nutri-

Score considers calorie density as a standalone signal and punishes items for having high

calorie densities. The GBS takes a different route as it has been adjusted for BMI and does

not consider calorie density as its own signal, effectively asking the question "If a consumer

is able to maintain their weight, is it ok for them to consume this shopping basket?". While

weight management is certainly an important goal when trying to compose a shopping

basket, it is also relatively easy to track for the consumer when stepping on a scale. The

GBS thus focuses on issues that are more difficult to keep track of, such as the nutrient

composition, which indirectly relates to weight management as fiber and protein tend to

provide more satiety per calorie than sugars.

The high degree of correlation with the AHEI is promising and not surprising considering that

the AHEI advises people to frequently eat fiber- and protein-rich foods while lowering sugar

and salt consumption. It is also unsurprising that the correlation is even better when using

the AHEI without assuming a protective effect of moderate alcohol consumption as the

beverage model penalizes drinks with higher calorie densities.

Our study has several limitations. Firstly, our grocery basket was based on all foods

consumed over 24h, as recalled by participants in the NHANES study. It represented one

person's daily diet. Although isolated grocery baskets in real life are unlikely to contain all of

one person's daily diet, over a period of time, combined grocery baskets can be

representative of a single person's or whole family's diet. Then, the greater number of GB

scores from a single or multiple supermarkets will improve the overall evaluation of the

health value of a family diet and the consumer can also be advised on how to improve the

diet.

A general limitation of working with the NHANES dataset is its cross-sectional nature. That

is, a person's nutrition might have changed significantly from the time of their dietary recalls

to their death or time of censoring. Furthermore, the fact that some signals are included in

the final model while others are not, should not be interpreted as a claim that the included

signals are somehow more predictive of mortality, let alone that there is more evidence for a

causal relation, i.e., that supplementing the respective nutrients will improve mortality. It

seems plausible that some of the signals used in the GBS function largely as proxies, such

as the fiber signal and the vitamin-C-energy-ratio signal being proxies for fruit and vegetable

or whole grains consumption which need to be considered when using the GBS to propose a

healthy diet.

Furthermore, it would have been desirable to have even more signals to get a more

comprehensive picture of someone's nutrition. However, due to the high degree of

correlation between many signals, including more signals would not only provide, at best,

minimal lift but also make the modeling more unstable due to variables stealing predictive

power from each other (15). While there are certainly other ways to do the modeling, and

judgment calls of some degree were necessary to make the signal selection, we believe that

the current model is capable of giving the consumer a useful evaluation of the quality of their

diet, especially if grocery basket scores from a single retailer, or groups of retailers, are

combined over a period of time.

The GBS is also relatively resistant to current attempts at gaming the system, e.g., by adding

soluble fiber or proteins to beverages as this would not improve the beverage model that

only focuses on calorie density. While having the degree of processing as a signal would

have been interesting, this is harder to do without requiring manual labeling. In addition,

ultra-processed foods are frequently high in salt, sugar, and saturated fats but relatively low

in protein and fiber, as illustrated by the correlation matrix in Figure 4, so the degree of

processing is at least indirectly represented in the GBS through surrogate signals (16). This

is supported by a recent study by Fang and colleagues, who found that adjusting for the

AHEI attenuated a previously observed association of ultra-processed food intake with

mortality (17). The annotation problem and the representation through surrogates also apply

to other potential signals like including whole grains, which have been suggested previously

(18).

Other limitations include the fact that this model was developed on a U.S. instead of a global

population which might limit its ability to generalize to other parts of the world. In addition,

the dietary information was self-reported. Furthermore, food items with missing nutritional

information were not considered during the validation, which might have introduced bias, as

well as the fact that for technical reasons Nutri-Score 2017 was used for the AHEI validation

instead of the more recent version 2023.

Lastly, another limitation of the GBS is that it does not model the negative health effects of

alcoholic beverages that go beyond their calorie density (19). The fact that adding the self-

reported alcohol consumption as a signal did not increase the performance of the model

when predicting mortality in NHANES might be due to inconsistencies in self-reported and

actual alcohol consumption that have previously been described (20).

538 A strength of the GBS is that it does not require manual annotation of food items prior to

539 deployment, as it leverages the Open Food Facts database and an imputation engine. Since

540 many retailers already have loyalty or digital shopping basket programs in place, this allows

541 for automated real-world implementation. The use of high-level signals yielding a fingerprint,

542 as opposed to very granular lists of food items people should consume, is aligned with the

543 quality of evidence in the nutritional epidemiology space and allows for flexibility when

544 composing a basket.

546 Most importantly, the GBS gives users an automated assessment of their entire shopping

547 basket without the need for manual tracking. Moreover, owing to its energy density-based

548 nature, GBS does not require absolute quantities or knowledge of the household size. Even

549 with shopping at multiple grocers and eating out, GBS can provide a reasonable score as

550 long as a representative subset of food intake is scored. As an outlook, the model could also

551 be used to derive personal shopping recommendations based on the strengths of the signals

552 for a given basket.

554 In conclusion, the Grocery Basket Score (GBS) based on a nutrient energy density model is

555 a methodology for scoring shopping baskets that goes beyond averaging the scores of

556 individual products. The GBS scoring methodology is aligned with the recommendations

557 formulated in nutritional guidelines, which is indicated by a high degree of correlation with

558 the AHEI. Product-level profiling systems such as the Nutri-Score also exhibit a correlation

559 with the GBS and can be used in combination.

References

References

1. Afshin A, Sur PJ, Fay KA, Cornaby L, Ferrara G, Salama JS, Mullany EC, Abate KH,

Abbafati C, Abebe Z, et al. Health effects of dietary risks in 195 countries, 1990–2017: a

systematic analysis for the Global Burden of Disease Study 2017. Lancet Elsevier;

2019;393:1958–72.

2. An R, Shi Y, Shen J, Bullard T, Liu G, Yang Q, Chen N, Cao L. Effect of front-of-

package nutrition labeling on food purchases: a systematic review. Public Health

2021;191:59–67.

3. Julia C, Etilé F, Hercberg S. Front-of-pack Nutri-Score labelling in France: an evidence-

based policy. Lancet Public Health 2018;3:e164.

4. van den Akker K, Bartelet D, Brouwer L, Luijpers S, Nap T, Havermans R. The impact of

the nutri-score on food choice: A choice experiment in a Dutch supermarket. Appetite

2022;168:105664.

5. Ten Haaf DSM, De Regt MF, Visser M, Witteman BJM, de Vries JHM, Eijsvogels TMH,

Hopman MTE. Insufficient protein intakes is highly prevalent among physically active

elderly. J Nutr Health Aging Springer Science and Business Media LLC; 2018;22:1112–

4.

6. McCullough ML, Willett WC. Evaluating adherence to recommended diets in adults: the

Alternate Healthy Eating Index. Public Health Nutr 2006;9:152–7.

7. Chiuve SE, Fung TT, Rimm EB, Hu FB, McCullough ML, Wang M, Stampfer MJ, Willett

WC. Alternative dietary indices both strongly predict risk of chronic disease. J Nutr

2012;142:1009–18.

8. Pestoni G, Karavasiloglou N, Braun J, Krieger J-P, Sych JM, Bopp M, Faeh D,

Gruebner O, Rohrmann S. Does diet map with mortality? Ecological association of

dietary patterns with chronic disease mortality and its spatial dependence in

Switzerland. Br J Nutr 2022;127:1037–49.

9.   Paulose-Ram R, Graber JE, Woodwell D, Ahluwalia N. The National Health and

Nutrition Examination Survey (NHANES), 2021-2022: Adapting Data Collection in a

COVID-19 Environment. Am J Public Health 2021;111:2149–56.

10. NCHS Data Linkage - Mortality Data - Public-Use Files [Internet]. 2022 [cited 2024 Jul

22]. Available from: https://www.cdc.gov/nchs/data-linkage/mortality-public.htm

11. FNDDS : USDA ARS [Internet]. [cited 2024 Jul 22]. Available from:

https://www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-

research-center/food-surveys-research-group/docs/fndds/

12. Open Food Facts [Internet]. Open Food Facts. [cited 2024 Apr 28]. Available from:

https://world.openfoodfacts.org/discover

13. Snetselaar LG, de Jesus JM, DeSilva DM, Stoody EE. Dietary Guidelines for

Americans, 2020-2025: Understanding the Scientific Process, Guidelines, and Key

Recommendations. Nutr Today 2021;56:287–95.

14. Healthy diet [Internet]. [cited 2024 Apr 29]. Available from: https://www.who.int/news-

room/fact-sheets/detail/healthy-diet

15. Kim JH. Multicollinearity and misleading statistical results. Korean J Anesthesiol

2019;72:558–69.

16. Machado PP, Steele EM, Levy RB, da Costa Louzada ML, Rangan A, Woods J, Gill T,

Scrinis G, Monteiro CA. Ultra-processed food consumption and obesity in the Australian

adult population. Nutr Diabetes 2020;10:39.

17. Fang Z, Rossato SL, Hang D, Khandpur N, Wang K, Lo C-H, Willett WC, Giovannucci

EL, Song M. Association of ultra-processed food consumption with all cause and cause

specific mortality: population based cohort study. BMJ 2024;385:e078476.

18. Kissock KR, Vieux F, Mathias KC, Drewnowski A, Seal CJ, Masset G, Smith J, Mejborn

H, McKeown NM, Beck EJ. Aligning nutrient profiling with dietary guidelines: modifying

the Nutri-Score algorithm to include whole grains. Eur J Nutr 2022;61:541–53.

19. Wood AM, Kaptoge S, Butterworth AS, Willeit P, Warnakula S, Bolton T, Paige E, Paul

DS, Sweeting M, Burgess S, et al. Risk thresholds for alcohol consumption: combined

analysis of individual-participant data for 599 912 current drinkers in 83 prospective

studies. Lancet Elsevier; 2018;391:1513–23.

20. Grüner Nielsen D, Andersen K, Søgaard Nielsen A, Juhl C, Mellentin A. Consistency

between self-reported alcohol consumption and biological markers among patients with

alcohol use disorder - A systematic review. Neurosci Biobehav Rev 2021;124:370–85.

33