Evaluating the association of *APOE* genotype and cognitive resilience in SuperAgers

Alaina Durant, BS[a]; Shubhabrata Mukherjee, MS, PhD[b]; Michael L. Lee, MPH, PhD[b]; Seo-Eun Choi, PhD[b]; Phoebe Scollard, MA[b,c]; Brandon S. Klinedinst, PhD[b]; Emily H. Trittschuh, PhD[d,e]; Jesse Mez, MS, MD[f]; Lindsay A. Farrer, PhD[f,g,h]; Katherine A. Gifford, PsyD[a,h]; Carlos Cruchaga, PhD[i,j]; Jason Hassenstab, PhD[k]; Adam C. Naj, PhD[l,m]; Li-San Wang, PhD[m]; Sterling C. Johnson, PhD[n,o]; Corinne D. Engelman, MSPH, PhD[n,o]; Walter A. Kukull, PhD[p]; C. Dirk Keene, MD, PhD[q]; Andrew J. Saykin, PyD[r,s]; Michael L. Cuccaro, PhD[t]; Brian W. Kunkle, MPH, PhD[t]; Margaret A. Pericak-Vance, PhD[t]; Eden R. Martin, PhD[t]; David A. Bennett, MD[u]; Lisa L. Barnes, PhD[u]; Julie A. Schneider, MD[u]; William S. Bush, PhD[v]; Jonathan L. Haines, PhD[v]; Richard Mayeux, MD, MSc[w]; Badri N. Vardarajan, MS, PhD[w]; Marilyn S. Albert, PhD[x]; Paul M. Thompson, PhD[y]; Angela L. Jefferson, PhD[a,z]; The Alzheimer's Disease Neuroimaging Initiative (ADNI)*; Alzheimer's Disease Genetics Consortium (ADGC); The Alzheimer's Disease Sequencing Project (ADSP); Paul K. Crane, MD, MPH[b]; Logan Dumitrescu, MS, PhD[a,z,aa]; Derek B. Archer, PhD[a,z,aa]; Timothy J. Hohman, PhD[a,z,aa]; Leslie S. Gaynor, PhD[a,bb]

[a]Vanderbilt Memory & Alzheimer's Center, Vanderbilt University Medical Center, 3319 West End Ave, Rm 819, Nashville, TN 37203, USA
[b]Department of Medicine, University of Washington, 1959 NE Pacific Street, Seattle, WA 98195-6420, USA
[c]University Bordeaux, Inserm, Bordeaux Population Health Research Center, UMR1219, 146 Rue Léo Saignat 11, Bordeaux, France 33076
[d]Department of Psychiatry and Behavioral Sciences, University of Washington School of Medicine, 1959 NE Pacific Street, Seattle, WA 98195-6560, USA
[e]VA Puget Sound Health Care System, GRECC, 1660 S Columbian Way, Seattle, WA 98108, USA
[f]Department of Neurology, Boston University Chobanian & Avedisian School of Medicine, Boston University Medical Center, 85 East Concord Street, 1st floor, Boston, MA 02118, USA
[g]Department of Biostatistics, Boston University School of Public Health, Crosstown Building, 801 Massachusetts Avenue 3rd Floor, Boston, MA 02118, USA
[h]Department of Medicine, Boston University Chobanian & Avedisian School of Medicine, 72 East Concord St., Boston, MA 02118, USA
[i]Department of Psychiatry, Washington University in St. Louis, 4444 Forest Park Avenue, St. Louis, MO 63108, USA
[j]NeuroGenomics and Informatics Center, Washington University in St. Louis, 4444 Forest Park Ave Room 5501, St. Louis, MO 63018, USA
[k]Department of Neurology and Department Psychological & Brain Sciences, Washington University in St Louis, 1 Brookings Drive, St. Louis, MO 63130, USA
[l]Department of Biostatistics, Epidemiology, and Informatics, University of Pennsylvania Perelman School of Medicine, 423 Guardian Drive, Philadelphia, PA 19104-6021, USA
[m]Penn Neurodegeneration Genomics Center, Department of Pathology and Laboratory Medicine, University of Pennsylvania Perelman School of Medicine, 3400 Spruce St., Philadelphia, PA 19104-4238, USA
[n]Alzheimer's Disease Research Center, University of Wisconsin-Madison, School of Medicine and Public Health, 600 Highland Ave J5/1 Mezzanine, Madison, WI 53792, USA
[o]Department of Population Health Sciences, University of Wisconsin-Madison, School of Medicine and Public Health, 500 Lincoln Dr., Madison, WI 53706, USA
[p]Department of Epidemiology, School of Public Health, University of Washington, 3980 15th Ave NE, Seattle, WA 98195, USA

qDepartment of Laboratory Medicine and Pathology, University of Washington, 1959 NE Pacific St, Seattle, WA 98195, USA☐

rDepartment of Radiology and Imaging Services, Indiana University School of Medicine, 550 University Blvd, Indianapolis, IN 46202, USA☐

sDepartment of Medical and Molecular Genetics, School of Medicine, Indiana University, 975 W Walnut St IB 130, Indianapolis, IN 46202, USA☐

tJohn P. Hussman Institute for Human Genomics, University of Miami Miller School of Medicine, 1501 NW 10th Ave, Miami, FL 33101, USA☐

uRush Alzheimer's Disease Center, Rush University Medical Center, 1750 W Harrison St, Chicago, IL 60612, USA☐

vCleveland Institute for Computational Biology, Department of Population and Quantitative Health Sciences, School of Medicine, Case Western Reserve University, 2109 Adelbert Rd, Cleveland, OH 44106, USA☐

wDepartment of Neurology, The Taub Institute for Research on Alzheimer's Disease and The Aging Brain, Columbia University Medical Center and The New York Presbyterian Hospital, 710 W 168th St #3, New York, NY 10032, USA☐

xDepartment of Neurology, The Johns Hopkins University School of Medicine, 550 N Broadway #415, Baltimore, MD 21205, USA☐

yKeck School of Medicine, University of Southern California, 1975 Zonal Ave, Los Angeles, CA 90033, USA☐

zDepartment of Neurology, Vanderbilt University Medical Center, 1211 Medical Center Drive, Nashville, TN 37232, USA

aaVanderbilt Genetics Institute, Vanderbilt University Medical Center, 1211 Medical Center Drive, Nashville, TN 37232, USA☐

bbDivision of Geriatric Medicine, Department of Medicine, Vanderbilt University Medical Center, 1161 21st Ave S, Nashville, TN 37232, USA

Corresponding Author: Leslie S. Gaynor, leslie.gaynor@vumc.org, Work Phone: (615) 421-0296, Address: Vanderbilt Memory and Alzheimer's Center, 3319 West End Ave, Rm 819, Nashville, TN 37203

Abstract

INTRODUCTION: "SuperAgers" are oldest-old adults (ages 80+) whose memory performance more closely resembles middle-aged adults. The present study examined *APOE* allele frequency in non-Hispanic Black (NHB) and non-Hispanic White (NHW) SuperAgers compared to controls and Alzheimer's disease dementia cases.

METHODS: In 18,080 participants from eight cohorts, harmonized clinical diagnostics and memory, executive function, and language domain scores were used to identify SuperAgers, cases, and controls across age-defined bins.

RESULTS: NHW SuperAgers had significantly lower frequency of *APOE*-ε4 alleles and higher frequency of *APOE*-ε2 alleles compared to all cases and controls, including oldest-old controls. Similar patterns were found in a small yet substantial sample of NHB SuperAgers; however, not all comparisons with controls reached significance.

DISCUSSION: We demonstrated strong evidence that *APOE* allele frequency relates to SuperAger status. Further research is needed with a larger sample of NHB SuperAgers to determine if mechanisms conferring resilience differ across race groups.

1 Background

"SuperAgers" is a term used to describe oldest-old (ages 80+) adults with episodic memory performance most closely resembling adults in their 50s to mid-60s.[1,2] Studies suggest that resilience to Alzheimer's disease (AD) pathological changes and neurodegeneration may explain SuperAgers' high memory scores.[3-6] Further research is needed to elucidate factors conferring resilience to AD-related brain changes and subsequent cognitive decline in SuperAgers. Moreover, research is needed to explore resiliency factors in non-Hispanic Black (NHB) SuperAgers, as this group is largely understudied.[7]

APOE-ε4 is the strongest genetic risk factor for late-onset AD.[8] The Northwestern SuperAging project reported lower *APOE*-ε4 allele frequency in SuperAgers (N = 10-12) compared to non-demented older adults.[2,9] In contrast, most studies report no differences in *APOE*-ε4 allele frequency between SuperAgers and oldest-old adults with typical memory performance, both groups having lower *APOE*-ε4 allele frequency compared to AD dementia cases.[4-6,10-12] Notably, these studies have small SuperAger samples (N = 25-64)[4-6,10-12] oftentimes drawn from the same cohort, thus limiting their generalizability and reliability. Further, these studies exclusively include NHW participants.[4-6,10-12] To our knowledge, only one study has been published with a NHB SuperAger sample (N = 61) and did not find a significant difference in *APOE*-ε4 allele frequency between SuperAgers and same-age controls.[7] Even fewer studies have explored the relationship of *APOE*-ε2, the protective *APOE* allele, and SuperAger status,[11-13] likely due to the low minor allele frequency of *APOE*-ε2. Studies of *APOE*-ε2 allele frequency and superior memory in the oldest-old have not found a significant relationship[11-13]; however, questions of statistical power, generalizability, and reliability remain.

The present study aims to explore *APOE*-ε4 and -ε2 allele frequency in SuperAgers compared to AD dementia cases and controls in a large, harmonized multicohort dataset from the Alzheimer's Disease Sequencing Project Phenotype Harmonization Consortium (ADSP-PHC). Using harmonized clinical diagnoses and cognitive domain scores (e.g., memory,

executive function, language), we classified NHW and NHB middle-aged, old, and oldest-old adults as cases, controls, or SuperAgers, and compared *APOE*-ε4 and -ε2 allele frequency of SuperAgers to cases and controls by age bin. Although prior literature suggests that there is not a relationship between optimal memory in oldest-old age and *APOE* genotype, this is likely due to a limitation of sample size. The ADSP-PHC has enabled us to complete, to our knowledge, the largest and most racially diverse study to date of *APOE* allele frequency and SuperAger status. We hypothesize that SuperAgers will possess a lower frequency of *APOE*-ε4 alleles and a higher frequency of *APOE*-ε2 alleles compared to both AD dementia cases and controls.

2 Methods

2.1 Study Population

The ADSP-PHC was assembled in 2021 to provide large-scale harmonization of ADSP cohorts, spanning markers of cognition, neuroimaging, fluid biomarkers, and neuropathology. Cohorts that are part of ADSP-PHC and were included in the present study are: Adult Changes in Thought (ACT),[14] Alzheimer's Disease Neuroimaging Initiative (ADNI),[15] Biomarkers of Cognitive Decline Among Normal Individuals (BIOCARD),[16] National Alzheimer's Coordinating Centers (NACC),[17] National Institute on Aging Alzheimer's Disease Family Based Study (NIA-AD FBS),[18] Religious Orders Study/Rush Memory and Aging Project/Minority Aging Research Study (ROS/MAP/MARS),[19,20,21] Knight Alzheimer's Disease Research Center (Knight ADRC),[22] and Wisconsin Registry for Alzheimer's Prevention (WRAP).[23]

Written informed consent was obtained from all participants in each cohort, and research was carried out with protocols approved by each site's institutional review board. These secondary analyses were approved by the Vanderbilt University Medical Center institutional review board.

2.2 Cognitive Domain Scores

Each cohort used different neuropsychological assessment tools to measure cognition which may not be on the same scale. We derived co-calibrated and harmonized cognitive

domain scores for four domains (i.e., memory, executive function, language, and visuospatial) using confirmatory factor analysis.[24] Consistent with previous definitions of SuperAgers, only memory, executive function, and language domains were used in present analyses.[1,2] To ensure inclusion of high-quality harmonization, cognitive domain scores with a standard error of measurement > 0.6 (estimated during the co-calibration and composite generation procedure) were excluded. Visit the data dictionary at https://vmacdata.org/adsp-phc/data/data-dictionary/cognition to explore the cognitive variables from each cohort that make up each cognitive domain.

For the purpose of this study, regression-based cognitive domain scores were created for participant classification. Age, years of education, sex, and cohort were regressed on cognitive domain score at each available timepoint. Residuals were used to capture cognitive performance not explained by demographic variables or cohort differences. Separate models were used for NHW and NHB participants.

2.3 Participant Classification

Participants were grouped by race (self or study reported), clinical status, and age (**Figure 1A**). Age bins were decided based on the age-related criteria of SuperAgers (ages 80+) and the middle-aged adults whose memory performance is compared to oldest-old adults to determine SuperAger status (ages 50-64).[25] Therefore, the age bins included middle-aged (ages 50-64), old (ages 65-79), and oldest-old (ages 80+) adults. Participant classification was determined separately for NHW and NHB adults. Given that longitudinal data were available for most participants, and some participants' ages spanned multiple age bins, participant classification was decided using a predetermined schema (**Figure 1B**).

Firstly, participants with a clinical diagnosis of AD dementia at least once were considered cases. Across all cohorts, a diagnosis of AD dementia followed standard published criteria (National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association, or NINCDS-ADRDA).[26-28] Given that

the present study focused on dementia cases likely due to AD, participants with comorbid neurological disorders (e.g., stroke, traumatic brain injury with significant loss of consciousness) or primary diagnoses of non-AD dementia (e.g., Dementia with Lewy Bodies, Vascular Dementia) at any timepoint were excluded. Age bin was determined based on age at first AD dementia diagnosis.

Next, SuperAgers were identified using previously published criteria.[1,2] Across all cohorts, SuperAgers were defined as oldest-old adults with (1) memory scores at or above the mean of middle-aged adults (ages 50-64), (2) executive function and language residual scores no more than 1 standard deviation below their same-aged peers (ages 80+) at the same visit as their superior memory performance, and (3) whose diagnosis remained "cognitively normal" for the duration of study participation.

Regarding classification as a control, given the importance of oldest-old (ages 80+) participants to our central analyses, we sequentially identified oldest-old, middle-aged, and then old controls. Criteria for controls included (1) memory, executive function, and language residual scores no lower than 1 standard deviation below their same-aged peers at a single visit, and (2) whose diagnosis remained "cognitively normal" for the duration of their study participation.

2.4 *APOE* Genotyping

APOE haplotypes were determined from the single nucleotide variants rs7412 and rs429358 for ACT, BIOCARD, NACC, NIA-AD FBS, Knight ADRC, and WRAP and from pyrosequencing of *APOE* codons 112 and 158 for ADNI and ROS/MAP/MARS.[29,30]

2.5 Statistical Analyses

Statistical analyses were performed using R Statistical Software (v4.2.3.).[31] Logistic regression models examined differences in *APOE-*ε4 and -ε2 allele frequency of SuperAgers compared to cases and controls at all age bins. *APOE* allele positivity was determined by allele presence (0 = no allele present, 1 = one or more alleles present). Models covaried for sex and years of education due to their known modifying effects on the relationship of *APOE* and late-life

cognition.[32,33] Sensitivity analyses included analyses removing individuals with *APOE*-ε2/ε4

genotype and adding *APOE* genotyping method as a covariate. Additionally, to determine

whether differences in sample size explained differences in the significance of comparisons

across NHW and NHB samples, we included sensitivity analyses randomly down-sampling the

NHW sample by participant classification and cohort. Correction for multiple comparisons was

applied using Benjamini-Hochberg false discovery rate (FDR) procedure.[34] Results tables

include odds ratios (OR), confidence intervals (CI), and FDR-corrected *p*-values.

3 Results

3.1 Participant Characteristics.

In total, 18,080 participants were included in the present analyses with a total of 78,549

datapoints (**Table 1**). Participants completed an average of 4 ± 4 visits over 5 ± 5 years. The

number of follow-up visits and length of follow-up varied by cohort due to differences in study

design.

Average baseline age varied by cohort (Age$_{[all\ cohorts]}$ = 72 ± 10); the youngest cohorts on

average, BIOCARD and WRAP, primarily recruited cognitively normal participants. Generally,

cohorts were highly educated (Years of Education$_{[all\ cohorts]}$ = 15 ± 3), mostly female (62.9%), and

mostly NHW (85.4%). The frequency of each *APOE* genotype differed by cohort, with higher

frequency of ε3/ε4 and ε4/ε4 genotypes in cohorts with a higher frequency of AD dementia

cases.

3.2 Participant Classification

SuperAgers made up 9% of all participants (N = 1,623). The two youngest cohorts,

BIOCARD and WRAP, did not contribute any SuperAgers. Cognitively unimpaired controls

comprised 42% of all participants (N = 7,628), and cases made up 49% of all participants (N =

8,829).

Table 2 displays participant characteristics of NHW and NHB SuperAgers, controls, and

cases in age-defined bins. On average, NHW SuperAgers (N = 1,412) were somewhat older,

had more years of education, and included more males than NHB SuperAgers (N = 211). There was not a difference in the proportion of participants in each participant classification (SuperAger, control, case) across racialized groups ($\chi^2_{[2]}$ = 3.79, p = 0.15). Comparing NHW and NHB participants across age-defined bins, all NHW bins had greater average years of education and a higher proportion of males than NHB bins.

3.3 *APOE* Allele Frequency

In NHW comparisons (**Table 3A**), SuperAgers had a significantly higher frequency of *APOE*-ε2 alleles (**Figure 2A;** middle-aged controls: OR = 1.38, 95%CI[1.13, 1.68], P_{FDR} = 0.002; old controls: OR = 1.24, 95%CI[1.05, 1.47], P_{FDR} = 0.015; oldest-old controls: OR = 1.28, 95%CI[1.03, 1.59], P_{FDR} = 0.035; middle-aged cases: OR = 4.55, 95%CI[3.30, 6.27], P_{FDR} < 0.001; old cases: OR = 4.02, 95%CI[3.26, 4.96], P_{FDR} < 0.001; oldest-old cases: OR = 2.03, 95%CI[1.68, 2.44], P_{FDR} < 0.001) and a significantly lower frequency of *APOE*-ε4 alleles (**Figure 2B**; middle-aged controls: OR = 0.37, 95%CI[0.31, 0.43], P_{FDR} < 0.001; old controls: OR = 0.55, 95%CI[0.47, 0.64], P_{FDR} < 0.001; oldest-old controls: OR = 0.81, 95%CI[0.67, 0.99], P_{FDR} = 0.044; middle-aged cases: OR = 0.18, 95%CI[0.15, 0.21], P_{FDR} < 0.001; old cases: OR = 0.09, 95%CI[0.08, 0.11], P_{FDR} < 0.001; oldest-old cases: OR = 0.32, 95%CI[0.27,0.37], P_{FDR} < 0.001) compared to all cases and controls. In contrast, NHB SuperAgers (**Table 3B**) had a significantly higher frequency of *APOE*-ε2 alleles only compared to cases (**Figure 2C;** middle-aged cases: OR = 3.59, 95%CI[1.60, 8.06], P_{FDR} = 0.005; old cases: OR = 4.56, 95%CI[2.75, 7.55], P_{FDR} < 0.001; oldest-old cases: OR = 2.73, 95%CI[1.75, 4.25], P_{FDR} < 0.001), and a significantly lower frequency of *APOE*-ε4 alleles compared to all cases and controls except oldest-old controls (**Figure 2D;** middle-aged controls: OR = 0.48, 95%CI[0.33, 0.71], P_{FDR} = 0.001; old controls: OR = 0.66, 95%CI[0.47, 0.94], P_{FDR} = 0.035; middle-aged cases: OR = 0.12, 95%CI[0.07, 0.22], P_{FDR} < 0.001; old cases: OR = 0.12, 95%CI[0.08, 0.18], P_{FDR} < 0.001; oldest-old cases: OR = 0.39, 95%CI[0.27, 0.57], P_{FDR} < 0.001).

3.4 Sensitivity Analyses

Analyses were repeated removing individuals with an *APOE*-ε2/ε4 genotype. Given their low frequency, very few participants were removed (**Supplementary Table 1;** NHW SuperAgers: N = 1,393; NHB SuperAgers: N = 200) and findings were preserved (**Supplementary Table 2**).

Findings were also preserved when *APOE* genotyping method was included as a covariate (**Supplementary Table 3**).

To determine the impact of sample size on the significance of NHB comparisons, we randomly down-sampled the NHW sample by participant classification and cohort (**Supplementary Table 4**). In the smaller sample, differences in *APOE*-ε2 and -ε4 allele frequency between NHW SuperAgers and controls were no longer significant, except for the comparison between NHW SuperAgers and middle-aged controls (**Supplementary Table 5**).

4 Discussion

Mechanisms conferring resilience to memory decline in oldest-old age are yet unknown. This is the largest published sample of SuperAgers and the largest study to date of the relationship of *APOE*-ε4 and -ε2 allele frequency and SuperAger status. Across 8 national aging cohorts, we identified 1,623 NHW and NHB SuperAgers with *APOE* genotyping using longitudinal harmonized cognitive and clinical data. As expected, we found that SuperAgers had a higher frequency of *APOE*-ε2 alleles and a lower frequency of *APOE*-ε4 alleles compared to individuals with AD dementia. Unlike most previous studies of *APOE* genotype in NHW SuperAgers, we found significant differences in *APOE* genotype compared to controls of all ages, including oldest-old controls. Specifically, the odds of having 1 ε4 allele was about 20% lower in NHW SuperAgers compared to same-age controls, and the odds of having 1 ε2 allele was about 30% higher in NHW SuperAgers compared to same-age controls. This finding held when ε2/ε4 carriers were removed from analyses and when *APOE* genotyping method was added as a covariate. Although NHB SuperAgers also had a higher frequency of *APOE*-ε2 alleles and a lower frequency of *APOE*-ε4 alleles compared to oldest-old controls, comparisons

did not rise to the level of significance. Sensitivity analyses down-sampling NHW SuperAgers suggest that sample size likely impacted the significance of comparisons with NHB SuperAgers.

APOE-ε4 is the strongest genetic risk factor for late-onset AD,[8] and has been shown to be related to increased entorhinal and hippocampal atrophy[35] and amnestic cognitive impairment.[36,37] Our data are supportive of these findings, such that AD dementia cases had a higher frequency of APOE-ε4 alleles compared to controls and SuperAgers.[38] Unlike most studies of genetic resilience in SuperAgers, we found that NHW SuperAgers had a lower frequency of APOE-ε4 alleles compared to oldest-old adults with typical memory performance. This was found in two previous studies published from the same cohort (N = 10-12),[2,9] but was not replicated in subsequent studies with larger samples of SuperAgers.[4-6,10-12] This finding is relatively unexpected. Across racialized groups, the effect of APOE-ε4 is most impressive prior to age 70.[39,40] Additionally, APOE-ε4 carriership is related to increased mortality.[41,42] In line with these studies, we found lower APOE-ε4 allele frequency in oldest-old compared to middle-aged and old controls. Despite NHW oldest-old controls being older than NHW SuperAgers on average, we found that SuperAgers had a significantly lower frequency of APOE-ε4 alleles compared to oldest-old controls, indicating that APOE-ε4 allele carriership influences memory even in adults who live past age 80. Moreover, dementia incidence increases exponentially with age,[43] suggesting that the oldest-old control group also represents exceptional aging. We found that NHW SuperAgers had a significantly lower frequency of APOE-ε4 alleles compared to same-age peers, thus underscoring the utility of the SuperAgers phenotype for uncovering factors conferring resilience in cognitive aging.

The protective APOE-ε2 allele is related to lower likelihood of late-onset AD dementia[13,44] and better cognitive performance in older adults even in the presence of AD neuropathology.[45] Unlike APOE-ε4, APOE-ε2 carriership was previously shown to affect cognition after age 80; more precisely, although oldest-old APOE-ε2 carriers were as likely as APOE-ε4 carriers to meet neuropathologic criteria for AD, they were less likely to be diagnosed

with dementia.[44] Nonetheless, no previous studies have found a relationship of *APOE*-ε2 allele frequency and SuperAger status.[11-13] The present study is the first to find that NHW SuperAgers had a significantly higher frequency of *APOE*-ε2 alleles compared to controls. Our results suggest that *APOE*-ε2 not only reduces the likelihood of dementia in oldest age but increases the likelihood one will possess optimal memory in oldest age. Future studies are needed to determine whether SuperAgers have similar levels of AD neuropathology compared to AD dementia cases as was previously found in oldest-old *APOE*-ε2 carriers without dementia.[44]

Genetic factors underlying the superior memory performance of NHB SuperAgers, a critically under-diagnosed and understudied group, are relatively unknown.[7] Previous research suggests that, while NHB individuals have a higher frequency of *APOE*-ε4 alleles compared to NHW individuals, *APOE*-ε4 carriership is associated with attenuated risk for late-onset AD, yet similar mortality risk, in NHB individuals compared to NHW individuals.[40,46-48] The effect is likely related to global population ancestry; more specifically, *APOE*-ε4 carriership is associated with a greater risk of AD in NHB older adults with decreased global African ancestry or increased global European ancestry.[40] Similar to *APOE*-ε4, NHB individuals have a higher frequency of *APOE*-ε2 alleles compared to NHW individuals.[46] Unlike *APOE*-ε4, researchers did not detect differences in the protective effect of *APOE*-ε2 alleles related to global population ancestry.[40] In fact, a recent study from the MARS cohort found that NHB older adults (ages 65+) with more *APOE*-ε2 alleles had slower cognitive decline over a 10-year study period.[49] Additionally, *APOE*-ε2 carriership was related to better survival in a sample of NHB and NHW individuals from NACC with and without AD neuropathology.[50] In the present study, NHB SuperAgers had a significantly lower frequency of *APOE*-ε4 alleles compared to cases and younger controls, and a significantly higher frequency of *APOE*-ε2 alleles compared only to cases. Importantly, the present study included a substantially smaller sample of NHB SuperAgers compared to NHW SuperAgers, although the sample was still far larger than has been previously reported. Sensitivity analyses suggest that sample size likely impacted the significance of comparisons

with NHB SuperAgers. It is also possible that the lack of significant difference in *APOE* allele carriership between NHB SuperAgers and oldest-old controls is partially explained by survivorship bias; recent research indicates that NHB adults have a higher probability of survival from ages 70 and 80 to 100 compared to NHW adults.[51] Thus, NHB SuperAgers and cognitively normal oldest-old adults may share environmental and genetic factors, including *APOE*-ε2 allele carriership, that support survival and reduced mortality risk in older age, resulting in a lack of significant difference in *APOE* genotype between these two, very similar, exceptional aging groups.[7] Still, more research and targeted recruitment of high-performing NHB oldest-old adults is necessary to determine the role of *APOE* genotype in their sustained optimal memory performance.

The present study has several strengths, including being the largest of its kind to explore the relationship of *APOE* genotype and optimal memory performance in both NHW and NHB SuperAgers. This is the largest study to date to identify differences in *APOE*-ε4 allele frequency based on SuperAger status, and the first study of SuperAgers to find a relationship between *APOE*-ε2 allele frequency and SuperAger status. Our study also has several limitations. Across cohorts, participants were highly educated, limiting generalizability to lower-educated or socioeconomically diverse populations. Additionally, studies have identified other genetic factors that may confer greater AD risk in NHB older adults compared to *APOE*, including *ABCA7*.[53,54] Future studies will need to consider other genetic factors that may also be relevant to exceptional aging in NHB older adults. Moreover, differences in the effect of genetic factors on AD risk have been associated with genetic ancestry, which was only available for a subset of participants in our sample.[40] Subsequent research leveraging advanced genetic analyses to consider admixture may further our understanding of genetic profiles underlying AD risk.

The present study reveals important information about genetic factors associated with SuperAger status. While significant findings were restricted to NHW comparisons, study results importantly direct our attention to the need for targeted recruitment of high-performing NHB

oldest-old adults to more definitively determine the role of genetics in sustained optimal memory performance in oldest-old age across racialized groups. There is an ongoing initiative that intends to answer complex questions about brain aging, resilience, and resistance in a well-characterized SuperAging cohort through longitudinal, multimodal data collection.[55,56] SuperAging research will contribute tremendously to our understanding of factors conferring resilience in oldest-old age, such as preserved regional brain volume, differences in lifestyle factors, and resistance to neuropathology.

References

Harrison, T. M., Mesulam, M. M., Rogalski, E. & Weintraub, S. Superior Memory and Higher Cortical Volumes in Unusually Successful Cognitive Aging. *Journal of the International Neuropsychological Society* **18**, 1081-1085 (2012). https://doi.org/10.1017/S1355617712000847
Rogalski, E. J. *et al.* Youthful Memory Capacity in Old Brains: Anatomic and Genetic Clues from the Northwestern SuperAging Project. *Journal of Cognitive Neuroscience* **25**, 29-36 (2013). https://doi.org/10.1162/jocn_a_00300
de Godoy, L. L. *et al.* Understanding brain resilience in superagers: a systematic review. *Neuroradiology* **63**, 663-683 (2021). https://doi.org/10.1007/s00234-020-02562-1
Dekhtyar, M. *et al.* Neuroimaging markers associated with maintenance of optimal memory performance in late-life. *Neuropsychologia* **100**, 164-170 (2017). https://doi.org/10.1016/j.neuropsychologia.2017.04.037
Gefen, T. *et al.* Morphometric and histologic substrates of cingulate integrity in elders with exceptional memory capacity. *Journal of Neuroscience* **35**, 1781-1791 (2015). https://doi.org/10.1523/JNEUROSCI.2998-14.2015
Harrison, T. M., Maass, A., Baker, S. L. & Jagust, W. J. Brain morphology, cognition, and β-amyloid in older adults with superior memory performance. *Neurobiology of aging* **67**, 162-170 (2018). https://doi.org/10.1016/j.neurobiolaging.2018.03.024
Trammell, A. R., Goldstein, F. C., Parker, M. W. & Hajjar, I. M. Characterization of African-American Super-Agers in the National Alzheimer's Coordinating Center cohort. *J Am Geriatr Soc* **72**, 1995-2005 (2024). https://doi.org/10.1111/jgs.18882
Corder, E. H. *et al.* Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science* **261**, 921-923 (1993). https://doi.org/10.1126/science.8346443
Rogalski, E. *et al.* Cognitive trajectories and spectrum of neuropathology in SuperAgers: The first 10 cases. *Hippocampus* **29**, 458-467 (2019). https://doi.org/10.1002/hipo.22828
Mapstone, M. *et al.* What success can teach us about failure: the plasma metabolome of older adults with superior memory and lessons for Alzheimer's disease. *Neurobiology of aging* **51**, 148-155 (2017). https://doi.org/10.1016/j.neurobiolaging.2016.11.007
Spencer, B. E. *et al.* Alzheimer's polygenic hazard score in SuperAgers: SuperGenes or SuperResilience? *Alzheimer's & Dementia: Translational Research & Clinical Interventions* **8**, e12321 (2022). https://doi.org/10.1002/trc2.1232
Garo-Pascual, M. *et al.* Brain structure and phenotypic profile of superagers compared with age-matched older adults: a longitudinal analysis from the Vallecas Project. *Lancet Healthy Longev* **4**, e374-e385 (2023). https://doi.org/10.1016/S2666-7568(23)00079-X
Corder, E. *et al.* Protective effect of apolipoprotein E type 2 allele for late onset Alzheimer disease. *Nature genetics* **7**, 180-184 (1994). https://doi.org/10.1038/ng0694-180
Kukull, W. A. *et al.* Dementia and Alzheimer disease incidence: a prospective cohort study. *Archives of neurology* **59**, 1737-1746 (2002). https://doi.org/10.1001/archneur.59.11.1737
Jack Jr, C. R. *et al.* The Alzheimer's disease neuroimaging initiative (ADNI): MRI methods. *Journal of Magnetic Resonance Imaging: An Official Journal of the International Society for Magnetic Resonance in Medicine* **27**, 685-691 (2008). https://doi.org/10.1002/jmri.21049
Sacktor, N. *et al.* The BIOCARD Index: A Summary Measure to Predict Onset of Mild Cognitive Impairment. *Alzheimer Disease & Associated Disorders* **31**, 114-119 (2017). https://doi.org/10.1097/WAD.0000000000000194
Beekly, D. L. *et al.* The National Alzheimer's coordinating center (NACC) database: an Alzheimer disease database. *Alzheimer Disease & Associated Disorders* **18**, 270-277 (2004). https://doi.org/lww/alz/88441/03-070R

Lee, J. H. *et al.* Analyses of the National Institute on Aging Late-Onset Alzheimer's Disease Family Study: Implication of Additional Loci. *Archives of neurology* **65**, 1518-1526 (2008). https://doi.org/10.1001/archneur.65.11.1518

Bennett, D. A. *et al.* Religious orders study and rush memory and aging project. *Journal of Alzheimer's disease* **64**, S161-S189 (2018). https://doi.org/10.3233/JAD-179939

Bennett, D. A. *et al.* Overview and findings from the Rush Memory and Aging Project. *Current Alzheimer research* **9**, 646-663 (2012). https://doi.org/10.2174/156720512801322663

Barnes, L. L., Shah, R. C., Aggarwal, N. T., Bennett, D. A. & Schneider, J. A. The Minority Aging Research Study: ongoing efforts to obtain brain donation in African Americans without dementia. *Current Alzheimer research* **9**, 734-745 (2012). https://doi.org/10.2174/156720512801322627

Fernandez, M. V. *et al.* Genetic and multi-omic resources for Alzheimer disease and related dementia from the Knight Alzheimer Disease Research Center. *Sci Data* **11**, 768 (2024). https://doi.org/10.1038/s41597-024-03485-9

Johnson, S. C. *et al.* The Wisconsin Registry for Alzheimer's Prevention: A review of findings and current directions. *Alzheimer's & dementia (Amsterdam, Netherlands)* **10**, 130-142 (2018). https://doi.org/10.1016/j.dadm.2017.11.007

Mukherjee, S. *et al.* Cognitive domain harmonization and cocalibration in studies of older adults. *Neuropsychology* **37**, 409 (2023). https://doi.org/10.1037/neu0000835

Harrison, T. M., Weintraub, S., Mesulam, M. M. & Rogalski, E. Superior Memory and Higher Cortical Volumes in Unusually Successful Cognitive Aging. *Journal of the International Neuropsychological Society* **18**, 1081-1085 (2012). https://doi.org/10.1017/S1355617712000847

McKhann, G. *et al.* Clinical diagnosis of Alzheimer's disease: Report of the NINCDS-ADRDA Work Group* under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology* **34**, 939-944 (1984). https://doi.org/10.1016/j.jalz.2011.03.005

McKhann, G. M. *et al.* The diagnosis of dementia due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers & Dementia: The Journal of the Alzheimer's Association* **7**, 263-269 (2011). https://doi.org/10.1016/j.jalz.2011.03.005

Dubois, B. *et al.* Research criteria for the diagnosis of Alzheimer's disease: revising the NINCDS-ADRDA criteria. *The Lancet Neurology* **6**, 734-746 (2007). https://doi.org/10.1016/S1474-4422(07)70178-3

Yu, L. *et al.* Neuropathologic features of TOMM40 '523 variant on late-life cognitive decline. *Alzheimer's & Dementia* **13**, 1380-1388 (2017). https://doi.org/10.1016/j.jalz.2017.05.002

Naj, A. C. *et al.* Common variants at MS4A4/MS4A6E, CD2AP, CD33 and EPHA1 are associated with late-onset Alzheimer's disease. *Nature Genetics* **43**, 436-441 (2011). https://doi.org/10.1038/ng.801

R: A language and environment for statistical computing. (R Foundation for Statistical Computing, Vienna, Austria, 2021).

Lee, M. *et al.* Education and Cardiovascular Health as Effect Modifiers of APOE ε4 on Dementia: The Atherosclerosis Risk in Communities Study. *Journals of Gerontology Series A, Biological Sciences & Medical Sciences* **77**, 1199-1207 (2022). https://doi.org/10.1093/gerona/glab299

Hohman, T. J. *et al.* Sex-Specific Association of Apolipoprotein E With Cerebrospinal Fluid Levels of Tau. *JAMA neurology* **75**, 989-998 (2018). https://doi.org/10.1001/jamaneurol.2018.0821

Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)* **57**, 289-300 (1995). https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Juottonen, K., Lehtovirta, M., Helisalmi, S., Riekkinen, P. J., Sr. & Soininen, H. Major decrease in the volume of the entorhinal cortex in patients with Alzheimer's disease carrying the apolipoprotein E ε4 allele. *J Neurology, Neurosurgy, and Psychiatry* **65**, 322-327 (1998). https://doi.org/10.1136/jnnp.65.3.322

Snowden, J. S. *et al.* Cognitive phenotypes in Alzheimer's disease and genetic risk. *Cortex; a journal devoted to the study of the nervous system and behavior* **43**, 835-845 (2007). https://doi.org/10.1016/s0010-9452(08)70683-x

Mukherjee, S. *et al.* Genetic data and cognitively defined late-onset Alzheimer's disease subgroups. *Molecular Psychiatry* **25**, 2942-2951 (2018). https://doi.org/10.1038/s41380-018-0298-8

Poirier, J. *et al.* Apolipoprotein E polymorphism and Alzheimer's disease. *The Lancet* **342**, 697-699 (1993). https://doi.org/10.1016/0140-6736(93)91705-q

Farrer, L. A. *et al.* Effects of age, sex, and ethnicity on the association between apolipoprotein E genotype and Alzheimer disease. A meta-analysis. APOE and Alzheimer Disease Meta Analysis Consortium. *Jama* **278**, 1349-1356 (1997). https://doi.org/10.1001/jama.1997.03550160069041

Belloy, M. E. *et al.* APOE Genotype and Alzheimer Disease Risk Across Age, Sex, and Population Ancestry. *JAMA neurology* **80**, 1284-1294 (2023). https://doi.org/10.1001/jamaneurol.2023.3599

Rosvall, L. *et al.* APOE-related mortality: effect of dementia, cardiovascular disease and gender. *Neurobiology of aging* **30**, 1545-1551 (2009). https://doi.org/10.1016/j.neurobiolaging.2007.12.003

McKay, G. J. *et al.* Variations in apolipoprotein E frequency with age in a pooled analysis of a large group of older people. *American journal of epidemiology* **173**, 1357-1364 (2011). https://doi.org/10.1093/aje/kwr015

Jorm, A. F. & Jolley, D. The incidence of dementia: a meta-analysis. *Neurology* **51**, 728-733 (1998). https://doi.org/10.1212/wnl.51.3.728

Berlau, D. J., Corrada, M. M., Head, E. & Kawas, C. H. APOE ε2 is associated with intact cognition but increased Alzheimer pathology in the oldest old. *Neurology* **72**, 829-834 (2009). https://doi.org/10.1212/01.wnl.0000343853.00346.a4

Shinohara, M. *et al.* APOE2 eases cognitive decline during Aging: Clinical and preclinical evaluations. *Annals of neurology* **79**, 758-774 (2016). https://doi.org/10.1002/ana.24628

Rajan, K. B. *et al.* Racial Differences in the Association Between Apolipoprotein E Risk Alleles and Overall and Total Cardiovascular Mortality Over 18 Years. *Journal American Geriatrics Society* **65**, 2425-2430 (2017). https://doi.org/10.1111/jgs.15059

Beydoun, M. A. *et al.* Race, APOE genotypes, and cognitive decline among middle-aged urban adults. *Alzheimers Research & Therapy* **13**, 120 (2021). https://doi.org/10.1186/s13195-021-00855-y

Tang, M. X. *et al.* The APOE-epsilon4 allele and the risk of Alzheimer disease among African Americans, whites, and Hispanics. *Jama* **279**, 751-755 (1998). https://doi.org/10.1001/jama.279.10.751

Nsor, N. A., Bourassa, K. J., Barnes, L. L. & Brown, C. L. The Effects of APOE Alleles, Cognitive Activities, and Social Activities on Cognitive Decline in African Americans. *The journals of gerontology. Series B, Psychological sciences and social sciences*, gbae172 (2024). https://doi.org/10.1093/geronb/gbae172

Shinohara, M. *et al.* APOE2 is associated with longevity independent of Alzheimer's disease. *eLife* **19**, e62199 (2020). https://doi.org/10.7554/eLife.62199

Ouellette, N. & Perls, T. Race and ethnicity dynamics in survival to 100 years in the United States. *J Intern Med* **297**, 2-21 (2025). https://doi.org/10.1111/joim.20031

Nemes, S., Jonasson, J. M., Genell, A. & Steineck, G. Bias in odds ratios by logistic regression modelling and sample size. *BMC Medical Research Methodology* **9**, 56 (2009). https://doi.org/10.1186/1471-2288-9-56

Logue, M. W., Dasgupta, S. & Farrer, L. A. Genetics of Alzheimer's Disease in the African American Population. *J Clin Med* **12**, 5189 (2023). https://doi.org/10.3390/jcm12165189

Reitz, C. *et al.* Variants in the ATP-binding cassette transporter (ABCA7), apolipoprotein E ε4,and the risk of late-onset Alzheimer disease in African Americans. *Jama* **309**, 1483-1492 (2013). https://doi.org/10.1001/jama.2013.2973

Rogalski, E. J. *et al.* Enrollment and Scientific Progress of the Multisite SuperAging Research Initiative. *Alzheimer's & Dementia* **19**, e078625 (2023). https://doi.org/10.1002/alz.078625

Rogalski, E. J. *et al.* The SuperAging Research Initiative: A multisite consortium focused on identifying factors promoting extraordinary cognitive aging. *Alzheimer's & Dementia* **18**, e066407 (2022). https://doi.org/10.1002/alz.066407

Acknowledgements/Conflicts/Funding Sources: The authors have nothing to disclose. The ADSP Phenotype Harmonization Consortium (ADSP-PHC) is funded by NIA (U24 AG074855, U01 AG068057 and R01 AG059716). This study was also funded by several additional funding sources, including K01 AG073584 (DBA) and K24 AG046373 (ALJ). The data contributed from the Adult Changes in Thought study was supported by NIA U19 AG066567. The data contributed from the Wisconsin Registry for Alzheimer's Prevention was supported by NIA R01 AG021155, R01 AG0271761, R01 AG037639, and R01 AG054047. Data collection and sharing for this project was funded (in part) by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (NIA U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California. Data contributed from MAP/ROS/MARS was supported by NIA R01 AG017917, P30 AG10161, P30 AG072975, R01 AG022018, R01 AG056405, UH2 NS100599, UH3 NS100599, R01 AG064233, R01 AG15819 and R01 AG067482, and the Illinois Department of Public Health (Alzheimer's Disease Research Fund). Data can be accessed at www.radc.rush.edu. The NACC database is funded by NIA/NIH Grant U24 AG072122. NACC data are contributed by the NIA-funded ADRCs: P30 AG062429 (PI James Brewer, MD, PhD), P30 AG066468 (PI Oscar Lopez, MD), P30 AG062421 (PI Bradley Hyman, MD, PhD), P30 AG066509 (PI Thomas Grabowski, MD), P30 AG066514 (PI Mary Sano, PhD), P30 AG066530 (PI Helena Chui, MD), P30 AG066507 (PI Marilyn Albert, PhD), P30 AG066444 (PI David Holtzman, MD), P30 AG066518 (PI Lisa Silbert, MD, MCR), P30 AG066512 (PI Thomas Wisniewski, MD), P30 AG066462 (PI Scott Small, MD), P30 AG072979 (PI David Wolk, MD), P30 AG072972 (PI Charles DeCarli, MD), P30 AG072976 (PI Andrew Saykin, PsyD), P30 AG072975 (PI Julie A. Schneider, MD, MS), P30 AG072978 (PI Ann McKee, MD), P30 AG072977 (PI Robert Vassar, PhD), P30 AG066519 (PI Frank LaFerla, PhD), P30 AG062677 (PI Ronald Petersen, MD, PhD), P30 AG079280 (PI Jessica Langbaum, PhD), P30 AG062422 (PI Gil Rabinovici, MD), P30 AG066511 (PI Allan Levey, MD, PhD), P30 AG072946 (PI Linda Van Eldik, PhD), P30 AG062715 (PI Sanjay Asthana, MD, FRCP), P30 AG072973 (PI Russell Swerdlow, MD), P30 AG066506 (PI Glenn Smith, PhD, ABPP), P30 AG066508 (PI Stephen Strittmatter, MD, PhD), P30 AG066515 (PI Victor Henderson, MD, MS), P30 AG072947 (PI Suzanne Craft, PhD), P30 AG072931 (PI Henry Paulson, MD, PhD), P30 AG066546 (PI Sudha Seshadri, MD), P30 AG086401 (PI Erik Roberson, MD, PhD), P30 AG086404 (PI Gary Rosenberg, MD), P20 AG068082 (PI Angela Jefferson, PhD), P30 AG072958 (PI Heather Whitson, MD), P30 AG072959 (PI James Leverenz, MD). NACC data can be accessed at naccdata.org. The recruitment and clinical characterization of research participants at Washington University were supported by NIH P30 AG066444 (JCM), P01 AG03991 (JCM), and P01 AG026276 (JCM). The National Institute on Aging-AD Family Based Study (NIA-AD FBS; https://www.neurology.columbia.edu/research/research-centers-and-programs/national-institute-aging-alzheimers-disease-family-based-study-nia-ad-fbs) collected the samples used in this study and is supported by National Institute on Aging (NIA) grant U24 AG056270. Additional families were contributed to the NIA-AD FBS through NIH grants: R01 AG028786, R01 AG027944, R01 AG027944, RF1 AG054074, U01 AG052410. This work was supported by access to equipment made possible by the Hope Center for Neurological Disorders, the Neurogenomics and Informatics Center (NGI: https://neurogenomics.wustl.edu/) and the Departments of Neurology and Psychiatry at Washington

University School of Medicine. Data contributed from the Wisconsin Registry for Alzheimer's Prevention were supported by R01 AG027161.

Consent Statement: All participants provided informed consent in their respective cohort studies.

Key Words: *APOE* genotype, cognitive aging, Alzheimer's disease, SuperAgers, cognitive resilience.

Figure 1. Flow Diagram for Participant Classification of SuperAgers, Cases, and Controls. (A) Flowchart depicting inclusion and exclusion criteria for identifying SuperAgers, AD dementia cases, controls. (B) Flowchart depicting selection order of SuperAgers, cases, and controls. Age range of participants indicated by line segment with arrows on each end. Age of participant classification is indicated by position of shorter, labeled line segments. Closed circles at the end of line segments indicate inclusion of age, such that age range is less-than-or-equal-to or greater-than-or-equal-to the age with which the circle aligns, while open circles indicate exclusion of age, such that age range is less-than or greater-than the age with which the circle aligns. Sequence of selection is indicated by line height, higher lines indicating earlier selection. Abbreviations: ADSP-PHC, Alzheimer's Disease Sequencing Project – Phenotype Harmonization Consortium; AD, Alzheimer's Disease; CN, Cognitively Normal; MEM, Memory; EXF, Executive Functioning; LAN, Language.

Table 1. Participant Characteristics by Cohort.

Participant Demographics	All	ACT	ADNI	BIOCARD	Knight ADRC	NACC	NIA-AD FBS	ROS, MAP, MARS	WRAP
No. participants	18080	2188	825	51	735	11851	645	1462	323
No. observations	78549	13016	3568	455	3314	42004	956	13931	1305
Visits, mean (SD)	4 (4)	6 (3)	4 (3)	9 (3)	5 (4)	4 (3)	1 (1)	10 (6)	4 (2)
Follow-up time, mean (SD), y	5 (5)	10 (6)	3 (3)	13 (4)	4 (5)	3 (4)	2 (3)	9 (6)	9 (5)
Baseline Age, mean (SD), y	72 (10)	73 (6)	74 (7)	53 (9)	74 (8)	72 (10)	74 (12)	77 (8)	53 (6)
NHW Race, No. (%)	15698 (87)	2116 (97)	753 (91)	51 (100)	635 (86)	10149 (86)	605 (94)	1086 (74)	303 (94)
Education, mean (SD), y	15 (3)	15 (3)	16 (3)	17 (2)	15 (3)	16 (3)	14 (3)	16 (4)	16 (3)
Female Sex, No. (%)	11213 (62)	1295 (59)	401 (49)	30 (59)	453 (62)	7319 (62)	406 (63)	1091 (75)	218 (67)
APOE genotype									
ε2/ε2, %	0.4	0.6	0.4	0	0.3	0.4	0.6	0.5	0.3
ε2/ε3, %	8.8	12.5	8.6	13.7	8.7	7.9	5.1	12.1	7.1
ε2/ε4, %	2.6	2.3	2.4	0	2.7	2.6	3.3	2.1	3.4
ε3/ε3, %	47.0	58.8	44.8	62.7	41.9	44.2	37.7	57.9	52.6
ε3/ε4, %	33.5	24.0	33.1	19.6	38.8	35.6	43.1	24.4	32.2
ε4/ε4, %	7.8	1.7	10.7	3.9	7.6	9.4	10.2	2.9	4.3
SuperAgers, No. (%)	1623 (9)	275 (13)	59 (7)	0 (0)	53 (7)	935 (8)	16 (2)	285 (19)	0 (0)
Controls, No. (%)	7628 (42)	1207 (55)	335 (41)	49 (96)	290 (39)	4605 (39)	309 (48)	510 (35)	323 (100)
Cases, No. (%)	8829 (49)	706 (32)	431 (52)	2 (4)	392 (53)	6311 (53)	320 (50)	667 (46)	0 (0)

Abbreviations: NHW, Non-Hispanic White; ACT, Adult Changes in Thought; ADNI, Alzheimer's Disease Neuroimaging Initiative; BIOCARD, Biomarkers of Cognitive Decline Among Normal Individuals; Knight ADRC, Knight Alzheimer's Disease Research Center at Washington University; NACC, National Alzheimer's Coordinating Centers; NIA-AD FBS, National Institute on Aging Alzheimer's Disease Family Based Study; ROS, Religious Orders Study; MAP, Memory and Aging Project; MARS, Minority Aging Research Study; WRAP, Wisconsin Registry for Alzheimer's Prevention.

Table 2. Characteristics of SuperAgers, Cases, and Controls by Race and Age Bin.

A

	SuperAgers	Middle-Aged Controls	Old Controls	Oldest-old Controls	Middle-Aged Cases	Old Cases	Oldest-old Cases
No. participants	1412	1622	3202	1213	1101	3528	3258
No. observations	11528	7067	12308	6980	2692	10858	15434
Visits, mean (SD)	8 (5)	4 (3)	4 (3)	6 (4)	2 (2)	3 (2)	5 (4)
Follow-up time, mean (SD), y	10 (6)	6 (5)	4 (4)	8 (6)	2 (2)	2 (3)	5 (6)
Baseline age, mean (SD), y	77 (7)	57 (5)	70 (4)	79 (7)	59 (5)	73 (4)	83 (6)
Education, mean (SD), y	16 (3)	16 (2)	16 (3)	15 (3)	15 (3)	15 (3)	15 (3)
Female Sex, No. (%)	953 (67)	1119 (69)	1964 (61)	661 (54)	618 (56)	1905 (54)	1989 (61)
APOE-ε2 Frequency, No. (%)	246 (17)	216 (13)	467 (15)	179 (15)	50 (5)	184 (5)	321 (10)
APOE-ε4 Frequency, No. (%)	276 (20)	647 (40)	981 (31)	270 (22)	639 (58)	2495 (71)	1425 (44)

B

	SuperAgers	Middle-Aged Controls	Old Controls	Oldest-old Controls	Middle-Aged Cases	Old Cases	Oldest-old Cases
No. participants	211	296	752	145	85	413	444
No. observations	1830	1024	3106	827	171	1300	2281
Visits, mean (SD)	9 (5)	3 (3)	4 (3)	6 (4)	2 (1)	3 (3)	5 (5)
Follow-up time, mean (SD), y	9 (6)	4 (4)	4 (4)	6 (5)	1 (1)	3 (3)	5 (6)
Baseline age, mean (SD), y	77 (6)	59 (4)	70 (4)	79 (6)	60 (4)	73 (4)	83 (6)

Education, mean (SD), y	15 (3)	15 (3)	15 (3)	13 (3)	14 (3)	14 (4)	13 (4)
Female Sex, No. (%)	177 (84)	222 (75)	564 (75)	115 (79)	58 (68)	283 (69)	331 (75)
APOE-ε2 Frequency, No. (%)	56 (27)	68 (23)	155 (21)	27 (19)	8 (9)	36 (9)	53 (12)
APOE-ε4 Frequency, No. (%)	54 (26)	125 (42)	261 (35)	39 (27)	65 (76)	307 (74)	216 (49)

(A) Non-Hispanic White; (B) Non-Hispanic Black

Table 3. Logistic Regression Model Results Comparing *APOE*-ε2 and -ε4 Allele Frequency among SuperAgers, Cases, and Controls.

A

	APOE-ε2		*APOE*-ε4	
	OR (CI)	P_{FDR}	OR (CI)	P_{FDR}
SuperAgers vs. Middle-Aged Controls	1.38 (1.13, 1.68)	*0.002*	0.37 (0.31, 0.43)	*<0.001*
SuperAgers vs. Old Controls	1.24 (1.05, 1.47)	*0.015*	0.55 (0.47, 0.64)	*<0.001*
SuperAgers vs. Oldest-Old Controls	1.28 (1.03, 1.59)	*0.035*	0.81 (0.67, 0.99)	*0.044*
SuperAgers vs. Middle-Aged Cases	4.55 (3.30, 6.27)	*<0.001*	0.18 (0.15, 0.21)	*<0.001*
SuperAgers vs. Old Cases	4.02 (3.26, 4.96)	*<0.001*	0.09 (0.08, 0.11)	*<0.001*
SuperAgers vs. Oldest-Old Cases	2.03 (1.68, 2.44)	*<0.001*	0.32 (0.27, 0.37)	*<0.001*

B

	APOE-ε2		*APOE*-ε4	
	OR (CI)	P_{FDR}	OR (CI)	P_{FDR}
SuperAgers vs. Middle-Aged Controls	1.19 (0.79, 1.80)	0.423	0.48 (0.33, 0.71)	*0.001*
SuperAgers vs. Old Controls	1.41 (0.99, 2.01)	0.079	0.66 (0.47, 0.94)	*0.035*
SuperAgers vs. Oldest-Old Controls	1.63 (0.93, 2.85)	0.115	1.18 (0.70, 1.98)	0.564
SuperAgers vs. Middle-Aged Cases	3.59 (1.60, 8.06)	*0.005*	0.12 (0.07, 0.22)	*<0.001*

SuperAgers vs. Old Cases	4.56 (2.75, 7.55)	*<0.001*	0.12 (0.08, 0.18)	*<0.001*
SuperAgers vs. Oldest-Old Cases	2.73 (1.75, 4.25)	*<0.001*	0.39 (0.27, 0.57)	*<0.001*

(A) Non-Hispanic White; (B) Non-Hispanic Black. Abbreviations: OR, Odds Ratio; CI, Confidence Interval (95%); P_{FDR}, FDR-corrected P-value.

A

B

Figure 1. Flow Diagram for Participant Classification of SuperAgers, Cases, and Controls.