Title: Quantifying the spatiotemporal dynamics of the first two epidemic waves of SARS-

CoV-2 infections in the United States

Short Title: Dynamics of the first two epidemic waves of SARS-CoV-2 infections in US

Rafael Lopes[1,2,*,#], Nicole A. Swartwood[3,*,#], Yu Lan[1,2], Melanie H. Chitwood[1,2], Fayette

Klaassen[3], Joshua A. Salomon[4], Nicolas A. Menzies[3], Joshua L. Warren[5,2], Nathan D.

Grubaugh[1,2], & Ted Cohen[1,2,†]

[1] Department of Epidemiology of Microbial Diseases, Yale School of Public Health, Yale

University, New Haven, Connecticut, USA

[2] Public Health Modeling Unit, Yale School of Public Health, Yale University, New Haven,

Connecticut, USA

[3] Department of Global Health and Population, Harvard T.H. Chan School of Public Health,

Harvard University, Boston, Massachusetts, USA

[4] Department of Medicine, Stanford University, Stanford, California, USA

[5] Department of Biostatistics, Yale School of Public Health, Yale University, New Haven,

Connecticut, USA

* Co-first authors

[†] Senior author

[#] Corresponding Authors:

Rafael Lopes, 350 George Street, New Haven, CT 06511.

Email: rafael.lopes@yale.edu Phone (+1) 475-300-8845

Nicole A. Swartwood, 90 Smith Street, Boston, MA 02120.

Email: nswartwood@hsph.harvard.edu. Phone (+1) 617-432-6171.

Abstract:

SARS-CoV-2 infection rates displayed striking temporal and spatial variation during the emergence of new variants globally and within the United States. While spatiotemporal "waves" of infection have been observed, quantitative assessments of their spread remain limited. Here, we estimate and compare the speed and spatial extent of the first two major infection waves in the United States, illustrating these dynamics through detailed visualizations. Our findings reveal that the origins of these waves coincide with large gatherings and the relaxation of masking mandates. Notably, the second wave spread more rapidly than the first, driven by multiple, non-contiguous origins of infection. This highlights the role of regional heterogeneity in epidemic dynamics and underscores the importance of localized public health measures in mitigating ongoing outbreaks.

Author Summary:

Over the pandemic of SARS-CoV-2, efforts to identify and visualize the disease progression were made. However, quantitative visualization of the infections spreading are limited until today. Here we developed tools to visualize the spatial and temporal spreading of SARS-CoV-2 first two waves of infections over the Contiguous United States. We generated novels figures and movies that captures the dynamics of spreading and developed a new mapping of incidence of SARS-CoV-2 that goes below county-level. These outputs can help public health understanding and control efforts of the disease.

Introduction

Person-to-person transmission of SARS-CoV-2 was confirmed in the United States (US) in late January 2020 and a national public health emergency was declared six weeks later (1,2). Spatial visualizations of observed and estimated COVID-19 cases and deaths displayed wave-like spread as new variants were introduced in the country (3,4). Previous investigations have examined the correlation between the spatiotemporal patterns of COVID-19 spread and human mobility (5–7). Other studies have estimated the effectiveness of public health measures such as lockdowns and travel restrictions in interrupting or modifying these patterns (7–9). However, efforts to quantify the speed and spatial extent of viral spread over distinct epidemic waves have been limited. Here, we apply a version of the Besag, York, and Mollié (BYM) spatial model (10,11) to temporally- and spatially-resolved estimates of SARS-CoV-2 infections derived from a previously described model (3). Based on this analysis we quantify the speed at which SARS-CoV-2 outbreaks spread during the two large waves of SARS-CoV-2 infections in the United States during March 2020 to December 2021.

Materials and Methods

Data sources

We used previously published daily estimates of SARS-CoV-2 infections at the county-level in the United States from a Bayesian nowcasting model that synthesized reported COVID-19 cases and deaths, accounting for both under ascertainment and time lags (3). Our dataset encompasses the period from March 2020 to December 2021. To estimate population denominators for per

capita analyses, we used 2019 population estimates from the US Census Bureau, which provided

population size estimates for all Census Block Groups (CBGs) in the United States (12).

Regularization of geographical units

The smallest consistently available unit of reporting for US COVID-19 cases, hospitalizations,

and deaths was at the county-level, which dictated the geographic resolution of the SARS-CoV-2

infection estimates. However, as the geographic size and population density of counties differs

systematically across the US (larger and less dense in the West than in the East), a county-level

analysis could lead to bias in estimates of the wave speed and expansion.

To improve our ability to detect spatial patterns in the data, we first created a spatially

regularized grid of infections per capita across the contiguous US. The grid is comprised of 7,665

unique hexagons, each enclosing 64.75 square kilometers (25 square miles). We employed an

area-weighted approach to distribute population estimates from CBGs to hexagons (**Figure 1A-**

C), assuming a constant population throughout the analysis period. We then distributed estimated

infections from counties to hexagons (12) based on the fraction of each county's population

contained in each hexagon and assuming that per capita infection rates were distributed equally

within a county. We combined these to produce daily estimates of SARS-CoV-2 infections per

capita for each hexagon (**Figure 1D-E**).

Assessment of infections per capita surfaces

We fit a modified version of the BYM model as implemented in R-INLA (10), called BYM2, to

estimate spatially smoothed rates of SARS-CoV-2 infections per capita across the hexagonal

grid. The BYM2 is a reparametrized version of the original Besag (11) model, which improves

the assignment of prior distributions for the corresponding model parameters and the subsequent

interpretation of parameters (10). Specifically, our model is given as

$$Z(A_i) = \mu + \theta(A_i) + \epsilon(A_i), \qquad (1)$$

where $Z(A_i)$ is the infections per capita in hexagon A_i, μ is the global intercept, $\theta(A_i)$ are

random effects that are assigned the BYM2 prior distribution, and $\epsilon(A_i)$ are residual error terms

assumed to be statistically independent. For more details on the model and on the priors used see

the **Supplementary Material.** We fit this model separately for each day over March 2020 to

December 2021. The collection of all the $\mu + \theta(A_i)$ (i.e., the global intercept plus the denoised

and spatially smoothed random effects) are the modeled infections per capita, or surfaces. We

then produced a daily sequence of these surfaces, which we used to calculate the speed of wave

expansion and to visualize changes over time. We opted for a separate model fit on each day to

avoid temporal oversmoothing effects, as the infection estimates were derived from a nowcasting

model that already applied temporal smoothing.

Definitions of wave and speed of expansion

We defined waves as one or more sets of contiguous hexagons with relatively high infections-

per-capita levels. To allow for multiple infection centers with the same wave, we did not limit

our definition to a single contiguous set. Using this framework, we categorized 'Wave 1' as the

period from September 2020 to February 2021 and 'Wave 2' as the period from July 2021 to

November 2021. To formally determine hexagons within a wave, we defined a level of per-capita

SARS-CoV-2 infection, above which we coded hexagons as being part of the infection wave. For

our main analysis we used a threshold of 165 daily infections per-capita to characterize a

hexagon as being within a wave. This value represents the 75th percentile of the infections-per-

capita value distribution during the study period (**Figure S1**). We conducted sensitivity analyses

using 85 and 300 infections per capita as alternative threshold values (**Supplementary Material**

Figure S2). We calculated the speed of wave expansion as the increase in area covered by the

wave each day (number of hexes recruited into a wave each day).

Statistical analysis

All analyses were performed using R Statistical Software (v4.3.0) (13)). We also used the **INLA**,

sf, **sp**, **spdep**, **areal**, **rgeoda**, and **magick** R packages (14–18).

Results

Spatiotemporal patterns of SARS-CoV-2 infections across the United States

As described in the **Material and Methods**, we categorized infections as belonging to Wave 1

(September 2020 to February 2021) or Wave 2 (July 2021 to December 2021) (**Figure 2A**).

Figure 2B-I shows the smoothed estimates of SARS-CoV-2 infections per capita from the

BYM2 model on eight dates, leading up to the peak of each of the two waves analyzed. Our

model estimated that Wave 1 originated with a set of hexes spanning central South Dakota,

eastern North Dakota, and northeastern Montana (**Figure 2B**) in mid-September of 2020. Then

the wave extended south and to both coasts from this point of origin, achieving a peak of over

1.5 million infections/day by mid-November 2020 (**Figure 2C-E**). Wave 2 originated in the

Ozarks (southern Missouri and northern Arkansas) (**Figure 2F**) in early July 2021, and then

expanded further south (**Figure 2G-I**). Secondary centers of infection appeared later in July in

the Pacific Northwest (**Figure 2G**), and the wave of infections subsequently spread throughout

the western United States (**Figure 2G-I**). This wave had a peak of 300 per capita infection/day

by early September 2021. Animations of the infection waves are provided in **Movie 1,** which

presents maps of the spatially smoothed infections across the full study period at a weekly

timestep.

Wave expansion and speed of expansion

Figure 3B-C shows contour plots of the speed of expansion for each wave. At its peak, each

wave involved the total contiguous United States (8,595,756 km^2), but the two waves differed in

the time needed to spread across the nation. Wave 1 expanded from an area of 327,780 km^2 to the

total United States in 64 days (September 8th–November 11th, 2020) (**Figure 3B**). Wave 2

expanded from an area of 246,047 km^2 to the total United States in 51 days (July 7th – August

27th, 2021) (**Figure 3C**).

Wave 1 reached its maximal speed of expansion (~ 580,000 km^2/day) 49 days after its formation

and 14 days prior to its peak. Wave 2 reached its maximal speed of expansion (~ 650,000

km^2/day) more quickly, in only 38 days after its formation and 25 days prior to its peak (**Figure**

3C). Sensitivity analyses using different thresholds for categorizing a hexagon as within a wave

resulted in consistent estimates for speed of wave expansion and time to reach peak expansion

speed (**Figure S2**). The shape and speed of wave patterns were also robust to the choice of

threshold (**Figure S3** and **Figure S4**). See **Supplementary Material Figure S5** displaying the

countour plot of the timing of each wave speed of expansion.

Discussion

We sought to quantify the expansion of SARS-CoV-2 infections across the United States during

the first two large waves of the epidemic (**Figures 2**, **Movie1**, and **Movie2**). The two waves had

unique origin sites within the United States. We estimated that Wave 1 originated in parts of

northeastern Montana, eastern North Dakota, and central South Dakota in September 2020

(**Figures 2B-E** and **Movie 2**). This estimated origin coincided with a motorcycle rally that

brought over 460,000 individuals to Sturgis, South Dakota in August 2020 and was later

epidemiologically implicated in elevated local COVID-19 rates and interstate spread of infection

(19,20). Wave 2 originated in the Ozarks (**Figures 2F-I** and **Movie 2**), during a time when there

was local re-opening of indoor music venues and relaxation of local masking policies (21,22).

Shortly following spread in the Ozarks, secondary centers of elevated SARS-CoV-2 infections

per capita were found in the Pacific Northwest. Previous work has implicated both domestic and

international travel in contributing to SARS-CoV-2 transmission across United States and likely

explains the rapid appearance of multiple wave centers for the second examined wave of the

epidemic (23).

While we found that Wave 2 had both a higher maximal speed of expansion and achieved this

peak speed more quickly than Wave 1, the overall similarity in our estimates of the speeds of

wave expansion for these first two waves is notable. This similarity in wave speed expansion is

more remarkable in light of the very different viral variants in circulation (Wave 1 was initially

due to wildtype SARS-CoV-2 and later the Alpha variant (24,25), while Wave 2 was due to the

spread of the Delta variant (23)), regional and secular changes in nonpharmaceutical intervention

policies, and the stark differences in the levels and types of infection- and vaccine-induced

population immunity at the time of these two waves (26). Such stability in the speed of wave

expansion suggests that underlying patterns of human mobility may have had a stronger impact

on the patterns of local epidemic expansion than viral variation, non-pharmaceutical

interventions, and host-immunity (7,20).

Our effort to provide quantitative estimates of speeds and locations of epidemic wave expansion

required several simplifying assumptions. Importantly, we elected to use estimates of infections

from a nowcasting model (rather than case notification data) as diagnosis and case reporting

varied markedly in quality and completeness over the epidemic (27). To overcome potential bias

associated with the irregular shapes and sizes of US counties (the smallest unit at which

estimates were possible), we distributed estimated infections on a hexagonal grid before

calculating the speed of wave expansion. This could have introduced bias if cases within counties

were not randomly distributed, though this effect would be modest. Finally, we arbitrarily

selected the threshold of infections per capita to define wave membership. However, our

sensitivity analyses found that the wave speeds were similar at alternative thresholds.

Factors that affect the magnitude and speed of epidemic waves is an area of substantial interest

for other viral diseases like influenza (e.g. Viboud et al. 2006 (28), Eggo et al. 2011 (29), Gog et

al. 2014 (30)). While our work on SARS-CoV-2 does not itself provide explanations for why

each of these first two epidemic waves spread in the manner that they did once they were

established, the development of methods to quantify the speed and extent of spread, such as

those we employ here, are a necessary first step. Further research to explore the manner in which

the pathogen, environment, and host characteristics affect the speeds and patterns of epidemic

expansion may help to predict wave expansion, and could provide valuable information for the

planning of spatially defined interventions.

Funding

This project is supported by Cooperative Agreement NU38OT000297 from the Centers for Disease

Control and Prevention (CDC) and the Council of State and Territorial Epidemiologists (CSTE),

SHEPheRD Contract 200-2016-91779 from the CDC, and the CDC Broad Agency Announcement

Contract 75D30122C14697. This work does not necessarily represent the views of the CDC or

CSTE.

Data availability:

Daily county estimates of per capita infections used in the analysis are available at Harvard

Dataverse (31). The code and the hexagonally distributed infections per capita estimates for this

analysis are available at: https://github.com/covidestim/waves.

Author contribution – CRediT Taxonomy

Conceptualization: RL, NAS, TC

Methodology: RL, NAS, JLW, TC

Investigation: RL, NAS

Visualization: RL, NAS, YL

Funding acquisition: NAM, JAS, NDG, TC

Supervision: JAS, NAM, JLW, NDG, TC

Writing – original draft: RL, NAS

Writing – review & editing: all authors

Competing interests

NDG is a paid consultant for BioNTech

Population (log10 scale)

Cumulative Infections (log10 scale)
(March 2020 – December 2021)

Figure 1: United States population and estimated cumulative SARS-CoV-2 infections per

capita distributed across the hexagonal grid. Panel **A, B,** and **C**: United States', New

England's and Connecticut's 2019 population Census estimates on the hexagonal grid. Panel **D,**

E, and **F**: United States', New England's, and Connecticut cumulative infections per 100,000

persons on the hexagonal grid (March 2020–December 2021).

Note: Numbers are given in a log 10 scale. Hexagons with no filling had no population reported

or infections counts ever estimated.

Figure 2: Estimated infections per capita of SARS-CoV-2 in the United States, March 2020–December 2021. Panel **A**: Time series of SARS-CoV-2 infection estimates for the United States, the gray shaded areas show the first two large waves of infections. Panels **B, C, D**, and **E**: Sequence of the spatially smoothed estimates of SARS-CoV-2 infections per capita associated with Wave 1 at 4 time points. Panels **F, G, H**, and **I**: Sequence of the spatially smoothed estimates of SARS-CoV-2 infections per capita associated with Wave 2 at 4 time points.

264 **Figure 3: SARS-CoV-2 infections-per-capita surfaces progression and speed of invasion for each wave.** Comparison of SARS-

265 CoV-2 infection Wave 1 and Wave 2 speed for 56- to 7- days prior to each wave's infection peak. **See Supplementary Material**

266 **Figure S5 for a countor plot of waves progression.**

References

1. List of Public Health Emergency Declarations [Internet]. [cited 2024 Dec 3]. Available from: https://aspr.hhs.gov:443/legal/PHE/Pages/default.aspx

2. Determination that a Public Health Emergency Exists [Internet]. [cited 2024 Dec 3]. Available from: https://aspr.hhs.gov:443/legal/PHE/Pages/2019-nCoV.aspx

3. Chitwood MH, Russi M, Gunasekera K, Havumaki J, Klaassen F, Pitzer VE, et al. Reconstructing the course of the COVID-19 epidemic over 2020 for US states and counties: Results of a Bayesian evidence synthesis model. PLOS Comput Biol [Internet]. 2022 Aug 30 [cited 2023 Oct 6];18(8):e1010465. Available from: https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1010465

4. Gamio L, Smith M, Yourish K, Almukhtar S. Watch How the Coronavirus Spread Across the United States. The New York Times [Internet]. 2020 Mar 21 [cited 2023 Dec 19]; Available from: https://www.nytimes.com/interactive/2020/03/21/us/coronavirus-us-cases-spread.html

5. Kissler SM, Kishore N, Prabhu M, Goffman D, Beilin Y, Landau R, et al. Reductions in commuting mobility correlate with geographic differences in SARS-CoV-2 prevalence in New York City. Nat Commun [Internet]. 2020 Sep 16 [cited 2024 Oct 17];11(1):4674. Available from: https://www.nature.com/articles/s41467-020-18271-5

6. Gatto M, Bertuzzo E, Mari L, Miccoli S, Carraro L, Casagrandi R, et al. Spread and dynamics of the COVID-19 epidemic in Italy: Effects of emergency containment measures. Proc Natl Acad Sci [Internet]. 2020 May 12 [cited 2024 Oct 14];117(19):10484–91. Available from: https://www.pnas.org/doi/10.1073/pnas.2004978117

7. Kraemer MUG, Yang CH, Gutierrez B, Wu CH, Klein B, Pigott DM, et al. The effect of human mobility and control measures on the COVID-19 epidemic in China. Science [Internet]. 2020 May [cited 2024 Oct 2];368(6490):493–7. Available from: https://www.science.org/doi/full/10.1126/science.abb4218

8. Pepe E, Bajardi P, Gauvin L, Privitera F, Lake B, Cattuto C, et al. COVID-19 outbreak response, a dataset to assess mobility changes in Italy following national lockdown. Sci Data [Internet]. 2020 Jul 8 [cited 2024 Oct 2];7(1):230. Available from: https://www.nature.com/articles/s41597-020-00575-2

9. Chinazzi M, Davis JT, Ajelli M, Gioannini C, Litvinova M, Merler S, et al. The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. Science [Internet]. 2020 Apr 24 [cited 2024 Oct 2];368(6489):395–400. Available from: https://www.science.org/doi/full/10.1126/science.aba9757

10. Riebler A, Sørbye SH, Simpson D, Rue H. An intuitive Bayesian spatial model for disease mapping that accounts for scaling. Stat Methods Med Res [Internet]. 2016 Aug 1 [cited 2024 Oct 2];25(4):1145–65. Available from: https://doi.org/10.1177/0962280216660421

11. Besag J, York J, Mollí A. Bayesian image restoration, with two applications in spatial statistics. Ann Inst Stat Math [Internet]. 1991 Mar [cited 2024 Oct 2];43(1):1–20. Available from: http://link.springer.com/10.1007/BF00116466

12. topojson/us-atlas [Internet]. TopoJSON; 2024 [cited 2024 Aug 7]. Available from: https://github.com/topojson/us-atlas

13. R: The R Project for Statistical Computing [Internet]. [cited 2024 Dec 3]. Available from: https://www.r-project.org/

14. Prener C, Revord C. areal: An R package for areal weighted interpolation [Internet]. Zenodo; 2020 [cited 2024 Aug 7]. Available from: https://zenodo.org/records/3822534

15. Pebesma E. Simple Features for R: Standardized Support for Spatial Vector Data. R J [Internet]. 2018 [cited 2024 Oct 5];10(1):439. Available from: https://journal.r-project.org/archive/2018/RJ-2018-009/index.html

16. magick: Advanced Graphics and Image-Processing in R [Internet]. [cited 2024 Oct 5]. Available from: https://ropensci.r-universe.dev/magick

17. Bivand R. spdep: Spatial Dependence: Weighting Schemes, Statistics [Internet]. 2002 [cited 2024 Oct 7]. p. 1.3-6. Available from: https://CRAN.R-project.org/package=spdep

18. Lindgren F, Rue H. Bayesian Spatial Modelling with R-INLA. J Stat Softw [Internet]. 2015 Feb 16 [cited 2024 Oct 14];63:1–25. Available from: https://doi.org/10.18637/jss.v063.i19

19. Firestone MJ. COVID-19 Outbreak Associated with a 10-Day Motorcycle Rally in a Neighboring State — Minnesota, August–September 2020. MMWR Morb Mortal Wkly Rep [Internet]. 2020 [cited 2024 May 22];69. Available from: https://www.cdc.gov/mmwr/volumes/69/wr/mm6947e1.htm

20. Dave D, McNichols D, Sabia JJ. The contagion externality of a superspreading event: The Sturgis Motorcycle Rally and COVID-19. South Econ J. 2021 Jan;87(3):769–807.

21. Barone E. TIME. 2021 [cited 2024 May 28]. How the Delta Variant Overtook Missouri: A Lesson for the Rest of the U.S. Available from: https://time.com/6085454/delta-variant/

22. Daly M. How a Tiny Town With an Anti-Mask Mayor Caused COVID Chaos. The Daily Beast [Internet]. 2021 Jul 23 [cited 2024 Oct 14]; Available from: https://www.thedailybeast.com/missouris-covid-surge-started-in-tourist-town-of-branson

23. Bolze A, Luo S, White S, Cirulli ET, Wyman D, Dei Rossi A, et al. SARS-CoV-2 variant Delta rapidly displaced variant Alpha in the United States and led to higher viral loads. Cell Rep Med [Internet]. 2022 Mar 15 [cited 2024 Mar 26];3(3):100564. Available from: https://www.sciencedirect.com/science/article/pii/S2666379122000714

24.   Alpert T, Brito AF, Lasek-Nesselquist E, Rothman J, Valesano AL, MacKay MJ, et al. Early introductions and transmission of SARS-CoV-2 variant B.1.1.7 in the United States. Cell [Internet]. 2021 May [cited 2023 May 11];184(10):2595-2604.e13. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0092867421004347

25.   Washington NL, Gangavarapu K, Zeller M, Bolze A, Cirulli ET, Barrett KMS, et al. Emergence and rapid transmission of SARS-CoV-2 B.1.1.7 in the United States. Cell [Internet]. 2021 May 13 [cited 2024 Nov 7];184(10):2587-2594.e7. Available from: https://www.cell.com/cell/abstract/S0092-8674(21)00383-4

26.   Klaassen F, Chitwood MH, Cohen T, Pitzer VE, Russi M, Swartwood NA, et al. Changes in Population Immunity Against Infection and Severe Disease From Severe Acute Respiratory Syndrome Coronavirus 2 Omicron Variants in the United States Between December 2021 and November 2022. Clin Infect Dis [Internet]. 2023 Aug 1 [cited 2023 Oct 16];77(3):355–61. Available from: https://doi.org/10.1093/cid/ciad210

27.   Pitzer VE, Chitwood M, Havumaki J, Menzies NA, Perniciaro S, Warren JL, et al. The Impact of Changes in Diagnostic Testing Practices on Estimates of COVID-19 Transmission in the United States. Am J Epidemiol [Internet]. 2021 Sep 1 [cited 2023 Oct 26];190(9):1908–17. Available from: https://doi.org/10.1093/aje/kwab089

28.   Viboud C, Bjørnstad ON, Smith DL, Simonsen L, Miller MA, Grenfell BT. Synchrony, Waves, and Spatial Hierarchies in the Spread of Influenza. Science [Internet]. 2006 Apr 21 [cited 2024 Feb 29];312(5772):447–51. Available from: https://www.science.org/doi/10.1126/science.1125237

29.   Eggo RM, Cauchemez S, Ferguson NM. Spatial dynamics of the 1918 influenza pandemic in England, Wales and the United States. J R Soc Interface [Internet]. 2011 Feb 6 [cited 2024 Dec 12];8(55):233–43. Available from: https://royalsocietypublishing.org/doi/10.1098/rsif.2010.0216

30.   Gog JR, Ballesteros S, Viboud C, Simonsen L, Bjornstad ON, Shaman J, et al. Spatial Transmission of 2009 Pandemic Influenza in the US. PLOS Comput Biol [Internet]. 2014 Jun 12 [cited 2024 Dec 12];10(6):e1003635. Available from: https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1003635

31.   Klaassen F. covidestim updated: Estimating SARS-CoV-2 infections and immunity over the entire pandemic [Internet]. Harvard Dataverse; 2024 [cited 2024 Oct 2]. Available from: https://dataverse.harvard.edu/dataset.xhtml?persistentId=doi:10.7910/DVN/G2ZXJG

 Supporting Information

372 **Figure S1 – Histogram of values for the risk surfaces and empirical cumulative density**
373 **function of the risk surface values.** The histogram show that the distribution of values is
374 concentrated around 0, which as expected due to for long periods we had low to no infections
375 occurring to a great number of places. The ECDF shows that a threshold of 165 infections per
376 capita will capture 75% of the mass of the values distribution.
377
378 **Figure S2 – Speed of invasion for different thresholds to the infection per capita surfaces**
379 **progression calculation.** Panel **A** is built with a threshold of 85 or more infections per capita,
380 panel **B** is built with a threshold of 300 infections per capita. As in **Figure 3C**, we observe a
381 maximal speed and a steep decrease after peak, and, as in **Figure 3C**, the second wave had a
382 higher speed of invasion and encompassed a larger area at peak than the first wave.
383
384 **Figure S3 – Infection per capita surface on a continuous scale of values.** As expected, the
385 wave-like pattern holds independently of the scale to be displayed, and as being an output of
386 spatial smooth model, the continuous scale gives a less defined border to the risk surface
387 expansion.
388
389 **Figure S4 - Infection per capita with a threshold equal to the mean of the risk values**
390 **distribution (85 infections per capita).** With a lower threshold showing on the map, the spread
391 process seems to happen faster.
392
393 **Figure S5 – Contour plot of the dates for the speed of expansion.** The contour plots shows the
394 contour surface at dates with 7 days spacing between each, up until the national curve peak.