Fusing Data from CT Deep Learning, CT Radiomics and Peripheral Blood Immune profiles to Diagnose Lung Cancer in a Cohort of Patients Experiencing Symptoms

Rami Mustapha[1]*, Balaji Ganeshan[2]*, Sam Ellis[3]*, Luigi Dolcetti[1]*, Mukunthan Tharmakulasingam[1], Karen DeSouza[1], Xiaolan Jiang[1], Courtney Savage[1], Sheena Lim[4], Emily Chan[3], Andrew Thornton[2], Luke Hoy[2], Raymond Endozo[2], Rob Shortman[2], Darren Walls[2], Shih-hsin Chen[2], Mark Rowley[5], Anthony CC Coolen[6,7], Ashley M Groves[2], Julia A Schnabel[1,3], Thida Win[4]^, Paul R Barber[1]^\$~, Tony Ng[1]^\$

Author affiliations:

[1] Comprehensive Cancer Centre, King's College London, London UK
[2] University College London, London UK
[3] School of Biomedical Engineering and Imaging Sciences, King's College London, London, UK
[4] East and North Hertfordshire NHS Trust, UK
[5] Saddle Point Science Ltd, York, UK
[6] Saddle Point Science Europe BV, 6525EC Nijmegen, The Netherlands
[7] Radboud University, 6525AJ Nijmegen, The Netherlands

* = Joint first author
^ = Joint last author
\$ = Joint senior author
~ = Corresponding Author

Corresponding Author:

    Dr Paul R Barber
    Comprehensive Cancer Centre
    New Hunt's House
    King's College London
    SE1 9RT
    paul.barber@kcl.ac.uk

1   Summary

Background: Lung cancer is the leading cause of cancer-related deaths. Diagnosis
at late stages is common due to the largely non-specific nature of presenting
symptoms contributing to high mortality. There is a lack of specific, minimally
invasive low-cost tests to screen patients ahead of the diagnostic biopsy.

Methods: 344 patients experiencing symptoms from the lung clinic of Lister hospital
suspected of lung cancer were recruited. Predictive covariates were successfully
generated on 170 patients from Computed Tomography (CT) scans using CT Texture
Analysis (CTTA) and Deep Learning Autoencoders (DLA) as well as from peripheral
blood data for immunity using high depth flow-cytometry and for exosome protein
components. Predictive signatures were formed by combining covariates using
Bayesian regression on a randomly chosen 128-patient training set and validated on
a 42-patient held-out set. Final signatures were generated by fusing the data sources
at different levels.

Findings: Immune, CTTA and DLA single modality signatures had overall AUCs of
0.69, 0.70 and 0.73 respectively. The final combined signature had a ROC AUC of
0.81. The overall sensitivity and specificity were 0.72 and 0.77 respectively.

Interpretation: Combining immune monitoring with CT scan data is an effective
approach to improving sensitivity and specificity of Lung cancer screening even in
patients experiencing symptoms.

Funding: CRUK [C1519/A27375], Wellcome Trust/EPSRC Centre for Medical
Engineering [WT203148/Z/16/Z], NIHR Clinical Research Facility at Guy's and St
Thomas' NHS Foundation Trust, NIHR Biomedical Research Centre.

Keywords: Lung cancer; Early Diagnosis; CT Scan; Blood test; Immune

1 Research in Context

2 Evidence before this study

Lung cancer is the leading cause of cancer related deaths and previous studies have
shown that early diagnosis is often difficult and one third of patients return to their GP
three or more times before a referral to a specialist according to a study in the British
Journal of Cancer in 2013. Screening is a possible solution and low dose CT scans
for patients with high-risk has been implemented, but this excludes non-smokers and
the young and can be inaccurate according to studies in 2022 and 2023. In addition,
the workload of screening on radiologists is resulting in delays in examinations as
recently reported by the Royal College of Radiologists in 2024. Some studies have
suggested the benefit of a blood test, to augment a CT scan, based on DNA
fragments or circulating tumour cells, for example (2021-2023).

A PubMed publications database search (search term: "blood"[Title] AND "CT"[Title]
AND "Lung"[Title] AND "cancer"[Title]) revealed 17 results on 8[th] May 2025, of which
2 were directly relevant involving combining a PET/CT scan with inflammatory blood
measurements, rejecting those where blood referred to blood volume or perfusion
imaging or ctDNA.

18 Added value of this study

We propose a blood test based on the immune response to the presence of cancer
(in contrast to other studies which are based on detecting nucleic acids) that is
combined with the automated analysis of CT scans (reducing radiologist workload)
by machine learning, which may provide a route to screening patients with symptoms
of lung diseases. The advantage of a blood test based on the immune response is
that immune cell-based detection of cancers can occur <u>when the tumours are</u>
<u>relatively small and at an earlier time point</u> than the shedding of nucleic acids. We
recruited 344 patients between October 2020 and November 2021 into the
LungExoDETECT study, with a blood test for immune profiling and routine CT scan.
After following their progress to determine their cancer diagnosis, we were able to
make a mathematical model that predicts diagnosis from 12 measurements based
on the blood and CT scan, and which was validated in a patient subset.

31 Implications of all the available evidence

Our study has important implications for the field: (1) We reinforce previous our
previous evidence that the response of the immune system to cancer can add
additional information for detecting cancer to that provided by the automated
analysis of CT scans. (2) Advanced Bayesian machine learning can produce simple
mathematical models that predict the presence of cancer and are highly interpretable
in terms of why a particular prediction has been made by the model by pinpointing
the particular component of the immune system that contributes to the prediction.

1 Introduction

2 Lung cancer and importance of early detection

Lung cancer is by far the leading cause of cancer-related deaths worldwide with an
estimated 1.8 million death in 2020 [1]. In the U.S.A., lung cancer accounts for about
two and a half times the number of cancer deaths than the second leading cause,
colorectal cancer. This is despite the fact that lung cancer only contributes to about
12% of newly diagnosed cancer cases, which is far less than prostate cancer and
breast cancer, the leading cancer diagnoses for men and women, respectively [2]. This
high mortality of lung cancer has been linked to diagnosis at late stages. 48% of lung
cancer cases are diagnosed with a distant metastasis and an expected 5-year survival
rate of 8% compared to the ≥60% rate when diagnosis happens at a localized stage
[3]. In England, 5-year survival for a stage I diagnosis is 63%, which drops down to
4.3% for a stage IV diagnosis [4]. Delayed diagnosis can be attributed to the largely
asymptomatic nature of early-stage lung cancer [5]. Other data have shown that patients
delaying their first general practitioner (GP) visit after manifestation of early symptoms
is another obstacle to a timely diagnosis [6]. Beyond that, a study from 2013 showed
that a third of patients with lung cancer have 3 or more visits with their GP with lung
cancer associated symptoms before special referral, which forms a sharp contrast to
the 3% observed in patients with breast cancer[7]. Similarly, the UK's National Cancer
Diagnosis Audit reported primary care delays of 60 and 90 days experienced by 17.9%
and 10.8% of patients with suspected lung cancer, respectively [8]. This highlights the
need for lung cancer screening, which to a certain extent has been implemented in the
form of low dose computed tomography (LDCT) for patients of high-risk. Despite this
practice leading to about a 20% reduction in mortality across multiple clinical trials,
there is still room for improvement [9, 10]. LDCT screening excludes patients that are
younger and those at lower risk as it has been shown to be not reliable as a screening
tool for the broader population, owing to its high chance of false positives [11, 12]. Lung
cancer screening also heavily relied on smoking history, which excludes the early
detection of lung cancer in non-smokers which represent around 25% of cases [13].
There is a clear need for a minimally invasive robust screening test for lung cancer
with similar sensitivity and specificity to a mammography for breast cancer, or the
faecal occult blood test currently in use for colorectal cancer. Furthermore, this needs
to be extended beyond screening, as numerous healthcare systems, including the
NHS in the UK, have put in place a rapid diagnostic pathway for patients already
symptomatic and suspected of lung cancer [14].

36 Role of radiologist and importance of alternative image analysis

37 The diagnostic pathway for lung cancer is heavily reliant on radiologist assessment at
38 multiple stages. According to the UK NHS National Optimal Lung Cancer Pathway
39 (NOLCP) [15] and National Institute for Health and Care Excellence (NICE) guidelines,
40 the typical pathway may include a chest radiograph (CXR), a CT scan, a PET-CT, a

CT guided biopsy, all of which require a radiologists intervention, bronchoscopy (EBUS-TBNA, EUS-FNA or navigational) and pleural procedures. This reliance is further exacerbated by the large amount of CT imaging data from lung cancer screening programmes. This workload combined with the shortage of radiologists in the UK and other European countries is manifesting in delays in examinations as recently reported by the Royal College of Radiologists [16, 17]. Implementing automated image analysis approaches like radiomics-based texture and deep learning analyses is a viable approach, especially given the availability of large screening datasets for training purposes. The low-dose radiation used in LDCT, to minimize patient exposure, does negatively impact the visibility of small or low-contrast lesions due to increased noise and reduced spatial resolution and artificial intelligence can alleviate this in many areas of CT screening [18,19]. In this study we exploit the latent vector of a deep autoencoder as a low dimensional representation of the lesion image characteristics and evaluate its predictive performance.

Liquid biopsy

Using a liquid biopsy for early detection and diagnosis of lung cancer has been heavily assayed to varying degrees of success [20]. Of note, Galleri® which is a circulating tumoral DNA (ctDNA) methylation-based multicancer early detection (MCED) test is seeing adoption across clinics. Early results seem very promising in the symptomatic setting, though these are not comparable across all cancer types. Of note, sensitivity for stage I lung cancer was 8.7% and 0% in the symptomatic and asymptomatic settings, respectively [21, 22]. Combining LDCT with minimally invasive blood test is one approach that can improve the shortcoming of either approach. This has proven useful in the BioMilk study that showed that double positivity LDCT and microRNA signature doubles the sensitivity of LDCT alone [23]. Similarly, other studies combined circulating proteins with LDCT achieving improved specificity [24]. However, there is still room for improvement using other blood biomarkers which include circulating tumour cells, microRNA, exosomes, tumour educated platelets, metabolites, tumour associated proteins, and autoantibodies. In addition, peripheral cell-based flow cytometry immune markers [25-27] (especially the immune checkpoint receptors) can potentially provide treatment targets for the cancers detected.

Immunology in lung cancer

Little research exists on the feasibility of immune profiling of circulating immune cells as a biomarker for early lung cancer. Peripheral blood mononuclear cells (PBMCs) are a heterogenous population of immune cells usually classified into subsets. The numbers, relative frequencies and functional status these subsets provide information on the state of the immune response and there have been recent advents in technology beyond classical multiparametric flow cytometry. Using patient PBMCs, single cell RNA sequencing was used to identify circulating anti-tumour CD8 cells, and Cytometry by time of flight (CyTOF) was used to correlate the frequency of HLADR expressing monocytes with response to ICB therapy [28, 29]. Numerous research groups,

including ours, have shown that changes at the tissue level of a solid cancer immune response are reflected as systemic changes in the immune response that can be detected at the peripheral level [28-30]. Lung cancer promotion after initial mutagenesis has been shown to be inflammation driven, with a recent link established to air pollution [31], further highlighting the potential of an immune based biomarker for early detection. Cancer specific T cell receptor repertoire has been shown to be effective at identifying early-stage lung cancer in an LDCT screened cohort. This proves that cancer specific immune cells can be reflected in the periphery [32].

Methods

Study design and data collection

The study entitled "Improving the Early Detection of Lung Cancer by Combining Exosomal Analysis of Hypoxia with Standard of Care Imaging (LungExoDETECT)" (https://clinicaltrials.gov/ct2/show/NCT04629079) is a prospective cohort study of patients referred to secondary care for the investigation of clinical symptoms or signs suspicious of lung cancer. The study analysis will determine whether the assay can detect clinical lung cancer at the time of imaging, and interval cancers during subsequent follow up. The study aimed to establish preliminary ROC AUC and sensitivity/specificity data for the "combined CT/blood risk stratification marker" and provide initial data on the potential association of the "combined CT/blood risk score" with the subsequent cancer progression and treatment response. The gold standard lung cancer diagnosis was determined by the National Optimal Lung Cancer Pathway (NOLCP).

The study included patients who have been referred to the Lung Cancer Clinic and Multi-Disciplinary Team (MDT) at The Lister, Hertford County, UK and New QEII Hospitals, Welwyn Garden City, UK for investigation of suspected lung cancer. Participants were recruited between October 2020 and November 2021 (344 patients). As part of the standard of care pathway they received a CT scan, at that time peripheral blood samples were collected. 174 cases were excluded, 162 due to lack of visible lesion or lesion <6mm and 12 cases with non-lung cancer. Sex/gender was self-reported by study participants and is reported but not included in multivariate analysis. The patient pathway and study design is illustrated in Figure 1 and patient characteristics are in Table 1.

Figure 1. Showing the patient pathway and study design. 344 patients referred to the clinic were recruited to the study. As part the standard of care pathway they received a CT scan, at which time we collected the peripheral blood samples. 174 cases were excluded from the analysis set due to lack of visible lesion, the lesion was smaller than 6mm or the diagnosis was a non-lung cancer diagnosis. The analysis was performed on the remaining patients.

Patient Characteristic	Non-cancer $n = 91$	Lung cancer $n = 79$	p-value*
Age			
Mean	71.9	74.9	0.152
Range	27.0 to 94.0	33.0 to 97.0	
Sex			
Male	43	38	0.646
Female	48	41	
Smoking status			
Ex-Smoker	33	31	
Non-Smoker	29	12	0.07
Smoker	17	21	
Unknown	12	15	
Cohort			
Training	64	64	0.1
Test	27	15	
Staging			
No Cancer	91		
Stage I		23	
Stage II		9	
Stage III		18	
Stage IV		29	

Table 1. Patient characteristics and distribution. The table shows the difference between cancer and non-cancer as well as staging for patients with cancer. *P-values were calculated to compare data from individuals with and without lung cancer for the following variables: mean ages using Student's unpaired two-tailed t-test whereas for sex distribution, smoking status and cohort using a χ2 test.

Sample and Data Processing

The details of CT scan acquisition, CT Texture Analysis (CTTA) (Supplementary Figure 1, Supplementary Figure 2), the Deep Learning Autoencoder (DLA), peripheral blood sampling and processing, flow cytometry and model (aka signature) generation are given in the Supplementary Methods. The total number of features generated by each modality are: 42 from CTTA, 32 from DLA, 45 from flow cytometry and 68 from exosome dot-blots (Supplementary Table S1, Supplementary Table S2, Supplementary Table S3).

Model Building and Performance Analysis

Separately, using the data from each analysis, a risk signatures (models) were generated using Bayesian multi-variate regression (BMR) which also performed covariate selection (supplementary material). Although 170 is a reasonable number of samples it is not significantly larger than the combined number of covariates. It is therefore reasonable to use all 170 for model building and through bootstrapping

training and test sets, gain indicators of performance. Recent literature indicates that
these BMR approaches will not overfit and therefore will generalise under the
assumption that the training set is representative[33, 34].

Of all the study samples, one quarter was randomly held out of the model training and
served as an internal test set. This was repeated 8 times to get a good estimate of
future performance independent of the training/test splitting. The performance of both
univariate biomarkers and model predicted outcomes were judged by the area under
the receiver operator characteristic curve (ROC AUC) with the cancer diagnosis result
from the clinic as the gold standard using the R 'pROC' package and the 'ci.auc'
function. Sensitivity and specificity were calculated at the Youden point of the ROC. A
Risk Score can be generated by combining covariate raw values in a linear model
according to their given weights and intercept. Final signatures were generated from
covariates that were significant in at least 50% of the 8 repeated training runs, and
assed on the full cohort. This offers credence to the robustness of the selected
features.

Statistics

For comparing groups, a Student's unpaired two-tailed t-test or a χ2 test was used for continuous or
discrete variables, as these are well used and understood tests for this purpose. Hazard Ratios, P-
values and confidence intervals from multivariate analyses were calculated from posterior parameter
probability distributions. In the Bayesian framework these are calculated using exact expansion
methods, that (whenever required by the data) go beyond quadratic (i.e. Fisher matrix based)
approximations, and are easily interpretable in line with other tests.

Ethics

24 All procedures were performed in compliance with relevant laws and institutional
25 guidelines and have been approved by the appropriate institutional committee(s), and
26 informed consent obtained from the subjects (REC reference: 19/EE/0357 20th Feb
27 2020).

Role of Funders

29 The funders of this study had no role or influence in study design, data collection, data
30 analysis, interpretation nor the writing of this manuscript.

Results

Patient Characteristics

33 The characteristics of the patients suspected of lung cancer and included in the final
34 analysis set are shown in Table 1. Of those suspected of having lung cancer 46.5%
35 (79 patients) were diagnosed with lung cancer over the follow up period. There were
36 no significant differences between the patients with and without cancer with regards
37 to sex, age, smoking status nor assignment to training or test set. The division of

patients between training (3/4) and test sets (1/4) was retrospective and at random, with the condition of having a balance of patients with and without cancer representative of the whole cohort (Supplementary material). Amongst the patients with lung cancer, the division across stage is shown in Table 1. Of note, 23 of the lung cancers (29.1%) were stage I, signifying that this is a good cohort for the potential development of a screening signature. 12 patients of the total recruited were diagnosed with a cancer that is not lung cancer (characteristics in Supplementary Table S4).

Generation of a Predictive Model from CTTA and DLA

The imaging data were analysed using CTTA and DLA, as explained in the methodology. The signatures are presented in Figure 2A. The DLA signature performed better than the texture analysis signature on both the training and test cohorts. On the training sets DLA had a range of AUC of 0.72-0.77 (mean ± 1 standard deviation), whereas CTTA had an AUC range of 0.67 to 0.73. On the test sets DLA had an AUC range 0.56 to 0.71, whereas CTTA had an AUC range of 0.57-0.79 (Figure 2A). Figure 2A also represents the result of combining CTTA and DLA analyses.

The CTTA features consistently selected by BMR across the repeated training splits were the mean intensity at spatial scale filter (SSF) = 0 (Mean_0) – which relates to the CT attenuation / density of the tumour, and the mean intensity of positive pixels at scale SSF=3 (mpp_3) – which relates to the average brightness at fine texture scale (corresponding to features of 3mm in size) within the tumour.

A preliminary exploration of the interpretation of the DLA latent space features selected by BMR was performed. This involved passing a lesion image into the encoder before manually varying the value of predictive features. By varying the value of a feature in this way, and passing the resulting latent space codes through the decoder, the impact of each feature in image space could be viewed. The 5 predictive DLA features selected by BMR (numbered 13 and 16) were observed to relate to lesion size, morphology (e.g. spiculation and sphericity), location within the lung (evidenced by changing distance and orientation of pleura), and degree of pleural attachment (Supplementary Figure 3). As quantitative evaluation of these interpretations would require large annotated datasets, including novel quantitative measures for characteristics such as pleural attachment, we leave this for future work.

B CTTA

Covariate	Beta_mean	Beta_sd	pval_mean	Count	Selected	Beta	pval	Weight
mean_0	0.605	0.125	<0.001	7	Yes	0.644	<0.001	0.026
mpp_3	0.525	0.138	<0.001	5	Yes	-0.521	<0.001	-0.026
kurtosis_4	0.424	0.13	0.001	1				
mean_3	0.408	0.107	<0.001	1				
skewness_0	0.444	0.114	<0.001	1				
skewness_4	0.356	0.119	0.003	1				
Constant								0.591

ROC AUC = 0.70

C DLA

Covariate	Beta_mean	Beta_sd	pval_mean	Count	Selected	Beta	pval	Weight
GAN32_13	0.394	0.106	<0.001	5	Yes	-0.349	<0.001	-0.396
GAN32_16	0.612	0.114	<0.001	4	Yes	0.521	<0.001	0.517
GAN32_10	0.542	0.112	<0.001	1				
GAN32_12	0.344	0.113	0.002	1				
GAN32_18	0.574	0.114	<0.001	3				
GAN32_20	0.405	0.115	0.001	2				
GAN32_22	0.32	0.111	0.004	1				
GAN32_28	0.486	0.117	<0.001	2				
Constant								0.2

ROC AUC = 0.73

2 **Figure 2. Risk generation by BMR from repeated data splits and comparing CT texture analysis**
3 **to deep learning autoencoder.** (A) Box plots showing the distributions (mean and inter-quartile
4 range, points overlaid) of AUCs obtained from 8 random training/test splits of the data. (B, C)
5 Outcome association statistics for CTTA and DLA of the 8 repeats, and the count of the number of
6 times the covariate was indicated as important. Covariates indicated more than 50% of the time were

selected for the final model generation; the final association (Beta), p-value for the association and weight in the final signature are given, as well as a ROC curve and AUC value for the final signature.

Generation of a Predictive Model from Peripheral Blood

The peripheral blood samples were analysed for their immune and their exosome protein content (dot-blot) as indicated in the methods. Separately, using the data from each analysis modality, a risk signature was generated (Figure 3A). The flow cytometry signature performed far better on the training and test cohort than the exosome protein content signature. On the training sets, the immune signature had an AUC range of 0.66-0.70, whereas the exosome derived signature had an AUC range of 0.50-0.60 (Figure 2A). Of note, the flow cytometry signature was driven by two significant features: a high proportion of type 2 dendritic cells indicated a non-cancer pathology, whereas a high proportion KIR3DL1 expressing CD8 T lymphocytes indicated lung cancer (Figure 3A). The exosome dot-blot results were not consistent, with the exception of some consistency of PD1, and was not used for further combined risk signature generation (Figure 3B).

Figure 3. Risk generation by BMR from peripheral blood samples comparing flow cytometry to plasma exosome dotblot analysis. (A, B) Outcome association statistics for Immune and Dotblot of the 8 repeats, and the count of the number of times the covariate was indicated as important. Covariates indicated more than 50% of the time were selected for the final model generation; the final association (Beta), p-value for the association and weight in the final signature are given, as well as a ROC curve and AUC value for the final signature. Associations for the dotblot data were so low that a final model was not generated.

1 Combining CT Analyses with Peripheral Blood Data

We proceeded to combine the data from the blood sample with the CTTA and DLA
individually, at the covariate level (early fusion). Both combined signatures from BMR
featured covariates from the immune and imaging data with the upregulation of
KIR3DL1 CD8 T cells being the strongest predictor of a lung cancer (Figure 4A). The
risk signature generated from the combination of flow cytometry and DLA performed
better on the training set with an AUC range of 0.77-0.83, which transferred well to
the test sets: 0.68-0.81 (Figure 2A). The risk signature generated from the
combination of peripheral immunity and CTTA on the training set had an AUC range
of 0.79-0.84, which also transferred well to the test sets: 0.66-0.82. Final models
were generated from the consistent covariates with AUCs of 0.81 and 0.71 (Figure
4).

A Immune CTTA Early Fusion

Covariate	Beta_mean	Beta_sd	pval_mean	Count	Selected	Beta	pval	Weight
mean_0	0.516	0.117	0.001	5	Yes	0.358	<0.001	0.014
cDC2	0.659	0.158	<0.001	7	Yes	-0.6	<0.001	-0.454
mpp_2	0.527	0.14	<0.001	1				
mpp_3	0.682	0.14	<0.001	1				
total_0	0.331	0.116	0.004	1				
KIR3DL1.CD8.T.lymphocytes	0.314	0.132	0.018	1				
Constant								0.196

ROC AUC = 0.71 (Sensitivity vs 1-Specificity)

B Immune DLA Early Fusion

Covariate	Beta_mean	Beta_sd	pval_mean	Count	Selected	Beta	pval	Weight
GAN32_16	0.579	0.116	<0.001	6	Yes	0.604	<0.001	0.6
GAN32_18	0.575	0.119	<0.001	4	Yes	0.299	0.001	0.299
KIR3DL1.CD8.T.lymphocytes	0.537	0.153	0.002	6	Yes	0.601	<0.001	1.242
cDC2	0.7	0.162	<0.001	8	Yes	-0.696	<0.001	-0.526
GAN32_10	0.411	0.118	0.001	2				
GAN32_11	0.625	0.139	<0.001	1				
GAN32_13	0.62	0.121	<0.001	2				
CD4CD8.double.positive.T.cells	0.378	0.108	<0.001	1				
CD4.Effector.memory	0.358	0.11	0.001	1				
CD4.Effector	0.45	0.128	<0.001	1				
Constant								1.009

ROC AUC = 0.81 (Sensitivity vs 1-Specificity)

14 **Figure 4. Risk generation by BMR from CT image analysis combined with flow cytometry,**
15 **comparing deep learning and texture analysis.** (A, B) Outcome association statistics for Immune-
16 CTTA and Immune-DLA fusion models of the 8 repeats, and the count of the number of times the
17 covariate was indicated as important. Covariates indicated more than 50% of the time were selected

1 for the final model generation; the final association (Beta), p-value for the association and weight in
2 the final signature are given, as well as a ROC curve and AUC value for the final signature.

Generation of a Combined Model by Data Fusion

We proceeded to combine all methods of CT and blood analyses using data fusion at different levels. Combination at the covariate level (early fusion) did not show a major improvement of the risk signature with an AUC range 0.77-0.83. The signature generation process included co-variates from all three modalities, but no CTTA covariates were selected in 50% of runs (Figure 5A).

We also generated predictive signatures by fusing the features included in individual signatures (feature level or intermediate fusion) and by fusing the individual signatures themselves (signature level or late fusion, Figures 5B,5C). These did not show better performance than the early fusion models.

An equivalent approach to data combination and signature generation was performed using an Elastic Net algorithm (glmnet R package) [35, 36] which produced some better signatures on the training sets, but struggled with overfitting which impacted the ROC AUCs on the test set. We are focusing on the results from BMR since they are more consistent and offer less overfitting. A summary of all the Elastic net signatures is in Supplementary Figure 4. Overcoming the problem of overfitting was the main reason for the use of the Bayesian regression, which will optimally choose the number of covariates to be predictive given the evidence in the training set. This is in contrast to the commonly used Elastic Net algorithm, which in its LASSO form will select an optimal set of parameters through regularisation but not information content.

A Immune CTTA DLA Early Fusion

Covariate	Beta_mean	Beta_sd	pval_mean	Count
GAN32_10	0.396	0.115	0.001	1
GAN32_11	0.579	0.135	<0.001	1
GAN32_13	0.613	0.127	<0.001	1
GAN32_16	0.532	0.112	<0.001	5
GAN32_18	0.597	0.119	<0.001	4
GAN32_22	0.386	0.114	0.001	1
mean_0	0.571	0.125	<0.001	2
mpp_3	0.635	0.145	<0.001	1
total_0	0.524	0.126	<0.001	1
KIR3DL1.CD8.T.lymphocytes	0.546	0.165	0.011	5
cDC2	0.694	0.161	<0.001	8

B Immune CTTA DLA Intermediate Fusion

Covariate	Beta_mean	Beta_sd	pval_mean	Count
GAN32_16	0.523	0.105	<0.001	4
GAN32_18	0.54	0.111	<0.001	3
GAN32_20	0.521	0.116	<0.001	1
mean_0	0.435	0.121	0.003	2
mpp_3	0.602	0.142	<0.001	1
cDC2	0.72	0.158	<0.001	7

C Immune CTTA DLA Late Fusion

Covariate	Beta_mean	Beta_sd	pval_mean	Count
dl_sig	0.726	0.118	<0.001	8
flow_sig	0.692	0.164	<0.001	5
texrad_sig	0.589	0.127	<0.001	2

Figure 5 Risk signature generation from fusion models. (A, B, C) Outcome association statistics for Immune-CTTA-DLA early, intermediate and late fusion models of the 8 repeats, and the count of the number of times the covariate was indicated as important.

Combined Signature Specificity and Sensitivity to Detect Lung Cancer at Different Stages and Across Other Cancers

Disease staging was intentionally excluded from the data sets for both signature generation and testing. As current methodologies struggle at detecting early-stage lung cancer [21, 22] where current available therapies are most effective, we assessed the ability of the final signature to detect cancer across all stages for the training and test sets (Figure 6A). The overall sensitivity was 72%. The sensitivity was high in most Stages at 76-78%, with a drop in Stage II but we have very limited numbers in

that Stage. Interestingly, the specificity was high at 77% despite the fact that these
are patients with advanced symptoms (Figure 6B) .

Data were analysed for the 11 patients who received a non-lung cancer diagnosis.
As the primary disease was not a lung cancer, the efficacy of the lung imaging data
was unclear, and hence we used the immune signature alone. The sensitivity of this
signature to detect cancer in these cohorts was 82%, with 9/11 cancers detected
(Figure 6C). The two missed cancers were both breast cancer.

A

Lung cancer	Total Cases	True Positive	False negative	Total Sensitivity	PPV	NPV
All Cases	79	57	22	0.722	0.86	0.79
Stage I	23	18	5	0.783		
Stage II	9	3	6	0.333		
Stage III	18	14	4	0.778		
Stage IV	29	22	7	0.759		

B

No Cancer	Non-Cancer Patients	True Negative	False Positive	Total Specificity
Final Signature	91	70	21	0.769

C

Immune Signature: Other Cancers	True Positive	False negative	Sensitivity
Colorectal	2	0	1
Lymphoma	1	0	1
Breast cancer	0	2	0
Thymoma	1	0	1
Chronic Lymphocytic Leukemia	2	0	1
Kidney cancer with Lung metastasis	1	0	1
Prostate	2	0	1

10 Figure 6 Tables of the performance of the final signature (the combined late-fusion model) when
11 thresholding the risk score according to Youden's criterion: sensitivity (A) and specificity (B). Table C
12 shows the performance of the immune signature with those patients who were diagnosed with a non-
13 lung cancer; the lung CT scan therefore not being relevant as a predictor of non-lung cancer.

Discussion

Immune Subpopulations

We have previously shown that early on-treatment changes in peripheral levels of memory CD8 T-cells can predict response in patients with head and neck cancer [30], and that pre-treatment levels of a subpopulation antigen presenting capable myeloid cells can predict treatment outcome. We therefore postulated that that the systemic immune response may also be indicative of diagnosis of cancer in this study and highlight the fact that the signature obtained from a simple blood sample had a stronger predictive power than a chest CT scan. One of the strongest predictors in the signatures was the upregulation of KIR+ CD8 T lymphocytes. KIR3DL1 is part of the Killer Cell Inhibitory Receptor (KIR) family. Its expression on CD8 T cells has been linked to effector CD8 T cells with reduced proliferative capabilities and poor IFNg production following TCR engagement [37, 38]. The inhibitory effect of KIR binding to HLA has been shown to not directly impact the T lymphocyte's ability to degranulate but rather to compromise activation induced transcription which is required for clonal expansion and cytokine production [39]. KIR expression has been linked to demethylation following chronic TCR stimulation which is consistent with cancer immunosurveillance [40, 41]. In our data set, the KIR+ CD8 population exhibited a low level of expression of CD107a (Lamp1) indicating that those cells did not engage in degranulation which is even more indicative of their exhausted functional state (Supplementary Figure 5, Supplementary Figure 6).

Although KIR expression cannot not be, strictly, considered an exclusive feature of tumour specific T lymphocytes (e.g. there is a number of works describing its implications in viral immunology, especially HIV), there are works that demonstrated that KIR+ lymphocytes are restricted to specific tumour infiltrating lymphocytes subsets in renal cell carcinoma[42] and melanoma[43] and the phenotype of this cells closely resemble that of CD28 negative memory T cells. The latter work, actually, offers an intriguing hypothesis, claiming that the inhibitory nature of KIRs (they interfere with the interaction with HLA), serves the purpose of preventing the excessive cytotoxic activity of CTLs that bear a TCR specificity to self-antigens (peripheral tolerance), still maintaining reactivity against tumoral cells that might downregulate class I expression (a characteristic mechanism of tumour immune surveillance evasion).

Moreover, recent works suggests an active regulatory function of KIR+CD8+ lymphocytes that, for example, seems to suppress anti-tumour immunity in patients with melanoma[44]. This peculiar subset of CD8 T-lymphocytes resemble the phenotype of mouse Ly49+CD8+ Tregs[45, 46] and appears to be particularly relevant in both auto-immune conditions, such as lupus and celiac disease in which they efficiently eliminate gliadin-specific CD4 T-lymphocytes, and SARS-CoV-2 [47], making them particularly relevant in the context of self-tolerance.

Of particular interest is that this signature could distinguish between a cancer and a non-cancer pathology. As most of the other pathologies were linked to an infection or an allergic reaction, the upregulation of cDC2 in those patients is only logical. cDC2 are a subtype of DCs that play a key role for antigen presentation for a type 2 helper response (TH2) [48] which is key in controlling humoral immunity. cDC2 have also been correlated with their ability to activate a TH17 response, which has been shown to be upregulated in chronic allergy [49, 50]. The cDC2 cells expressed high levels of HLADR and CD38, indicating that they were indeed functional and pro-inflammatory. This subpopulation of myeloid cells did not express PDL2 indicating that it was not functionally suppressed (Supp. Fig. 6).

In recent years, the immune-surveillance theory and the efficacy of anticancer immunotherapies have been progressively more, and closely connected to the concept of immunometabolism and the microbiome profile of patients with cancer[51], making clear that immune evasion (a contributing factor to tumorigenesis) may be affected directly by metabolic pathways e.g. lactate metabolism[52], and these metabolic dysregulations can, in turn, affect immune cell function and tumour cell progression[53]. The KIR3DL1 CD8 T cells could represent a hypofunctional T cell subset[54] that contributes to tumorigenesis.

It is now recognised that T and NK quiescence, activation, and exhaustion are regulated by cell metabolic states that work in conjunction with immunological stimuli[55]. In this perspective it would be of pivotal interest including in future models modalities that consider metabolomic parameters that could strengthen, because of causative correlations, our exquisitely immunological observations. The other consistent immunological parameter that links with the cancer pathology is reduced peripheral blood cDC2. Further work is required to investigate whether this is correlated with an increased degree of infiltration by cDC2 as previously reported in head and neck cancers[56].

Deep Learning Radiomics and Combined Model Generation

There has been significant recent activity in the development of techniques to extract novel information from medical images and other patient sample data and exploit it for clinical use[57]. Links between imaging and the immune system have also been made[58-62]. Grossman et al.[58] evaluated radiomic texture and shape features in NSCLC and found correlations with immune response, inflammation and survival, and a combined signature that combined clinical, genetic and radiomics achieved a test set concordance index of 0.73 against OS.

Deep Learning networks have shown great promise in several fields, including cancer diagnosis, for extracting significant features from complex data, but their often black-box nature is one barrier to clinical use[63]. Autoencoders are able to learn semantically meaningful representations of images in an unsupervised manner and offer some advantages: training on an external data set and transparency through the exploration of the latent space parameters. Examples have been published in using epigenetic

data to determine subtypes of lung cancer[64], and image-based lesion detection and segmentation[65].

The method for combining radiomics with blood data, needs consideration (i.e. data fusion[66]) as do the algorithms used to generate the predictive models. A model formed from many covariates can easily have exceptional performance on the training set, but this often does not translate to test and validation data sets (the problem of overfitting). Many studies use parameter regularisation such as LASSO or Elastic Net algorithms for signature generation and covariate selection. In particular, a study investigated the feasibility of preoperative 18F-fluorodeoxyglucose (FDG) PET/CT radiomics with machine learning (LASSO) to predict microsatellite instability (MSI) status in patients with colorectal cancer (CRC), with a test set performance (ROC AUC) of 0.867 from a published set of radiomics features[67]. Methods more advanced than regularisation offer a greater potential for generalisation to test and validation sets. For instance, neural networks and XGBoost models have achieved high ROC AUC of 0.75-0.8 in multimodal data sets[68-70]. Bayesian methods offer robust models that are generalisable and more interpretable and can have automatic feature selection that aid interpretability[71, 72].

While a completely independent validation cohort with deep radiomic and immune profiles is currently outside the scope of this manuscript, we have repeated the randomisation of the patient cohort and machine learning 8 times, and the same two immune variables (reduced cDC2 levels and increased KIR3DL1+ CD8 T cells) have been retained in the model in the majority of these 8 independent model runs, lending credence to the selected predictive features.

A ROC AUC of 0.8 is within the realms of what may be clinically useful depending on the context. Choosing the optimal point on the ROC curve, to define operational sensitivity and specificity, may not be an obvious one for the clinical situation. Indeed, the multi-cancer Grail tests[21, 73] have a similar AUC. The Galleri test has a specificity of around 99% but a sensitivity as low as 17% (depending on cancer and stage)[74].

The 6mm clinical cutoff lesion size may limit the applicability of this model and we may obtain better results in early cancers is this is relaxed. This would however require a different benchmark to the optimal NOLCP pathway and longer follow-up to obtain definitive diagnosis which was not in the scope of this study. Further improvement may be obtained by retraining the deep learning autoencoder on a dataset with a greater match to the problem to avoid any domain mismatch between the LUNA16 dataset and the study acquired images. There may be other sources of bias in the training and test datasets that cannot be controlled for in this study, such as a lack of heterogeneity or cancer subtypes.

This present study has data collected in a prospective and standardised manner and used Bayesian machine learning techniques that emphasise reproducibility. Furthermore, ours is a "white box" method that generates an optimal set of parameters and weights that are available for interpretation. We explore a combination of deep

1   learning radiomics and pre-determined standardized features in the form of the well-

2   published filtration-histogram based texture analysis approach (using the

3   commercially available TexRAD software) for feature selection from CT images.

Contributors

Contributions to this manuscript are as follows: Data curation: RM, BG, SE, LD, MT, XJ, CS, SL, EC, AT, LH, RE, RS, DW, SC, PB; Formal analysis: RM, BG, SE, EC, AC, PB; Investigation: RM, SE, LD, XJ, CS, SC, PB; Methodology: RM, BG, SE, LD, XJ, CS, AC, JS, TN, PB; Resources: RM, LD, KD, SL; Validation: RM, LD, MR, AC, TN, PRB; Visualisation: RM, PB; Software: SE, MT, MR, AC, PB; Supervision: BG, AC, AG, JS, TW, PB, TN; Writing – original draft: RM, BG, SE, KD, JAS, TW, PB, TN; Writing – review & editing: LD, PB, TN; Conceptualisation: AG, JS, TW, TN; Funding acquisition: AG, JS, TW, TN; Project administration: TN.

All authors read and approve the final version of the manuscript. RM and PB had access to the underlying data for the multivariate analysis.

Data Sharing Statement

Model generation was performed using the MsaWrapper R package (https://github.com/paulbarber/Lung_EDx_Manuscript_Processing and https://github.com/paulbarber/msaWrapper) and Saddle Point Signature (commercial software distributed by Saddle Point Science Ltd).

Anonymised data used for this analysis is available at reasonable request (to comply with patient consent) from: kcl.figshare.com/account/articles/28070390

Declaration of Interests

BG is the co-founder/co-inventor of TexRAD texture analysis software used in this study for CT Texture analysis and a shareholder (not an employee) of Feedback Plc., a UK-based company which owns the TexRAD texture analysis software. PB and TN are both currently employees of GlaxoSmithKline (GSK) but this study has not received any funding support from GSK. TC is CEO of Saddle Point Europe BV. MR is CEO of Saddle Point Science Ltd.

All remaining authors have declared no conflicts of interest.

Acknowledgements

We are grateful to all participating staff at The Lister, Hertford County, UK and New QEII Hospitals, Welwyn Garden City, UK who were responsible for patient recruitment and sample collection.

This work was funded by the CRUK grant [C1519/A27375]. SE, EC and JAS were further supported by core funding from the Wellcome Trust/EPSRC Centre for Medical

Engineering [WT203148/Z/16/Z] and by the National Institute for Health and Care Research (NIHR) Clinical Research Facility at Guy's and St Thomas' NHS Foundation Trust. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care. Work was undertaken at UCLH/UCL, which received a proportion of the funding from the UK's Department of Health's NIHR Biomedical Research Centre's funding scheme.

References

Sung H, Ferlay J, Siegel RL et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. CA Cancer J Clin 2021; 71 (3): 209-249.

Siegel RL, Miller KD, Wagle NS, Jemal A. Cancer statistics, 2023. Ca-Cancer J Clin 2023; 73 (1): 17-48.

Leiter A, Veluswamy RR, Wisnivesky JP. The global burden of lung cancer: current status and future trends. Nature Reviews Clinical Oncology 2023; 20 (9): 624-639.

Cancer Research UK, Lung cancer statistics, https://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type/lung-cancer#heading-Zero accessed May. 2024.

Malalasekera A, Nahm S, Blinman PL et al. How long is too long? A scoping review of health system delays in lung cancer. European Respiratory Review 2018; 27 (149): 180045.

Walter FM, Rubin G, Bankhead C et al. Symptoms and other factors associated with time to diagnosis and stage of lung cancer: a prospective cohort study. Br J Cancer 2015; 112 Suppl 1 (Suppl 1): S6-13.

Lyratzopoulos G, Abel GA, Mcphail S et al. Measures of promptness of cancer diagnosis in primary care: secondary analysis of national audit data on patients with 18 common and rarer cancers. British Journal of Cancer 2013; 108 (3): 686-690.

Swann R, McPhail S, Witt J et al. Diagnosing cancer in primary care: results from the National Cancer Diagnosis Audit. Br J Gen Pract 2018; 68 (666): e63-e72.

National Lung Screening Trial Research T, Aberle DR, Berg CD et al. The National Lung Screening Trial: overview and study design. Radiology 2011; 258 (1): 243-253.

de Koning HJ, van der Aalst CM, de Jong PA et al. Reduced Lung-Cancer Mortality with Volume CT Screening in a Randomized Trial. N Engl J Med 2020; 382 (6): 503-513.

Wood DE, Kazerooni EA, Aberle D et al. NCCN Guidelines(R) Insights: Lung Cancer Screening, Version 1.2022. J Natl Compr Canc Netw 2022; 20 (7): 754-764.

Dingillo G, Bassiri A, Badrinathan A et al. Lung Cancer in Young Patients is Associated With More Advanced Disease but Better Overall Survival. J Surg Res 2023; 292: 307-316.

Woodard GA, Jablons DM. Shades of Gray: Do Never Smokers Benefit From Lung Cancer Screening Programs? J Thorac Oncol 2024; 19 (8): 1135-1137.

Dolly SO, Jones G, Allchorne P et al. The effectiveness of the Guy's Rapid Diagnostic Clinic (RDC) in detecting cancer and serious conditions in vague symptom patients. Br J Cancer 2021; 124 (6): 1079-1087.

National Optimal Lung Cancer Pathway (NOLCP). https://rmpartners.nhs.uk/wp-content/uploads/2024/09/national-optimal-lung-cancer-pathway_v4_01jan2024.pdf: NHS England; 2024.

The Royal College of Radiologists, Clinical oncology census reports 2022. https://www.rcr.ac.uk/news-policy/policy-reports-initiatives/clinical-oncology-census-reports/ accessed April. 2024.

Lung Cancer Europe: 2020 Position Paper. Disparities and Challenges in Access to Lung Cancer Diagnosis and Treatment Across Europe. Accessed: April.

Cellina M, Cacioppa LM, Ce M et al. Artificial Intelligence in Lung Cancer Screening: The Future Is Now. Cancers (Basel) 2023; 15 (17).

Kang E, Min J, Ye JC. A deep convolutional neural network using directional wavelets for low-dose X-ray CT reconstruction. Med Phys 2017; 44 (10): e360-e375.

Casagrande GMS, Silva MO, Reis RM, Leal LF. Liquid Biopsy for Lung Cancer: Up-to-Date and Perspectives for Screening Programs. Int J Mol Sci 2023; 24 (3).

Nicholson BD, Oke J, Virdee PS et al. Multi-cancer early detection test in symptomatic patients referred for cancer investigation in England and Wales (SYMPLIFY): a large-scale, observational cohort study. Lancet Oncol 2023; 24 (7): 733-743.

Schrag D, Beer TM, McDonnell CH, 3rd et al. Blood-based tests for multicancer early detection (PATHFINDER): a prospective cohort study. Lancet 2023; 402 (10409): 1251-1260.

Pastorino U, Boeri M, Sestini S et al. Baseline computed tomography screening and blood microRNA predict lung cancer risk and define adequate intervals in the BioMILD trial. Ann Oncol 2022; 33 (4): 395-405.

Ostrin EJ, Bantis LE, Wilson DO et al. Contribution of a Blood-Based Protein Biomarker Panel to the Classification of Indeterminate Pulmonary Nodules. J Thorac Oncol 2021; 16 (2): 228-236.

Valadez-Cosmes P, Maitz K, Kindler O et al. Identification of Novel Low-Density Neutrophil Markers Through Unbiased High-Dimensional Flow Cytometry Screening in Non-Small Cell Lung Cancer Patients. Front Immunol 2021; 12: 703846.

Fortunato O, Huber V, Segale M et al. Development of a Molecular Blood-Based Immune Signature Classifier as Biomarker for Risks Assessment in Lung Cancer Screening. Cancer Epidemiol Biomarkers Prev 2022; 31 (11): 2020-2029.

Lee SW, Kim YJ, Rho KN et al. Immunophenotypic analysis on circulating T cells for early diagnosis of lung cancer. Biomark Res 2024; 12 (1): 161.

Krieg C, Nowicka M, Guglietta S et al. High-dimensional single-cell analysis predicts response to anti-PD-1 immunotherapy. Nat Med 2018; 24 (2): 144-+.

Pauken KE, Shahid O, Lagattuta KA et al. Single-cell analyses identify circulating anti-tumor CD8 T cells and markers for their enrichment. J Exp Med 2021; 218 (4).

Barber PR, Mustapha R, Flores-Borja F et al. Predicting progression-free survival after systemic therapy in advanced head and neck cancer: Bayesian regression and model development. Elife 2022; 11.

Hill W, Lim EL, Weeden CE et al. Lung adenocarcinoma promotion by air pollutants. Nature 2023; 616 (7955): 159-167.

Li M, Zhang C, Deng S et al. Lung cancer-associated T cell repertoire as potential biomarker for early detection of stage I lung cancer. Lung Cancer 2021; 162: 16-22.

Massa E, Coolen ACC. Observable asymptotics of regularized Cox regression models with standard Gaussian designs: a statistical mechanics approach. Journal of Physics A: Mathematical and Theoretical 2025; 58 (10): 105001.

Massa E, Coolen ACC. Practical debiasing with the Covariant Prior in the proportional regime when p < n. Journal of Physics Communications 2025; 9 (5): 055001.

Friedman J, Tibshirani R, Hastie T. Regularization Paths for Generalized Linear Models via Coordinate Descent. Journal of Statistical Software 2010; 33 (1).

Simon N, Friedman J, Tibshirani R, Hastie T. Regularization Paths for Cox's Proportional Hazards Model via Coordinate Descent. Journal of Statistical Software 2011; 39 (5).

Anfossi N, Doisne JM, Peyrat MA et al. Coordinated expression of Ig-like inhibitory MHC class I receptors and acquisition of cytotoxic function in human CD8+ T cells. J Immunol 2004; 173 (12): 7223-7229.

Arlettaz L, Degermann S, De Rham C et al. Expression of inhibitory KIR is confined to CD8+ effector T cells and limits their proliferative capacity. Eur J Immunol 2004; 34 (12): 3413-3422.

Henel G, Singh K, Cui D et al. Uncoupling of T-cell effector functions by inhibitory killer immunoglobulin-like receptors. Blood 2006; 107 (11): 4449-4457.

McLane LM, Abdel-Hakeem MS, Wherry EJ. CD8 T Cell Exhaustion During Chronic Viral Infection and Cancer. Annu Rev Immunol 2019; 37: 457-495.

Li G, Yu M, Weyand CM, Goronzy JJ. Epigenetic regulation of killer immunoglobulin-like receptor expression in T cells. Blood 2009; 114 (16): 3422-3430.

Guerra N, Guillard M, Angevin E et al. Killer inhibitory receptor (CD158b) modulates the lytic activity of tumor-specific T lymphocytes infiltrating renal cell carcinomas. Blood 2000; 95 (9): 2883-2889.

Bakker ABH, Phillips JH, Figdor CG, Lanier LL. Killer Cell Inhibitory Receptors for MHC Class I Molecules Regulate Lysis of Melanoma Cells Mediated by NK cells, γδ T Cells, and Antigen-Specific CTL1. The Journal of Immunology 1998; 160 (11): 5239-5245.

Lu BY, Lucca LE, Lewis W et al. Circulating tumor-reactive KIR(+)CD8(+) T cells suppress anti-tumor immunity in patients with melanoma. Nat Immunol 2025; 26 (1): 82-91.

Kim H-J, Wang X, Radfar S et al. CD8[+] T regulatory cells express the Ly49 Class I MHC receptor and are defective in autoimmune prone B6-Yaa mice. Proceedings of the National Academy of Sciences 2011; 108 (5): 2010-2015.

Xia Q, Huang H, Davis MM. A High-Throughput Strategy for T-Cell Receptor Cloning and Expression. Methods Mol Biol 2022; 2574: 251-264.

Koh C-H, Lee S, Kwak M et al. CD8 T-cell subsets: heterogeneity, functions, and therapeutic potential. Experimental & Molecular Medicine 2023; 55 (11): 2287-2299.

Kumar S, Jeong Y, Ashraf MU, Bae YS. Dendritic Cell-Mediated Th2 Immunity and Immune Disorders. Int J Mol Sci 2019; 20 (9).

Izumi G, Nakano H, Nakano K et al. CD11b(+) lung dendritic cells at different stages of maturation induce Th17 or Th2 differentiation. Nat Commun 2021; 12 (1): 5029.

Zhao J, Lloyd CM, Noble A. Th17 responses in chronic allergic airway inflammation abrogate regulatory T-cell-mediated tolerance and contribute to airway remodeling. Mucosal Immunol 2013; 6 (2): 335-346.

Zhou CB, Zhou YL, Fang JY. Gut Microbiota in Cancer Immune Response and Immunotherapy. Trends Cancer 2021; 7 (7): 647-660.

Tong H, Jiang Z, Song L et al. Dual impacts of serine/glycine-free diet in enhancing antitumor immunity and promoting evasion via PD-L1 lactylation. Cell Metabolism 2024; 36 (12): 2493-2510.e2499.

Anderson KG, Stromnes IM, Greenberg PD. Obstacles Posed by the Tumor Microenvironment to T cell Activity: A Case for Synergistic Therapies. Cancer Cell 2017; 31 (3): 311-325.

Zhang X, Lu X, Moog C et al. KIR3DL1-Negative CD8 T Cells and KIR3DL1-Negative Natural Killer Cells Contribute to the Advantageous Control of Early Human Immunodeficiency Virus Type 1 Infection in HLA-B Bw4 Homozygous Individuals. Frontiers in Immunology 2018; Volume 9 - 2018.

Chapman NM, Boothby MR, Chi H. Metabolic coordination of T cell quiescence and activation. Nat Rev Immunol 2020; 20 (1): 55-70.

Kirchner J, Plesca I, Rothe R et al. Type I conventional dendritic cells and CD8(+) T cells predict favorable clinical outcome of head and neck squamous cell carcinoma patients. Front Immunol 2024; 15: 1414298.

Chetan MR, Gleeson FV. Radiomics in predicting treatment response in non-small-cell lung cancer: current status, challenges and future perspectives. Eur Radiol 2021; 31 (2): 1049-1058.

Grossmann P, Stringfield O, El-Hachem N et al. Defining the biological basis of radiomic phenotypes in lung cancer. Elife 2017; 6.

Yang Q, Gong H, Liu J et al. A 13-gene signature to predict the prognosis and immunotherapy responses of lung squamous cell carcinoma. Sci Rep 2022; 12 (1): 13646.

Wiesweg M, Mairinger F, Reis H et al. Machine learning reveals a PD-L1-independent prediction of response to immunotherapy of non-small cell lung cancer by gene expression context. Eur J Cancer 2020; 140: 76-85.

Nunez NG, Berner F, Friebel E et al. Immune signatures predict development of autoimmune toxicity in patients with cancer treated with immune checkpoint inhibitors. Med 2023; 4 (2): 113-129 e117.

Duruisseaux M, Martinez-Cardus A, Calleja-Cervantes ME et al. Epigenetic prediction of response to anti-PD-1 treatment in non-small-cell lung cancer: a multicentre, retrospective analysis. Lancet Respir Med 2018; 6 (10): 771-781.

Singh A, Sengupta S, Lakshminarayanan V. Explainable Deep Learning Models in Medical Image Analysis. J Imaging 2020; 6 (6).

Wang Z, Wang Y. Extracting a biologically latent space of lung cancer epigenetics with variational autoencoders. BMC Bioinformatics 2019; 20 (Suppl 18): 568.

Astaraki M, Toma-Dasu I, Smedby Ö, Wang C. Normal Appearance Autoencoder for Lung Cancer Detection and Segmentation. Cham: Springer International Publishing; 2019:249-256.

Lipkova J, Chen RJ, Chen B et al. Artificial intelligence for multimodal data integration in oncology. Cancer Cell 2022; 40 (10): 1095-1110.

Kim S, Lee JH, Park EJ et al. Prediction of Microsatellite Instability in Colorectal Cancer Using a Machine Learning Model Based on PET/CT Radiomics. Yonsei Med J 2023; 64 (5): 320-326.

Jiang J, Jin Z, Zhang Y et al. Robust Prediction of Immune Checkpoint Inhibition Therapy for Non-Small Cell Lung Cancer. Front Immunol 2021; 12: 646874.

Tian P, He B, Mu W et al. Assessing PD-L1 expression in non-small cell lung cancer and predicting responses to immune checkpoint inhibitors using deep learning on computed tomography images. Theranostics 2021; 11 (5): 2098-2107.

Ahn BC, So JW, Synn CB et al. Clinical decision support algorithm based on machine learning to assess the clinical response to anti-programmed death-1 therapy in patients with non-small-cell lung cancer. Eur J Cancer 2021; 153: 179-189.

Shalabi A, Inoue M, Watkins J et al. Bayesian clinical classification from high-dimensional data: Signatures versus variability. Stat Methods Med Res 2018; 27 (2): 336-351.

Hozumi H, Shimizu H. Bayesian network enables interpretable and state-of-the-art prediction of immunotherapy responses in cancer patients. PNAS Nexus 2023; 2 (5): pgad133.

Klein EA, Beer TM, Seiden M. The Promise of Multicancer Early Detection. Comment on Pons-Belda et al. Can Circulating Tumor DNA Support a Successful Screening Test for Early Cancer Detection? The Grail Paradigm. Diagnostics 2021, 11, 2171. Diagnostics 2022; 12 (5): 1243.

Pons-Belda OD, Fernandez-Uriarte A, Diamandis EP. Multi Cancer Early Detection by Using Circulating Tumor DNA—The Galleri Test. Reply to Klein et al. The Promise of Multicancer Early Detection. Comment on "Pons-Belda et al. Can Circulating Tumor DNA Support a Successful Screening Test for Early Cancer Detection? The Grail Paradigm. Diagnostics 2021, 11, 2171". Diagnostics 2022; 12 (5): 1244.

Figure Legends

Figure 1. Showing the patient pathway and study design. 344 patients referred to the clinic were recruited to the study. As part the standard of care pathway they received a CT scan, at which time we collected the peripheral blood samples. 174 cases were excluded from the analysis set due to lack of visible lesion, the lesion was smaller than 6mm or the diagnosis was a non-lung cancer diagnosis. The analysis was performed on the remaining patients.

Figure 2. Risk generation by BMR from repeated data splits and comparing CT texture analysis to deep learning autoencoder. (A) Box plots showing the distributions (mean and inter-quartile range, points overlaid) of AUCs obtained from 8 random training/test splits of the data. (B, C) Outcome association statistics for CTTA and DLA of the 8 repeats, and the count of the number of times the covariate was indicated as important. Covariates indicated more than 50% of the time were selected for the final model generation; the final association (Beta), p-value for the association and weight in the final signature are given, as well as a ROC curve and AUC value for the final signature.

Figure 3. Risk generation by BMR from peripheral blood samples comparing flow cytometry to plasma exosome dotblot analysis. (A, B) Outcome association statistics for Immune and Dotblot of the 8 repeats, and the count of the number of times the covariate was indicated as important. Covariates indicated more than 50% of the time were selected for the final model generation; the final association (Beta), p-value for the association and weight in the final signature are given, as well as a ROC curve and AUC value for the final signature. Associations for the dotblot data were so low that a final model was not generated.

Figure 4. Risk generation by BMR from CT image analysis combined with flow cytometry, comparing deep learning and texture analysis. (A, B) Outcome association statistics for Immune-CTTA and Immune-DLA fusion models of the 8 repeats, and the count of the number of times the covariate was indicated as important. Covariates indicated more than 50% of the time were selected for the final model generation; the final association (Beta), p-value for the association and weight in the final signature are given, as well as a ROC curve and AUC value for the final signature.

Figure 5 Risk signature generation from fusion models. (A, B, C) Outcome association statistics for Immune-CTTA-DLA early, intermediate and late fusion models of the 8 repeats, and the count of the number of times the covariate was indicated as important.

Figure 6 Tables of the performance of the final signature (the combined late-fusion model) when thresholding the risk score according to Youden's criterion: sensitivity (A) and specificity (B). Table C shows the performance of the immune signature with those patients who were diagnosed with a non-lung cancer; the lung CT scan therefore not being relevant as a predictor of non-lung cancer.