HaptiKart: An engaging videogame reveals elevated proprioceptive vs. visual bias in individuals with autism spectrum disorder

Short Title: *Proprioceptive vs. visual bias and autism spectrum disorder*

Daniel E. Lidstone, Ph.D.[1,2*], Mohit Singhala, Ph.D.[3], Liam J. Wang[4], Jeremy D. Brown, Ph.D.[5], Stewart H. Mostofsky, M.D.[2,6,7]

[1]School of Behavioral Sciences and Education, Penn State Harrisburg, Middletown, Pennsylvania, United States of America

[2]Center for Neurodevelopmental and Imaging Research, Kennedy Krieger Institute, Baltimore, Maryland, United States of America

[3]Department of Otolaryngology-Head & Neck Surgery, University of Pittsburgh, Pittsburgh, Pennsylvania, United States of America

[4]Electrical Engineering and Computer Science Department, University of Michigan, Ann Arbor, Michigan, United States of America

[5]Department of Mechanical Engineering, Johns Hopkins University, Baltimore, Maryland, United States of America

[6]Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, Maryland, United States of America

[7]Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, Maryland, United States of America

* Corresponding author

E-mail: dbl5689@psu.edu (DE)

Abstract

An overreliance on proprioceptive (intrinsic) sensory input from the body, compared to visual (extrinsic) input from the environment, may underpin core features of autism spectrum disorder (ASD). We developed an engaging videogame ("HaptiKart") as a tool to examine differences in proprioceptive vs. visual bias in children and adults with ASD and whether bias correlates with age, core autism features, and intellectual ability. Eighty-one participants (33 ASD, 48 typically-developing, TD) aged 8 to 31 years played "HaptiKart," a driving videogame with a force-feedback steering wheel that provided "steering assist" during gameplay. In separate trials, proprioceptive and visual feedback were selectively delayed, and differences in driving error between the conditions were used to calculate perceptual bias scores. Effects of autism diagnosis and age on perceptual (proprioceptive vs. visual) bias scores were examined, controlling for sex, as were associations of perceptual bias with autism symptom severity (ADOS-2, SRS-2), attention-deficit symptom severity (Conners4 ADHD Total Scores) ratings, and IQ (general ability index, GAI). The ASD group exhibited significantly higher proprioceptive bias than did the TD group ($p=0.002$). There was a trend for decreasing proprioceptive bias with age, but no significant diagnosis-by-age interaction. Increased proprioceptive bias correlated with higher autism severity and with lower IQ, but not ADHD symptoms. HaptiKart provides a highly scalable approach for measuring sensory-motor bias, revealing that individuals with ASD show elevated proprioceptive bias, correlating with autism severity. HaptiKart's sensory-motor bias measure may thereby serve as a digital biomarker for addressing autism heterogeneity in ways that can improve targeted intervention.

Author Summary

In this study, we explored how people with autism spectrum disorder (ASD) may rely more on body-based (proprioceptive) senses than on visual information when interacting with their surroundings. Using a custom video game, *HaptiKart*, we designed a fun and engaging driving task to measure this sensory preference. Players drove a virtual car using a steering wheel that could be set to delay either visual or proprioceptive feedback, allowing us to observe how each type of feedback affects driving accuracy.

Our results showed that individuals with ASD had a stronger preference for proprioceptive feedback than those without ASD. This sensory preference was linked to greater autism symptom severity and lower IQ, suggesting that heightened reliance on body-centered feedback may contribute to learning and skill differences often seen in autism. These findings support the potential of *HaptiKart* as a simple, accessible tool to help clinicians better understand and address sensory biases in ASD, tailoring interventions to improve learning from visual information.

Introduction

Acquiring social-communicative and motor skills from others through observation and imitation requires efficient dynamic visual-motor integration (VMI). Evidence from children with autism spectrum disorder (ASD) shows significant differences in dynamic VMI in non-social (e.g., ball catching) and social (e.g., motor imitation) behaviors, which may affect the implicit learning of motor and social skills [1,2]. Studies suggest a shared neural basis underlying both motor and social-communicative differences in children with ASD, demonstrated by differences in tasks that rely on dynamic VMI, such as ball catching, motor imitation, and visual-motor tracking [3–6]. Moreover, differences in dynamic VMI can be detected as early as six months of age, potentially serving as an early indicator of ASD [7]. This suggests that differences in brain mechanisms responsible for dynamic VMI may be present at birth or at least emerge early in children who later develop the core autism phenotype and receive an ASD diagnosis.

Autism-associated challenges with dynamic visual-motor integration (VMI) may lead to a preference for intrinsic proprioceptive feedback over extrinsic visual feedback in daily tasks. This bias could support skill acquisition in static environments but impede learning through observation in dynamic settings. Research shows that dynamic visual input perturbations disrupt motor control less in children with ASD compared to typically developing peers, indicating a stronger reliance on proprioceptive input [8–10]. Importantly, this proprioceptive vs. visual bias seems unique to ASD, as it is not seen in conditions like ADHD [11]. Most studies focus on children under 12 [9,10,12], suggesting that these biases may diminish in adolescence, potentially influenced by interventions that promote learning from dynamic visual cues, such as Social Skills Groups. A prolonged proprioceptive vs. visual bias in childhood may also impact intellectual development reliant on implicit learning from visual stimuli. Those with lower intellectual abilities might depend more on proprioceptive feedback for environmental exploration and learning. However, the literature has yet to explore whether increased proprioceptive vs. visual bias correlates with intellectual differences in individuals with ASD. Identifying this bias in children can help clinicians tailor interventions to enhance learning from dynamic visual feedback, such as social and motor behavior observation and imitation.

The relative weighting of proprioceptive vs. visual feedback is a simple measure that may serve as a useful digital biomarker of a particular autism subtype with utility for aiding diagnosis, parsing autism heterogeneity, and guiding intervention. However, previous computational methods to examine proprioceptive vs. visual bias have relied on inaccessible tools (i.e., expensive, non-portable robots) and long-duration lab assessments that limit clinical utility and accessibility. Furthermore, proprioceptive vs. visual bias has primarily been examined using tasks, such as ballistic reaching, where proprioception is the primary feedback modality [10,11,13–15]. Thus, it remains unclear whether autism-associated proprioceptive vs. visual bias is also present in visual-dominant tasks.

To address limitations in prior computational methods for examining proprioceptive vs. visual bias, we developed "HaptiKart," an arcade-style driving videogame (**Fig 1**).

Fig 1. HaptiKart videogame. A child participant playing the HaptiKart videogame.

This platform delivers configurable visual feedback on the car's position and proprioceptive
feedback through a force-feedback steering wheel. We assessed driving errors in two
conditions—delayed proprioceptive feedback and delayed visual feedback. Importantly, this
approach to delivering proprioceptive and visual error was implicit, as it was challenging for
participants to recognize whether the visual or proprioceptive feedback had changed explicitly.
Additionally, the order of the conditions was counterbalanced within each group to control for
potential order effects.

This first study of HaptiKart had three aims: (1) to examine the effects of ASD diagnosis on
perceptual (proprioceptive vs. visual) bias, (2) to examine the effects of age, as well as age-by-
diagnosis on perceptual bias, (3) to examine associations with ASD clinical severity, as well as
with severity of commonly co-occurring ADHD symptoms and with intellectual ability. Our
corresponding hypotheses were: (1) individuals with ASD would show elevated proprioceptive
vs. visual bias relative to TD individuals, (2) proprioceptive vs. visual bias would show a
significant decrease with age with steeper age-related declines in the ASD vs. TD groups, and (3)
proprioceptive vs. visual bias would correlate significantly with core autism features, but not
attentional (ADHD) traits. We also hypothesized that, in addition to the association between
elevated proprioceptive vs. visual bias and core autism features, this bias would also be linked to
lower intellectual ability. Specifically, we propose that disrupted visual input from the external
environment, beyond just social stimuli, may more generally affect cognitive development.

Results

Elevated Proprioceptive vs. Visual Bias in Individuals with ASD Relative to TD Controls

We examined ASD vs. TD differences proprioceptive vs. visual bias by comparing driving errors between the counterbalanced proprioceptive delay and visual delay conditions (**Fig 2**).

Fig 2. **Experimental paradigm and sample data.** Schematic showing the paradigm used to examine proprioceptive vs. visual bias. The delayed visual and proprioceptive feedback conditions were counterbalanced within each group. Trials from two sample participants.

The findings from ANCOVA did not show a significant diagnosis-by-age effect (p = 0.68) or effect of age (p = 0.06) but did show a significant effect of diagnosis (p = 0.002) on bias scores (**Fig 3**).

Fig 3. **Group differences in perceptual bias scores**. Children with ASD show significantly greater proprioceptive vs. visual bias as compared to TD children.

Bias scores in the ASD group were more positive than the TD group (ASD bias score: -4.24 ± 5.83 vs. TD bias score: -8.34 ± 5.90), showing elevated proprioceptive vs. visual bias. Further, the biological sex covariate did not have a significant effect on bias scores ($p = 0.68$). Collectively, individuals with an ASD diagnosis showed increased preference for proprioceptive vs. visual feedback that was not explained by age or biological sex.

Proprioceptive vs. Visual Bias Correlates with Core Autism Features and IQ, but not ADHD Features

Correlations between proprioceptive vs. visual bias and core autism features revealed significant correlation with clinical ratings of autism symptoms within the ASD group (ADOS-CS, $r = 0.48$, $p = 0.035$, $p_{adj} = 0.046$) and parent rated autism symptoms at the whole group level (SRS-2 total score, $r = 0.52$, $p = 0.0003$, $p_{adj} = 0.0006$) (**Fig 4A-B**), such that elevated symptoms predicted elevated proprioceptive vs. visual bias. In contrast, parent-rated ADHD symptom severity was not significantly correlated with bias scores (Conners4 total score, $r = 0.17$, $p = 0.38$, $p_{adj} = 0.38$) (**Fig 4F**). Bias scores were also significantly correlated with GAI, such that elevated proprioceptive vs. visual bias was associated with lower GAI ($r = 0.54$, $p = 0.000002$, $p_{adj} = 0.000008$) (**Fig 4C**).

Fig 4. Correlations between perceptual bias scores, age, and autism-associated features:
Simple linear regression showing a significant correlation between elevated proprioceptive vs.
visual bias and: (A) elevated clinically rated ASD symptoms (ADOS-CS (Module 3) scores,
p=0.03), (B) elevated parent-rated ASD symptoms (SRS-2 total scores, p < 0.001), and (C) lower
intellectual ability (GAI, p < 0.001). (D-E) Correlations between age and bias scores were not
significant for either group (p>0.1) but showed a negative trend-level association at the whole
cohort level (age, p=0.08). (F) There was no correlation between parent-rated ADHD symptoms
and bias scores (Conners4 total score, p=0.38).

Trend for Decreasing Proprioceptive vs. Visual Bias with Increasing Age Across Both ASD and TD Cohorts

ANCOVA revealed a trend-level main effect of age on bias scores (p = 0.06). Simple linear
regression (bias scores ~ age) revealed a trend for an effect of age on perceptual bias across the
entire cohort, with decreasing proprioceptive vs. visual bias with increasing age (r = -0.2, p =
0.08) (**Fig 4E**). However, there was no significant interaction of diagnosis and age (p = 0.68),
such that this trend for increasing reliance on visual error vs. proprioceptive error from childhood
to adulthood was similar in both diagnostic groups (ASD: r = -0.26, p=0.13; TD: r = -0.17, p =
0.25) (**Fig 4D**).

Discussion

This study reveals two main findings that support the stated hypotheses: (1) children with ASD show significantly elevated proprioceptive vs. visual bias relative to TD children, (2) elevated proprioceptive vs. visual bias correlates with greater core autism clinical severity and with lower intellectual ability but not with ADHD clinical severity. While associations between age and proprioceptive vs. visual bias were not significant, trend-level associations with age were observed such that elevated proprioceptive vs. visual bias was observed in younger vs. older participants. Examination of diagnosis-by-age effects on proprioceptive vs. visual bias did not support our hypothesis of elevated bias in younger but not older individuals with ASD vs. TD. These findings suggest that ASD-associated proprioceptive vs. visual bias persists across development and does not appear to normalize to TD levels with increasing age. However, including younger participants (<8 years of age) in future studies will provide a deeper insight into autism-associated differences in developmental trajectories of proprioceptive vs. visual bias.

The finding that children with ASD show elevated proprioceptive vs. visual bias in the current study supports a degree of task invariance of the measure as it has been previously observed in proprioception-dominant tasks such as ballistic reaching [11,14,15] and now with HaptiKart, a more visual task. Consistent with prior studies, the current study showed elevated proprioceptive vs. visual bias correlated with parent-rated core autism features [14]. The finding that elevated proprioceptive vs. visual bias correlated with core autism symptoms but not ADHD symptoms suggests that proprioceptive vs. visual bias may be a promising digital phenotypic marker of autism.

Our findings indicate a strong correlation between preference for proprioceptive vs. visual feedback and general intellectual ability (GAI), suggesting this bias may be more pronounced in individuals with severe ASD symptoms. While previous studies have linked intellectual ability and motor skills in ASD, these correlations are expected due to the complexity of standardized motor tests, which can be challenging for those with autism and intellectual differences [16,17]. Notably, our study highlights a robust association between elevated proprioceptive vs. visual bias and IQ, focusing solely on sensory preferences rather than motor skills. This sensory-motor preference could partially explain intellectual differences in individuals with ASD and may serve as a target for therapies aimed at improving skill learning through visual observation and imitation. The HaptiKart task, requiring minimal instructions ("keep the car on the track centerline") and being child-friendly, shows promise for reliably assessing sensory-motor preferences in the diverse ASD population. The strong link between elevated proprioceptive vs. visual bias and lower IQ also suggests that intervention approaches that aim to increase reliance on dynamic visual feedback and stimuli, particularly in early development, may enhance cognitive development.

HaptiKart's ability to assess sensory-motor preferences makes it a valuable asset for clinicians and therapists. By identifying elevated proprioceptive vs. visual bias, HaptiKart can help tailor interventions that focus on increasing reliance on visual feedback. This approach aligns with evidence-based autism therapies, such as Applied Behavior Analysis (ABA) and Naturalistic

Developmental Behavioral Interventions, where therapists help guide children towards dynamic
visual stimuli to teach social skills (e.g., eye/facial/limb movements) and language skills (e.g.,
mouth movement).

Therapists can also use HaptiKart's quick assessments of sensory bias to develop therapy plans
to modify sensory bias and as an objective outcome measure of treatment response following
intervention. For children showing elevated proprioceptive vs. visual bias, digital interventions
can be designed to enhance visual feedback reliance by introducing inaccuracies in
proprioceptive feedback, such as gradually reducing the gain of proprioceptive feedback. By
fine-tuning visual and proprioceptive feedback, therapists can create diverse therapy goals
tailored to individual needs. HaptiKart could be used to deliver interventions aimed at reducing
anomalous sensory-motor bias at-home, reducing costs, and increasing access of services to
families living in underserved and rural communities. Further, HaptiKart offers an objective
measure of sensory bias, enabling therapists to monitor progress and adjust treatments as needed.
Its digital format also ensures that these interventions are accessible across different cultures,
unaffected by language barriers.

HaptiKart's quick measure of proprioceptive vs. visual bias can also provide clinicians and
therapists information on how best to instruct a child on a new skill. For instance, children with
elevated proprioceptive vs. visual bias may benefit from a "hands-on" manual guidance approach
when learning new skills (e.g., a therapist manually guides a child's arm towards the target) as
compared to using visual demonstration of the desired action. Therefore, HaptiKart has strong
potential as an assessment and digital intervention tool that can be easily integrated within
clinical settings to aid development of intervention plans to address anomalous sensory-motor
biases.
254
While allowing for more targeted therapeutic intervention, future iterations of HaptiKart's
assessment of proprioceptive/tactile vs. visual bias may be extended to examine other types of
sensory-motor biases such as proprioceptive vs. auditory and tactile vs. visual that can inform
clinical practice. Further, an adapted version of HaptiKart could be extended to toddlers (1-3
years) to better understand typical developmental trajectories of sensory-motor bias with the goal
of improving early identification and support for affected children during the most sensitive
developmental period.
262
One limitation of our study is the lack of data on participants' prior experiences with intervention
services, which could influence sensory preferences. Future research should explore how such
experiences relate to proprioceptive and visual biases. Additionally, we did not include a control
group, limiting our ability to assess the specificity of these biases to ASD. Future studies should
include clinical control groups for comparison. Another limitation is the single-session design,
which did not allow us to assess the test-retest reliability of HaptiKart's measures. Establishing
this reliability is crucial for clinical utility. Some participants were excluded for hand removal
from the wheel; future iterations of HaptiKart will include sensors to capture this behavior.
271
272 In summary, our study using HaptiKart demonstrated significant proprioceptive vs. visual bias in
273 children with ASD compared to typically developing controls. The brief assessment format
274 (approximately 5 minutes) has strong potential for clinical and home applications to track
275 sensory preferences throughout development and guide interventions aimed at enhancing

preference for dynamic visual feedback while reducing reliance on proprioceptive feedback in individuals with ASD.

Materials and Methods

Participants

A total of 81 participants 8-31 years old (33 ASD and 48 TD) participated in the study and met criteria for inclusion. Groups did not differ in age (ASD: 14.66 ± 6.28 years; TD: 14.49 ± 5.64 years; p = 0.9) or sex (ASD: 69% male, TD: 50% male; χ^2=3.11, p = 0.07) (**Table 1**). Not included in these participant totals is one outlier in the ASD group with a bias score exceeding 3 standard deviations (SD) from the ASD group mean. An additional ASD participant was removed due to an outlier IQ score approximating 3 SD below the ASD mean. Additional participants not included in the study were eight participants (6 ASD and 2 TD children) that removed their hands from the wheel during gameplay. These participants had to be excluded as there was no objective way to determine at what interval their hands were removed from the wheel during gameplay.

All participants in the ASD group received a clinical diagnosis of ASD based on DSM-5 criteria, as confirmed by a pediatric neurologist with decades of experience working with individuals with ASD across both clinical and research settings (SHM). Child participants with ASD (8 to 17 years; n=25) were required to have a prior diagnosis of ASD, score within the autism spectrum range on the Autism Diagnostic Observation Schedule, 2nd edition (ADOS-2), and score within the clinical range on the Social Responsiveness Scale, 2nd edition (SRS-2). Adult participants (>17 years; n=8) were required to have a prior clinical diagnosis of autism or ASD and score within the clinical range on the SRS-2. Further, the ADOS-2 Comparison Score (ADOS-CS) and SRS-2 were combined to analyze the clinical severity of ASD.

Additional data collected on subsets of participants included IQ General Ability Index (GAI) measured using the Wechsler Intelligence Scale for Children 5th Edition (WISC-V) (participants < 12 years old) or Wechsler Abbreviated Scale of Intelligence 2nd Edition (WASI-II) (participants >13 years old). For participants receiving the WISC-V, GAI was calculated using the Verbal Comprehension Index and Fluid Reasoning Index and converted to a composite score. For the participants completing the WASI-II, the full-scale IQ score calculated from Verbal Comprehension and Perceptual Reasoning subscales was used to calculate GAI. ADHD severity was assessed using the parent version of the Conners 4th edition (Conners4) total score, derived from the inattention and hyperactivity/impulsivity subscales of the Conners4 and standardized using T-scores [18].

Ethics approval was received from the Johns Hopkins University School of Medicine Institutional Review Board before the study (IRB00269589). Written informed consent was obtained from all adult participants, legal guardians of each child participant, and verbal assent from all children.

314 **Table 1: Participant demographics for all participants included in analysis.**

	TD (n = 48)	ASD (n = 33)	Test Statistic
Age (years)	14.49 ± 5.64	14.66 ± 6.28	p = 0.90
Sex (% male)	50%	70%	χ^2=3.11, p = 0.07
GAI (TD: 38, ASD: 29)	116.11 ± 15.08	108.66 ± 16.85	p = 0.06
SRS-2 Total Score (TD: 22, ASD: 20)	45.45 ± 4.81	70.30 ± 11.39	p < 0.0001
Conners4 Total Score (TD: 16, ASD: 13)	42.56 ± 4.03	62.92 ± 8.23	p < 0.0001
ADOS-2 CS (Module 3) (ASD: 20)	-	6.85 ± 1.50	-
ADOS-2 Total Score (Modules 3 and 4) (ASD: 26)	-	11.15 ± 3.07	-

315 TD: Typically developing
316 ASD: Autism spectrum disorder
317 GAI: General Ability Index
318 SRS-2: Social Responsiveness Scale, 2nd Edition
319 ADOS-2 CS: Autism Diagnostic Observation Schedule, 2nd Edition Comparison Score
320

Experimental Design

HaptiKart Procedures

323
Participants played the custom-built HaptiKart videogame in a quiet, light-controlled room using
a Logitech G29 steering wheel. The game was run on an MSI GS66 Stealth laptop and displayed
on a Dell monitor adjusted to eye level. The wheel was affixed to a height-adjustable stand, with
participants instructed to maintain their hands at a standardized 10 o'clock (left hand) and 2
o'clock (right hand) position (**Fig 1**). They were directed to "drive the car on the centerline"
without awareness of any visual or steering perturbations. To ensure data reliability, participants
who exhibited challenging behavior—specifically, removing their hands from the wheel during
gameplay—were excluded from the analysis, as there was no objective measure for when hand
release occurred. Six children with ASD and two TD children displayed this behavior and were
not included in the analyses of participant characteristics or sensory preferences.

Measurement of Driving Error

335
336 During gameplay, errors were measured continuously as the mean of two error types: (1) the
337 absolute distance of the car relative to the track centerline (position error, PE) that was expressed
338 as a percentage of maximum lateral error (6 meters) and (2) the absolute orientation of the car
339 relative to the track centerline (heading error, HE) that was expressed a percentage of the

maximum orientation error (+/- 90-degrees) (**Fig 5**). The absolute relative position and heading errors were averaged across all laps within each block to produce a single error metric (Mean Total Error Percentage (%) = 0.5 x [ABS(PE)/6] + [ABS(HE)/90]).

Fig 5. Error calculation. Continuous calculation of heading and position error used for proprioceptive feedback.

Visual-error feedback

The visual-error feedback (**Fig 6A**) was presented as the car's position relative to the track centerline. While other forms of error feedback could have been utilized—such as color changes or symbols indicating the direction and magnitude of error—the continuous feedback of the kart's position relative to the centerline was most akin to the ongoing nature of proprioceptive feedback from the steering wheel. This similarity is crucial for quantifying proprioceptive vs. visual bias because it allows for a direct comparison between the sensory modalities. By using a feedback mechanism that mimics the continuous flow of proprioceptive-error feedback, we can better assess how children with ASD integrate these two types of feedback. This approach ensures that any performance differences between the feedback delay conditions are more accurately attributed to differences in sensory bias rather than differences in the nature of the feedback provided.

Proprioceptive-error feedback (force feedback wheel)

The two error types (HE and PE) were used to provide proprioceptive-error feedback, felt as a corrective torque or "steering assist" applied to the wheel, that continuously corrected steering by guiding the car toward the track centerline. The gain constant (k) on the "steering assist" was multiplied by each error type (HE and PE) to adjust the resting angle of the centering spring of

369   the steering wheel (**Eq. 1, Fig 6B**). The gain constant was selected to provide some "steering
370   assistance" while still requiring steering input from the user to stay on the centerline.
371

$$Centering\ Spring\ Angle = k * (Heading\ Error + Position\ Error) \tag{1}$$

A. Visual-Error Feedback

B. Proprioceptive-Error Feedback

374   **Fig 6. Visual and proprioceptive-error feedback.** (A) Visual display seen by the participant
375   during gameplay. (B) Method of delivering corrective proprioceptive-error feedback by using
376   driving errors to update wheel centering spring angle (perceived as "steering assist"). Right plot
377   in panel B shows opposing relationship between driving error and the corrective proprioceptive-
378   error feedback.

Temporally Delayed Feedback

The paradigm to examine proprioceptive vs. visual bias involved selectively delaying proprioceptive and visual feedback in separate conditions to measure relative preferences for each feedback modality. The selected visual delay was 250 milliseconds (ms), and the proprioceptive delay was 300 ms. The delays were selected such that they would be reliably detected by young children [19]. Previous findings in children as young as five years old show that visual-proprioceptive asynchrony detection thresholds continue to increase up to 300 ms [19]. Pilot testing in a small number of children revealed that a visual delay of 300 ms may be too challenging given the visual dominance of the driving task, so a lesser 250 ms visual delay was selected.

Experimental Paradigm

The paradigm used to measure proprioceptive vs. visual feedback was completed as part of a larger paradigm that included several sensory conditions, completed in the same order for all participants, and separated by washout blocks. The washout blocks provided veridical visual and proprioceptive feedback to mitigate crossover effects between conditions with varied sensory feedback. Below, we describe a "brief" paradigm (~5-min) within the larger paradigm that was designed to examine proprioceptive vs. visual bias while controlling for both condition order (i.e., counterbalancing) and crossover effects (i.e., using washout blocks).

Following several different sensory feedback conditions that all participants completed in a set order, each participant completed the brief paradigm to examine proprioceptive vs. visual bias. The brief paradigm included three conditions: (1) visual delay (*DV+P*; 6 laps), (2) washout (3 laps), and (3) proprioceptive delay (*V+DP*; 6 laps) (6 laps) (**Fig 2**). To account for potential order effects on the measurement of proprioceptive vs. visual bias, the order of the delay conditions was counterbalanced within each group such that an equal proportion of participants completed the visual delay condition first and the other half completed the proprioceptive delay condition first.

Calculation of Proprioceptive vs. Visual Bias

Proprioceptive vs. visual bias was calculated as the difference in mean total error between the delayed proprioceptive feedback (*V+DP*) and delayed visual feedback (*DV+P*) conditions, as shown in Equation (2), where a positive score (>0) indicates proprioceptive vs. visual bias and a negative score (<0) indicates visual vs. proprioceptive bias (**Fig 7**).

$$Bias\ score = V{+}DP\ error\ \text{-}\ DV{+}P\ error \qquad (2)$$

420 **Fig 7**. **Calculation of perceptual bias scores**. Visualization of the two delay conditions and
421 interpretation of the bias score.

Statistical analysis

To examine the interaction effects between bias scores and age, an ANCOVA was conducted to
assess the interaction effect of diagnosis and age on bias scores while controlling for biological
sex as a covariate (bias score ~ diagnosis*age + sex). Overall main effects of diagnosis and age
on bias scores were also evaluated. Simple linear regression was used to examine the correlation
between bias scores and core autism symptoms (ADOS-CS and SRS-2 total score), ADHD
symptoms (Conners4 total score), intellectual ability (GAI), and age. Significance for all tests
was set at an alpha level of 0.05, and false discovery rate correction was applied to the p-values
in the correlation analysis. All statistical analyses were conducted using RStudio version
2021.09.1. All assumptions for ANCOVA were met including homogeneity of variance
(Levene's Test: $p = 0.7$), and normality of residuals (Shapiro-Wilk; $p = 0.10$). Homogeneity of
slopes was not assessed separately, as it is part of the hypothesis being tested.

Acknowledgements

439 We sincerely thank our participants and their families for their time, effort, and support, which
440 made this study possible.

References

1. Lidstone DE, Mostofsky SH. Moving Toward Understanding Autism: Visual-Motor Integration, Imitation, and Social Skill Development. Pediatr Neurol. 2021;122: 98–105. doi:10.1016/j.pediatrneurol.2021.06.010

2. Mostofsky SH, Ewen JB. Altered connectivity and action model formation in autism is autism. Neuroscientist. 2011;17: 437–448. doi:10.1177/1073858410392381

3. Ament K, Mejia A, Buhlman R, Erklin S, Caffo B, Mostofsky S, et al. Evidence for Specificity of Motor Impairments in Catching and Balance in Children with Autism. J Autism Dev Disord. 2015;45: 742–751. doi:10.1007/s10803-014-2229-0

4. De Francesco S, Morello L, Fioravanti M, Cassaro C, Grazioli S, Busti Ceccarelli S, et al. A multimodal approach can identify specific motor profiles in autism and attention-deficit/hyperactivity disorder. Autism Research. 2023. doi:10.1002/aur.2989

5. Lidstone DE, Miah FZ, Poston B, Beasley JF, Mostofsky SH, Dufek JS. Children with Autism Spectrum Disorder Show Impairments During Dynamic Versus Static Grip-force Tracking. Autism Research. 2020;13: 2177–2189. doi:10.1002/aur.2370

6. Tunçgenç B, Pacheco C, Rochowiak R, Nicholas R, Rengarajan S, Zou E, et al. Computerized Assessment of Motor Imitation as a Scalable Method for Distinguishing Children With Autism. Biol Psychiatry Cogn Neurosci Neuroimaging. 2021;6: 321–328. doi:10.1016/j.bpsc.2020.09.001

7. Landa RJ, Haworth JL, Nebel MB. Ready, Set, Go! Low Anticipatory Response during a Dyadic Task in Infants at High Familial Risk for Autism. Front Psychol. 2016;7: 721. doi:10.3389/fpsyg.2016.00721

8. Gepner B, Mestre D, Masson G, de Schonen S. Postural effects of motion vision in young autistic children. Neuroreport. 1995;6: 1211–1214. doi:10.1097/00001756-199505300-00034

9. Greffou S, Bertone A, Hahler EM, Hanssens JM, Mottron L, Faubert J. Postural hypo-reactivity in autism is contingent on development and visual environment: A fully immersive virtual reality study. J Autism Dev Disord. 2012;42: 961–970. doi:10.1007/s10803-011-1326-6

10. Hirai M, Sakurada T, Izawa J, Ikeda T, Monden Y, Shimoizumi H, et al. Greater reliance on proprioceptive information during a reaching task with perspective manipulation among children with autism spectrum disorders. Sci Rep. 2021;11. doi:10.1038/s41598-021-95349-0

11. Izawa J, Pekny SE, Marko MK, Haswell CC, Shadmehr R, Mostofsky SH. Motor learning relies on integrated sensory inputs in ADHD, but over-selectively on proprioception in autism spectrum conditions. Autism Research. 2012;5: 124–136. doi:10.1002/aur.1222

12. Takamuku S, Ohta H, Kanai · Chieko, De AF, Hamilton C, Gomi H. Seeing motion of
controlled object improves grip timing in adults with autism spectrum condition: evidence
for use of inverse dynamics in motor control. Exp Brain Res. 2021;239: 1047–1059.
doi:10.1007/s00221-021-06046-3

13. Glazebrook CM, Gonzalez D, Hansen S, Elliott D. The role of vision for online control of
manual aiming movements in persons with autism spectrum disorders. Autism. 2009;13:
411–433. doi:10.1177/1362361309105659

14. Haswell CC, Izawa J, Dowell LR, Mostofsky SH, Shadmehr R. Representation of internal
models of action in the autistic brain. Nat Neurosci. 2009;12: 970–2. doi:10.1038/nn.2356

15. Marko MK, Crocetti D, Hulst T, Donchin O, Shadmehr R, Mostofsky SH. Behavioural
and neural basis of anomalous motor learning in children with autism. Brain. 2015;138:
784–797. doi:10.1093/brain/awu394

16. Green D, Charman T, Pickles A, Chandler S, Loucas T, Simonoff E, et al. Impairment in
movement skills of children with autistic spectrum disorders. Dev Med Child Neurol.
2009;51: 311–316. doi:10.1111/j.1469-8749.2008.03242.x

17. Surgent OJ, Walczak M, Zarzycki O, Ausderau K, Travers BG. IQ and Sensory Symptom
Severity Best Predict Motor Ability in Children With and Without Autism Spectrum
Disorder. J Autism Dev Disord. 2021;51: 243–254. doi:10.1007/s10803-020-04536-x

18. Conners CK. Conners third edition (Conners 3). Los Angeles, CA: Western Psychological
Services. 2008.

19. Jaime M, Longard J, Moore C. Developmental changes in the visual-proprioceptive
integration threshold of children. J Exp Child Psychol. 2014;125: 1–12.
doi:10.1016/j.jecp.2013.11.004