Identifying Opportunities for Fluid Balance Optimization in Critically Ill Children

Denise Hasson, MD[1]; Ami Shah, MD[1]; Chloe G. Braun, MD[2]; Ulka Kothari, MD[3]; Steve Drury, MS[4]; Heda Dapul, MD[1]; Julie C. Fitzgerald, MD, PhD, MSCE[5]; Celeste Dixon, MD[5]; Andrew Barbera, MD[2]; James Odum, MD[2]; Nina Terry, BSN[6]; Scott L. Weiss, MD, MSCE[7]; Susan D. Martin, MD[6]; Adam C. Dziorny, MD, PhD [4,6]

[1] Dept of Pediatrics, Hassenfeld Children's Hospital at NYU Langone Health, NY, NY

[2] Dept of Pediatrics, University of Alabama at Birmingham, Birmingham, AL

[3] Dept of Pediatrics, NYU Long Island School of Medicine, Mineola, NY

[4] University of Rochester Health Lab, University of Rochester Medical Center, Rochester, NY

[5] Dept of Anesthesiology & Critical Care Medicine, Children's Hospital of Philadelphia, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA

[6] Dept of Pediatrics, Golisano's Children's Hospital, University of Rochester School of Medicine, Rochester, NY

[7] Division of Critical Care, Dept of Pediatrics, Nemour's Children's Hospital, Wilmington, DE

Corresponding Author:

Denise Hasson, MD

Denise.Hasson@nyulangone.org

430 E 34th St

NY, NY 10016

646-341-2376

Running headline: Fluid Balance Optimization in PICU Patients

Abstract

Introduction: Fluid overload (FO), a state of pathologic positive cumulative fluid balance (CFB), is common in Pediatric Intensive Care Units (PICU) and associated with morbidity and mortality. Because different PICUs may have unique needs, barriers, and limitations to accurately report fluid balance (FB) and reduce FO, understanding the drivers of positive FB is needed. We hypothesize CFB >5% and >10% is common within initial days of PICU admission, but that reasons for high %CFB will vary across sites, as will barriers to accurate FB recording and opportunities to improve FB recording and management.

Methods: Concurrent mixed methods study utilizing a retrospective observational cohort design and prospective interview and survey design performed at four tertiary pediatric ICUs. FB data was extracted from the electronic health record. A federated data collection framework allowed for rapid data aggregation. The primary outcome was %CFB on PICU days 1 and 2, defined as total intake minus total output divided by PICU admission weight. Chi-square test and Wilcoxon rank sum tests compared results across and within sites.

Results: Amongst 3,071 PICU encounters, day 2 CFB >5% varied from 39% to 54% (p=0.03) and day 2 CFB >10% varied from 16% to 25% (p=0.04) across sites. Urine occurrence recordings and patients receiving >100% Holliday-Segar fluids on Day 1 differed across sites (p<0.001). Sites discussed overall FB and specific FB goals on rounds with differing frequency (42-73% and 19-39%, respectively), but they reported similar barriers to accurate FB reporting and achievable opportunities to improve FB measurements.

Conclusion: Day 2 CFB >5% and >10% was common among PICU encounters but proportion of patients varied significantly across PICUs. Individual ICUs have different drivers of FO that must be targeted to improve FB management.

Key Words: fluid overload; fluid management; fluid accumulation; acute kidney injury; federated data collection

Introduction

Fluid overload (FO), a state of pathologic positive cumulative fluid balance (CFB), is common in children with critical illness and is associated with prolonged mechanical ventilation, increased risk of acute kidney injury (AKI), increased intensive care unit (ICU) length of stay (LOS), morbidity, and mortality.[1] Although early literature associating FO with morbidity and mortality focused on critically ill children requiring renal replacement therapy (RRT),[2–5] most pediatric ICU (PICU) patients with FO have similar associations with poor outcomes.[6] A 2018 meta-analysis found that %CFB >5% within 24 hours of PICU admission was associated with nine-fold greater odds of mortality, while peak %CFB >10% during PICU admission was associated with 15-fold greater mortality.[1] Awareness of this excess mortality has led to awareness of the need for unified definitions[1] and increased interest in strategies to prevent, mitigate, and reduce positive %CFB.

Recent international multicenter studies on the impact of volume and temporality of this positive %CFB showed that small amounts (≥5%) early in the ICU course (Day 1 and Day 2) are associated with the aforementioned negative ICU outcomes.[1,6] One approach toward solving this problem is to develop a clearer understanding of the key drivers of fluid exposure.[6] Studies have begun to identify reasons behind positive %CFB, such as fluid creep (i.e., the administration of non-nutritive, non-resuscitative fluids)[7] and have found associations between this metric and negative PICU outcomes.[8,9] However, we lack a holistic, multidisciplinary, and multicenter approach to investigating this problem, including identification of other key drivers of fluid exposure, barriers to accurate fluid balance (FB) reporting, and targeted fluid deresuscitation.

Nuances in FB management vary across institutions and even between units within the same institution. Unique challenges to optimizing fluid sensitive care include accurate FB reporting, lack of structured discussion and goal setting around FB, and varied opportunities and actions to reduce %CFB. The objective of this study was to quantify the frequency of positive

%CFB amongst critically ill patients in our multicenter collaborative, and qualitatively identify areas for improvement of fluid sensitive care. We hypothesize that CFB >5% and >10% is common within initial days of PICU admission, but that reasons for high %CFB will vary across sites, as will barriers to accurate FB recording and opportunities to improve FB recording and management.

Materials and Methods

Study Design

This is a concurrent mixed methods study conducted within a collaborative composed of four tertiary care centers (with four PICUs and one pediatric cardiac ICU) spanning 148 ICU beds. We performed a quantitative retrospective observational cohort study to measure FB metrics and ICU outcomes from all new pediatric ICU admissions in the 6 months prior to qualitative data collection, between April 2022 and January 2023. All sites use the same Electronic Health Record (EHR), Epic (Epic Systems Corporation, Verona, WI). We also performed a prospective observational survey using qualitative methods, including interviews and surveys with bedside nurses to understand current ICU-specific practices and barriers. This study followed procedures in accordance with ethical standards as delineated by the Helsinki Declaration of 1975. Each site's institutional review board deemed the project exempt from IRB review.

We used a federated data collection framework to collect and aggregate clinical data.[10] We wrote data extraction queries, analysis scripts, survey data dictionaries, and semi-structured interview scripts at a single site and disseminated each to all sites to promote process standardization. This allowed for site-specific understanding of FB management while still allowing for comparison and collaboration. No row-level (i.e., individual patient level) data were shared between sites.

Baseline Measures and Definitions

We collected metrics including baseline demographics, admission weight, height, total intake and output by route and type, urine occurrence counts (e.g. urine, stool, or mixed output that was not weighed or measured as an exact volume), number of patients with invasive mechanical ventilation orders (as a surrogate for use of invasive mechanical ventilation), and ICU LOS and hospital disposition. Total intake and output included all measures noted to be "intake" or "output" within the EHR. We harmonized input and output types among sites where necessary. Consistent with prior studies, we attributed fluid measures to ICU days as follows: Day 0 was defined as admission through the next 7 AM, and each subsequent ICU day (1 through 7) included all fluid measures from 7 AM through the next 7 AM or ICU discharge.[6,11]

We calculated %CFB as:[12]

$$\%CFB = \frac{\text{cumulative fluid in (L) - cumulative fluid out (L)}}{\text{PICU admission weight (kg)}} \times 100$$

We calculated the ratio of fluids administered to Holliday-Segar (H-S) maintenance fluid rates as: (the sum of all IV fluids including parenteral nutrition) / (4x first 10 kg weight + 2x number of kg between 10-20 + 1x number of kg >20).[13]

Interview and Survey Based Assessment of ICU FB Practice

We followed a multistep survey creation framework to identify site-specific barriers to facilitators to improve accurate FB reporting.[14] We wrote a question script based on goals we hoped to achieve (**Supplementary Methods**), then we interviewed 3-5 nurses at Sites A and B. These answers informed the questions and responses that we incorporated into the Research Electronic Data Capture (REDCap) Nurse Survey (**Supplementary Methods**). After drafting specific questions with informed multiple-choice answers, we emailed surveys to bedside nurses at each site and allowed them a month to complete this survey. We shared data dictionaries and implemented the same survey in each site-specific REDCap instance. Nurses

were asked to choose the three best answers for questions 1 and 2, and the single best answer for question 3 of the REDCap survey.

We measured the frequency and content of discussions related to FB, recognizing that providers cannot address an issue they 1) are unaware of and 2) do not discuss. We focused on discussions during ICU rounds, as this is when daily plans and goals are established in collaboration with a multidisciplinary team. Sites A, B, and C interviewed available nurses 1-4 hours after rounds (**Supplementary Methods**) 2-3 times weekly for 4-6 weeks to account for the varied practices of different providers. Site D employed a research assistant that participated on daily PICU rounds (Monday-Friday) to observe and record the frequency of FB discussions for 6 weeks. We recorded observation results in a standard REDCap instance or shared Excel spreadsheet (Microsoft, Redmond, WA, USA) and aggregated results across sites.

Statistical Analysis

We performed clinical data analyses at each individual site using analysis scripts written in R (https://www.r-project.org, R Foundation for Statistical Computing, Vienna, Austria). Results are presented by institution. We reported categorical data in frequencies (%) over a given period (e.g., one week or 7 days) and aggregate data as median (interquartile range, IQR). We compared counts across and within sites using Chi-square and compared distributions within sites using Wilcoxon rank sum tests. Baseline fluid metric comparisons by site excluded patients with urine "occurrences," as such occurrences suggest that urine or stool volume was not recorded, thus biasing toward a more positive %CFB than what may have occurred. However, we also performed sensitivity analyses including encounters with urine occurrences without calculating or extrapolating missing data. We reported descriptive statistics of survey results at each site and aggregated them across all sites.

Results

Fluid Balance Metric Comparisons by Site

Over 6 months, we included 3,071 ICU encounters and 2,850 hospitalizations encompassing 2,632 unique patients (**Table 1**). Excluding patients with urine occurrences charted, CFB >5% was common early in the ICU course, occurring in 34% (N=697/2,064) of ICU encounters on Day 1 and 47% (N=670/1,437) of ICU encounters on Day 2. Ten percent (N=205/2,064) of ICU encounters had >10% CFB on Day 1 and 20% (N=292/1,437) had >10% CFB on Day 2. Total number (%) of ICU encounters with Day 2 CFB >5% and >10% significantly differed by site (**Table 1**, p=0.03 and p=0.04, respectively). Day 1 CFB >5% and >10% did not differ between sites (**Table 1**, p=0.15 and p=0.79, respectively), with sites having 5% or less difference between incidence of CFB >5% and 3% or less difference for CFB >10%. However, when patients with urine occurrences were included, day 1 CFB >5% also differed by site (**Supplementary Table S1**).

Instead of measuring all total IVF volume measured (excluding enteral nutrition and blood but including parenteral nutrition, as some EHRs do not differentiate the latter from IVF), we chose to measure total number (%) of ICU encounters where patients' total IVF was >100% of H-S fluid volume. Sites varied in the number of ICU encounters where >100% of H-S fluids were prescribed on Day 1 but not on Day 2 (**Table 1**, p < 0.001 and p=0.21, respectively). All sites had at least 20% of encounters where >100% H-S fluids were prescribed on both days, but on day 1, multiple sites approached 30% of encounters with >100% H-S fluids prescribed. Given the importance of accurate FB recording to these metrics, we measured the number of encounters that had a urine occurrence documented. The frequency of encounters with urine occurrences varied by site and ranged from 15.5% to 50.7% (**Table 1**, p <0.001).

Table 1: Baseline Fluid Metric Comparisons by Site, Not Including Urine Occurrences

Fluid Metric	Site A	Site B	Site C	Site D	Chi Square

	(N=427)	(N=553)	(N=1356)	(N=735)	Test
Day 1 CFB >5%**	61 / 178 (34%)	133 / 427 (31%)	344 / 950 (36%)	159 / 509 (31%)	X2=5.32 P=0.15
Day 1 CFB >10%**	14 / 178 (7.9%)	45 / 427 (11%)	96 / 950 (10%)	50 / 509 (9.8%)	X2=1.06 P=0.79
Day 2 CFB >5%**	48 / 89 (54%)	113 / 287 (39%)	351 / 732 (48%)	158 / 329 (48%)	**X2=8.75 P=0.03**
Day 2 CFB >10%**	22 / 89 (25%)	46 / 287 (16%)	166 / 732 (23%)	58 / 329 (18%)	**X2=8.31 P=0.04**
Urine Occurrences	667 / 1315 (51%)	292 / 1618 (18%)	694 / 4479 (16%)	364 / 1912 (19%)	**X2=775.8 P < 0.001**
Day 1 H-S fluids >100%	66 / 335 (20%)	124 / 433 (29%)	341 / 1096 (31%)	181 / 646 (28%)	**X2=16.5 P < 0.001**
Day 2 H-S fluids >100%	40 / 190 (21%)	65 / 235 (28%)	188 / 709 (27%)	73 / 327 (22%)	X2=4.58 P=0.21

Cumulative fluid balance (CFB), urine occurrence measurements, and IVF intake above H-S calculation for each site. Site-specific sample size in header row reflects total ICU admissions over the study period. Each cell reports number of patient-days (%) meeting fluid metric. Cell-specific denominator reflects total number of ICU patients present on that day of ICU stay, or during any day if no day is specified. H-S fluids metric only includes patients with IVF intake. We report row-wise chi-square p-value to examine the relation between fluid metric and site. ** indicates only those samples without urine occurrences. Blue indicates favorable differences; red indicates unfavorable differences. P-values <0.05 represented with bolding. CFB-cumulative fluid balance, H-S- Holliday-Segar.

Mechanical ventilation orders were more common in those encounters that had >100% of H-S fluids prescribed at all sites on Day 1 (A: 28.5% vs 13.3%, p<0.001; B: 40.3% vs 23.7%, p=0.001; C: 49.8% vs 23.4%, p<0.001; D: 43.9% vs 21.0%, p<0.001). Mechanical ventilation orders were more common in those with >100% of H-S fluids prescribed at sites C and D on Day 2 but not sites A and B (A: 25.0% vs 15.4%, p=0.11; B: 34.0% vs 23.4%, p=0.07; C: 43.0% vs 15.8%, p<0.001; D: 33.8% vs 12.5%, p<0.001) at each site (**Table 2**). The lack of difference seen for sites A and B on Day 2 could have been partially due to smaller sample sizes on this day; however, it is worth noting that Site A had a lower number of encounters with >100% of H-S fluids prescribed in both groups compared to the other sites.

Table 2: Frequency of Encounters with >100% Holliday-Segar Fluids by Mechanical Ventilation Status, by Site

	Site A	Site B	Site C	Site D

	+MV	-MV	+MV	-MV	+MV	-MV	+MV	-MV
ICU Day 1, N (%)	**40/140 (28.6%)**	**26/195 (13.3%)**	**52/129 (40.3%)**	**72/304 (23.7%)**	**159/319 (49.8%)**	**182/777 (23.4%)**	**87/198 (43.9%)**	**94/448 (21.0%)**
ICU Day 2, N (%)	28/112 (25.0%)	12/78 (15.4%)	32/94 (34.0%)	33/141 (23.4%)	**120/279 (43.0%)**	**68/430 (15.8%)**	**51/151 (33.8%)**	**22/176 (12.5%)**

Measured the association of MV with >100% H-S fluids on each ICU day, by site, using Chi Square Test. Bolded values indicate comparisons are statistically significant with a P-value < 0.001. MV-mechanical ventilation, ICU-intensive care unit.

In unadjusted analyses excluding patients with urine occurrences, ICU LOS was significantly longer in patients with Days 1 and 2 CFB >5% at all sites (**Table 3**, p < 0.01). ICU LOS was significantly longer in patients with Days 1 and 2 CFB >10% at three of four sites, associated with a median of 1.5-2.5 greater ICU days (**Table 3)**.

Table 3: Intensive Care Unit Length of Stay by % Cumulative Fluid Balance Metric, by Site

	Site A		Site B		Site C		Site D	
	> CFB	< CFB	> CFB	< CFB	> CFB	< CFB	> CFB	< CFB
ICU Day 1, 5 % cut-off	**4.5 (2.2,8.6)**	**1.0 (0.6,3.6)**	**2.2 (1.6,4.2)**	**1.6 (0.9,2.8)**	**2.9 (1.7,6.2)**	**2.1 (1.1,4.8)**	**2.6 (1.4,5.9)**	**1.3 (0.8,2.6)**
ICU Day 1, 10% cut-off	3.2 (2.3,5.8)	1.9 (0.8,4.9)	3.5 (2.0,9.2)	1.8 (1.0,2.8)	**4.1 (2.4,6.9)**	**2.3 (1.1,4.9)**	**3.1 (1.9,7.1)**	**1.5 (0.9,3.0)**
ICU Day 2, 5 % cut-off	4.7 (2.5,8.3)	3.7* (1.6,4.9)	**2.8 (1.8,5.2)**	**1.9 (1.3,3.3)**	3.8 (2.1,7.4)	2.5 (1.6,5.1)	3.3 (1.9,7.1)	2.0 (1.3,3.6)
ICU Day 2, 10% cut-off	4.7 (2.5,6.9)	4.0 (1.8,6.3)	4.0 (2.2,9.2)	1.9 (1.5,3.4)	**4.3 (2.8,7.7)**	**2.7 (1.6,5.4)**	**4.9 (2.7,8.9)**	**2.2 (1.4,4.1)**

Median (interquartile range) length of stay in days for patients who did not meet cut-offs of 5% and 10% CFB compared to those who did, by site, excluding urine occurrences. Wilcoxon rank sum tests determined P values, with P values <0.01 represented by bolding, P value <0.05 indicated with *. CFB-cumulative fluid balance, LOS-length of stay.

Barriers and Opportunities to Improve Accurate FB Recording

All sites had the same top two Most Frequent Barriers to Accurate FB Recording (**Figure 1a, Supplementary Table S2**): "ambulatory patients not saving urine/stool," and "patients not keeping track of oral intake." Site D respondents reported that "standardizing charting of

medication flushes could be helpful," while the other three sites reported "families throwing diapers away before being weighed as an important barrier." All four sites had two of the same four Achievable Opportunities to Accurate FB Recording (**Figure 1b, Supplementary Table S2**): "standardizing recording of flush volumes" and "collaborating with other units to add fluids into flowsheets." Three of four sites shared the other top two opportunities: "standardizing flush volumes per medication" and "consistent zero-ing of bed scales for daily weights." When asked the most Feasible Method for FB Recording, all sites agreed that "setting a fluid balance goal on rounds and discussing ways to meet it" would be more feasible than "recording strict I/O without urine occurrence measurements" or "obtaining daily weights" (**Figure 1c, Supplementary Table S2**).

Setting FB Goals

Sites surveyed nurses on a median of 8.5 (8.0-46.5) days (total of 84 days) over a median of 6 (4.25-9.25) weeks (total of 26 weeks) (**Table 4**). All sites discussed FB on 62% (N=578) of patients' rounds, with Site C discussing FB most frequently (78%, N=93), p<0.001. A FB goal was only set on 23% (N=214) of patient rounds recorded, with Site B setting a goal most frequently (39%, N=26), p=0.005. Sites A and C discussed a FB goal just over 20% of the time (23%, N=52 and 21%, N=103). Sites A-C recorded whether changes were made to the FB plan, such as decreasing maintenance fluids or starting/changing diuretics. Changes were only made for 18% of patients (N=79/439), with sites making changes on 14-36% of patients with varying frequency (p<0.001).

Table 4: Assessment of Fluid Balance Practices on Rounds

	Site A (N=252)	Site B (N=67)	Site C (N=120)	Site D (N=494)	Total (N=933)	P value

Number of Days/Weeks Surveyed	9 d/ 7 wk	8 d/ 5 wk	8 d/ 4 wk	59 d/ 10 wk	84 d/ 26 wk	
Discussed Fluid Balance, N (%)	102 (40)	46 (69)	93 (78)	337 (68)	578 (62)	<0.001
Set a Fluid Balance Goal, N (%)	52 (21)	26 (39)	33 (28)	103 (21)	214 (23)	0.005
Made Fluid Balance Changes, N (%)	35 (14)	24 (36)	20 (17)	n/a	79 (8.5)	<0.001

Results of rounds' observations or post-rounds interviews of nurses to report discussions about fluid balance practices and goals. Site A included data from both pediatric and cardiac intensive care units. Site D did not gather data on whether there were changes to the plan to address fluid balance goals. d-days, wk-week(s). We report row-wise association of fluid balance discussion and site using Chi-square test.

Discussion

Across four tertiary PICUs, CFB >5% and >10% on PICU Days 1 and 2 were very common, with similar rates on Day 1 but varied rates on Day 2 across sites. Sites prescribed total IVF in excess of H-S calculations in 20-30% of ICU encounters, with varying frequency on Day 1 but not on Day 2. Sites routinely discussed FB on rounds, but setting goals and making changes based on said goals occurred <40% of the time. A major limitation to the study of FB is accuracy of FB measuring and recording. We identified that up to 50% of encounters can have urine occurrences charted, leading to inaccurate FB calculations. More importantly, though, we identified many barriers to accurate FB recording and opportunities to improve FB recording, with commonality between sites, but with enough variation to warrant site-specific targeted management.

The rates of positive %CFB we measured are very similar to rates reported by others. In a multicenter international collaborative study, Selewski et al identified CFB >5% on Day 1 and 2 in 38.1% and 53.3% of patients and CFB >10% on Day 1 and 2 in 11.7% and 25.1% of patients.[6] Another two-center study found Day 3 CFB >10% in 24.6% of patients,[8] similar to Day 2 %CFB at 2 of our sites. Both these studies identified strong associations between these FB cut-offs and fluid in excess of hydration requirements with fewer ventilator-free days, ICU-free days, and 28-day ICU mortality.[6,8] Although these aggregate data greatly add to the known

epidemiology of positive %CFB, especially as they pertain to outcomes, they do not discuss intra-unit variability. Specifically, they suggest that next steps should identify key drivers of such high %CFB, as this can inform next steps to reduce this incidence.[6]

Targeting "fluid creep," the administration of non-nutritive, non-resuscitation fluid in excess of maintenance fluid requirements, is a worthy strategy for FO prevention given its association with increased morbidity and mortality.[8,9] Barhight et al found that each 10mL/kg of fluid given in excess of hydration increases the odds of death by 1%.[8] A single center PICU was able to increase ventilator- and PICU-free days by restricting maintenance fluids to 40% compared to 70-80% in mechanically ventilated children.[15] Our data suggest that some sites reduce this "fluid creep" better than others, but that nearly a quarter of patients may be receiving more fluid than needed. This number is lower than others have reported, especially given certain populations are at higher risk for FO than others,[8,9] but we believe it is still too high given its association with adverse outcomes. Our team of bedside nurses have identified that setting a total fluid goal on rounds, and discussing ways to meet it, would be the most feasible method for accurate FB recording. Additionally, our data suggest that more work must be done to improve goal setting, and that providers must be cognizant of making adjustments to ensure those goals are met.

It is possible that a major barrier to improving FO incidence may be the difficulty in clinically recognizing positive %CFB. Even 15% CFB can be missed routinely by PICU providers.[7] Providers must then rely on more objective recorded data to know a patient's %CFB. Although advances in the EHR have brought extreme amounts of data to the clinicians' fingertips, it is dependent on and limited by the accuracy of the input data. Simple calculations can be run by the EHR to alert clinicians to unseen FO, but these are made inaccurate by modifiable barriers such as urine occurrences and unclear processes for charting fluids such as medication flushes, as well as non-modifiable barriers such as insensible fluid losses.[16] Although using daily weights can account for insensible losses,[4] there are barriers to accurate

daily weights in PICU patients, and our data suggest this may be difficult to incorporate into unit workflow. Technical barriers exist to building custom EHR-based scripts that perform complex but clinically useful calculations, such as %CFB. Overcoming these barriers may promote earlier clinical recognition of FO.

There are many strengths to this study. We established a robust multicenter collaborative to aggregate the large amount of granular data necessary for the study of FB without time-intensive manual chart review. Our FB metrics consider inaccuracies that occur when urine and stool are collected as occurrence counts (rather than measured volumes). We have sought to go beyond FB epidemiology, to investigate reasons for positive %CFB and to improve FB recognition by addressing barriers to accurate FB measurement and recording. By comparing practices between sites, we provide feasible solutions that address the nuances of FB management across sites. Our sites have size and geographic diversity, enhancing generalizability. We have leveraged the expertise of a multidisciplinary group of providers, most notably, our bedside nurses, who often know the most about how much fluid goes in and from which sources.

There are also study limitations. Our baseline data was retrospective, and we did not collect data on severity of illness to adjust for in our analyses. It is possible that patients with urine occurrences may reflect patients who are less critically ill and, therefore, in whom these inaccuracies are less clinically relevant. To account for this, we performed our retrospective analyses both including and excluding these encounters. To minimize the number of site-specific modifications and parameters within our shared queries and analysis platform, we were unable to collect granular respiratory support data, and instead collected ventilator orders as a surrogate for mechanical ventilation. We also did not share row-level data among sites, which limited our ability to compute multivariable regression models across sites. As with any survey, there likely was selection bias in those subjects who chose to respond. We attempted to

overcome this limitation by including multiple sites to recruit a varied nursing population, but biases may have been present within each site.

Next steps involve measuring acute kidney injury (AKI) frequency to better understand the interplay between AKI and FB in our population. We plan to implement site-specific opportunities to improve FB recording and management to reduce CFB amongst the collaboratives' ICUs. We will also be implementing novel EHR-based calculations across sites to assist all team members in identifying positive %CFB earlier.

Conclusion

Across our multisite collaborative, CFB >5% and >10% was common within the first 2 PICU days but varied across sites. Workflow practices to address positive %CFB varied significantly across sites despite similar barriers to accurate FB recording and opportunities to reduce %CFB. Such opportunities will be the targets of upcoming prospective interventional trials to reduce %CFB and assess ability to improve ICU outcomes.

Supplementary Materials

Supplementary Methods (PDF)

Supplementary Tables (PDF)

Supplementary Table S1. Baseline Fluid Metric Comparisons by Site, Including Urine Occurrences

Supplementary Table S2. Barriers and Opportunities for Accurate Fluid Balance Recording

Acknowledgements

We would like to acknowledge the work of Denise Dauterman, who aided in the creation of the fluid balance score, as well as the ICU nurses who completed our surveys and questionnaires and not only provide excellent patient care, but always share their knowledge and insights that have gone into improving our management of FB. We would also like to acknowledge Sheel Chitre and Adrian La Duca (NYU), Amina Khan (CHOP), and Heather Dutton (UAB) for their assistance with clinical data extraction.

Author Contributions

DH, SM, and ACD conceived of the study and along with AS, CGB, UK, HD, JCF, CD, AB, JO, and SLW contributed to the overall design. ACD and SD wrote software queries and analysis scripts, which were executed at each site by UK, ACD, SD, and JO. All authors contributed to qualitative data collection. ACD and DH completed the analysis and created the figures and tables. DH drafted the manuscript. All authors provided revisions and approved of the final manuscript version.

Statements and Declarations

Ethical considerations

This study followed procedures in accordance with ethical standards as delineated by the Helsinki Declaration of 1975. Each site's institutional review board either approved this study with a waiver of informed consent or deemed the project exempt from IRB review.

Consent to participate: Waived as above

Consent for publication: Not applicable

Declaration of conflicting interest

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article. Dr. Dziorny is supported by NIH NIDDK (1K23DK138299) and AHRQ (1R21HS030123). Dr. Fitzgerald is supported by NIH NIDDK funding and her institution has received prior funding from BioPorto for a research study. All other authors confirm they have no disclosures to report.

Funding Statement

Data Availability

Data can be made available upon request to corresponding author.

References

1. Alobaidi R, Morgan C, Basu RK, et al. Association Between Fluid Balance and Outcomes in Critically Ill Children: A Systematic Review and Meta-analysis. *JAMA Pediatr*. 2018;172(3):257-268. doi:10.1001/jamapediatrics.2017.4540

2. Sutherland SM, Zappitelli M, Alexander SR, et al. Fluid overload and mortality in children receiving continuous renal replacement therapy: the prospective pediatric continuous renal replacement therapy registry. *Am J Kidney Dis*. 2010;55(2):316-325. doi:10.1053/j.ajkd.2009.10.048

3. Goldstein SL, Currier H, Graf Cd null, Cosio CC, Brewer ED, Sachdeva R. Outcome in children receiving continuous venovenous hemofiltration. *Pediatrics*. 2001;107(6):1309-1312. doi:10.1542/peds.107.6.1309

4. Selewski DT, Cornell TT, Lombel RM, et al. Weight-based determination of fluid overload status and mortality in pediatric intensive care unit patients requiring continuous renal replacement therapy. *Intensive Care Med*. 2011;37(7):1166-1173. doi:10.1007/s00134-011-2231-3

5. Gillespie RS, Seidel K, Symons JM. Effect of fluid overload and dose of replacement fluid on survival in hemofiltration. *Pediatr Nephrol*. 2004;19(12):1394-1399. doi:10.1007/s00467-004-1655-1

6. Selewski DT, Gist KM, Basu RK, et al. Impact of the Magnitude and Timing of Fluid Overload on Outcomes in Critically Ill Children: A Report From the Multicenter International Assessment of Worldwide Acute Kidney Injury, Renal Angina, and Epidemiology (AWARE) Study. *Crit Care Med*. 2023;51(5):606-618. doi:10.1097/CCM.0000000000005791

7. Al-Lawati ZH, Sur M, Kennedy CE, Akcan Arikan A. Profile of Fluid Exposure and Recognition of Fluid Overload in Critically Ill Children. *Pediatr Crit Care Med*. 2020;21(8):760-766. doi:10.1097/PCC.0000000000002337

8. Barhight MF, Nelson D, Chong G, Basu RK, Sanchez-Pinto LN. Non-resuscitation fluid in excess of hydration requirements is associated with higher mortality in critically ill children. *Pediatr Res*. 2022;91(1):235-240. doi:10.1038/s41390-021-01456-z

9. Hudkins MR, Miller-Smith L, Evers PD, Muralidaran A, Orwoll BE. Nonresuscitation Fluid Accumulation and Outcomes After Pediatric Cardiac Surgery: Single-Center Retrospective Cohort Study. *Pediatr Crit Care Med*. 2023;24(12):1043-1052. doi:10.1097/PCC.0000000000003373

10. Mandl KD, Kohane IS. Federalist principles for healthcare data networks. *Nat Biotechnol*. 2015;33(4):360-363. doi:10.1038/nbt.3180

11. Menon S, Krallman KA, Arikan AA, et al. Worldwide Exploration of Renal Replacement Outcomes Collaborative in Kidney Disease (WE-ROCK). *Kidney Int Rep*. 2023;8(8):1542-1552. doi:10.1016/j.ekir.2023.05.026

12. Selewski DT, Barhight MF, Bjornstad EC, et al. Fluid assessment, fluid balance, and fluid overload in sick children: a report from the Pediatric Acute Disease Quality Initiative (ADQI)

conference. *Pediatr Nephrol.* Published online November 7, 2023. doi:10.1007/s00467-023-06156-w

13. Holliday MA, Segar WE. The maintenance need for water in parenteral fluid therapy. *Pediatrics.* 1957;19(5):823-832.

14. Artino AR, La Rochelle JS, Dezee KJ, Gehlbach H. Developing questionnaires for educational research: AMEE Guide No. 87. *Med Teach.* 2014;36(6):463-474. doi:10.3109/0142159X.2014.889814

15. Charaya S, Angurana SK, Nallasamy K, Jayashree M. Restricted versus Usual/Liberal Maintenance Fluid Strategy in Mechanically Ventilated Children: An Open-Label Randomized Trial (ReLiSCh Trial). *Indian J Pediatr.* Published online October 18, 2023. doi:10.1007/s12098-023-04867-4

16. Dziorny A, Kothari U, Drury S, et al. Improving Fluid Overload Among Critically Ill Children: Federated Data Analysis for Multi-Center Quality Improvement. (Accepted) presented at: Pediatric Academic Societies Annual Meeting; May 2024; Toronto, ON.

A Most Frequent Barriers to Accurate FB Capture

B Achievable Opportunities to Improve Accurate FB Capture

C Most Feasible Method for Accurate FB Capture