Computational modeling of the temporal influences between cues, craving and use in addiction: A dynamical system analysis based on ecological momentary assessment

Running title: Craving and dynamical systems

Christophe Gauld [1,2,3,†], Damien Depannemaecker [4,†,*], Fuschia Serre [3,5,6,X], Marc Auriacombe [3,5,6,X,*]

Affiliations
[1] Department of Child Psychiatry, CHU de Lyon, F-69000 Lyon, France
[2] Institut des Sciences Cognitives Marc Jeannerod, UMR 5229 CNRS & Université Claude Bernard Lyon 1, F-69000 Lyon, France
[3] SANPSY, CNRS, UMR 6033, F-33076 Bordeaux, France
[4] Aix Marseille Université, Institut National de la Santé et de la Recherche Médicale, Institut de Neurosciences des Systèmes (INS) UMR1106, Marseille 13005, France
[5] University of Bordeaux, F-33076 Bordeaux, France
[6] Pôle interétablissement d'addictologie, CHU de Bordeaux et CH Charles Perrens, F-33076 Bordeaux, France

[†] : These authors contributed equally as first author.
[X] : These authors contributed equally as last author.

* Corresponding authors
Marc Auriacombe
e-mail: marc.auriacombe@u-bordeaux.fr
Sanpsy-addiction, Pôle Addictologie, CHCP, 121 rue de la Béchade, 33076 Bordeaux Cedex, France
Tel: +33 556 561 738
Damien Depannemaecker
e-mail: damien.depannemaecker@univ-amu.fr
Institut de neuroscience des systèmes, Aix-Marseille Université, 27 bd, Jean Moulin 13005 Marseille, France

Main text words: 4,693 / 5,000
Abstract words: 280/300
References: 56
Figures: 5
Tables: 4

Abstract (275 / 300 words)

Substance Use Disorders (SUD) can be conceptualized as a prospective link from cues to craving and

use. To explore the nonlinear interplay between craving and cues, this study applied dynamical

systems theory (DST) to ecological momentary assessment (EMA) data. Optimized linear Seasonal

Auto-Regressive Integrated Moving Average with eXogenous variable (SARIMAX) models were

used to phenotype patients with SUD (alcohol, tobacco, cannabis, opiates, and cocaine), considering

the potential for complex interactions between cue exposure and craving intensity in daily life. These

phenotypic profiles were replicated in computational DST models to analyze the nonlinear

interactions between cues, craving, and use. The study involved 211 individuals and 8,260

observations, with 154 patients fitting the SARIMAX model for the influence of cues on craving, and

57 patients fitting the SARIMAX model for a possible influence of craving on cues. Two DST models

were adjusted to replicate the complex temporal dynamics of SUD based on these two directions of

influence. The first DST model (adjusted to the influence of cues on craving) showed that an increase

in cues leads to a rise in craving, which then diminishes both cues and craving itself, with use patterns

following craving's trajectory. This patient profile is driven by a phenomenon of "maximum cue

saturation". The second DST model (adjusted to the influence of craving on cues) demonstrated that

an increase in craving heightens cue perception, further increasing craving and leading to use, with

use peaking and then reducing craving. This patient profile is characterized by a phenomenon of

"maximum use saturation". Both models underscore craving as an essential modulator between cues

and use, opening new therapeutic avenues.

Keywords: substance use disorder; craving; ecological momentary assessment; dynamical system;

biomarker; addiction.

 Introduction

Substance Use Disorder (SUD) is characterized by recurrent relapses despite efforts to reduce use or

abstain [1]. One of the main objectives of SUD treatment is to reduce the frequency of relapses over

time. As an important symptom of SUD [1–8], the fluctuating phenomenon of craving can be captured

by daily-life monitoring systems such as ecological momentary assessment (EMA) [9]. EMA, adapted

and validated to study craving in SUD [10], allows a real-time measurement of individual momentary

states in the natural environment through repeated assessments overtime. Based on EMA research, a

human model of addiction has been suggested across several substances and behaviors: the cues –

craving – use model [4, 11, 12]. In this model, an increase in the number of reported cues is associated

with an increase in the intensity of self-reported craving in the following EMA survey, itself

associated with an increased likelihood of use in the following EMA survey. This craving – use link

has been considered unidirectional [11, 13–15] using mixed models like hierarchical linear models

(HLM), which allows for the analysis of data with a nested structure, accounting for variability at

multiple levels of observation. However, HLM has some limitations such as the difficulty in

accounting for cross-effects, i.e., integrating complex interactions between levels. Furthermore, they

are only linear: the effects (e.g., craving) may be directly proportional to the causes (e.g., cues) – as

opposed to non-linear models where the relationships between variables can be disproportionate,

interactive, producing effects which are not predictable simply by addition or multiplication. Finally,

HLM cannot account for multidirectionality, e.g., that cues and craving could influence each other in

a loop. Due to the important intra-individual variability and its intrinsic non-linearity, the evolutionary

dynamics of a specific patient with a SUD are particularly complex to model.

To go further, modeling the evolution of SUD variables and their interactions requires finding

alternatives to traditional statistical methods [2]. A growing literature based on dynamical systems

theory (DST) seeks to capture the temporal dynamics of psychiatric conditions (e.g., [16–36]),

including SUD. DST accounts for the evolution of clinical manifestations over time without

abolishing the complexity of the relationships between variables and their multidirectionality,

allowing the prediction of different types of temporal evolution. This theory is based on dynamic

objects (e.g., tipping points, transitions, steady states, strange attractors, limit cycles or oscillations)

and dynamic concepts (e.g., emergence, instability, or variations related to fluctuations in initial

conditions – referring to the "butterfly effect", etc.). However, most psychiatric studies include

biological factors (e.g., [37]) and few studies have empirically included psychiatric clinical

manifestations (e.g., [20]).

Based on EMA data, considering the possibility that, in addition to a unidirectional prospective link

from cues to craving, an increase in craving could change perception of cues [38], DST models allow

the representation of dynamics and mutual influences between variables. . The main goal of this study

was to model alternative evolutions of craving and cues, to explore their influence on substance use.

Such results could lead to proposing a classification of individuals with SUD based on dynamics'

profiles, as well as providing information on the attractive states (i.e., the states to which the system

gravitates) – allowing to understand what stabilizes the patient in a given (healthy or pathological)

state.

Methods

This study is based on two successive stages: i) the use of optimized linear statistical models

(SARIMAX) to identify, based on EMA data, two patient profiles according to the direction of

influence between cues and craving; ii) the use of a computational model (DST) to analyze the

complex interplay between cues and craving according to these profiles. In line with the classic cues-

craving-use model (in which an increase in cues can linearly contribute to an increase in craving), we

hypothesized that a state of craving could make an individual more "sensitive" or more "receptive" to

cues, with an effect on use, creating a feedback loop: cues lead to craving, which increases cue

perception, leading to more craving, and ultimately, more use. This reflects a non-linear dynamic

where continuous mutual interactions bring about a dynamic process. The modeling focuses on the

cues-craving relationship, as use is the outcome.

Data collection in EMA

Participants

Data were extracted from two EMA studies with similar protocol [10], which comprised individuals

aged 18 years and above, who were beginning addiction outpatient treatment (in Bordeaux and

Bayonne, France), were diagnosed with at least one DSM-5 Substance Use Disorder for tobacco,

alcohol, cannabis, cocaine or opiates, and were free from active psychosis or severe cognitive

impairment [39]. Informed consent was obtained, and the studies were approved by the French

Regulation and ethical committee (CPP SOOM III/DC-2009/01; CNIL/DR-2015-408; CPP IDF X:

11-2019/IR-RCB: 2018-A00952-53; CNIL/MR003). Participants received standard comprehensive

care, consisting of individual behavioral treatment focused on relapse prevention and psychosocial

support combined, when available, with pharmacotherapy. After establishing a target quit date

according to the patient's personal goals, full abstinence or use reduction was encouraged as an

outcome, but with no negative consequences for the patient if failure to achieve this goal.

Assessments

Each participant underwent an initial interview to collect sociodemographic variables and substance

use-related information using a validated French version of the Addiction Severity Index (ASI) [40,

41], modified to consider tobacco addiction. The Mini International Neuropsychiatric Interview–Plus

(MINI) structured interview [42] was adapted to evaluate the current DSM-5 SUD status within the

past year. When multiple SUD co-occurred, the primary substance was determined according to the

main problematic substance reported by the individual, and on which the treatment was focused. The

number of endorsed SUD criteria defined MINI addiction severity (**Table 1**).

Ecological momentary assessment

The EMA protocol was previously described in detail [10]. EMA assessments began within the first

month after treatment initiation. Following a training session, each participant received a tablet to

carry for 14 days. Each tablet was programmed to administer four electronic surveys per day,

inquiring about primary substance use, craving, and the presence of environmental cues. The signal

schedule was adjusted to accommodate participants' usual sleep patterns. Financial compensation was

provided based on the number of surveys completed, with a maximum of 100 euros for participants

who completed 75% or more of the assessments. Methodological details and comparisons with other

studies using this validated protocol [11, 11, 43–45] are presented in **Supplementary Material 1**.

Data of interest

At each electronic assessment, participants were asked to rate the maximum intensity level of craving

they had felt since the previous assessment on a 7-point scale (1 = no desire to 7 = extreme desire).

This item was referred to as "craving". Participants were also asked if they had encountered cues

related to their primary substance, selected from a personalized list for each individual. This item was

labeled "cues" and included specific objects or contexts typically associated with substance use, such

as materials used for consumption (e.g., a bottle), the sight or smell of the substance, or places

associated with use (e.g., a bar). It also encompassed personal cues, identified during a personal

interview, that referenced specific circumstances, objects, individuals, emotions, or environmental

contexts linked to their substance use (e.g., "being with my friend Sam") [11]. Although these refer to

"cue perceptions", for reading convenience, we used the term "cues" below. Finally, participants were

asked if they had used the primary substance. This item was referred to as "use". All three questions

inquired about participants' experiences since the previous assessment.

Due to a restricted number of assessments occurring on the starting and ending day of the study, and

their overlap with investigator contact, only data collected between days 2 and 13 were analyzed

(n=12 days) (**Table 2**).

Statistical and computational models

Aims of the models

First, we used optimized linear SARIMAX models with craving and cues to determine the mutual

influence profiles of patients. We aimed to identify: i) a *SARIMAX-1* model, referring to the influence

of cues on craving, hereinafter named "cues-to-craving" model; ii) a *SARIMAX-2* model, referring to

the influence of craving on cues, hereinafter named "craving-to-cues" model.

Secondly, we used the SARIMAX model 1 to determine the specific parameter values of a model

based on the DST (*DST-model 1*), and the SARIMAX model 2 to determine the specific parameter

values of another model based on the DST (*DST-model 2*). We qualitatively observed the complex

evolution of each of the two variables within the two DST models. "Use" was treated as an output of

the model, with its values deliberately left unconstrained to follow the parameterized dynamics of the

cues-craving structure in both DST models. This approach allowed us to capture the intricate

interactions between cues, craving, and use.

By combining linear SARIMAX models with DST models, we aimed to create a framework that

captures both the linear and nonlinear dynamics of craving and cues. The SARIMAX models first

classified patients based on their interaction profiles. These profiles served to establish parameter

values for the DST models, which further analyze the nonlinear interactions between craving and

cues. This approach allowed us to utilize linear models for the initial sequence of variables without

necessarily providing a parameter characterization. Subsequently, the DST models determine these

parameters to capture intricate feedback loops between craving, cues, and use. Though SARIMAX

models were constrained by assumptions of linearity and stationarity, this hybrid approach could offer

an interesting tool for understanding complex craving dynamics across varied profiles. As we will

describe later, the comparison of SARIMAX and DST models is a methodological challenge due to

differences in scope and (non)linearity. Specifically, in the DST framework, the parameter ranges are

reconstructed to fit the data, rather than being directly calculated from empirical measurements. This

approach emphasizes conceptual alignment with circadian-scale dynamics rather than exact value

matching.

To ensure the reliability of our method, we conducted a simulation study by generating two groups

based on distinct craving-to-cues and cues-to-craving dynamics, according to the DST model

parameter distributions). We then verified our ability to recover these groups using SARIMAX

modeling, applying separate models to identify each directional influence (**Supplementary Material**

2).

SARIMAX models

SARIMAX models are linear algorithms for forecasting data, considering past values (AR –

AutoRegressive) and data noise (MA – Moving Average) to predict future values [46].

Stationarity and seasonality assessments of the EMA data. Stationarity is a condition of validity for

this time-series analysis using the Auto-Regressive Moving Average (AR-MA) method. The

stationarity of the time-series data for craving and cues was assessed using the Augmented Dickey-

Fuller (ADF) test [47]. If the time series was not stationary, an integration factor (I(d)) was added to

the model, allowing differentiation to make the data stationary (i.e., resulting in a "AR-I-MA" model).

Seasonality (classically denoted S in the model, providing a "S-ARIMA" model) was considered

given the repeated daily nature of the data. Finally, an exogenous variable (denoted X in the model,

resulting in a "SARIMA-X" model) could be added to model an additional effect on the time series of

the first variable (i.e., cues for the craving time series, and craving for the cues time series) [48, 49].

Model fitting. For each patient, two SARIMAX models were fit to the data: one where craving was

modeled as the dependent variable influenced by cues (*SARIMAX-model 1* – "cues-to-craving"

model), and another where cues were the dependent variable influenced by craving (*SARIMAX-model*

2 – "craving-to-cues" model). Different combinations of autoregressive and moving average terms

were tested for each patient. Only the parameter values for the equations governing y (craving) and z

(cues) were adjusted to align with each of the two SARIMAX models. The parameter values for the

equation of x (use) were deliberately left unconstrained, allowing to observe its dynamics as an output

variable within the models.

Parameter optimization. The y and z parameters of the two SARIMAX models were optimized

individually for each patient. A grid search was performed across potential values of the AR and MA

component, identifying the combination that minimized the Akaike Information Criterion (AIC). This

criterion balances model fit and complexity to prevent overfitting. The choice to use the AIC over the

Bayesian Information Criterion (BIC) was motivated by the objective of optimizing predictive

accuracy while accommodating the complexity of individual temporal dynamics, as AIC imposes a

less stringent penalty on model parameters, thus enhancing the model's sensitivity to subtle within-

subject variations. Two AIC were provided per patient (one for each SARIMAX model of a patient).

Model selection. For each patient, the preferred SARIMAX model (either *SARIMAX-model 1* or

SARIMAX-model 2) was determined based on the lower AIC. We assume that each patient aligns with

only one specific model, excluding the other. Following the identification of which participants best

fit each model, we compared the two selected subgroups based on socio-demographic characteristics,

substance types, addiction severity, psychiatric diagnoses, and EMA variable characteristics (**Table**

3). This comparison aimed to 1) clinically describe the profiles of patients displaying different

dynamic patterns (cues to craving versus craving to cues), and 2) methodologically confirm the

absence of significant differences between the groups, ensuring that any observed distinctions in

model dynamics were not attributable to these underlying characteristics. Details on these models are

provided in **Supplementary Material 3**.

Sensitivity analysis. We then performed a sensitivity analysis on the sub-datasets of alcohol and

cocaine to assess the stability of the models across these substance-specific data.

Auto-correlation analysis. Finally, we proposed studying the temporal dynamics between craving and

cues through auto-correlation and cross-correlation analyses, allowing to identify individual temporal

structures, key time lags, and descriptive statistics revealing variability in craving and cues across the

sample.

DST models

Dynamical system modeling. A set of differential equations was used to model the dynamical

relationship between craving and cues. The parameter values were searched, in order to fit with each

of the two SARIMAX models, by successive simulations to obtain a qualitative understanding of the

system's behavior. The model was integrated over the 12-day observation window and extended to a

theoretically infinite duration to visualize the temporal interaction of one variable on the other.

Details of the DST model. The detail of the DST model has been described elsewhere [50]. In a

nutshell, it is based on four differential equations, producing a nonlinear representation of

participant's internal factors (i.e., subjective phenomenological experiences and/or biological

elements) (y) and of environmental noise (z). Based on the empirical data collected in EMA regarding

SUD, the "patient's internal factor" corresponds to the intensity of craving (y), and the

"environmental noise" corresponds cues (more precisely, to the salience of perceived cues) (z). Two

other variables allow modulating: i) the long-term temporality of patients' conditions (f); ii) and the

intensity of behavior (x), corresponding here to substance use (passively evolving according to the

two cues-craving models). Thirteen parameters are described within these four variables, with their

clinical translation, presented in **Supplementary Material 4**. Details of interactions between

parameters within each variable are described elsewhere [50], but the four differential equations and

the thirteen parameters of the model are presented below[1]:

$$\dot{X} = \begin{bmatrix} \dot{x} \\ \dot{y} \\ \dot{z} \\ \dot{f} \end{bmatrix} = \begin{bmatrix} \frac{S_{\max}}{1+\exp\left(\frac{R_s-y}{\lambda_s}\right)} - x \\ \frac{P}{1+\exp\left(\frac{R_b-y}{\lambda_b}\right)} + fL - xy - z \\ S(\alpha x + \beta y)\zeta(t) - z \\ y - \lambda_f f \end{bmatrix}$$

As an example, **Figure 1** presents two simulations of the DST model with different parameter values,

by considering three random variables. We observe that the simulated patient trajectories in the

model's "relief" differ when the parameter values are varied.

[FIGURE 1]

Construction of the two DST models. We built two DST models with specific parameter values. Given

the short time scale considered (tens of days), we fixed the f variable – originally representing

predisposing factors over months and years. The parameter values in the y (craving) and z (cues)

equations were adjusted to align with each of the two SARIMAX models. In other words, patient

---

[1] x: symptom intensity, representing use; y: participant's internal factors, representing craving intensity; z: environmental noise, representing the salience of perceived cues. $\tau_{x,y,z,f}$: time scales of the four equations; $S_{\square\square\square}$: maximal symptom level; $R_\square$ and $R_\square$: sensitivity, with $R_\square$ for symptom sensitivity and $R_\square$ for internal element (potentiation) sensitivity; $\lambda_\square$ and $\lambda_\square$: slope of symptom/internal element curves, with $\lambda_\square$ affecting symptom intensity based on patient state γ, and $\lambda_\square$ affecting internal elements based on environment z; P: maximal rate of internal elements, aggregating biological, psychological, and phenomenological factors, shaping phenotype from biological signals; L: level of predisposing factors contributing to a permanent shift in internal elements; λ_f: scaling factor for slow fluctuations affecting L; S: overall environmental sensitivity; α and β: weights of x and y effects on environmental perception.

profiles aligned with each of the two linear SARIMAX models (either cues-to-craving or craving-to-

cues) were fitted to a nonlinear DST model, parameterized to capture their dynamics. For instance, the

parameter L (representing the "vulnerability level" in the y equation) may need to be set high to reflect

the increasing influence of craving on cues. Such parameterization of vulnerability would constitute

an interesting result: for craving to affect sensitivity to cues, a patient must have a high level of

vulnerability (e.g., genetic predispositions, psychological vulnerability, neurological comorbidities,

etc.). Thus, from the dynamical model, we identified a subset of the parameter space, representing the

possible dynamical repertoire that matches the circadian evolution (i.e., changes observed over a 24-

hour day-night cycle) observed in the data. This selection was based on the features associated with

temporality and rhythmicity, central to interpreting model comparisons. This specific parameter

tuning, grounded in an informed and iterative fit process, represents the sequential emergence and

fluctuation patterns of craving and cues in each model, offering a close alignment with observed

dynamics – rather than a merely contingent outcome.

All the analyzes were performed with Python [51]. The code is now added to the open access

repository: https://github.com/ChristopheGauld/Dynamical_systems_for_craving.

297   **Results**

Sample description and assessments

As described in **Table 1**, 211 individuals were included in the study. The mean age was 38.2 years old

and 36% were female (n=36). Primary substance was: 35% alcohol, 28% tobacco, 18% cannabis, 16%

opiates and 3% cocaine. The flow-chart of participant inclusion is presented in **Supplementary**

Material 5.

Table 1. Characteristics of the Substance Use Disorder (SUD) sample (n=211 participants): primary

substance, demographic variables and substance use-related information. ASI: Addiction Severity

Index; MINI: Mini International Neuropsychiatric Interview–Plus; SD: standard deviation.

	Sample characteristics	All participants (n=211)	
		% (n)	Mean (SD)
Primary substance use		37 (3,087)	
Primary substance	Alcohol	35 (73)	–
	Tobacco	28 (58)	–
	Cannabis	18 (39)	–
	Opiates	16 (34)	–
	Cocaine	3 (7)	–
Demographic variables	Sex (female)	36 (78)	–
	Employed	56 (116)	–
	Age	–	38.2 (11.4)
	Education (years)	–	12.5 (2.74)
	Pharmacological treatment	79 (167)	–
Substance use-related information	ASI addiction severity (0-9)	–	6.6 (0.76)
	Mood disorder on the MINI (current diagnosis)	34 (72)	–
	Anxiety disorder on the MINI (current diagnosis)	45 (93)	–

EMA assessments

Among the 10,128 EMA electronic assessments planned and delivered, 8,260 (82%) were completed

Majority of EMA assessments reported episodes of craving (70% with non-null intensity), and 37%

reported primary substance use (**Table 2**).

Table 2. Ecological Momentary Assessment (EMA) of intensity of craving, number of cues, and

reports of primary substance use on the Substance Use Disorder (SUD) sample of participants

(n=211). SD: standard deviation.

EMA variables	All participants (n=211 – 8,260 observations)	
	% (n)	Mean (SD)
Craving episodes (intensity ≥ 1)	70 (5,810)	–
Craving intensity (1-7)	–	3.6 (2.10)
Number of cues	–	4.3 (3.4)
Use	37 (3,087)	–

SARIMAX models

The ADF test confirmed that the time-series does not have a time-dependent structure ($p = 0.0003$ for craving and $p \ll 0.001$ for cues, indicating that both series are stationary), supporting stationarity for each patient and for each variable (cues, craving and use). No detrending was necessary. No seasonality, which could have been present here due to typical day-by-day fluctuations, was found in the data. SARIMAX models could thus have been named "ARMAX", but we keep the initial term to ensure understanding.

After testing the two SARIMAX models for each patient and after individual optimization, we found 154 patients who fit better (according to the AIC) with the SARIMAX-model 1 ("cues-to-craving") and 57 patients who fit better with the SARIMAX-model 2 ("craving-to-cues"). As expected in the analysis seeking to maximize the fit of a patient with one of the two models, no patient belonged to both models. Details on the AIC of the 1 and 2 SARIMAX models and on the preferred model according to the lowest AIC are given in **Supplementary Material 3**. **Table 3** provides the characteristics of the patients who best fit each of the two SARIMAX models. There were no significant differences in the distribution of sociodemographic characteristics, addiction severity, psychiatric diagnoses, or substance types between patients fitting each model, except for the EMA variables.

Table 3. Socio-demographic characteristics, substances types, addiction severity, psychiatric diagnoses and EMA variable characteristics of the two temporal profile (SARIMAX) of patients with Substance Use Disorder (SUD) (n=211). SARIMAX: Seasonal Auto-Regressive Integrated Moving Average with eXogenous variable. "Model 1": SARIMAX-model 1 ("cues-to-craving"); "Model 2": SARIMAX-model 2 ("craving-to-cues").

	Model 1 (N= 154)				Model 2 (N=57)					
	%	n	Mean	SD	%	n	Mean	SD	Test	p*
Sample characteristics										
Sex (female)	36.4	56			38.6	22			$\chi^2=0.089$	0.765
Employed	55.6	84			57.1	32			$\chi^2= 0.038$	0.845
Age			37.6	11.1			39.7	11.3	F= 1.53	0.217

Education (years)		12.4	2.6		12.5	3.1	F= 0.021	0.886

Education (years)			12.4	2.6		12.5	3.1	F= 0.021	0.886
Primary substance									
Alcohol	30.5	47			45.6	26			
Tobacco	34.4	53			8.7	5			
Cannabis	20.1	31			14.0	8			
Opiates	13.0	20			24.6	14			
Cocaine	1.9	3			7.0	4			
Addiction severity									
ASI ISR (0-9)			6.6	0.8			6.5	0.7	F= 0.088	0.767
Current pharmacological treatment for addiction	80.5	124			75.4	43			χ^2=0.651	0.419
MINI current diagnosis									
Mood disorder	32.0	48			42.9	24			χ^2= 2.11	0.146
Anxiety disorder	42.0	63			52.6	30			χ^2= 1.89	0.170
EMA variables	**6004 observations**				**2256 observations**					
Frequency of craving episodes (intensity > 1)			73.3	32.4			63.7	27.7	F= 3.98	**0.0472**
Craving intensity (1-7)			3.7	1.6			3.2	1.3	F= 4.25	**0.0405**
Number of cues			4.9	2.5			2.8	1.7	F= 35.91	**<.0001**
Frequency of primary substance use			42.4	35.1			26.6	29.9	F= 9.08	**0.0029**

The results of the sensitivity analysis and of the auto-correlation analysis between craving and cues

are presented in **Supplementary Material 3**.

DST models

Using successive simulations based on different parameterization of the z (cues) and y (craving)

variables of the DST model, we sought to represent in a non-linear way (unlike SARIMAX models)

the mutual interactions between craving and cues. We did not seek to modify the other two variables

(f variable for long-term temporality or x variable for use). The parameter values of the equations of y

(craving) and z (cues) equations are modified in order to fit with each of the two SARIMAX models.

We defined two sets of parameters: the first set of parameters of the *DST-model 1,* corresponds to the

dynamical system representation of the 154 patients who fit with the *SARIMAX-model 1*; the second

set of parameters of the *DST-model 2,* corresponds to the dynamical system representation of the 57

patients who fit with the *SARIMAX-model 2*.

The details of the parameter values of the two DST models are provided in **Table 4**. Five parameter

values were necessarily modified to recreate the two SARIMAX-models: *L, R_b, P , τ_x,* and τ_y (in red in

Table 4). Firstly, the value of the *L* parameter (corresponding to the vulnerability level in the *y*

equation) was barely modified (*L* = 1.01 in the DST-model 1 and *L* = 1.00 in the DST-model 2) to

obtain these two profiles of influence of craving on cues. Secondly, the value of the parameter R_b

(referring to the sensitivity of the internal elements) was necessarily modified (R_b = 1.04 in the DST-

model 1 and R_b = 0.904 in the DST-model 2): for craving to have an influence on sensitivity to cues

(DST-model 2), a patient must have a lower level of sensitivity. Thirdly, the value of the parameter *P*

(representing potentiation, or the maximal rate of internal elements amplifying symptoms) was

necessarily increased for DST-model 1 (*P* = 10) compared to DST-model 2 (*P* = 1), indicating that

craving (which drives DST-model 1) serves as an internal mechanism that amplifies use. Finally, τ_x

and τ_y refer solely to the time scales specific to each variable (e.g., τ_x • *dx/dt*) without clinical

interpretation, and their increase in DST-model 1 does not alter the interpretation of the other

parameter values.

Table 4. Parameter values of the two dynamical system models fitted with linear models, for the

Substance Use Disorder (SUD) sample of participants (n=211). DST: Dynamical System Theory. The

details of the parameters and their clinical translation are given in **Supplementary Material 4**.

DST models	Parameters of x, y and z variables													
	x				*y*						*z*			
	S_{max}	R_s	λ_s	τ_x	B_{max}	R_b	λ_b	P	L	τ_y	S	α	β	τ_z
DST-model 1	10	1	0.1	14	10	1.04	0.05	10	1.01	14	1.0	0.5	0.5	1
DST-model 2	10	1	0.1	2	10	0.904	0.05	1	1.00	2	1.0	0.5	0.5	1

x: symptom intensity, representing use; *y*: participant's internal factors, representing craving intensity;
z: environmental noise, representing the salience of perceived cues. $\tau_{x,y,z,f}$: time scales of the four
equations; $S_{\square\square}$: maximal symptom level; $R_\square$ and $R_\square$: sensitivity, with $R_\square$ for symptom
sensitivity and $R_\square$ for internal element (potentiation) sensitivity; $\lambda_\square$ and $\lambda_\square$: slope of
symptom/internal element curves, with $\lambda_\square$ affecting symptom intensity based on patient state γ, and
$\lambda_\square$ affecting internal elements based on environment z; P: maximal rate of internal elements,
aggregating biological, psychological, and phenomenological factors, shaping phenotype from
biological signals; L: level of predisposing factors contributing to a permanent shift in internal
elements; λ_f: scaling factor for slow fluctuations affecting L; S: overall environmental sensitivity; α
and β: weights of x and y effects on environmental perception.

Dynamics of craving and cues on the *DST-model 1* and on the *DST-model 2* are represented in **Figure 2** and **Figure 3,** respectively. Within these two figures, although there was no fixed scale due to the nonlinear interactions between relative dynamics, it was important to consider that the baseline represented an absence of interaction, visualized by an increase in a variable (e.g., the baseline for craving represented the level corresponding to the absence of craving, when it did not spontaneously increase or respond to another variable).

In **Figure 2** (DST-model 1 for *SARIMAX-model 1* patients, N=154), when an increase in cues (visible as negative events in the *z* variable) preceded craving, there was a relatively rapid increase in craving, which in turn reduced the cues, followed by a slower decrease in craving (**Figure 2**, bottom). Craving peaked in alignment with peak cues. Craving only returned to its baseline level after cues also returned to baseline. Use progressively increased throughout the duration of the craving rise (e.g., from 8 p.m. to midnight on the 1st day) and only began to decrease in temporal conjunction with the relatively sudden drop in craving. This temporal overlap between the decreases in craving and use limited our ability to determine the direction of influence between the two variables. However, it is noteworthy that the decrease in cues preceded this joint decrease in craving and use (e.g., at 4 p.m. on the 1st day). The return to baseline was therefore attributed to cue saturation, suggesting modulation by craving.

[FIGURE 2]

In **Figure 3** (*DST-model 2* for *SARIMAX-model 2* patients, N=57), when an increase in craving preceded cues, this prior increase in craving randomly influenced cue perception, resulting in greater simultaneity between these variables compared to the previous model (**Figure 3**, bottom). Craving decreased during the period of noisy cue perception, with both craving and cues returning to baseline simultaneously. Use increased gradually throughout the duration of the craving increase and, as before, only began to decrease with the relatively sudden drop in craving. Cue reporting returned to baseline in temporal conjunction with this decrease in use and craving (unlike the earlier return seen

407 in the previous model). The return to the baseline state was therefore due to saturation of use,

408 implying a modulation of craving.

410   **[FIGURE 3]**

412 Finally, the simulation study demonstrates the robustness of our modeling approach in reliably

413 distinguishing the craving-to-cues and cues-to-craving groups, confirming the SARIMAX method's

414 effectiveness in capturing these distinct directional influences based on different parameter

415 distributions of the DST models (**Supplementary Material 2**).

 Discussion

The main goal of this study was to understand the multiple and complex influences of craving and

cues on substance use. To achieve this, two SARIMAX models were tested for each patient, resulting

in 154 patients aligning better with *SARIMAX-model 1* ("cues-to-craving") and 57 with *SARIMAX-*

model 2 ("craving-to-cues"). Parameter sets were defined through a simulation analysis to propose

two DST models, representing the evolutionary and interactional dynamics of cues, craving, and use

for patients corresponding to each SARIMAX model. **Figure 4** illustrates the sequence of interactions

between the variables in *DST-model 1*, from which two main results could be drawn.

The first finding offers insights into the mechanism of cessation in these patients. Specifically,

cessation of use is not due to the saturation of use itself (i.e., the point where further increase is

impossible due to reaching maximum capacity). Instead, cessation results from a specific sequence

involving cues and craving. In this model, a peak in cues initially occurs. Once this peak is reached,

we observe a gradual reduction in use, mediated by craving levels. This decrease in cues returns to the

baseline level, ultimately leading to cessation of use. In other words, cessation is triggered by the

preceding peak in cues, modulated by craving levels.

The second finding of this model addresses the complexity of the interaction between craving and use.

As observed in practice, craving leads to an increase in use [2]. In this model, cues precede craving,

but craving peaks simultaneously with the peak in cues. A high level of cues and craving is followed

by a gradual decrease in cues to a baseline threshold, resulting in cessation of use. The direct

interaction between cues and use remains unclear – specifically, whether cues directly saturate use,

given that cues precede use in this model. However, modulation by craving clarifies the interactions

among these three variables. The model suggests that cues and craving rise in tandem, leading to an

increase in use. At the peak of craving, cues also reach their maximum. After this peak, craving

begins to decrease, which in turn reduces cues to the baseline, eventually leading to cessation of use.

Importantly, use does not interrupt the craving peak (since a reduction in craving is necessary once it

reaches its peak). The conclusion here is that use plays a secondary role: it follows craving, which

controls the peak in cues, and it also follows the reduction in cue perception.

[FIGURE 4]

Figure 5 illustrates the sequence of interactions between the variables in *DST-model 2*. The primary

result from this model pertains to the role of use in the interplay between craving, cues, and use.

In this model, craving increases and simultaneously enhances the salience of cues. These cues do not

exceed a relatively low threshold, resulting in a "noisy" appearance. This scattered pattern may reflect

the intermittent encounter of cues by patients, consistent with the DST-model 2, where cues

fluctuations do not align directly with craving peaks but rather create a baseline disturbance, testifying

to a strong impact of craving on use without the presence of clearly defined prior cues – consistent

with previous experimental data [44]. At the same time, craving drives an increase in use. This

increase is linked not only to heightened craving but also to greater cue engagement. Interestingly, the

decrease in cues does not drive the reduction in use or craving, as no maximum threshold of cues is

reached in this scenario (unlike in *DST-model 1*). Once use has peaked, it must necessarily decline. At

this point, craving returns to its baseline, appearing to be influenced by the reduction in use. When use

reaches maximum saturation, there is a sharp decrease in craving. The dynamics of this model are

thus as follows: initially, craving may lead to a peak in use, while simultaneously amplifying cue

perception. Once use hits its peak, it may sharply reduce craving, suggesting a complex feedback loop

between use and craving.

[FIGURE 5]

To summarize, when cues precede craving, the process appears to be primarily driven by a

phenomenon of "maximum cue saturation", which initially increases craving and use, then leads them

to decrease as cues subside. When craving precedes cues, the process seems primarily driven by a

phenomenon of "maximum use saturation", which causes craving and cues to decrease once use

reaches its peak. In both profiles, craving serves as a critical modulator between cues and use. These

findings align with animal models demonstrating spontaneous exhaustion, or saturation [52].

The literature includes numerous simulation studies on psychiatry and DST (e.g., [16–35]); however,

to our knowledge, no DST study has yet been conducted in SUD. Most existing studies focus on a

single (phenotypic) variable (e.g., [32]) and have not integrated phenotype, internal factors, and

environment together – three elements essential for a nuanced understanding of clinical practice.

Furthermore, only eight studies have applied empirical data within DST models in psychiatry [20, 31,

53–58], and none in the context of SUD. Finally, the integration of a linear statistical method with

non-linear computational analysis appears to be unprecedented.

This study's innovative approach offers interesting insights into the complex interplay of

environmental, subjective, and behavioral variables, enabling modeling of clinical decision-making

strategies essential for understanding SUD mechanisms. Such modeling opens therapeutic avenues,

such as accelerating the peak of use to reduce its intensity, which may prompt an earlier reduction in

craving. This focus on dynamic markers aimed at reducing use is particularly novel. Additionally,

EMA could enable more detailed simulation visualizations of patient progression, support the study of

phenotype variability, provide insights into attractors, structure longitudinal cohorts, and potentially

refine nosology with dynamic factors [59–61].

This study encompasses limitations. Firstly, the list of cues was pre-identified with the patient.

However, in this study we were not trying to model the type of cues, but rather change in number

(quantity) of cues, and their interactions with craving. Secondly, we are trying to model rapid

temporal sequences between variables, based on data that is aggregated over 4-hour periods; within

these windows, it is not possible to know the mutual influence between variables. More frequent

EMA measurements would provide better precision. Thirdly, at the linear level, the present study

focused on analyzing the interaction between cues and craving as a priority, considering that the use

behavior was an outcome variable in relation to cue perception and craving. This choice raises the

concern that, in linear models, the analysis of cues might merely reflect use patterns, and that future

(SARIMAX) analyzes may integrate the use as a (exogenous) influence variable. Moreover, certain

statistical choices were necessarily made for these linear models (e.g., favoring AIC over BIC),

methods that could be compared through simulations, for instance. However, beyond this potential

extension to linear models, the use of DST models allows us to highlight the interactions between use,

craving, and cues. Fourthly, although subgroup analyses by SUD type (alcohol, tobacco, cannabis,

opiates, and cocaine) could provide valuable insights, incorporating them could overcomplicate the

already substantial findings. Models were developed by combining different substance groups based

on evidence that cues influence craving, and craving, in turn, affects substance use and relapse across

various types of SUD [2–4]. Fifthly, a known limitation of the SARIMAX models is that they

evaluate each patient as aligning with one model, though a patient's condition may not be stable over

time. Furthermore, SARIMAX models have difficulty handling nonlinear relationships and complex

data patterns, requiring significant fine-tuning to achieve optimal performance and assuming

stationarity in the data. We sought to address these limitations by verifying stationarity, performing a

grid search to optimize the models, and subsequently employing nonlinear models for greater

precision. In parallel, the nonlinear models based on DST also present certain methodological

constraints [29]. For instance, it could be necessary to refine the model to modelize symptoms and

behaviors of different types (and not only "use" or "craving"), different set of parameters or

differential equations could have achieved relatively similar dynamics and the qualitative

interpretation of nonlinear phenomena, although constrained by clinical perspectives, remains

challenging. Different parameter sets could yield similar data fits, highlighting the critical need for

robust model validation.

517   **Conclusion**

In conclusion, this study introduces an innovative approach that combines EMA data, temporal

statistical models (SARIMAX), and computational models (DST) to examine the nonlinear and

interconnected relationships between cues, craving, and use in substance use disorders (SUD). The

results could have important implications for the temporal phenotyping of patients, helping to refine

the complex patterns of influence among SUD variables and supporting the identification of

temporally anchored markers and mechanisms in clinical practice – thus opening critical therapeutic

avenues for precision addiction treatments. Moving forward, validating the robustness of the model

will remain essential, further reinforcing the reliability of these findings and their potential to advance

tailored approaches in addiction treatment.

Acknowledgements

The authors express theirs thanks to all participants for their contribution and are grateful to all the

interviewers of the Addiction Team of SANPSY lab 2009-2022.

Conflict of Interest

None.

References

1.   Auriacombe M, Serre F, Denis C, Fatseas M. Diagnosis of addictions. Chapter 11. The
Routledge Handbook of the Philosophy and Science of Addiction2018. p. 132–144.
2.   Vafaie N, Kober H. Association of Drug Cues and Craving With Drug Use and Relapse: A
Systematic Review and Meta-analysis. JAMA Psychiatry. 2022;79:641–650.
3.   Serre F, Fatseas M, Denis C, Swendsen J, Auriacombe M. Predictors of craving and substance
use among patients with alcohol, tobacco, cannabis or opiate addictions: Commonalities and
specificities across substances. Addict Behav. 2018;83:123–129.
4.   Serre F, Fatseas M, Swendsen J, Auriacombe M. Ecological momentary assessment in the
investigation of craving and substance use in daily life: a systematic review. Drug Alcohol Depend.
2015;148:1–20.
5.   Cavicchioli M, Vassena G, Movalli M, Maffei C. Is craving a risk factor for substance use
among treatment-seeking individuals with alcohol and other drugs use disorders? A meta-analytic
review. Drug Alcohol Depend. 2020;212:108002.
6.   Cleveland HH, Knapp KS, Brick TR, Russell MA, Gajos JM, Bunce SC. Effectiveness and
Utility of Mobile Device Assessment of Subjective Craving during Residential Opioid Dependence
Treatment. Subst Use Misuse. 2021;56:1284–1294.
7.   Bujarski S, Roche DJO, Sheets ES, Krull JL, Guzman I, Ray LA. Modeling naturalistic
craving, withdrawal, and affect during early nicotine abstinence: a pilot EMA study. Exp Clin
Psychopharmacol. 2015;23:81–89.
8.   Sayette MA. The Role of Craving in Substance Use Disorders: Theoretical and
Methodological Issues. Annu Rev Clin Psychol. 2016;12:407–433.
9.   Stone AA, Shiffman S. Ecological Momentary Assessment (Ema) in Behavioral Medicine.
Annals of Behavioral Medicine. 1994;16:199–202.
10.   Serre F, Fatseas M, Debrabant R, Alexandre J-M, Auriacombe M, Swendsen J. Ecological
momentary assessment in alcohol, tobacco, cannabis and opiate dependence: a comparison of
feasibility and validity. Drug Alcohol Depend. 2012;126:118–123.
11.   Fatseas M, Serre F, Alexandre J-M, Debrabant R, Auriacombe M, Swendsen J. Craving and
substance use among patients with alcohol, tobacco, cannabis or heroin addiction: a comparison of
substance- and person-specific cues. Addiction. 2015;110:1035–1042.
12.   Auriacombe M, Fatséas M, Daulouède J-P, Tignol J. Le craving et nouvelle clinique de
l'addiction : une perspective simplifiée et opérationnelle. Annales Médico-Psychologiques, Revue
Psychiatrique. 2018;176:746–749.
13.   Ellis JD, Mun CJ, Epstein DH, Phillips KA, Finan PH, Preston KL. Intra-individual
variability and stability of affect and craving among individuals receiving medication treatment for
opioid use disorder. Neuropsychopharmacology. 2022;47:1836–1843.
14.   Burgess-Hull AJ, Panlilio LV, Preston KL, Epstein DH. Trajectories of craving during
medication-assisted treatment for opioid-use disorder: Subtyping for early identification of higher
risk. Drug Alcohol Depend. 2022;233:109362.
15.   Panlilio LV, Stull SW, Kowalczyk WJ, Phillips KA, Schroeder JR, Bertz JW, et al. Stress,
craving and mood as predictors of early dropout from opioid agonist therapy. Drug Alcohol Depend.
2019;202:200–208.
16.   Kelso JAS. Dynamic patterns: The self-organization of brain and behavior. Cambridge, MA,
US: The MIT Press; 1995.
17.   Boldrini M, Placidi G, Marazziti D. Applications of Chaos Theories to Psychiatry: A Review
and Future Perspectives. CNS Spectrums. 1998;3:22–29.
18.   Ayers S. The Application of Chaos Theory to Psychology. Theory & Psychology.
1997;7:373–398.
19.   Barton S. Chaos, self-organization, and psychology. Am Psychol. 1994;49:5–14.
20.   Bonsall MB, Geddes JR, Goodwin GM, Holmes EA. Bipolar disorder dynamics: affective
instabilities, relaxation oscillations and noise. J R Soc Interface. 2015;12:20150670.
21.   Bystritsky A, Nierenberg AA, Feusner JD, Rabinovich M. Computational non-linear
dynamical psychiatry: a new methodological paradigm for diagnosis and course of illness. J Psychiatr
Res. 2012;46:428–435.

22.   Chang S-S, Chou T. A Dynamical Bifurcation Model of Bipolar Disorder Based on Learned
Expectation and Asymmetry in Mood Sensitivity. Computational Psychiatry. 2018;2:205–222.
23.   Cramer AOJ, Van Der Sluis S, Noordhof A, Wichers M, Geschwind N, Aggen SH, et al.
Dimensions of Normal Personality as Networks in Search of Equilibrium: You Can't like Parties if
you Don't like People. Eur J Pers. 2012;26:414–431.
24.   Cramer, Borkulo CD van, Giltay EJ, Maas HLJ van der, Kendler KS, Scheffer M, et al. Major
Depression as a Complex Dynamic System. PLOS ONE. 2016;11:e0167490.
25.   Cui J, Lichtwarck-Aschoff A, Olthof M, Li T, Hasselman F. From Metaphor to Computation:
Constructing the Potential Landscape for Multivariate Psychological Formal Models. Multivariate
Behavioral Research. 2023;58:743–761.
26.   Cui J, Hasselman F, Lichtwarck-Aschoff A. Unlocking nonlinear dynamics and multistability
from intensive longitudinal data: A novel method. 2023. 1 December 2023.
27.   Demic S, Cheng S. Modeling the Dynamics of Disease States in Depression. PLoS One.
2014;9:e110358.
28.   Durstewitz D, Huys QJM, Koppe G. Psychiatric Illnesses as Disorders of Network Dynamics.
Biol Psychiatry Cogn Neurosci Neuroimaging. 2021;6:865–876.
29.   Gauld C, Depannemaecker D. Dynamical systems in computational psychiatry: A toy-model
to apprehend the dynamics of psychiatric symptoms. Frontiers in Psychology. 2023;14.
30.   Globus GG, Arpaia JP. Psychiatry and the new dynamics. Biological Psychiatry.
1994;35:352–364.
31.   Gottschalk A, Bauer MS, Whybrow PC. Evidence of Chaotic Mood Variation in Bipolar
Disorder. Archives of General Psychiatry. 1995;52:947–959.
32.   Haken H, Tschacher W. How to Modify Psychopathological States? Hypotheses Based on
Complex Systems Theory. Nonlinear Dynamics Psychology and Life Sciences. 2017;21:19–34.
33.   Haslbeck J, Ryan O, Robinaugh D, Waldorp L, Borsboom D. Modeling Psychopathology:
From Data Models to Formal Theories. 2019. December 2019. https://doi.org/None.
34.   Hayes AM, Yasinski C, Barnes JB, Bockting CLH. Network destabilization and transition in
depression: New methods for studying the dynamics of therapeutic change. Clin Psychol Rev.
2015;41:27–39.
35.   Hayes AM, Laurenceau J-P, Feldman G, Strauss JL, Cardaciotto L. Change is Not Always
Linear: The Study of Nonlinear and Discontinuous Patterns of Change in Psychotherapy. Clin Psychol
Rev. 2007;27:715–723.
36.   Witkiewitz K, Marlatt GA. Modeling the complexity of post-treatment drinking: it's a rocky
road to relapse. Clin Psychol Rev. 2007;27:724–738.
37.   Foo JC, Noori HR, Yamaguchi I, Vengeliene V, Cosa-Linan A, Nakamura T, et al.
Dynamical state transitions into addictive behaviour and their early-warning signals. Proceedings of
the Royal Society B: Biological Sciences. 2017;284:20170882.
38.   Serre F, Fatseas M, Swendsen J, Auriacombe M. Does craving intensity influence cue
exposure reports? An ecological momentary assessment study in patients with alcohol, tobacco,
cannabis and heroin use disorder. Drug and Alcohol Dependence. 2015;C:e201.
39.   Fatseas M, Serre F, Swendsen J, Auriacombe M. Effects of anxiety and mood disorders on
craving and substance use among patients with substance use disorder: An ecological momentary
assessment study. Drug Alcohol Depend. 2018;187:242–248.
40.   Denis C, Fatséas M, Beltran V, Serre F, Alexandre J-M, Debrabant R, et al. Usefulness and
validity of the modified Addiction Severity Index: A focus on alcohol, drugs, tobacco, and gambling.
Subst Abus. 2016;37:168–175.
41.   McLellan AT, Kushner H, Metzger D, Peters R, Smith I, Grissom G, et al. The Fifth Edition
of the Addiction Severity Index. J Subst Abuse Treat. 1992;9:199–213.
42.   Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al. The Mini-
International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured
diagnostic psychiatric interview for DSM-IV and ICD-10. J Clin Psychiatry. 1998;59 Suppl 20:22-
33;quiz 34-57.
43.   Fatseas M, Serre F, Swendsen J, Auriacombe M. Effects of anxiety and mood disorders on
craving and substance use among patients with substance use disorder: An ecological momentary
assessment study. Drug and Alcohol Dependence. 2018;187:242–248.

44.   Serre F, Gauld C, Lambert L, Baillet E, Beltran V, Daulouede J, et al. Predictors of substance
use during treatment for addiction: A network analysis of ecological momentary assessment data.
Addiction. 2024:add.16658.
45.   Chirokoff V, Dupuy M, Abdallah M, Fatseas M, Serre F, Auriacombe M, et al. Craving
dynamics and related cerebral substrates predict timing of use in alcohol, tobacco, and cannabis use
disorders. Addiction Neuroscience. 2023;9:100138.
46.   Shumway RH, Stoffer DS. ARIMA Models. In: Shumway RH, Stoffer DS, editors. Time
Series Analysis and Its Applications: With R Examples, Cham: Springer International Publishing;
2017. p. 75–163.
47.   Mushtaq R. Augmented Dickey Fuller Test. 2011.
48.   Vagropoulos SI, Chouliaras GI, Kardakos EG, Simoglou CK, Bakirtzis AG. Comparison of
SARIMAX, SARIMA, modified SARIMA and ANN-based models for short-term PV generation
forecasting. 2016 IEEE International Energy Conference (ENERGYCON), 2016. p. 1–6.
49.   Alharbi FR, Csala D. A Seasonal Autoregressive Integrated Moving Average with Exogenous
Factors (SARIMAX) Forecasting Model-Based Time Series Approach. Inventions. 2022;7:94.
50.   Gauld C, Depannemaecker D. Dynamical systems in computational psychiatry: A toy-model
to apprehend the dynamics of psychiatric symptoms. Front Psychol. 2023;14:1099257.
51.   Van Rossum G, Drake FL. Python 3 Reference Manual. Scotts Valley, CA: CreateSpace;
2009.
52.   Ahmed SH. Validation crisis in animal models of drug addiction: Beyond non-disordered
drug use toward drug addiction. Neuroscience & Biobehavioral Reviews. 2010;35:172–184.
53.   Pezard L, Nandrino JL, Renault B, ElMassioui F, Allilaire JF, Muller J, et al. Depression as a
dynamical disease. Biol Psychiatry. 1996;39:991–999.
54.   Tschacher W, Scheier C, Hashimoto Y. Dynamical analysis of schizophrenia courses.
Biological Psychiatry. 1997;41:428–437.
55.   van de Leemput IA, Wichers M, Cramer AOJ, Borsboom D, Tuerlinckx F, Kuppens P, et al.
Critical slowing down as early warning for the onset and termination of depression. Proceedings of
the National Academy of Sciences. 2014;111:87–92.
56.   Tschacher W, Haken H. Causation and chance: Detection of deterministic and stochastic
ingredients in psychotherapy processes. Psychotherapy Research. 2020;30:1075–1087.
57.   Odgers CL, Mulvey EP, Skeem JL, Gardner W, Lidz CW, Schubert C. Capturing the ebb and
flow of psychiatric symptoms with dynamical systems models. Am J Psychiatry. 2009;166:575–582.
58.   Heiby E, Pagano I, Blaine D, Nelson K, Heath R. Modeling Unipolar Depression as a Chaotic
Process. Psychological Assessment. 2003;15:426–434.
59.   Wichers M, Schreuder MJ, Goekoop R, Groen RN. Can we predict the direction of sudden
shifts in symptoms? Transdiagnostic implications from a complex systems perspective on
psychopathology. Psychol Med. 2019;49:380–387.
60.   Nelson B, McGorry PD, Wichers M, Wigman JTW, Hartmann JA. Moving From Static to
Dynamic Models of the Onset of Mental Disorder: A Review. JAMA Psychiatry. 2017;74:528–534.
61.   Salvi JD, Rauch SL, Baker JT. Behavior as Physiology: How Dynamical-Systems Theory
Could Advance Psychiatry. AJP. 2021;178:791–792.
684

Figure 1. Phase space representation of the co-evolution of the cues, craving and use for two

randomly chosen patient from the dataset. The relations between these variables are complex and no

trivial linear interdependence appear.

Figure 2. Dynamic representation of the temporal profile of patients with Substance Use Disorder

(SUD) belonging to the *SARIMAX-model 1* ("cues-to-craving") (n=154), for 4 days. **Top**: Space phase

of the model, i.e., the conceptual framework illustrating all possible states of the system and the

interactions between variables over time. This 3D representation illustrates the dynamic interactions

between craving, cues, and use, providing a visual model of how fluctuations in one variable

correspond with variations in the others. Although such a depiction may not be immediately readable,

it remains essential for capturing the complex, nonlinear relationships inherent in dynamic systems

and is commonly used to elucidate interdependencies across multiple variables. **Bottom**: Temporal

representation of the model (4 times 12 hours, hours are noted every 4 hours and over 24 hours for the

sake of clarity). Since the model equations require reversing the sign of the cues, the increase in cue

salience in clinical practice, as described in the manuscript, is here visualized as a decline in the curve

(i.e., an increase of the cues is visible as negative events in the z variables). SARIMAX: Seasonal

Auto-Regressive Integrated Moving Average with eXogenous variable; DST: Dynamical System

Theory.

Figure 3. Dynamic representation of the temporal profile of patients with Substance Use Disorder

(SUD) belonging to the *SARIMAX-2* model ("craving-to-cues") (n=211), for 4 days. **Top**: Space phase

of the model, i.e., the conceptual framework illustrating all possible states of the system and the

interactions between variables over time. This 3D representation illustrates the dynamic interactions

between craving, cues, and use, providing a visual model of how fluctuations in one variable

correspond with variations in the others. Although such a depiction may not be immediately readable,

it remains essential for capturing the complex, nonlinear relationships inherent in dynamic systems

and is commonly used to elucidate interdependencies across multiple variables. **Bottom**: Temporal

representation of the model (4 times 12 hours, hours are noted every 4 hours and over 24 hours for the

sake of clarity). Since the model equations require reversing the sign of the cues, the increase in cue

salience in clinical practice, as described in the manuscript, is here visualized as a decline in the curve

(i.e., an increase of the cues is visible as negative events in the z variables). SARIMAX: Seasonal

Auto-Regressive Integrated Moving Average with eXogenous variable; DST: Dynamical System

Theory.

Figure 4. Temporal sequence of the interaction of the three variables of use, craving and cues in the

DST-model 1 ("cues-to-craving"). These relationships are described here in a linear manner in order to

facilitate understanding but should be interpreted in a non-linear manner (in particular with mutual

and reciprocal interactions) within the working framework of the DST. The arrow indicates the trigger

variable in this model.

Figure 5. Temporal sequence of the interaction of the three variables of use, craving and cues in the

DST-model 2 ("craving-to-cues"). These relationships are described here in a linear manner in order to

facilitate understanding but should be interpreted in a non-linear manner (in particular with mutual

and reciprocal interactions) within the working framework of the DST. The arrow indicates the trigger

variable in this model.

Tables

Table 1. Characteristics of the Substance Use Disorder (SUD) sample (n=211 participants): primary

substance, demographic variables and substance use-related information. ASI: Addiction Severity

Index; MINI: Mini International Neuropsychiatric Interview–Plus; SD: standard deviation.

	Sample characteristics	All participants (n=211)	
		% (n)	Mean (SD)
Primary substance use		37 (3,087)	
Primary substance	Alcohol	35 (73)	–
	Tobacco	28 (58)	–
	Cannabis	18 (39)	–
	Opiates	16 (34)	–
	Cocaine	3 (7)	–
Demographic variables	Sex (female)	36 (78)	–
	Employed	56 (116)	–
	Age	–	38.2 (11.4)
	Education (years)	–	12.5 (2.74)
	Pharmacological treatment	79 (167)	–
Substance use-related information	ASI addiction severity (0-9)	–	6.6 (0.76)
	Mood disorder on the MINI (current diagnosis)	34 (72)	–
	Anxiety disorder on the MINI (current diagnosis)	45 (93)	–

738 **Table 2**. Ecological Momentary Assessment (EMA) of intensity of craving, number of cues, and

739 reports of primary substance use on the Substance Use Disorder (SUD) sample of participants

740 (n=211). SD: standard deviation.

EMA variables	All participants (n=211 – 8,260 observations)	
	% (n)	Mean (SD)
Craving episodes (intensity ≥ 1)	70 (5,810)	–
Craving intensity (1-7)	–	3.6 (2.10)
Number of cues	–	4.3 (3.4)
Use	37 (3,087)	–

742 **Table 3.** Socio-demographic characteristics, substances types, addiction severity, psychiatric

743 diagnoses and EMA variable characteristics of the two temporal profile (SARIMAX) of patients with

744 Substance Use Disorder (SUD) (n=211). SARIMAX: Seasonal Auto-Regressive Integrated Moving

745 Average with eXogenous variable. "Model 1": SARIMAX-model 1 ("cues-to-craving"); "Model 2":

746 SARIMAX-model 2 ("craving-to-cues").

	Model 1 (N= 154)				Model 2 (N=57)					
	%	n	Mean	SD	%	n	Mean	SD	Test	p*
Sample characteristics										
Sex (female)	36.4	56			38.6	22			χ^2=0.089	0.765
Employed	55.6	84			57.1	32			χ^2= 0.038	0.845
Age			37.6	11.1			39.7	11.3	F= 1.53	0.217
Education (years)			12.4	2.6			12.5	3.1	F= 0.021	0.886
Primary substance										
Alcohol	30.5	47			45.6	26				
Tobacco	34.4	53			8.7	5				
Cannabis	20.1	31			14.0	8				
Opiates	13.0	20			24.6	14				
Cocaine	1.9	3			7.0	4				
Addiction severity										
ASI ISR (0-9) [§]			6.6	0.8			6.5	0.7	F= 0.088	0.767
Current pharmacological treatment for addiction	80.5	124			75.4	43			χ^2=0.651	0.419
MINI current diagnosis										
Mood disorder	32.0	48			42.9	24			χ^2= 2.11	0.146
Anxiety disorder	42.0	63			52.6	30			χ^2= 1.89	0.170
EMA variables [#]	6004 observations				2256 observations					
Frequency of Craving episodes (intensity > 1)			73.3	32.4			63.7	27.7	F= 3.98	**0.0472**
Craving intensity (1-7)			3.7	1.6			3.2	1.3	F= 4.25	**0.0405**
Number of cues			4.9	2.5			2.8	1.7	F= 35.91	**<.0001**
Frequency of primary substance use			42.4	35.1			26.6	29.9	F= 9.08	**0.0029**

748 **Table 4**. Parameter values of the two dynamical system models fitted with linear models, for the

749 Substance Use Disorder (SUD) sample of participants (n=211). DST: Dynamical System Theory. The

750 details of the parameters and their clinical translation are given in **Supplementary Material 4**.

DST models	Parameters of x, y and z variables													
	x				y						z			
	S_{max}	R_s	λ_s	τ_x	B_{max}	R_b	λ_b	P	L	τ_y	S	α	β	τ_z
DST-model 1	10	1	0.1	14	10	1.04	0.05	10	1.01	14	1.0	0.5	0.5	1
DST-model 2	10	1	0.1	2	10	0.904	0.05	1	1.00	2	1.0	0.5	0.5	1

752 **Figures**
753

756 **Figure 1.**

757
758

761   **Figure 2**

765 **Figure 3**

768   **Figure 4**

771 **Figure 5**

Funding source

Funding for the EMA studies was provided by Research Grant AAP-Recherche-CRA (20091301018) from the Aquitaine Regional Council (MA) and IRESP-19-ADDICTIONS-16 from IReSP and Alliance Aviesan for CUSEMA study (FS). For this study, MA and FS received financial support from the French government in the framework of the University of Bordeaux's IdEx "Investments for the Future" program / GPR BRAIN_2030. Office and staff support was provided by CH Charles Perrens. The funding sponsors had no role in the design of the study or interpretation of the data. They were not involved in the preparation, review, or approval of this manuscript.

Data availability

The data that support the findings of the current manuscript are available from the corresponding author, upon reasonable request.

Authors' contributions

Marc Auriacombe and Fuschia Serre were the principal investigators, obtaining funding and supervised the study. Marc Auriacombe provided access to participants. Damien Depannemaecker and Christophe Gauld conceptualized and undertook analysis, interpretation of data and drafting of manuscript. Fuschia Serre participated in patient recruitment, data collection and database management. Marc Auriacombe and Fuschia Serre sustained the elaboration of the methodology and writing. All authors undertook the critical revision of the manuscript for important intellectual content and all authors significantly contributed to the manuscript and approved the final version.