Computational modeling of the temporal influences between cues, craving and use in addiction: A dynamical system analysis based on ecological momentary assessment data

Running title: Craving and dynamical systems

Christophe Gauld [1,2,3,†,*], Damien Depannemaecker [4,†], Marc Auriacombe [3,5,6,X,*], Fuschia Serre [3,5,6,X]

Affiliations
[1] Department of Child Psychiatry, CHU de Lyon, F-69000 Lyon, France
[2] Institut des Sciences Cognitives Marc Jeannerod, UMR 5229 CNRS & Université Claude Bernard Lyon 1, F-69000 Lyon, France
[3] Sleep, Addiction and Neuropsychiatry (SANPSY), Centre National de la Recherche Scientifique, (CNRS), UMR 6033, F-33076 Bordeaux, France
[4] Aix Marseille Université, Institut National de la Santé et de la Recherche Médicale, Institut de Neurosciences des Systèmes (INS) UMR1106, F-13005 Marseille, France
[5] University of Bordeaux, F-33076 Bordeaux, France
[6] Pôle interétablissement d'addictologie, CHU de Bordeaux et CH Charles Perrens, F-33076 Bordeaux, France

[†] : These authors contributed equally as first author.
[X] : These authors contributed equally as last author.

* Corresponding authors
Marc Auriacombe
e-mail: marc.auriacombe@u-bordeaux.fr
Sanpsy-addiction, Pôle Addictologie, CHCP, 121 rue de la Béchade, 33076 Bordeaux Cedex, France
Tel: +33 556 561 738

Main text words: 6009
Abstract words: 269
References: 61
Figures: 5
Tables: 4

Abstract

Substance Use Disorders (SUD) can be conceptualized as a prospective link from cues to craving and

use. To explore the nonlinear relationships between craving and cues, this study applied dynamical

systems theory (DST) to ecological momentary assessment (EMA) data. Optimized linear Seasonal

Auto-Regressive Integrated Moving Average with eXogenous variable (SARIMAX) models were used

to phenotype patients with SUD (alcohol, tobacco, cannabis, opiates, and cocaine), considering the

potential for complex interactions between cue exposure and craving intensity in daily life. These

phenotypic profiles were replicated in computational DST models to analyze the nonlinear interactions

between cues, craving, and use. The study involved 211 individuals and 8,260 observations, with 154

patients fitting the SARIMAX model for the influence of cues on craving, and 57 patients fitting the

SARIMAX model for a possible influence of craving on cues. Two DST models were adjusted to

replicate the complex temporal dynamics of SUD based on these two directions of influence. The first

DST model (adjusted to the influence of cues on craving) showed that an increase in cues leads to a rise

in craving, which then diminishes both cues and craving itself, with use patterns following craving's

trajectory. This patient profile is driven by a phenomenon of "maximum cue saturation". The second

DST model (adjusted to the influence of craving on cues) demonstrated that an increase in craving was

followed by an increase in cue reporting, leading to use, with use peaking and then reducing craving.

This patient profile is characterized by a phenomenon of "maximum use saturation". Both models

highlight craving as an essential modulator between cues and use, opening new therapeutic avenues.

Keywords: substance use disorder; craving; ecological momentary assessment; dynamical system;

biomarker; addiction.

Introduction

Substance Use Disorder (SUD) is characterized by recurrent relapses despite efforts to reduce use or abstain [1]. One of the main objectives of SUD treatment is to reduce the frequency of relapses over time. As an important symptom of SUD [1–8], the fluctuating phenomenon of craving can be captured by daily-life monitoring systems such as ecological momentary assessment (EMA) [9]. EMA, adapted and validated to study craving in SUD [10], allows a real-time measurement of individual momentary states in the natural environment through repeated assessments overtime. Based on EMA research, a human model of addiction has been suggested across several substances and behaviors: the cues – craving – use model [1, 4, 11]. In this model, an increase in the number of reported cues is associated with an increase in the intensity of self-reported craving in the following EMA survey, itself associated with an increased likelihood of use in the following EMA survey. This craving – use link has been considered unidirectional [11–14] using mixed models like hierarchical linear models (HLM), which allows for the analysis of data with a nested structure, accounting for variability at multiple levels of observation. However, HLM has some limitations such as the difficulty in accounting for cross-effects, i.e., integrating complex interactions between levels. Furthermore, they are only linear: the effects (e.g., craving) may be directly proportional to the causes (e.g., cues) – as opposed to non-linear models where the relationships between variables can be disproportionate, interactive, producing effects which are not predictable simply by addition or multiplication. Finally, HLM cannot account for multidirectionality, e.g., that cues and craving could influence each other in a loop. Due to the important intra-individual variability and its intrinsic non-linearity, the evolutionary dynamics of a specific patient with a SUD are particularly complex to model.

To go further, modeling the evolution of SUD variables and their interactions requires finding alternatives to traditional statistical methods [2]. A growing literature based on dynamical systems theory (DST) seeks to capture the temporal dynamics of psychiatric conditions (e.g., [15–35]), including SUD. DST accounts for the evolution of clinical manifestations over time without abolishing the complexity of the relationships between variables and their multidirectionality, allowing the prediction of different types of temporal evolution. This theory is based on dynamic objects (e.g., tipping points,

transitions, steady states, strange attractors, limit cycles or oscillations) and dynamic concepts (e.g.,

emergence, instability, or variations related to fluctuations in initial conditions – referring to the

"butterfly effect", etc.). However, most psychiatric studies include biological factors (e.g., [36]) and

few studies have empirically included psychiatric clinical manifestations (e.g., [19]).

Based on EMA data, considering the possibility that, in addition to a unidirectional prospective link

from cues to craving, an increase in craving could change perception of cues [37], DST models allow

the representation of dynamics and mutual influences between variables. The main goal of this study

was to model alternative evolutions of craving and cues, to explore their influence on substance use.

Such results could lead to proposing a classification of individuals with SUD based on dynamics'

profiles, as well as providing information on the attractive states (i.e., the states to which the system

gravitates) – allowing to understand what stabilizes the patient in a given (healthy or pathological) state.

Methods

This study is based on two successive stages: i) the use of optimized linear statistical models

(SARIMAX) to identify, based on EMA data, two patient profiles according to the direction of influence

between cues and craving; ii) the use of a computational model (DST) to analyze the complex

relationships between cues and craving according to these profiles. In line with the classic cues-craving-

use model (in which an increase in cues can linearly contribute to an increase in craving), we

hypothesized that a state of craving could make an individual more "sensitive" or more "receptive" to

cues, with an effect on use, creating a feedback loop: cues lead to craving, which increases cue

perception, leading to more craving, and ultimately, more use. This reflects a non-linear dynamic where

continuous mutual interactions bring about a dynamic process. The modeling focuses on the cues-

craving relationship, as use is the outcome.

Data collection in EMA

Participants

Data were extracted from two EMA studies with similar protocol [10, 43], which comprised individuals

aged 18 years and above, who were beginning addiction outpatient treatment (in Bordeaux and

Bayonne, France), were diagnosed with at least one DSM-5 Substance Use Disorder for tobacco,

alcohol, cannabis, cocaine or opiates, and were free from active psychosis or severe cognitive

impairment [38]. Informed consent was obtained, and the studies were approved by the French

Regulation and ethical committee (CPP SOOM III/DC-2009/01; CNIL/DR-2015-408; CPP IDF X: 11-

2019/IR-RCB: 2018-A00952-53; CNIL/MR003). All methods were performed in accordance with the

relevant guidelines and regulations. Participants received standard comprehensive care, consisting of

individual behavioral treatment focused on relapse prevention and psychosocial support combined,

when available, with pharmacotherapy. After establishing a target quit date according to the patient's

personal goals, full abstinence or use reduction was encouraged as an outcome, but with no negative

consequences for the patient if failure to achieve this goal.

Assessments

Each participant underwent an initial interview to collect sociodemographic variables and substance

use-related information using a validated French version of the Addiction Severity Index (ASI) [39,

40], modified to consider tobacco addiction. The Mini International Neuropsychiatric Interview–Plus

(MINI) structured interview [41] was adapted to evaluate the current DSM-5 SUD status within the past

year. When multiple SUD co-occurred, the primary substance was determined according to the main

problematic substance reported by the individual, and on which the treatment was focused.

Ecological momentary assessment

The EMA protocol was previously described in detail [10]. EMA assessments began within the first

month after treatment initiation. Following a training session, each participant received a tablet to carry

for 14 days. Each tablet was programmed to administer four electronic surveys per day, inquiring about

primary substance use, craving, and the presence of cues. The signal schedule was adjusted to

accommodate participants' usual sleep patterns. Financial compensation was provided based on the

number of surveys completed, with a maximum of 100 euros for participants who completed 75% or

more of the assessments. Methodological details and comparisons with other studies using this validated

protocol [11, 42–44] are presented in **Supplementary Material 1**.

Data of interest

At each electronic assessment, participants were asked to rate the maximum intensity level of desire to

use their primary substance they had felt since the previous assessment on a 7-point scale (1 = no desire

to use 7 = extreme desire to use). This item was referred to as "craving". Participants were also asked

if they had encountered cues related to their primary substance, selected from a personalized list for

each individual. This item was labeled "cues" and included specific objects or contexts typically

associated with substance use, such as materials used for consumption (e.g., a bottle), the sight or smell

of the substance, or places associated with use (e.g., a bar). It also encompassed personal cues, identified

during a personal interview, that referenced specific circumstances, objects, individuals, emotions, or

environmental contexts linked to their substance use (e.g., "being with my friend Sam") [11]. The

"cues" variable is a count measure that ranges from 0 upwards, with no upper limit, reflecting the total

number of cues reported since the previous assessment. Although these refer to "cue perceptions", for

reading convenience, we used the term "cues" below. Finally, participants were asked if they had used

the primary substance, with this variable being binary (0 = no, 1 = yes), focusing solely on whether

substance use occurred since the prior assessment. This item was referred to as "use". All three

questions inquired about participants' experiences since the previous assessment.

Due to a restricted number of assessments occurring on the starting and ending day of the study, and

their overlap with investigator contact, only data collected between days 2 and 13 were analyzed (n=12

days) (**Table 2**).

Statistical and computational models

Aims of the models

First, as detailed in **Supplementary Material 2**, we used optimized linear SARIMAX models with

craving and cues to determine the mutual influence profiles of patients. We aimed to identify: i) a

SARIMAX-1 model*,* referring to the influence of cues on craving, hereafter named "cues-to-craving"

model; ii) a *SARIMAX-2* model, referring to the influence of craving on cues, hereinafter named

"craving-to-cues" model.

Secondly, we used the SARIMAX model 1 to determine the specific parameter values of a model based

on the DST (*DST-model 1*), and the SARIMAX model 2 to determine the specific parameter values of

another model based on the DST (*DST-model 2*). We qualitatively observed the complex evolution of

each of the two variables within the two DST models. "Use" was treated as an output of the model, with

its values deliberately left unconstrained to follow the parameterized dynamics of the cues-craving

structure in both DST models. This approach allowed us to capture the intricate interactions between

cues, craving, and use.

By combining linear SARIMAX models with DST models, we aimed to create a framework that

captures both the linear and nonlinear dynamics of craving and cues. The SARIMAX models first

classified patients based on their interaction profiles. These profiles served to establish parameter values

for the DST models, which further analyze the nonlinear interactions between craving and cues. This

approach allowed us to utilize linear models for the initial sequence of variables without necessarily

providing a parameter characterization. Subsequently, the DST models determine these parameters to

capture intricate feedback loops between craving, cues, and use. Though SARIMAX models were

constrained by assumptions of linearity and stationarity, this hybrid approach could offer an interesting

tool for understanding complex craving dynamics across varied profiles. As we will describe later, the

comparison of SARIMAX and DST models is a methodological challenge due to differences in scope

and (non)linearity. Specifically, in the DST framework, the parameter ranges are reconstructed to fit

the data, rather than being directly calculated from empirical measurements. This approach emphasizes

conceptual alignment with circadian-scale dynamics rather than exact value matching.

To ensure the reliability of our method, we conducted a simulation study by generating two groups

based on distinct craving-to-cues and cues-to-craving dynamics, according to the DST model parameter

distributions. We then verified our ability to recover these groups using SARIMAX modeling, applying

separate models to identify each directional influence. Finally, based on the bifurcation analysis

described below, we checked whether the parameter values of the DST model could be recovered for

each of the two groups, ensuring consistency between the simulated data and the underlying DST model

parameters (**Supplementary Material 3**).

SARIMAX models

SARIMAX models, extensively detailed in **Supplementary Material 2**, are linear algorithms for

forecasting data, considering past values (AR – AutoRegressive) and data noise (MA – Moving

Average) to predict future values [45].

Stationarity and seasonality assessments of the EMA data. Stationarity is a condition of validity for this

time-series analysis using the Auto-Regressive Moving Average (AR-MA) method. The stationarity of

the time-series data for craving and cues was assessed using the Augmented Dickey-Fuller (ADF) test

[46]. If the time series was not stationary, an integration factor ($I(d)$) was added to the model, allowing

differentiation to make the data stationary (i.e., resulting in a "AR-I-MA" model). Seasonality

(classically denoted S in the model, providing a "S-ARIMA" model) was considered given the repeated

daily nature of the data. Finally, an exogenous variable (denoted X in the model, resulting in a

"SARIMA-X" model) could be added to model an additional effect on the time series of the first

variable (i.e., cues for the craving time series, and craving for the cues time series) [47, 48].

Model fitting. For each patient, two SARIMAX models were fit to the data: one where craving was

modeled as the dependent variable influenced by cues (*SARIMAX-model 1 –* "cues-to-craving" model),

and another where cues were the dependent variable influenced by craving (*SARIMAX-model 2 –*

"craving-to-cues" model). Different combinations of autoregressive and moving average terms were

tested for each patient. Only the parameter values for the equations governing y (craving) and z (cues)

were adjusted to align with each of the two SARIMAX models. The parameter values for the equation

of x (use) were deliberately left unconstrained, allowing to observe its dynamics as an output variable

within the models.

Parameter optimization. The y and z parameters of the two SARIMAX models were optimized

individually for each patient. A grid search was performed across potential values of the AR and MA

component, identifying the combination that minimized the Akaike Information Criterion (AIC). This

criterion balances model fit and complexity to prevent overfitting. The choice to use the AIC over the

Bayesian Information Criterion (BIC) was motivated by the objective of optimizing predictive accuracy

while accommodating the complexity of individual temporal dynamics, as AIC imposes a less stringent

penalty on model parameters, thus enhancing the model's sensitivity to subtle within-subject variations.

Two AIC were provided per patient (one for each SARIMAX model of a patient).

Model selection. For each patient, the preferred SARIMAX model (either *SARIMAX-model 1* or

SARIMAX-model 2) was determined based on the lower AIC. We assume that each patient aligns with

only one specific model, excluding the other. Following the identification of which participants best fit

each model, we compared the two selected subgroups based on socio-demographic characteristics,

substance types, addiction severity, psychiatric diagnoses, and EMA variable characteristics (**Table 3**).

This comparison aimed to 1) clinically describe the profiles of patients displaying different dynamic

patterns (cues to craving versus craving to cues), and 2) methodologically confirm the absence of

significant differences between the groups, ensuring that any observed distinctions in model dynamics

were not attributable to these underlying characteristics. Details on these models are provided in

Supplementary Material 2.

Sensitivity analysis. We then performed a sensitivity analysis on the sub-datasets of alcohol and cocaine

to assess the stability of the models across these substance-specific data.

Auto-correlation analysis. Finally, we proposed studying the temporal dynamics between craving and

cues through auto-correlation and cross-correlation analyses, allowing to identify individual temporal

structures, key time lags, and descriptive statistics revealing variability in craving and cues across the

sample.

DST models

Dynamical system modeling. A set of differential equations was used to model the dynamical

relationship between craving and cues. The model was integrated over the 12-day observation window

and extended to a theoretically infinite duration to visualize the temporal interaction of one variable on

the other.

Details of the DST model. The detail of the DST model has been described elsewhere [28]. In a nutshell,

it is based on four differential equations, producing a nonlinear representation of participant's internal

factors (i.e., subjective phenomenological experiences and/or biological elements) (*y*) and of

environmental noise (*z*). Based on the empirical data collected in EMA regarding SUD, the "patient's

internal factor" corresponds to the intensity of craving (*y*), and the "environmental noise" corresponds

cues (more precisely, to the reporting of cues) (*z*). Two other variables allow modulating: i) the long-

term temporality of patients' conditions (*f*); ii) and the intensity of behavior (*x*), corresponding here to

substance use (passively evolving according to the two cues-craving models). Thirteen parameters are

described within these four variables, with their clinical translation, presented in **Supplementary**

Material 4. Details of interactions between parameters within each variable are described elsewhere

[28], but the four differential equations and the thirteen parameters of the model are presented below[1]:

$$\dot{X} = \begin{bmatrix} \dot{x} \\ \dot{y} \\ \dot{z} \\ \dot{f} \end{bmatrix} = \begin{bmatrix} \frac{S_{\max}}{1+\exp\left(\frac{R_s - y}{\lambda_s}\right)} - x \\ \frac{P}{1+\exp\left(\frac{R_b - y}{\lambda_b}\right)} + fL - xy - z \\ S(\alpha x + \beta y)\zeta(t) - z \\ y - \lambda_f f \end{bmatrix}$$

269   As an example, **Figure 1** presents two simulations of the DST model with different parameter values,

270   by considering three random variables. We observe that the simulated patient trajectories in the model's

271   "relief" differ when the parameter values are varied.

273   **[FIGURE 1]**

---

[1] *x*: symptom intensity, representing use; *y*: participant's internal factors, representing craving intensity; *z*: environmental noise, representing the reporting of cues. $\tau_{x,y,z,f}$: time scales of the four equations; $S_{\max}$: maximal symptom level; R_s and R_β: sensitivity, with R_s for symptom sensitivity and R_β for internal element (potentiation) sensitivity; λ_s and λ_β: slope of symptom/internal element curves, with λ_s affecting symptom intensity based on patient state γ, and λ_β affecting internal elements based on environment z; P: maximal rate of internal elements, aggregating biological, psychological, and phenomenological factors, shaping phenotype from biological signals; L: level of predisposing factors contributing to a permanent shift in internal elements; λ_f: scaling factor for slow fluctuations affecting L; S: overall environmental sensitivity; α and β: weights of x and y effects on environmental perception.

Construction of the two DST models. We built two DST models with specific parameter values. Given

the short time scale considered (tens of days), we fixed the *f* variable – originally representing

predisposing factors over months and years. The parameter values in the *y* (craving) and *z* (cues)

equations were adjusted to align with each of the two SARIMAX models.

To identify the regions of interest in the parameter space, we performed a bifurcation analysis.

Bifurcation analysis is a well-known method in the field of dynamical systems to study how the

qualitative behavior of a system changes as parameters are varied, identifying transitions such as the

emergence of cycles, stability, or new equilibrium states [49, 50]. We focused on the parameters

associated with the dynamics of the internal state (craving), specifically λ_b, R_b, and *L*. This allowed to

determine the ranges of values that enable the existence of a limit cycle, corresponding to the observed

circadian oscillations. Similarly, we identified the regions of the parameter space that allow for the

existence of a stable fixed point, representing a situation where the craving variable does not oscillate

spontaneously but remains sensitive to perturbations induced by external inputs related to cues.

To explain it differently, we identified a subset of the parameter space using bifurcation analysis,

representing the possible dynamical repertoire that matches the circadian evolution (i.e., changes

observed over a 24-hour day-night cycle) observed in the data. This selection was based on the features

associated with temporality and rhythmicity, central to interpreting model comparisons. The parameter

adjustment process relied on an iterative hands-on fit, enabling the reproduction of sequential patterns

of emergence and fluctuation between craving and cues in each model. This empirical manual

exploration of the parameters of the system, to align with the perceived temporal dynamics, was

conducted without automated method (e.g., simulation-based inference, to be applied in future works).

The simulations shown here were chosen as the best matching observation within the range identified

previously.

Then, for each subject, the best-fitting linear SARIMAX model (either cues-to-craving or craving-to-

cues) was selected and subsequently used to fit a nonlinear DST model, parameterized to capture their

dynamics. The parameter adjustment process was conducted through successive simulations, allowing

to iteratively refine the model to obtain a qualitative understanding of the system's behavior while

ensuring it aligned with the observed dynamics in the SARIMAX models. For instance, the parameter

L (representing the "vulnerability level" in the *y* equation) may need to be set high to reflect the

increasing influence of craving on cues. Such parameterization of vulnerability would constitute an

interesting result: for craving to affect sensitivity to cues, a patient must have a high level of

vulnerability (e.g., genetic predispositions, psychological vulnerability, neurological comorbidities,

etc.).

All the analyzes were performed with Python [51].

Code availability: The code is added to the open access repository:

https://github.com/ChristopheGauld/Dynamical_systems_for_craving.

313   **Results**

Sample description and assessments

As described in **Table 1**, 211 individuals were included in the study. The mean age was 38.2 years old

and 36% were female (n=36). Primary substance was: 35% alcohol, 28% tobacco, 18% cannabis, 16%

opiates and 3% cocaine. The flow-chart of participant inclusion is presented in **Supplementary**

Material 5.

Table 1. Characteristics of the sample (n=211 participants): primary substance, demographic variables,

substance use-related information and psychiatric comorbidity. ASI: Addiction Severity Index; MINI:

Mini International Neuropsychiatric Interview–Plus; SD: standard deviation.

	Sample characteristics	All participants (n=211)	
		% (n)	Mean (SD)
Primary substance	Alcohol	35 (73)	–
	Tobacco	28 (58)	–
	Cannabis	18 (39)	–
	Opiates	16 (34)	–
	Cocaine	3 (7)	–
Demographic variables	Sex (female)	36 (78)	–
	Employed	56 (116)	–
	Age	–	38.2 (11.4)
	Education (years)	–	12.5 (2.74)
	Pharmacological treatment	79 (167)	–
Substance use-related information and psychiatric comorbidity	ASI addiction severity (0-9)	–	6.6 (0.76)
	Mood disorder on the MINI (current diagnosis)	34 (72)	–
	Anxiety disorder on the MINI (current diagnosis)	45 (93)	–

EMA assessments

Among the 10,128 EMA electronic assessments planned and delivered, 8,260 (82%) were completed.

The majority of EMA assessments reported episodes of craving (70% with non-null intensity), and 37%

reported primary substance use (**Table 2**).

Table 2. Ecological Momentary Assessment (EMA) report of craving episodes and intensity, number

of cues, and reports of primary substance use of participants (n=211). SD: standard deviation.

EMA variables	All participants (n=211 – 8,260 observations)	
	% (n)	Mean (SD)
Craving episodes (intensity ≥ 1)	70 (5,810)	–
Craving intensity (1-7)	–	3.6 (2.10)
Number of cues	–	4.3 (3.4)
Use	37 (3,087)	–

SARIMAX models

The ADF test confirmed that the time-series does not have a time-dependent structure ($p = 0.0003$ for craving and $p \ll 0.001$ for cues, indicating that both series are stationary), supporting stationarity for each patient and for each variable (cues, craving and use). No detrending was necessary. No seasonality, which could have been present here due to typical day-by-day fluctuations, was found in the data. SARIMAX models could thus have been named "ARMAX", but we keep the initial term to ensure understanding.

After testing the two SARIMAX models for each patient and after individual optimization, we found 154 patients who fit better (according to the AIC) with the SARIMAX-model 1 ("cues-to-craving") and 57 patients who fit better with the SARIMAX-model 2 ("craving-to-cues"). As expected in the analysis seeking to maximize the fit of a patient with one of the two models, no patient belonged to both models. Details on the AIC of the 1 and 2 SARIMAX models and on the preferred model according to the lowest AIC are given in **Supplementary Material 2**. **Table 3** provides the characteristics of the patients who best fit each of the two SARIMAX models. There were no significant differences in the distribution of sociodemographic characteristics, addiction severity, psychiatric diagnoses, or substance types between patients fitting each model; however there were differences for some of the EMA variables (see **Table 3**).

Table 3. Socio-demographic characteristics, substances types, addiction severity, psychiatric diagnoses and EMA variable characteristics of the two temporal profile (SARIMAX) of patients with Substance Use Disorder (SUD) (n=211). SARIMAX: Seasonal Auto-Regressive Integrated Moving Average with eXogenous variable. "Model 1": SARIMAX-model 1 ("cues-to-craving"); "Model 2": SARIMAX-model 2 ("craving-to-cues").

	Model 1 (N= 154)				Model 2 (N=57)					
	%	n	Mean	SD	%	n	Mean	SD	Test	p*
Sample characteristics										
Sex (female)	36.4	56			38.6	22			χ^2=0.089	0.765
Employed	55.6	84			57.1	32			χ^2= 0.038	0.845
Age			37.6	11.1			39.7	11.3	F= 1.53	0.217

	% / Mean	N / SD	% / Mean	N / SD	Test	p
Education (years)		12.4	2.6		12.5	3.1	F= 0.021	0.886
Primary substance								**<.001**
Alcohol	30.5	47		45.6	26			
Tobacco	34.4	53		8.7	5			
Cannabis	20.1	31		14.0	8			
Opiates	13.0	20		24.6	14			
Cocaine	1.9	3		7.0	4			
Addiction severity								
ASI ISR (0-9)		6.6	0.8		6.5	0.7	F= 0.088	0.767
Current pharmacological treatment for addiction	80.5	124		75.4	43		χ^2=0.651	0.419
MINI current diagnosis								
Mood disorder	32.0	48		42.9	24		χ^2= 2.11	0.146
Anxiety disorder	42.0	63		52.6	30		χ^2= 1.89	0.170
EMA variables	**6,004 observations (38.99/patient)**			**2,256 observations (39.58/patient)**				
Frequency of craving episodes (intensity > 1)		73.3	32.4		63.7	27.7	F= 3.98	**0.0472**
Craving intensity (1-7)		3.7	1.6		3.2	1.3	F= 4.25	**0.0405**
Number of cues		4.9	2.5		2.8	1.7	F= 35.91	**<.0001**
Frequency of primary substance use		42.4	35.1		26.6	29.9	F= 9.08	**0.0029**

The results of the sensitivity analysis and of the auto-correlation analysis between craving and cues are presented in **Supplementary Material 2**.

DST models

Using successive simulations based on different parameterization of the z (cues) and y (craving) variables of the DST model, we sought to represent in a non-linear way (unlike SARIMAX models) the mutual interactions between craving and cues. We did not seek to modify the other two variables (f variable for long-term temporality or x variable for use). The parameter values of the equations of y (craving) and z (cues) equations are modified in order to fit with each of the two SARIMAX models. We defined two sets of parameters: the first set of parameters of the *DST-model 1,* corresponds to the dynamical system representation of the 154 patients who fit with the *SARIMAX-model 1*; the second set

of parameters of the *DST-model 2,* corresponds to the dynamical system representation of the 57 patients

who fit with the *SARIMAX-model 2*.

The details of the parameter values of the two DST models are provided in **Table 4**. Five parameter

values were necessarily modified to recreate the two SARIMAX-models: L, R_b, P, τ_x, and τ_y (in red in

Table 4). Firstly, the value of the L parameter (corresponding to the vulnerability level in the y equation)

was barely modified ($L = 1.01$ in the DST-model 1 and $L = 1.00$ in the DST-model 2) to obtain these

two profiles of influence of craving on cues. Secondly, the value of the parameter R_b (referring to the

sensitivity of the internal elements) was necessarily modified ($R_b = 1.04$ in the DST-model 1 and $R_b =$

0.904 in the DST-model 2): for craving to have an influence on sensitivity to cues (DST-model 2), a

patient must have a lower level of sensitivity. Thirdly, the value of the parameter P (representing

potentiation, or the maximal rate of internal elements amplifying symptoms) was necessarily increased

for DST-model 1 ($P = 10$) compared to DST-model 2 ($P = 1$), indicating that craving (which drives

DST-model 1) serves as an internal mechanism that amplifies use. Finally, τ_x and τ_y refer solely to the

time scales specific to each variable (e.g., $\tau_x \bullet dx/dt$) without clinical interpretation, and their increase

in DST-model 1 does not alter the interpretation of the other parameter values.

Table 4. Parameter values of the two dynamical system models fitted with linear models (n=211). DST:

Dynamical System Theory. The details of the parameters and their clinical translation are given in

Supplementary Material 4.

DST models	Parameters of x, y and z variables													
	x				*y*						*z*			
	S_{max}	R_s	λ_s	τ_x	B_{max}	R_b	λ_b	P	L	τ_y	S	α	β	τ_z
DST-model 1	10	1	0.1	14	10	1.04	0.05	10	1.01	14	1.0	0.5	0.5	1
DST-model 2	10	1	0.1	2	10	0.904	0.05	1	1.00	2	1.0	0.5	0.5	1

x: symptom intensity, representing use; *y*: participant's internal factors, representing craving intensity;
z: environmental noise, representing the reporting of cues. $\tau_{x,y,z,f}$: time scales of the four equations; S_{max}:
maximal symptom level; R_s and R_β: sensitivity, with R_s for symptom sensitivity and R_β for internal
element (potentiation) sensitivity; λ_s and λ_β: slope of symptom/internal element curves, with λ_s affecting
symptom intensity based on patient state γ, and λ_β affecting internal elements based on environment z;
P: maximal rate of internal elements, aggregating biological, psychological, and phenomenological
factors, shaping phenotype from biological signals; L: level of predisposing factors contributing to a
permanent shift in internal elements; λ_f: scaling factor for slow fluctuations affecting L; S: overall
environmental sensitivity; α and β: weights of x and y effects on environmental perception.

Dynamics of craving and cues on the *DST-model 1* and on the *DST-model 2* are represented in **Figure**

2 and **Figure 3,** respectively. Within these two figures, although there was no fixed scale due to the

nonlinear interactions between relative dynamics, it was important to consider that the baseline

represented an absence of interaction, visualized by an increase in a variable (e.g., the baseline for

craving represented the level corresponding to the absence of craving, when it did not spontaneously

increase or respond to another variable).

In **Figure 2** (DST-model 1 for *SARIMAX-model 1* patients, N=154), when an increase in cues (visible

as positive events in the z variable) preceded craving, there was a relatively rapid increase in craving,

which in turn reduced the cues, followed by a slower decrease in craving (**Figure 2**, bottom). Craving

peaked in alignment with peak cues. Craving only returned to its baseline level after cues also returned

to baseline. Use progressively increased throughout the duration of the craving rise (e.g., from 8 p.m.

to midnight on the 1st day) and only began to decrease in temporal conjunction with the relatively sudden

drop in craving. This temporal overlap between the decreases in craving and use limited our ability to

determine the direction of influence between the two variables. However, it is noteworthy that the

decrease in reporting of cues preceded this joint decrease in craving and use (e.g., at 4 p.m. on the 1st

day). The return to baseline may therefore be attributed to cue saturation, with a craving modulation

(highlighted by the observation that the return to baseline occurred only after a reduction in craving).

411   **[FIGURE 2]**

In **Figure 3** (*DST-model 2* for *SARIMAX-model 2* patients, N=57), when an increase in craving occurred

before cues, this prior craving increase influenced cue reporting, leading to greater simultaneity between

the two variables compared to the previous model (**Figure 3**, bottom). Craving decreased during the

period of noisy cue reporting, with both craving and cues returning to baseline simultaneously. Use

increased gradually throughout the duration of the craving increase and, as before, only began to

decrease with the relatively sudden drop in craving. Cue reporting returned to baseline in temporal

conjunction with this decrease in use and craving (unlike the earlier return seen in the previous model).

The return to the baseline state was therefore due to saturation of use, implying a modulation of craving.

422   **[FIGURE 3]**

Finally, the simulation study demonstrates the robustness of our modeling approach in reliably

distinguishing the craving-to-cues and cues-to-craving groups, confirming the SARIMAX method's

effectiveness in capturing these distinct directional influences based on different parameter distributions

of the DST models. Additionally, for each SARIMAX model, we verified the alignment of the

recovered simulations with the dynamics initially generated by the DST model, utilizing the same

bifurcation analysis as in the main analysis. This step further validated that the parameter values of the

DST model accurately reflected the distinct craving-to-cues and cues-to-craving dynamics

(**Supplementary Material 3**).

The main goal of this study was to contribute to understanding the multiple and complex influences of

craving and cues on substance use using dynamical system analysis based on ecological momentary

assessment data. To achieve this, two SARIMAX models were tested for each patient, resulting in 154

patients aligning better with *SARIMAX-model 1* ("cues-to-craving") and 57 with *SARIMAX-model 2*

("craving-to-cues"). Parameter sets were defined through a simulation analysis to propose two DST

models, representing the evolutionary and interactional dynamics of cues, craving, and use for patients

corresponding to each SARIMAX model.

Figure 4 illustrates the sequence of interactions between the variables in *DST-model 1*, from which two

main results could be drawn. The first finding offers insights into the mechanism of cessation of use in

these patients. Specifically, cessation of use was not due to the saturation of use itself (i.e., the point

where further increase is impossible due to reaching maximum capacity). Instead, cessation resulted

from a specific sequence involving cue reporting and craving. In this model, a peak in cue reporting

initially occurs. Once this peak is reached, we observe a gradual reduction in use, mediated by craving

levels. This decrease in cue reporting returns to the baseline level, ultimately leading to cessation of

use. In other words, cessation may have been triggered by the preceding peak in cue reporting,

modulated by craving levels. The second finding of this model addresses the complexity of the

interaction between craving and use. As observed in practice and documented in many studies, craving

leads to an increase in use [2]. In this model, cue reporting precedes craving, but craving peaks

simultaneously with the peak in cue reporting. A high level of cue reporting and craving is followed by

a gradual decrease in cue reporting to a baseline threshold, resulting in cessation of use. The direct

interaction between cue reporting and use remains unclear – specifically, whether cue reporting directly

saturates use, given that cue reporting precedes use in this model. However, modulation by craving

clarifies the interactions among these three variables. The model suggests that cue reporting and craving

rise in tandem, leading to an increase in use. At the peak of craving, cues also reach their maximum.

After this peak, craving begins to decrease, which in turn reduces cue reporting to the baseline,

eventually leading to cessation of use. Importantly, use does not interrupt the craving peak (since a

reduction in craving is necessary once it reaches its peak). The conclusion here is that use plays a

secondary role: it follows craving, which controls the peak in cue reporting, and it also follows the

reduction in cue reporting.

[FIGURE 4]

Figure 5 illustrates the sequence of interactions between the variables in *DST-model 2*. The primary

result from this model pertains to the role of use in the relationships between craving, cues, and use.

In this model, craving increases and simultaneously (or immediately after) enhances cue reporting.

These reporting of cues do not exceed a relatively low threshold, resulting in a "noisy" appearance. This

scattered pattern may reflect the intermittent encounter or perception of cues by patients, reflecting cue

reporting, consistent with the DST-model 2, where cue reporting fluctuations do not align directly with

craving peaks but rather create a baseline disturbance, testifying to a strong impact of craving on use

without the presence of clearly defined prior cue reporting – consistent with recent experimental data

[43]. At the same time, craving drives an increase in use. This increase is linked not only to heightened

craving but also to greater cue reporting. Interestingly, the decrease in cue reporting does not drive the

reduction in use or craving, as no maximum threshold of cue reporting is reached in this scenario (unlike

in *DST-model 1*). Once use has peaked, it must necessarily decline. At this point, craving returns to its

baseline, appearing to be influenced by the reduction in use. When use reaches maximum saturation,

there is a sharp decrease in craving. The dynamics of this model may thus be as follows: initially,

craving may lead to a peak in use, while simultaneously amplifying cue reporting that could be mediated

by an increased perception of cues. Once use hits its peak, it may sharply reduce craving, suggesting a

complex feedback loop between use and craving.

[FIGURE 5]

To summarize, when cue reporting precedes craving, the process appears to be primarily driven by a

phenomenon of "maximum cue reporting saturation", which initially increases craving and use, then

leads them to decrease as cue reporting subsides. When craving precedes cue reporting, the process

seems primarily driven by a phenomenon of "maximum use saturation", which causes craving and cue

reporting to decrease once use reaches its peak. In both profiles, craving serves as a critical modulator

between cue reporting and use. These findings align with animal models demonstrating spontaneous

exhaustion, or saturation in use [52].

The literature includes numerous simulation studies on psychiatry and DST (e.g., [15–34]); however,

to our knowledge, no DST study has yet been conducted in SUD. Most existing studies focus on a single

(phenotypic) variable (e.g., [31]) and have not integrated phenotype, internal factors, and environment

together – three elements essential for a nuanced understanding of clinical practice. Furthermore, only

eight studies have applied empirical data within DST models in psychiatry [19, 30, 53–58], and none

in the context of SUD. Finally, the integration of a linear statistical method with non-linear

computational analysis appears to be unprecedented.

This study's method provides meaningful perspectives on the relationship of environmental, subjective,

and behavioral variables. Such a dynamic systems approach allows for rigorous and time-sensitive

modeling of clinical reasoning processes. This focus on dynamic markers aimed at reducing use is

particularly novel. Additionally, based on EMA, DST could help address several other questions in the

field of SUD, such as providing more detailed simulation visualizations of patient progression, studying

phenotype variability, improving the understanding of dynamic attractors, structuring data from

longitudinal cohorts, and potentially refining nosology with dynamic factors [59–61].

This study encompasses limitations. Firstly, the list of cues was pre-identified with the patient.

However, in this study we were not trying to model the type of cues but rather change in number

(quantity) of cues, and their interactions with craving. Also, what is actually measured by EMA was the

reporting of cues when asked, that implies that cues were available and perceived. Only perceived cues

can be reported. Whitin these data, we cannot determine if change in cue reporting was driven by change

in the number of cues available to the patient, that may change with time, or by a change in state (such

as craving variation), that influences the quality of cue perception, independently of cue presence.

Secondly, we were trying to model rapid temporal sequences between variables, based on data that is

aggregated over 4-hour periods; within these windows, it is not possible to know the mutual influence

between variables. More frequent EMA measurements would provide better precision. Thirdly, at the

linear level, the present study focused on analyzing the interaction between cue reporting and craving

as a priority, considering that the use behavior was an outcome variable in relation to cue perception

and craving. This choice raises the concern that, in linear models, the analysis of cue reporting might

merely reflect cue presence that may be consequential to use patterns, and that future (SARIMAX)

analyzes may integrate the use as an (exogenous) influence variable. Moreover, certain statistical

choices were necessarily made for these linear models (e.g., favoring AIC over BIC), methods that

could be compared through simulations, for instance. However, beyond this potential extension to linear

models, the use of DST models allows us to highlight the interactions between use, craving, and cues.

Fourthly, although subgroup analyses by SUD type (alcohol, tobacco, cannabis, opiates, and cocaine)

could provide valuable insights, incorporating them could overcomplicate the already substantial

findings. Models were developed by combining different substance groups based on evidence that cues

influence craving, and craving, in turn, affects substance use and relapse across various types of SUD

[2–4]. Fifthly, a known limitation of the SARIMAX models is that they evaluate each patient as aligning

with one model, though a patient's condition may not be stable over time, and may change from one

model to the other, and there is the possibility that the two models could co-exist for some individuals.

Furthermore, SARIMAX models have difficulty handling nonlinear relationships and complex data

patterns, requiring significant fine-tuning to achieve optimal performance and assuming stationarity in

the data. We sought to address these limitations by verifying stationarity, performing a grid search to

optimize the models, and subsequently employing nonlinear models for greater precision. In parallel,

the nonlinear models based on DST also present certain methodological constraints [28]. For instance,

it could be necessary to refine the model to modelize symptoms and behaviors of different types (and

not only "use" or "craving"), different set of parameters or differential equations could have achieved

relatively similar dynamics and the qualitative interpretation of nonlinear phenomena, although

constrained by clinical perspectives, remains challenging. Different parameter sets could yield similar

data fits, highlighting the critical need for robust model validation.

Conclusion

In conclusion, this study introduces an innovative approach that combines EMA data, temporal statistical models (SARIMAX), and computational models (DST) to examine the nonlinear and interconnected relationships between cues, craving, and use in substance use disorders (SUD). The results could have important implications for the temporal phenotyping of patients, helping to refine the complex patterns of influence among SUD variables and supporting the identification of temporally anchored markers and mechanisms in clinical practice – thus opening critical therapeutic avenues for precision addiction treatments. Moving forward, validating the robustness of the model will remain essential, further reinforcing the reliability of these findings and their potential to advance tailored approaches in addiction treatment.

554 **Acknowledgements**

555 The authors express theirs thanks to all participants for their contribution and are grateful to all the

556 interviewers of the Addiction Team of SANPSY lab 2009-2022.

558 **Conflict of Interest**

References

1. Auriacombe M, Serre F, Denis C, Fatseas M. Diagnosis of addictions. Chapter 11. The Routledge Handbook of the Philosophy and Science of Addiction2018. p. 132–144.

2. Vafaie N, Kober H. Association of Drug Cues and Craving With Drug Use and Relapse: A Systematic Review and Meta-analysis. JAMA Psychiatry. 2022;79:641–650.

3. Serre F, Fatseas M, Denis C, Swendsen J, Auriacombe M. Predictors of craving and substance use among patients with alcohol, tobacco, cannabis or opiate addictions: Commonalities and specificities across substances. Addict Behav. 2018;83:123–129.

4. Serre F, Fatseas M, Swendsen J, Auriacombe M. Ecological momentary assessment in the investigation of craving and substance use in daily life: a systematic review. Drug Alcohol Depend. 2015;148:1–20.

5. Cavicchioli M, Vassena G, Movalli M, Maffei C. Is craving a risk factor for substance use among treatment-seeking individuals with alcohol and other drugs use disorders? A meta-analytic review. Drug Alcohol Depend. 2020;212:108002.

6. Cleveland HH, Knapp KS, Brick TR, Russell MA, Gajos JM, Bunce SC. Effectiveness and Utility of Mobile Device Assessment of Subjective Craving during Residential Opioid Dependence Treatment. Subst Use Misuse. 2021;56:1284–1294.

7. Bujarski S, Roche DJO, Sheets ES, Krull JL, Guzman I, Ray LA. Modeling naturalistic craving, withdrawal, and affect during early nicotine abstinence: a pilot EMA study. Exp Clin Psychopharmacol. 2015;23:81–89.

8. Sayette MA. The Role of Craving in Substance Use Disorders: Theoretical and Methodological Issues. Annu Rev Clin Psychol. 2016;12:407–433.

9. Stone AA, Shiffman S. Ecological Momentary Assessment (Ema) in Behavioral Medicine. Annals of Behavioral Medicine. 1994;16:199–202.

10. Serre F, Fatseas M, Debrabant R, Alexandre J-M, Auriacombe M, Swendsen J. Ecological momentary assessment in alcohol, tobacco, cannabis and opiate dependence: a comparison of feasibility and validity. Drug Alcohol Depend. 2012;126:118–123.

11. Fatseas M, Serre F, Alexandre J-M, Debrabant R, Auriacombe M, Swendsen J. Craving and substance use among patients with alcohol, tobacco, cannabis or heroin addiction: a comparison of substance- and person-specific cues. Addiction. 2015;110:1035–1042.

12. Ellis JD, Mun CJ, Epstein DH, Phillips KA, Finan PH, Preston KL. Intra-individual variability and stability of affect and craving among individuals receiving medication treatment for opioid use disorder. Neuropsychopharmacology. 2022;47:1836–1843.

13. Burgess-Hull AJ, Panlilio LV, Preston KL, Epstein DH. Trajectories of craving during medication-assisted treatment for opioid-use disorder: Subtyping for early identification of higher risk. Drug Alcohol Depend. 2022;233:109362.

14. Panlilio LV, Stull SW, Kowalczyk WJ, Phillips KA, Schroeder JR, Bertz JW, et al. Stress, craving and mood as predictors of early dropout from opioid agonist therapy. Drug Alcohol Depend. 2019;202:200–208.

15. Kelso JAS. Dynamic patterns: The self-organization of brain and behavior. Cambridge, MA, US: The MIT Press; 1995.

16. Boldrini M, Placidi G, Marazziti D. Applications of Chaos Theories to Psychiatry: A Review and Future Perspectives. CNS Spectrums. 1998;3:22–29.

17. Ayers S. The Application of Chaos Theory to Psychology. Theory & Psychology. 1997;7:373–398.

18. Barton S. Chaos, self-organization, and psychology. Am Psychol. 1994;49:5–14.

19. Bonsall MB, Geddes JR, Goodwin GM, Holmes EA. Bipolar disorder dynamics: affective instabilities, relaxation oscillations and noise. J R Soc Interface. 2015;12:20150670.

20. Bystritsky A, Nierenberg AA, Feusner JD, Rabinovich M. Computational non-linear dynamical psychiatry: a new methodological paradigm for diagnosis and course of illness. J Psychiatr Res. 2012;46:428–435.

21. Chang S-S, Chou T. A Dynamical Bifurcation Model of Bipolar Disorder Based on Learned Expectation and Asymmetry in Mood Sensitivity. Computational Psychiatry. 2018;2:205–222.

22. Cramer AOJ, Van Der Sluis S, Noordhof A, Wichers M, Geschwind N, Aggen SH, et al.

Dimensions of Normal Personality as Networks in Search of Equilibrium: You Can't like Parties if
you Don't like People. Eur J Pers. 2012;26:414–431.
23.   Cramer, Borkulo CD van, Giltay EJ, Maas HLJ van der, Kendler KS, Scheffer M, et al. Major
Depression as a Complex Dynamic System. PLOS ONE. 2016;11:e0167490.
24.   Cui J, Lichtwarck-Aschoff A, Olthof M, Li T, Hasselman F. From Metaphor to Computation:
Constructing the Potential Landscape for Multivariate Psychological Formal Models. Multivariate
Behavioral Research. 2023;58:743–761.
25.   Cui J, Hasselman F, Lichtwarck-Aschoff A. Unlocking nonlinear dynamics and multistability
from intensive longitudinal data: A novel method. 2023. 1 December 2023.
26.   Demic S, Cheng S. Modeling the Dynamics of Disease States in Depression. PLoS One.
2014;9:e110358.
27.   Durstewitz D, Huys QJM, Koppe G. Psychiatric Illnesses as Disorders of Network Dynamics.
Biol Psychiatry Cogn Neurosci Neuroimaging. 2021;6:865–876.
28.   Gauld C, Depannemaecker D. Dynamical systems in computational psychiatry: A toy-model
to apprehend the dynamics of psychiatric symptoms. Frontiers in Psychology. 2023;14.
29.   Globus GG, Arpaia JP. Psychiatry and the new dynamics. Biological Psychiatry.
1994;35:352–364.
30.   Gottschalk A, Bauer MS, Whybrow PC. Evidence of Chaotic Mood Variation in Bipolar
Disorder. Archives of General Psychiatry. 1995;52:947–959.
31.   Haken H, Tschacher W. How to Modify Psychopathological States? Hypotheses Based on
Complex Systems Theory. Nonlinear Dynamics Psychology and Life Sciences. 2017;21:19–34.
32.   Haslbeck J, Ryan O, Robinaugh D, Waldorp L, Borsboom D. Modeling Psychopathology:
From Data Models to Formal Theories. 2019. December 2019. https://doi.org/None.
33.   Hayes AM, Yasinski C, Barnes JB, Bockting CLH. Network destabilization and transition in
depression: New methods for studying the dynamics of therapeutic change. Clin Psychol Rev.
2015;41:27–39.
34.   Hayes AM, Laurenceau J-P, Feldman G, Strauss JL, Cardaciotto L. Change is Not Always
Linear: The Study of Nonlinear and Discontinuous Patterns of Change in Psychotherapy. Clin Psychol
Rev. 2007;27:715–723.
35.   Witkiewitz K, Marlatt GA. Modeling the complexity of post-treatment drinking: it's a rocky
road to relapse. Clin Psychol Rev. 2007;27:724–738.
36.   Foo JC, Noori HR, Yamaguchi I, Vengeliene V, Cosa-Linan A, Nakamura T, et al.
Dynamical state transitions into addictive behaviour and their early-warning signals. Proceedings of
the Royal Society B: Biological Sciences. 2017;284:20170882.
37.   Serre F, Fatseas M, Swendsen J, Auriacombe M. Does craving intensity influence cue
exposure reports? An ecological momentary assessment study in patients with alcohol, tobacco,
cannabis and heroin use disorder. Drug and Alcohol Dependence. 2015;C:e201.
38.   Fatseas M, Serre F, Swendsen J, Auriacombe M. Effects of anxiety and mood disorders on
craving and substance use among patients with substance use disorder: An ecological momentary
assessment study. Drug Alcohol Depend. 2018;187:242–248.
39.   Denis C, Fatséas M, Beltran V, Serre F, Alexandre J-M, Debrabant R, et al. Usefulness and
validity of the modified Addiction Severity Index: A focus on alcohol, drugs, tobacco, and gambling.
Subst Abus. 2016;37:168–175.
40.   McLellan AT, Kushner H, Metzger D, Peters R, Smith I, Grissom G, et al. The Fifth Edition
of the Addiction Severity Index. J Subst Abuse Treat. 1992;9:199–213.
41.   Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al. The Mini-
International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured
diagnostic psychiatric interview for DSM-IV and ICD-10. J Clin Psychiatry. 1998;59 Suppl 20:22-
33;quiz 34-57.
42.   Fatseas M, Serre F, Swendsen J, Auriacombe M. Effects of anxiety and mood disorders on
craving and substance use among patients with substance use disorder: An ecological momentary
assessment study. Drug and Alcohol Dependence. 2018;187:242–248.
43.   Serre F, Gauld C, Lambert L, Baillet E, Beltran V, Daulouede J, et al. Predictors of substance
use during treatment for addiction: A network analysis of ecological momentary assessment data.
Addiction. 2024:add.16658.

44.   Chirokoff V, Dupuy M, Abdallah M, Fatseas M, Serre F, Auriacombe M, et al. Craving
dynamics and related cerebral substrates predict timing of use in alcohol, tobacco, and cannabis use
disorders. Addiction Neuroscience. 2023;9:100138.
45.   Shumway RH, Stoffer DS. ARIMA Models. In: Shumway RH, Stoffer DS, editors. Time
Series Analysis and Its Applications: With R Examples, Cham: Springer International Publishing;
2017. p. 75–163.
46.   Mushtaq R. Augmented Dickey Fuller Test. 2011.
47.   Vagropoulos SI, Chouliaras GI, Kardakos EG, Simoglou CK, Bakirtzis AG. Comparison of
SARIMAX, SARIMA, modified SARIMA and ANN-based models for short-term PV generation
forecasting. 2016 IEEE International Energy Conference (ENERGYCON), 2016. p. 1–6.
48.   Alharbi FR, Csala D. A Seasonal Autoregressive Integrated Moving Average with Exogenous
Factors (SARIMAX) Forecasting Model-Based Time Series Approach. Inventions. 2022;7:94.
49.   Strogatz SH. Nonlinear Dynamics And Chaos: With Applications To Physics, Biology,
Chemistry, And Engineering. 1st edition. Cambridge, Mass: CRC Press; 2000.
50.   Izhikevich EMM. Dynamical Systems in Neuroscience: The Geometry of Excitability and
Bursting. 24383rd edition. Cambridge, Massachusetts London, England: The MIT Press; 2010.
51.   Van Rossum G, Drake FL. Python 3 Reference Manual. Scotts Valley, CA: CreateSpace;
2009.
52.   Ahmed SH. Validation crisis in animal models of drug addiction: Beyond non-disordered
drug use toward drug addiction. Neuroscience & Biobehavioral Reviews. 2010;35:172–184.
53.   Pezard L, Nandrino JL, Renault B, ElMassioui F, Allilaire JF, Muller J, et al. Depression as a
dynamical disease. Biol Psychiatry. 1996;39:991–999.
54.   Tschacher W, Scheier C, Hashimoto Y. Dynamical analysis of schizophrenia courses.
Biological Psychiatry. 1997;41:428–437.
55.   van de Leemput IA, Wichers M, Cramer AOJ, Borsboom D, Tuerlinckx F, Kuppens P, et al.
Critical slowing down as early warning for the onset and termination of depression. Proceedings of
the National Academy of Sciences. 2014;111:87–92.
56.   Tschacher W, Haken H. Causation and chance: Detection of deterministic and stochastic
ingredients in psychotherapy processes. Psychotherapy Research. 2020;30:1075–1087.
57.   Odgers CL, Mulvey EP, Skeem JL, Gardner W, Lidz CW, Schubert C. Capturing the ebb and
flow of psychiatric symptoms with dynamical systems models. Am J Psychiatry. 2009;166:575–582.
58.   Heiby E, Pagano I, Blaine D, Nelson K, Heath R. Modeling Unipolar Depression as a Chaotic
Process. Psychological Assessment. 2003;15:426–434.
59.   Wichers M, Schreuder MJ, Goekoop R, Groen RN. Can we predict the direction of sudden
shifts in symptoms? Transdiagnostic implications from a complex systems perspective on
psychopathology. Psychol Med. 2019;49:380–387.
60.   Nelson B, McGorry PD, Wichers M, Wigman JTW, Hartmann JA. Moving From Static to
Dynamic Models of the Onset of Mental Disorder: A Review. JAMA Psychiatry. 2017;74:528–534.
61.   Salvi JD, Rauch SL, Baker JT. Behavior as Physiology: How Dynamical-Systems Theory
Could Advance Psychiatry. AJP. 2021;178:791–792.
709

710   **Figures**

712
713

716   **Figure 1.** Phase space representation of the co-evolution of the cues, craving and use for two randomly

717   chosen patient from the dataset. The relations between these variables are complex and no trivial linear

718   interdependence appear.

Figure 2. Dynamic representation of the temporal profile of patients with Substance Use Disorder

(SUD) belonging to the *SARIMAX-model 1* ("cues-to-craving") (n=154), for 4 days. **Top**: Space phase

of the model, i.e., the conceptual framework illustrating all possible states of the system and the

interactions between variables over time. This 3D representation illustrates the dynamic interactions

between craving, cues, and use, providing a visual model of how fluctuations in one variable correspond

with variations in the others. Although such a depiction may not be immediately readable, it remains

essential for capturing the complex, nonlinear relationships inherent in dynamic systems and is

commonly used to elucidate interdependencies across multiple variables. **Bottom**: Temporal

representation of the model (4 times 12 hours, hours are noted every 4 hours and over 24 hours for the

sake of clarity). SARIMAX: Seasonal Auto-Regressive Integrated Moving Average with eXogenous

variable; DST: Dynamical System Theory.

Figure 3. Dynamic representation of the temporal profile of patients with Substance Use Disorder

(SUD) belonging to the *SARIMAX-2* model ("craving-to-cues") (n=211), for 4 days. **Top**: Space phase

of the model, i.e., the conceptual framework illustrating all possible states of the system and the

interactions between variables over time. This 3D representation illustrates the dynamic interactions

between craving, cues, and use, providing a visual model of how fluctuations in one variable correspond

with variations in the others. Although such a depiction may not be immediately readable, it remains

essential for capturing the complex, nonlinear relationships inherent in dynamic systems and is

commonly used to elucidate interdependencies across multiple variables. **Bottom**: Temporal

representation of the model (4 times 12 hours, hours are noted every 4 hours and over 24 hours for the

749 sake of clarity). SARIMAX: Seasonal Auto-Regressive Integrated Moving Average with eXogenous

750 variable; DST: Dynamical System Theory.

755 **Figure 4.** Temporal sequence of the interaction of the three variables of use, craving and cues in the

756 *DST-model 1* ("cues-to-craving"). These relationships are described here in a linear manner in order to

757 facilitate understanding but should be interpreted in a non-linear manner (in particular with mutual and

758 reciprocal interactions) within the working framework of the DST. The arrow indicates the trigger

759 variable in this model.

Figure 5. Temporal sequence of the interaction of the three variables of use, craving and cues in the

DST-model 2 ("craving-to-cues"). These relationships are described here in a linear manner in order to

facilitate understanding but should be interpreted in a non-linear manner (in particular with mutual and

reciprocal interactions) within the working framework of the DST. The arrow indicates the trigger

variable in this model.

Table 1. Characteristics of the sample (n=211 participants): primary substance, demographic variables,

substance use-related information and psychiatric comorbidity. ASI: Addiction Severity Index; MINI:

Mini International Neuropsychiatric Interview–Plus; SD: standard deviation.

	Sample characteristics	All participants (n=211)	
		% (n)	Mean (SD)
Primary substance	Alcohol	35 (73)	–
	Tobacco	28 (58)	–
	Cannabis	18 (39)	–
	Opiates	16 (34)	–
	Cocaine	3 (7)	–
Demographic variables	Sex (female)	36 (78)	–
	Employed	56 (116)	–
	Age	–	38.2 (11.4)
	Education (years)	–	12.5 (2.74)
	Pharmacological treatment	79 (167)	–
Substance use-related information and psychiatric comorbidity	ASI addiction severity (0-9)	–	6.6 (0.76)
	Mood disorder on the MINI (current diagnosis)	34 (72)	–
	Anxiety disorder on the MINI (current diagnosis)	45 (93)	–

777 **Table 2**. Ecological Momentary Assessment (EMA) report of craving episodes and intensity, number

778 of cues, and reports of primary substance use of participants (n=211). SD: standard deviation.

EMA variables	All participants (n=211 – 8,260 observations)	
	% (n)	Mean (SD)
Craving episodes (intensity ≥ 1)	70 (5,810)	–
Craving intensity (1-7)	–	3.6 (2.10)
Number of cues	–	4.3 (3.4)
Use	37 (3,087)	–

Table 3. Socio-demographic characteristics, substance types, addiction severity, psychiatric diagnoses and EMA variable characteristics of the two temporal profile (SARIMAX) of patients with Substance Use Disorder (SUD) (n=211). SARIMAX: Seasonal Auto-Regressive Integrated Moving Average with eXogenous variable. "Model 1": SARIMAX-model 1 ("cues-to-craving"); "Model 2": SARIMAX-model 2 ("craving-to-cues").

	Model 1 (N= 154)				Model 2 (N=57)					
	%	n	Mean	SD	%	n	Mean	SD	Test	p*
Sample characteristics										
Sex (female)	36.4	56			38.6	22			χ^2=0.089	0.765
Employed	55.6	84			57.1	32			χ^2= 0.038	0.845
Age			37.6	11.1			39.7	11.3	F= 1.53	0.217
Education (years)			12.4	2.6			12.5	3.1	F= 0.021	0.886
Primary substance										**<.001**
Alcohol	30.5	47			45.6	26				
Tobacco	34.4	53			8.7	5				
Cannabis	20.1	31			14.0	8				
Opiates	13.0	20			24.6	14				
Cocaine	1.9	3			7.0	4				
Addiction severity										
ASI ISR (0-9)			6.6	0.8			6.5	0.7	F= 0.088	0.767
Current pharmacological treatment for addiction	80.5	124			75.4	43			χ^2=0.651	0.419
MINI current diagnosis										
Mood disorder	32.0	48			42.9	24			χ^2= 2.11	0.146
Anxiety disorder	42.0	63			52.6	30			χ^2= 1.89	0.170
EMA variables	6,004 observations (38.99/patient)				2,256 observations (39.58/patient)					
Frequency of craving episodes (intensity > 1)			73.3	32.4			63.7	27.7	F= 3.98	**0.0472**
Craving intensity (1-7)			3.7	1.6			3.2	1.3	F= 4.25	**0.0405**
Number of cues			4.9	2.5			2.8	1.7	F= 35.91	**<.0001**
Frequency of primary substance use			42.4	35.1			26.6	29.9	F= 9.08	**0.0029**

786 **Table 4**. Parameter values of the two dynamical system models fitted with linear models (n=211). DST:

787 Dynamical System Theory. The details of the parameters and their clinical translation are given in

788 **Supplementary Material 4**.

DST models	Parameters of x, y and z variables												
	x				y						z			
	S_{max}	R_s	λ_s	τ_x	B_{max}	R_b	λ_b	P	L	τ_y	S	α	β	τ_z
DST-model 1	10	1	0.1	14	10	1.04	0.05	10	1.01	14	1.0	0.5	0.5	1
DST-model 2	10	1	0.1	2	10	0.904	0.05	1	1.00	2	1.0	0.5	0.5	1

Funding source

Funding for the EMA studies was provided by Research Grant AAP-Recherche-CRA (20091301018) from the Aquitaine Regional Council (MA) and IRESP-19-ADDICTIONS-16 from IReSP and Alliance Aviesan for CUSEMA study (FS). For this study, MA and FS received financial support from the French government in the framework of the University of Bordeaux's IdEx "Investments for the Future" program / GPR BRAIN_2030. Office and staff support was provided by CH Charles Perrens. The funding sponsors had no role in the design of the study or interpretation of the data. They were not involved in the preparation, review, or approval of this manuscript.

Data availability

The data that support the findings of the current manuscript are available from the corresponding author, upon reasonable request.

Authors' contributions

Marc Auriacombe and Fuschia Serre were the principal investigators, obtaining funding and supervised the study. Marc Auriacombe provided access to participants. Damien Depannemaecker and Christophe Gauld conceptualized and undertook analysis, interpretation of data and drafting of manuscript. Fuschia Serre participated in patient recruitment, data collection and database management. Marc Auriacombe and Fuschia Serre sustained the elaboration of the methodology and writing. All authors undertook the critical revision of the manuscript for important intellectual content and all authors significantly contributed to the manuscript and approved the final version.

5   **Figure 1.**

Figure 2

Figure 3

Figure 4

Figure 5